

Yu-Hung Yeh^a, Dario Panzeri^{a,1}, Yasuhiro Kadota^{b,1,2}, Yi-Chun Huang^a, Pin-Yao Huang^a, Chia-Nan Tao^a, Milena Roux^{b,3}, Hsiao-Chiao Chien^a, Tzu-Chuan Chin^a, Po-Wei Chu^a, Cyril Zipfel^b, and Laurent Zimmerli^{a,4}

^aDepartment of Life Science and Institute of Plant Biology, National Taiwan University, Taipei 106, Taiwan.

^bThe Sainsbury Laboratory, Norwich Research Park, Norwich NR4 7UH, United Kingdom.

¹These authors contributed equally to this work.

²Present address: RIKEN Center for Sustainable Resource Science, Plant Immunity Research Group, Suehiro-cho 1-7-22 Tsurumi-ku, Yokohama 230-0045, Japan.

³Present address: University of Copenhagen, Department of Biology, Ole Maaløes Vej 5, Copenhagen, 2200, Denmark.

⁴Corresponding author e-mail: lauzim2@ntu.edu.tw

The *Arabidopsis* Malectin-Like/LRR-RLK IOS1 is Critical for BAK1-Dependent and BAK1-Independent Pattern-Triggered Immunity

Running title: IOS1 role in *Arabidopsis* Innate Immunity

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantcell.org) is: Laurent Zimmerli (lauzim2@ntu.edu.tw).

Synopsis: The RLK IOS1 is critical for priming of *Arabidopsis thaliana* innate immunity and is required for optimal function of BAK1-dependent and BAK1-independent cell surface immune receptors.

ABSTRACT

Plasma membrane-localized pattern recognition receptors (PRRs) such as FLAGELLIN SENSING2 (FLS2), EF-TU RECEPTOR (EFR) and CHITIN ELICITOR RECEPTOR KINASE 1 (CERK1) recognize microbe-associated molecular patterns (MAMPs) to activate pattern-triggered immunity (PTI). A reverse genetics approach on genes responsive to the priming agent beta-aminobutyric acid (BABA) revealed IMPAIRED OOMYCETE SUSCEPTIBILITY1 (IOS1) as a critical PTI player. *Arabidopsis thaliana ios1* mutants were hyper-susceptible to *Pseudomonas syringae* bacteria. Accordingly, *ios1* mutants showed defective PTI responses, notably delayed up-regulation of the PTI-marker gene *FRK1*, reduced callose deposition and mitogen-activated protein kinase activation upon MAMP treatment. Moreover, *Arabidopsis* lines over-expressing *IOS1* were more resistant to bacteria and had a primed PTI response. *In vitro* pull-down, bimolecular fluorescence complementation, co-immunoprecipitation and mass spectrometry analyses supported the existence of complexes between the membrane-localized IOS1 and BRASSINOSTEROID INSENSITIVE1-ASSOCIATED KINASE1 (BAK1)-dependent PRRs FLS2 and EFR, as well as with the BAK1-independent PRR CERK1. IOS1 also associated with BAK1 in a ligand-independent manner, and positively regulated FLS2-BAK1 complex formation upon MAMP treatment. In addition, IOS1 was critical for chitin-mediated PTI. Finally, *ios1* mutants were defective in BABA-induced resistance and priming. This work reveals IOS1 as a novel regulatory protein of FLS2-, EFR- and CERK1-mediated signaling pathways that primes PTI activation.

INTRODUCTION

Plants possess multilayered recognition systems that detect pathogens at various stages of infection and proliferation. Recognition of microbial invasion is essentially based upon the host's ability to distinguish between 'self' and 'non-self' components. Early microbial pathogens detection is performed by cell surface-localized pattern-recognition receptors (PRRs) that sense pathogen- or microbe-associated molecular patterns (PAMPs or MAMPs) (Monaghan and Zipfel, 2012). Major examples of MAMPs are lipopolysaccharides (LPS) present in the envelope

of Gram-negative bacteria, eubacterial flagellin, eubacterial elongation factor Tu (EF-Tu), peptidoglycans from Gram-positive bacteria, methylated bacterial DNA fragments and fungal cell wall-derived chitins (Girardin et al., 2002; Cook et al., 2004; Boller and Felix, 2009). MAMP recognition promptly triggers the activation of pattern-triggered immunity (PTI) (Tsuda and Katagiri, 2010). Early PTI responses such as calcium influx, production of reactive oxygen species (ROS), and activation of mitogen-activated protein kinases (MAP kinases), induce transcriptional reprogramming mediated by plant WRKY transcription factors as well as calmodulin-binding proteins (Boller and Felix, 2009; Tena et al., 2011). In addition, *Arabidopsis thaliana* plants close stomata in a MAMP-dependent manner when in contact with bacteria (Melotto et al., 2006; Singh et al., 2012a). Callose deposition and PTI marker gene up-regulation are usually observed later (Zipfel and Robatzek, 2010). Activation of PTI leads to broad resistance to pathogens (Nicaise et al., 2009; Tsuda and Katagiri, 2010; Zeng et al., 2010; Desclos-Theveniau et al., 2012). Virulent bacterial pathogens inject proteins some of which suppress PTI (Deslandes and Rivas, 2012; Feng and Zhou, 2012). Often, recognition of microbial effectors by plant intracellular nucleotide-binding site and leucine-rich repeat (NLR) proteins activates effector-triggered immunity (ETI). ETI is a rapid and robust response, usually associated with a hypersensitive reaction (Maekawa et al., 2011; Gassmann and Bhattacharjee, 2012).

In *Arabidopsis*, the most extensively studied PRRs are the leucine-rich repeat receptor-like kinases (LRR-RLKs) FLAGELLIN SENSING2 (FLS2) and EF-TU receptor (EFR). FLS2 and EFR recognize bacterial flagellin (or the derived peptide flg22) and EF-Tu (or the derived peptides elf18/elf26), respectively (Gomez-Gomez and Boller, 2000; Zipfel et al., 2006). Upon ligand binding, FLS2 and EFR rapidly associate with another LRR-RLK, BRI1-ASSOCIATED RECEPTOR-LIKE KINASE/SOMATIC EMBRYOGENESIS RECEPTOR-LIKE KINASE3 (BAK1/SERK3), forming a ligand-inducible complex that triggers downstream PTI responses (Chinchilla et al., 2007; Heese et al., 2007; Roux et al., 2011). In addition to associating with FLS2, BAK1 recognizes the C-terminus of the FLS2-bound flg22 thus acting as a co-receptor (Sun et al., 2013). BAK1-LIKE1/SOMATIC EMBRYOGENESIS RECEPTOR-LIKE KINASE4 (BKK1/SERK4) also cooperates with BAK1 to regulate PRR-mediated signaling pathway (Roux et al., 2011). Interestingly, the BAK1-INTERACTING RECEPTOR KINASE2 (BIR2) prevents BAK1 interaction with FLS2 before elicitation. Importantly, BIR2 is released from BAK1 upon

MAMP perception, allowing FLS2-BAK1 association and PTI activation (Halter et al., 2014). While BAK1 and other SERKs are the primary regulators downstream of FLS2 and EFR, the perception of the fungal MAMP chitin and signaling through CHITIN ELICITOR RECEPTOR KINASE1 (CERK1) does not require BAK1 (Shan et al., 2008; Kemmerling et al., 2011; Ranf et al., 2011). Although CERK1 was considered as the major PRR for chitin (Miya et al., 2007; Wan et al., 2008; Wan et al., 2012), recent data suggest that the LYSIN MOTIF RECEPTOR KINASE5 (LYK5) is the primary receptor for chitin (Cao et al., 2014). Upon chitin elicitation, CERK1 and LYK5 form a complex to activate plant innate immunity (Cao et al., 2014). CERK1 is also involved in the recognition of peptidoglycans (Willmann et al., 2011). Other proteins downstream of PRRs modulate the PTI response. Typically, the receptor-like cytoplasmic kinase (RLCK) BOTRYTIS-INDUCED KINASE1 (BIK1) plays a critical role in mediating early flagellin signaling from the FLS2/BAK1 receptor complex and regulates responses induced by elf18, Pep1 and chitin, and thus acts as a convergent point downstream of multiple PRRs (Lu et al., 2010a; Zhang et al., 2010). Other RLCKs such as the PTI COMPROMISED RECEPTOR-LIKE CYTOPLASMIC KINASE 1 (PCRK1) and PCRK2 function downstream of multiple PRRs (Sreekanta et al., 2015; Kong et al., 2016). In addition, the BRASSINOSTEROID-SIGNALING KINASE1 (BSK1) associates with unstimulated FLS2 (Shi et al., 2013). The DENN (Differentially Expressed in Normal and Neoplastic cells) domain protein STOMATAL CYTOKINESIS-DEFECTIVE1 (SCD1) is also necessary for some FLS2- and EFR-mediated responses and associates in a ligand-independent manner with FLS2 *in vivo* (Korasick et al., 2010). Furthermore, lectin receptor kinases (LecRKs) such as LecRK-VI.2 and LecRK-V.5 modulate early PTI signaling (Desclos-Theveniau et al., 2012; Singh et al., 2012a; Singh and Zimmerli, 2013; Huang et al., 2014).

In addition to PTI and ETI, other resistance responses such as systemic acquired resistance and induced systemic resistance are activated after pathogen challenges (Durrant and Dong, 2004; Van Wees et al., 2008). Organic and inorganic compounds can also induce systemic resistance in plants. The non-protein amino acid beta-aminobutyric acid (BABA) is a potent inducer of resistance against abiotic stress (Jakab et al., 2005; Zimmerli et al., 2008), nematodes (Oka et al., 1999), insects (Hodge et al., 2005), and microbial pathogens (Jakab et al., 2001; Zimmerli et al., 2001; Cohen, 2002; Ton and Mauch-Mani, 2004; Po-Wen et al., 2013). BABA-induced resistance is associated with a faster activation of defense mechanisms upon stress perception, a

phenomenon known as priming (Conrath et al., 2006; Navarova et al., 2012). Although accumulation of defense signaling components before stress exposure (Beckers et al., 2009; Singh et al., 2012a) and epigenetic modifications (Jaskiewicz et al., 2011; Luna et al., 2012; Rasmann et al., 2012; Slaughter et al., 2012; Po-Wen et al., 2013) are suggested to be critical for priming, the identity of signaling components involved in priming is still largely unknown.

In an effort to identify novel critical players in *Arabidopsis* immunity and priming, we used a reverse genetic approach by testing mutants of genes whose expression levels are induced by the priming agent BABA (Tsai et al., 2011). Three independent insertion lines in the malectin-like/LRR-RLK *IMPAIRED OOMYCETE SUSCEPTIBILITY 1 (IOS1)* (Hok et al., 2011) were found to be hyper-susceptible to bacterial pathogens. IOS1 is known to contribute to disease provoked by filamentous (hemi)biotrophs and to attenuate abscisic acid (ABA) responses in *Arabidopsis* (Hok et al., 2011; 2014). Through loss- and gain-of-function analyses and biochemical approaches, we show that IOS1 is an important modulator of *Arabidopsis* PTI that associates with the LRR-RLKs FLS2, EFR and BAK1 in a ligand-independent manner, notably controlling the complex formation between FLS2 and BAK1. IOS1 also associates with the LysM-domain RLK CERK1 and controls chitin-mediated PTI.

RESULTS

IOS1 is Required for Resistance to Hemi-Biotrophic Bacteria

To identify *Arabidopsis* genes involved in immunity to bacteria, we followed a reverse genetic analysis of genes up-regulated by the priming agent BABA (Tsai et al., 2011). One of these genes is the Malectin-like/LRR-RLK *IOS1* (At1g51800) (Hok et al., 2011; 2014). For our analyses, we used *ios1-1*, a transcriptional knockout *Ds* transposon insertion line in *Ler-0* background (GT_5_22250) recently isolated (Hok et al., 2011), and *ios1-2* (Salk_137388) and *ios1-3* (SAIL_343_B11), two independent T-DNA insertion lines in Col-0 background still producing truncated *IOS1* transcripts (see Supplemental Figure 1 online). To test whether IOS1 is required for anti-bacterial immunity, the 3 insertion lines were dip-inoculated with the virulent hemi-biotrophic bacteria *Pseudomonas syringae* pv. *tomato* DC3000 (*Pst* DC3000) or

Pseudomonas syringae pv. *maculicola* ES4326 (*Psm* ES4326). At 3 days post-inoculation (dpi), *ios1-1*, *ios1-2* and *ios1-3* developed stronger symptoms than wild-type (WT) plants, as illustrated by increased chlorosis and necrosis formation (Figure 1A). This phenotype was associated with significantly higher bacterial titers (Figure 1B). Typically, the susceptibility phenotype of *ios1-2* to *Pst* DC3000 is similar to the mutant *bak1-5* (see Supplemental Figure 2 online). We also evaluated the susceptibility of *ios1* mutants to the *Pst* DC3000 *hrcC* mutant, a strain defective in delivering type-III effectors that cannot repress the PTI response, and consequently is mostly non-virulent on *Arabidopsis* (Brooks et al., 2004). All 3 *ios1* mutants tested allowed more growth of *Pst* DC3000 *hrcC* than WT plants upon syringe-infiltration (see Supplemental Figure 3 online), suggesting a defective PTI response in *ios1* mutants.

To test whether IOS1 is also required for immunity to pathogens other than bacteria, susceptibility of *ios1* mutants to the necrotrophic fungal pathogens *Botrytis cinerea* and *Alternaria brassicicola* was evaluated by droplet-inoculation. Mutants *ios1-1*, *ios1-2* and *ios1-3* were as susceptible as WT plants to both pathogens (see Supplemental Figure 4A and 4B online), suggesting that IOS1 is critical for immunity to virulent hemi-biotrophic bacteria, but not to necrotrophic fungi such as *B. cinerea* and *A. brassicicola*.

The role of IOS1 in anti-bacterial immunity was further evaluated by analyzing the susceptibility to *Pst* DC3000 of transgenic *Arabidopsis* lines over-expressing *IOS1* mRNA (OE1 and OE3) (see Supplemental Figure 5 online). Both *IOS1*-OE lines were significantly less susceptible to *Pst* DC3000 (Figure 1C). Notably, although *ios1* mutants did not show any defect in stomatal innate immunity (see Supplemental Figure 6 online), over-expression of *IOS1* inhibited the bacteria-mediated re-opening of stomata (Figure 1D). Together, these data are consistent with a positive role of IOS1 in anti-bacterial immunity.

IOS1 is Critical for Late PTI Responses

To analyze whether IOS1 is involved in PTI responses, we first monitored *IOS1* mRNA expression levels by qRT-PCR after treatments of seedlings with 100 nM flg22 or elf18. Both MAMPs induced *IOS1* transcripts accumulation at 1 h after treatment (see Supplemental Figure 7 online). To evaluate the role of IOS1 in late PTI responses, we measured callose deposition in *ios1* mutants after infiltration with the bacterial MAMPs flg22 or elf26. Aniline blue staining and

image analysis indicated lower levels of callose deposition in *ios1-1* and *ios1-2* than in WT leaves (Figure 2A). These results suggest that IOS1 is critical for PTI-induced callose deposition. To further evaluate late PTI responses, we monitored expression levels of the PTI marker gene *FRK1* (Asai et al., 2002; Xiao et al., 2007; Boudsocq et al., 2010) after treatment with flg22 or elf18. At 1 h after MAMP treatment, both *ios1-1* and *ios1-2* mutants demonstrated lower *FRK1* up-regulation levels (Figure 2B). Furthermore, we analyzed late PTI responses in the *IOS1*-OE lines. Interestingly, these lines did not exhibit constitutive callose deposition, while more callose deposits were observed in OE1 and OE3 upon elicitation with the MAMPs flg22 or elf18 (Figure 2C). Similarly, constitutive up-regulation of the PTI marker gene *FRK1* was not observed in *IOS1*-OE lines, but *FRK1* expression levels were potentiated in the OE1 and OE3 lines upon flg22 or elf18 treatments (Figure 2D). These data suggest that over-expression of *IOS1* primes late PTI responses and that IOS1 positively regulates several late PTI responses.

IOS1 Modulates Several Early PTI Responses

To test whether IOS1 is required for early PTI events, we analyzed ROS production in response to 10 nM flg22 or elf26 for 30 min in WT, *ios1-1*, *ios1-2* and *IOS1*-OE leaves. Both mutants and OE lines displayed WT levels of ROS production, while MAMP-mediated ROS production was strongly reduced in *bak1-4* (Figure 3A and 3B; see Supplemental Figure 8A and 8B online). Treatments with MAMPs rapidly activate *Arabidopsis* MAP kinases MPK3 and MPK6 (Nuhse et al., 2000). Notably, both *ios1-1* and *ios1-2* mutants demonstrated a weaker activation of MPK3 and MPK6 than WT following treatment with flg22 or elf18 (Figure 3C; see Supplemental Figure 8C and 8D online). On the other hand, MPK3 and MPK6 activation was stronger than WT in the OE1 and OE3 transgenic lines (Figure 3D; see Supplemental Figure 8E online). Together these results suggest that IOS1 is required for full MPK activation, but not for the ROS burst after MAMP perception. This observation is consistent with these responses being uncoupled (Segonzac et al., 2011; Xu et al., 2014a). However, since we do not provide a kinetic analysis, we cannot exclude a slower or faster MPK response in *ios1* mutants or OE lines, respectively.

IOS1 Likely Localizes to the Plasma Membrane

IOS1 is a predicted trans-membrane RLK (Hok et al., 2011). We analyzed IOS1 subcellular localization by transiently expressing IOS1-GFP fusion protein driven by the cauliflower mosaic virus 35S promoter in *Arabidopsis* mesophyll protoplasts. The fluorescence signal was mainly confined to the cell surface with a pattern similar to the plasma membrane marker pm-rk CD3-1007 (Nelson et al., 2007), while the control protoplasts expressing GFP alone showed a nuclear/cytoplasmic localization (Figure 4A). These data suggest that similarly to the PRRs FLS2 and EFR (Robatzek et al., 2006; Haweker et al., 2010), IOS1 is likely localized at the plasma membrane.

IOS1 Associates with FLS2, EFR and BAK1 in a Ligand-Independent Manner

IOS1 acts up-stream of MPK in flg22- and elf26- or elf18-triggered PTI signaling cascades. We thus evaluated whether IOS1 associates with PRRs such as FLS2 or EFR. We first used pull-down analysis to show that a Trx-6xHis-tagged IOS1 kinase domain (KD) interacted with MBP-tagged FLS2 and EFR *in vitro* (Figure 4B). Next, interactions were evaluated by bimolecular fluorescence complementation (BiFC) assays (Walter *et al.*, 2004) in *Arabidopsis* protoplasts. To test whether our experimental conditions were appropriate, we first analysed the interactions between BAK1 and FLS2 or EFR that occur only upon elicitation (Chinchilla *et al.*, 2007; Heese *et al.*, 2007; Roux et al., 2011). As expected, the YFP signal was clearly observed after flg22 or elf18 treatment (Figure 4C; see Supplemental Figure 9A online). YFP fluorescence was detected before and after elicitation with flg22 or elf18 when testing IOS1 interaction with FLS2 or EFR respectively (Figure 4C; see Supplemental Figure 9A online). Similarly, IOS1 interacted with BAK1 in a ligand independent manner (Figure 4C; see Supplemental Figure 9A online). The low temperature and salt-responsive protein 6B (LTI6b/RCI2B) fused to GFP, which is known to localize at the plasma membrane (Cutler et al., 2000), was used as a negative control. LTI6b is known to dimerize (Huang et al., 2014), and YFP fluorescence was indeed observed when *Arabidopsis* protoplasts were transfected with LTI6b-YFP^N and LTI6b-YFP^C (Figure 4D; see Supplemental Figure 9B online), indicating that both constructs were functional. Importantly, no YFP fluorescence at the plasma membrane was observed when testing IOS1 interaction with LTI6b, even after elicitation with flg22 or elf18 (Figure 4D; see Supplemental Figure 9B online). Although we cannot exclude artifacts inherent to over-expression in protoplasts, these data

suggest that IOS1 interacts at the plasma membrane with the PRRs FLS2 and EFR and the co-receptor BAK1 in a ligand-independent manner.

To test whether IOS1 associates with FLS2 *in vivo*, we transiently co-expressed FLS2-HA₃ with GFP epitope-tagged IOS1 in *Arabidopsis* protoplasts. Equal amount of IOS1 were immunoprecipitated with GFP-Trap beads and analyzed for presence of FLS2-HA₃ using anti-HA immunoblotting. FLS2 could be detected in mock- and flg22-treated samples (Figure 5A). Similarly, we analyzed the possible association of IOS1 with EFR and BAK1 before and after elicitation with elf18 (for EFR) or flg22 and elf18 (for BAK1). For that purpose, IOS1-GFP was transiently co-expressed with EFR-HA₃ or BAK1-HA₃ in *Arabidopsis* protoplasts, and IOS1 was immunoprecipitated with GFP-Trap beads. EFR and BAK1 could also be detected in the IOS1 immunoprecipitate before and after MAMP treatment (Figure 5A). As a negative control, we tested the association of IOS1 with LTI6b by immunoprecipitating equal amount of LTI6b with GFP-trap beads and by analyzing FLS2-HA₃, EFR-HA₃ and BAK1-HA₃ presence using anti-HA immunoblotting. FLS2, EFR and BAK1 could not be detected, suggesting that they do not associate with GFP at the plasma membrane (Figure 5A). These observations suggest that IOS1 associates with FLS2, EFR and BAK1 in a ligand-independent manner. Of note, IOS1 homo-dimerized independently of flg22 treatment (see Supplemental Figure 10 online), as previously reported for FLS2 (Sun et al., 2012). In addition, the GFP fusion of IOS1 does not affect its function as it can complement the defective MPK3 and MPK6 activation observed in *ios1-1* and *ios1-2* mutants (see Supplemental Figure 11 online).

To test whether IOS1-GFP associates with FLS2 in *Arabidopsis* as well, we performed co-immunoprecipitation experiments using transgenic lines over-expressing IOS1-GFP. IOS1-GFP was immunoprecipitated with anti-GFP magnetic beads, and analyzed for the presence of endogenous BAK1 and FLS2 using anti-BAK1 and anti-FLS2 immunoblotting. As negative controls, anti-GFP magnetic beads were incubated with protein extracts of untransformed Col-0 and transgenic plants expressing LTI6b fused to GFP. Signals for FLS2 and BAK1 upon LTI6b-GFP immunoprecipitation were largely weaker than those observed upon IOS1-GFP, suggesting that FLS2 and BAK1 do not aspecifically bind to anti-GFP magnetic beads, nor do they interact with GFP itself at the plasma membrane (Figure 5B). In contrast, we could detect a clear association of IOS1-GFP with native FLS2 and BAK1 (Figure 5B). Treatment with flg22 did not affect significantly or reproducibly the associations of IOS-GFP with FLS2 and BAK1 (Figure

5B). Moreover, we found IOS1 as part of the *in vivo* EFR complex in an unbiased manner while searching for EFR-associated proteins *in planta* by proteomics analysis. In these experiments, anti-GFP immunoprecipitates were prepared from untreated and elf18-treated transgenic *efr-1*/EFRp:EFR-eGFP seedlings, as well as from untreated *efr-1* null mutant or Col-0 seedlings, in order to reveal proteins that non-specifically bind to GFP beads. Liquid chromatography-tandem mass spectrometry (LC-MS/MS) analysis identified 8 different peptides matching IOS1 in the EFR-eGFP immunoprecipitates, but none in the negative controls (see Supplemental Table 1 online). The IOS1 peptides were found in both untreated and elf18-treated samples, corroborating the fact that IOS1 associates with the PRRs FLS2 and EFR in a ligand-independent manner *in vivo*.

IOS1 is Required for Optimal flg22-Induced FLS2-BAK1 Association, But Functions Independently of BIK1

To test whether associations between IOS1 and both FLS2 and BAK1 impact other biochemical events within the FLS2 complex, we analyzed ligand-induced FLS2-BAK1 association (Chinchilla et al., 2007; Heese et al., 2007). Toward this goal, BAK1 was first immunoprecipitated from *ios1-2* plants treated or not with 100 nM flg22 and associated FLS2 was revealed by anti-FLS2 immunoblotting. After flg22 treatment, the mutant *ios1-2* displayed significantly less FLS2 coimmunoprecipitated with BAK1 than the WT control (Figures 6A and B). By contrast, a significant increase in coimmunoprecipitated FLS2 was observed in *Arabidopsis* over-expressing *IOS1* treated with 2 different concentrations of flg22 (Figures 6C and D). These data show that the active kinase IOS1 (see Supplemental Figure 12 online), positively regulates the association of BAK1 with FLS2. However, flg22-mediated phosphorylation of BIK1, which is a direct substrate of FLS2 (Lu et al., 2010a; Zhang et al., 2010), was not affected in *ios1-2* and the OE3 line (Figure 7A-D). Together, these results indicate that IOS1 modulates FLS2-BAK1 association upon elicitation but is not critical for BIK1 phosphorylation. We further evaluated IOS1 dependency to BAK1 and BIK1 by analyzing callose deposition in lines over-expressing *IOS1* in *bak1-5* and *bik1* mutant backgrounds. *bak1-5* and *bik1* mutants are largely defective in flg22-mediated callose deposition (Figure 7E; Zhang et al., 2010). While over-expression of *IOS1* strongly primed callose deposition in the Col-0 WT

control, *bak1-5* mutation completely abolished *IOS1*-mediated priming of callose deposition (Figure 7E). However, lines over-expressing *IOS1* in the *bik1* mutant background still displayed a large increase in callose deposits after flg22 treatment (Figure 7E). Collectively, these results suggest that *IOS1* functions in a BAK1-dependent, but BIK1-independent manner in the FLS2 complex.

Defective Chitin Responses in *ios1* Mutants

To evaluate whether *IOS1* function is uniquely linked with BAK1, we analyzed MPK3/6 activities upon elicitation with the MAMP chitin. Fungal chitin recognition is mediated by LysM-domain RLKs such as CERK1 (Miya et al., 2007; Wan et al., 2008; 2012), and BAK1 is not required for chitin perception and signaling (Shan et al., 2008; Kemmerling et al., 2011; Ranf et al., 2011). Reduced MPK3/6 activities were observed after chitin treatment in the *ios1-1* and *ios1-2* mutants (Figure 8A), suggesting that *IOS1* also plays a role in PRR complexes that do not recruit BAK1. To further investigate whether *IOS1* is necessary for the chitin-mediated PTI response, callose deposition in *ios1-1* and *ios1-2* mutants was analyzed. Both mutants accumulated less callose than WT *Arabidopsis* at 16 h after chitin treatment (Figure 8B), indicating that *IOS1* is necessary for chitin-mediated callose deposition. The fungal pathogen *B. cinerea* produces the MAMP chitin (Windram et al., 2012) and *A. brassicicola* is commonly used in chitin perception studies (Miya et al., 2007), but *ios1* mutants demonstrated WT resistance to both pathogens (see Supplemental Figure 4 online). We thus tested the resistance response of the OE1 and OE3 lines towards these necrotrophic pathogen. Both OE lines harbored smaller *B. cinerea*-mediated disease lesions (Figure 8C), while they did not show increased resistance towards *A. brassicicola* (see Supplemental Figure 13 online). Taken together, these data suggest that *IOS1* is critical for chitin-mediated PTI and plays a positive role in *Arabidopsis* resistance to some, but not all pathogens that produce the MAMP chitin.

***IOS1* Associates with CERK1**

Possible association of *IOS1* with CERK1 was tested by Co-IP in *Arabidopsis* protoplasts. The previously described CERK1 dimerization (Liu et al., 2012) was observed indicating that the

CERK1-GFP and CERK1-HA₃ constructs were functional (Figure 9A). Association of IOS1 with CERK1 was then performed by immunoprecipitating equal amount of IOS1 with GFP-Trap beads and by analyzing CERK1-HA₃ presence using anti-HA immunoblotting. CERK1 could clearly be detected before and after elicitation with chitin (Figure 9A), suggesting ligand-independent association. We also co-expressed CERK1-HA₃ with empty vector (EV)-GFP. No signal was observed in this negative control, suggesting that the association between CERK1-HA₃ and IOS1-GFP is not due to a direct binding of CERK1-HA₃ with GFP proteins or GFP-Trap beads (Figure 9A). As an additional negative control, we tested the association of CERK1 with LTI6b after elicitation with chitin by immunoprecipitating equal amount of LTI6b with GFP-trap beads and by analyzing CERK1-HA₃ presence using anti-HA immunoblotting. CERK1-HA₃ could not be detected or at very low levels (Figure 9A). Together, these data suggest specific association of IOS1 with CERK1 at the plasma membrane.

These observations suggest that IOS1 is part of the CERK1 receptor complex. To further clarify IOS1-CERK1 complex, direct interaction of IOS1 with CERK1 was analyzed by BiFC in *Arabidopsis* protoplasts. The dimerization of CERK1 was first used to demonstrate that CERK1-YFP^N and CERK1-YFP^C constructs are functional (Liu et al., 2012). As expected, the YFP signal was observed when both CERK1-YFP^N and CERK1-YFP^C were co-transfected in *Arabidopsis* protoplasts before and after chitin treatment indicating dimerization of CERK1 (Figure 9B). Similarly, a clear YFP signal was visible independently of chitin treatment with the CERK1-YFP^N and IOS1-YFP^C constructs (Figure 9B), suggesting that both proteins can directly interact in *Arabidopsis* protoplasts. Collectively, these results suggest that in addition to FLS2 and EFR, IOS1 also interacts with CERK1.

IOS1 is Necessary for BABA-Induced Resistance and Priming

Since over-expression of the BABA-responsive *IOS1* primes *Arabidopsis* PTI (Figures 2C and 2D; Figure 3D), we tested whether IOS1 is required for induced resistance to *Pst* DC3000 and *Psm* ES4326 triggered by the priming agent BABA (Zimmerli et al., 2000; Tsai et al., 2011). While BABA treatments protected both Col-0 and *Ler*-0 WT against *Pst* DC3000 and *Psm* ES4326 infection, *ios1-1* and *ios1-2* mutants demonstrated a defective BABA-induced resistance towards these hemi-biotrophic bacteria (Figure 10A). BABA is known to prime the PTI response

in *Arabidopsis* (Singh et al., 2012a; Po-Wen et al., 2013). We thus tested the role of IOS1 in BABA-induced priming of PTI responses. Notably, the priming effect of BABA on flg22-induced callose deposition and *FRK1* expression was largely abolished in *ios1-1* and *ios1-2*, in comparison to WT (Figure 10B and C). IOS1 positively modulates flg22-mediated FLS2-BAK1 association (Figure 6) and BABA treatment up-regulates *IOS1* expression (Tsai et al., 2011). We thus asked whether BABA affects FLS2-BAK1 association upon flg22 elicitation. No clear increase in FLS2-BAK1 association upon flg22 treatment was observed in BABA-treated Col-0 plants (see Supplemental Figure 14 online).

Since BABA inhibits bacteria-mediated stomatal re-opening (Tsai et al., 2011), we tested whether IOS1 is involved in this phenomenon. While BABA inhibited bacteria-mediated stomatal re-openings in Col-0 and *Ler-0* WT, it did not in *ios1-1* and *ios1-2* mutants (Figure 10D). Taken together, these data suggest a positive role for IOS1 in BABA-induced resistance and BABA-mediated priming of PTI, including strengthening of stomatal innate immunity.

DISCUSSION

PRRs are critical to elicit PTI responses and to restrict pathogen ingress (Boller and Felix, 2009; Nicaise et al., 2009; Zhang and Zhou, 2010; Huang and Zimmerli, 2014). To date, all known plant PRRs are modular transmembrane proteins containing ligand-binding ectodomains that function as part of multi-protein complexes (Böhm et al., 2014; Zipfel, 2014). In this work, we analyzed the role of the *Arabidopsis* malectin-like/LRR-RLK IOS1 in innate immunity and priming with genetic and biochemical approaches. The results support the following conclusions:

IOS1 is Necessary for Full Activation of PTI in *Arabidopsis*

Our reverse genetic approach identified 3 independent *IOS1* insertion mutants with hyper-susceptibility to virulent hemi-biotrophic *Pst* DC3000 and *Psm* ES4326 bacteria, but with WT sensitivity to the necrotrophic fungal pathogens *B. cinerea* and *A. brassicicola*. These observations suggest that IOS1 is critical for resistance to hemi-biotrophic bacteria, but not to necrotrophic fungi. However, lines over-expressing *IOS1* were more resistant to *B. cinerea*,

suggesting a role for IOS1 in *Arabidopsis* resistance to this necrotrophic pathogen. A partially defective PTI response in *ios1* mutants may not be sufficient to produce a visible increased susceptibility phenotype upon infection by *B. cinerea*. Necrotrophic fungal pathogens such as *B. cinerea* produce toxins, cell wall degrading enzymes and ROS to promote disease and macerate plant tissue (Prins et al., 2000), possibly hiding the effect of a defective PTI. By contrast, a stronger PTI in IOS1 OE lines may restrict early *B. cinerea* infection at least partially independently of toxins, cell wall degrading enzymes and ROS produced by *B. cinerea*, leading to increased resistance. Redundancy may also explain the lack of increased sensitivity in *ios1* loss-of-function mutants. Lines over-expressing *IOS1* demonstrated a WT resistance to *A. brassicicola* suggesting that *Arabidopsis* resistance to this necrotrophic fungus occurs independently of IOS1. The mutant *ios1-1* is known to be more resistant to the filamentous oomycete pathogens *Hyaloperonospora arabidopsidis* and *Phytophthora parasitica* (Hok et al., 2011; 2014). IOS1 could be a direct or indirect target of oomycete effectors necessary for pathogen virulence. IOS1 absence in *ios1-1* would not allow *H. arabidopsidis* or *P. parasitica* to fully repress *Arabidopsis* PTI (Hok et al., 2014). Taken together, these observations suggest that IOS1 is involved in *Arabidopsis* immunity to various microbial pathogens.

Increased susceptibility of *ios1* mutants to virulent bacteria was correlated with a defective PTI response. Typically, bacteria- and MAMP-induced callose depositions were dramatically reduced in *ios1* mutants. In addition, up-regulation of the PTI-responsive gene *FRK1* was delayed in plants with a defective IOS1. Coherently, *Arabidopsis* over-expressing *IOS1* demonstrated increased accumulation of callose and potentiated expression levels of *FRK1* upon MAMP elicitation. By contrast, both *FRK1* expression and callose deposition were not affected in the *ios1-1* mutant after inoculation with filamentous pathogens (Hok et al., 2014). These discrepancies may be explained by early (this work) versus late time point analyses (Hok et al., 2014). MPK3/6 activation was reduced in *ios1* mutants and augmented in *IOS1* over-expression lines, suggesting that IOS1 acts upstream of MPK3/6 in PTI signaling. These observations point to the fact that IOS1 is necessary for full activation of both early and late PTI responses. Similarly, *LecRK-VI.2* is necessary for full activation of some early and late PTI responses (Singh et al., 2012a). However, while *Arabidopsis* over-expressing *LecRK-VI.2* demonstrate a constitutive PTI response, *IOS1* over-expression lines showed a strengthened PTI only upon elicitation by bacteria or MAMPs, suggesting a different mechanism of action for these two

positive regulators of PTI. Importantly, PTI-mediated ROS production was at WT levels in *ios1* mutants and in *IOS1*-OE lines, suggesting that IOS1 may not regulate all aspects of the PTI response. The PRR-associated kinase BIK1 directly regulates apoplastic ROS production during PTI (Kadota et al., 2014; Li et al., 2014; Kadota et al., 2015). So, the apparent absence of IOS1 regulation of ROS production could be explained by the fact that IOS1 acts largely in a BIK1-independent manner (Figures 7). The *ios1* mutants demonstrated WT bacteria-mediated stomatal closure while *IOS1*-OE lines harbored a strengthened stomatal immunity. Redundancy may explain stomatal innate immunity discrepancy between *ios1* mutants and *IOS1*-OE lines (compare Figure 1D and Supplemental Figure 6 online). Other malectin-like/LRR-RLKs may indeed play a redundant role in stomatal closure (Hok et al., 2011), thus masking the possible function of IOS1 in this early PTI response. Stomata of the *ios1-1* mutant are hyper-responsive to ABA (Hok et al., 2014), and ABA signaling is critical for stomatal immunity (Melotto et al., 2006; Desclos-Theveniau et al., 2012), but we observed WT stomatal closure in *ios1-1* after *Pst* DC3000 inoculation. As suggested by Hok et al. (2014), IOS1 may use different signalings for the activation of PTI in response to bacteria and in the down-regulation of ABA upon infection with filamentous pathogens. Taken together, these data reveal IOS1 as a major positive regulator of *Arabidopsis* PTI against bacteria, acting upstream of MPK3/6 in FLS2- and EFR-dependent defense signaling pathways. Noteworthy, typical concentrations of MAMPs were used for each PTI assay resulting in the use of various concentrations of MAMPs in different experiments. We thus cannot fully exclude dose-dependent effects.

IOS1 Associates with FLS2, EFR and BAK1 in a Ligand-Independent Manner

Having genetically demonstrated the importance of IOS1 in bacteria-, flg22- and elf26/elf18-triggered PTI upstream of MPK3/6, and also considering that IOS1 is a LRR-RLK with 2 LRR motifs, a transmembrane domain, and a complete extracellular malectin-like domain (Hok et al., 2011), we further investigated whether IOS1 is part of PRR complexes recognizing bacterial MAMPs. We first showed that the KD of IOS1 associates *in vitro* with the KDs of FLS2 and EFR using a pull-down approach. In addition, *in vivo* association of IOS1 with FLS2, EFR and/or the regulatory LRR-RLK BAK1 were evaluated by BiFC and CoIP analyses. We performed BiFC assays and CoIP experiments in *Arabidopsis* protoplasts and found that IOS1

constitutively associates with FLS2 and EFR and that elicitation with flg22 or elf18 does not significantly affect the association. The constitutive IOS1 and FLS2 association was further confirmed in *Arabidopsis* using transgenic lines over-expressing IOS1-GFP, while IOS1 was also found to be part of unstimulated and stimulated EFR complexes by *in planta* proteomics analysis on EFR-associated proteins (see Supplemental Table 1 online). In addition to associating with PRRs, IOS1 interacts with the BRASSINOSTEROID SIGNAL-KINASE 3 in *Arabidopsis* (Xu et al., 2014b). In *Arabidopsis*, only few proteins are known to be present in PRR complexes before elicitation by MAMPs. Notably, the cytoplasmic kinases BIK1/PBLs and BSK1 interact constitutively with FLS2 and are released upon elicitation (Lu et al., 2010a; Zhang et al., 2010; Shi et al., 2013). Both PCRK1 and PCRK2 associate with FLS2 (Kong et al., 2016), and heterotrimeric G proteins directly interact with FLS2 to regulate PTI (Liang et al., 2016). Additionally, the DENN domain protein SCD1 that negatively regulates innate immunity associates in a ligand-independent manner with FLS2 *in vivo* (Korasick et al., 2010). Furthermore, the ubiquitin E3 ligases PUB12/13 interact with BAK1 prior elicitation and ubiquitinate FLS2 upon flg22-induced FLS2/BAK1 complex formation, leading to FLS2 degradation (Lu et al., 2011) and BIR2 negatively regulates *Arabidopsis* PTI by association before elicitation with BAK1 (Halter et al., 2014). This work thus reveals a novel component of FLS2 and EFR protein complexes.

IOS1 Positively Regulates FLS2-BAK1 Complex Formation

Since the malectin-like LRR-RLK IOS1 constitutively associates with the PRRs FLS2 and EFR and the regulatory LRR-RLK BAK1, and since *ios1* mutants demonstrate a defective PTI, we hypothesized that IOS1 affects early events at PRR complexes. The flg22-mediated association of FLS2 and BAK1 was indeed reduced in *ios1-2* and increased in the OE3 line when compared to WT Col-0 controls. By contrast, the positive regulator of PTI LecRK-VI.2 does not modulate flg22-mediated association of FLS2 and BAK1 (Singh et al., 2012a). The heteromerization between FLS2 and BAK1 occurs within seconds (Schulze et al., 2010) with BAK1 acting as co-receptor for flg22 (Sun et al., 2013), indicating that both LRR-RLKs most likely exist in close proximity at the plasma membrane, as recently suggested in the case of BAK1 and BRI1 (Bücherl et al., 2013). We thus propose that the plasma membrane-localized IOS1 is required for

promoting rapid FLS2-BAK1 complex formation upon flg22 binding. Importantly, the flg22-mediated association between FLS2 and BAK1 was not completely abolished in *ios1-2*. Other players such as other malectin-like LRR-RLKs may generate the partial FLS2-BAK1 association observed upon flg22 elicitation in *ios1-2*. IOS1 constitutively interacts with both FLS2 and BAK1, however FLS2 and BAK1 complex formation only occurs after flg22 treatment. In addition, FLS2-FLS2 and IOS1-IOS1 homo-dimerization could be observed independently of elicitation (Sun et al., 2012; see Supplemental Figure 10 online). IOS1 monomers or dimers may thus bind both FLS2 and BAK1 in different complexes before PTI elicitation. Upon flg22 treatment, IOS1 may participate in the formation of a new complex that integrates both FLS2 and BAK1. Contrarily to IOS1, BIR2 negatively regulates FLS2-BAK1 complex formation (Halter et al., 2014). Thus BIR2 may directly or indirectly antagonize IOS1.

Treatments with flg22 induce rapid phosphorylation of BIK1, which further increases phosphorylation of FLS2 and BAK1 (Lu et al., 2010a; Zhang et al., 2010). BIK1 phosphorylation occurs within minutes after flg22 treatment and is thus considered a good marker of PRRs activities. Surprisingly, BIK1 phosphorylation was at WT levels in flg22-treated *ios1-2* mutant and in the OE3 line. Since MPK3/6 activities were altered in *ios1* mutants and in *IOS1*-OE lines, a BIK1-independent signaling cascade that affects MPK3/6 activities must be present in *ios1-2* and in *IOS1*-OE lines. This observation is in agreement with Zhou and colleagues who demonstrated that BIK1 and the closely related PBL1 are not required for flg22-induced MAPK activation (Feng et al., 2012). Therefore, other receptor-like cytoplasmic kinases could play a role in regulating different branches of PTI signaling (Lu et al., 2010b). In addition, *Arabidopsis* over-expressing *IOS1* in the *bik1* mutant background still demonstrated a strong priming of callose deposition. By contrast, augmented callose deposition after flg22 treatment was strongly abolished in lines over-expressing *IOS1* in the *bak1-5* background. Taken together, these data suggest that an altered FLS2-BAK1 association in *ios1-2* impacts MPK3/6 activation independently of BIK1 phosphorylation.

IOS1 is Necessary for Chitin-Mediated PTI Responses and Associates with CERK1

IOS1 plays a critical role in BAK1-dependent PRR complexes such as FLS2 and EFR in a BIK1-independent manner. We next asked whether IOS1 is also necessary for a full PTI response

activated by PRR complexes functioning in a BAK1-independent manner. In contrary to FLS2 that associates with the BAK1 co-receptor to sense flagellin (Sun et al., 2013), the LysM-domain RLK CERK1 that is part of PRR complexes that recognize chitin (Miya et al., 2007; Wan et al., 2008; 2012; Cao et al., 2014), and peptidoglycans (Willmann et al., 2011), functions in a BAK1-independent manner. Notably, BAK1 is not required for chitin perception and signaling (Shan et al., 2008; Kemmerling et al., 2011; Ranf et al., 2011). Both *ios1-1* and *ios1-2* mutants demonstrated defective MPK3/6 activation and reduced callose deposition after chitin treatment. In addition, CoIP and BiFC analyses suggested direct association of IOS1 with CERK1. We thus propose that IOS1 acts at the CERK1 receptor complex to positively regulate chitin-mediated PTI responses. IOS1 thus regulates the PTI response at both BAK1-dependent and BAK1-independent PRR complexes. Similarly, BIK1 is involved in both FLS2/EFR and CERK1 complexes to activate downstream PTI responses (Lu et al., 2010a; Zhang et al., 2010). Notably, BIK1 regulates CERK1-mediated chitin responses, including the accumulation of ROS and the induction of defense genes (Zhang et al., 2010). Since CERK1 does not associate with BAK1 upon elicitation, IOS1 should modulate the chitin-mediated PTI response through other regulatory mechanisms than PRR association with BAK1 as observed for FLS2 (Figure 6). CERK1 is also involved in bacterial resistance (Gimenez-Ibanez et al., 2009), and is a critical member of the PRR complex that recognizes bacterial peptidoglycans (Willmann et al., 2011), suggesting that IOS1 plays a role in at least 3 PRR complexes recognizing bacterial MAMPs. This may explain the strong phenotypes of *ios1* mutants and IOS1-OE lines observed upon bacterial infection. By contrast, the role of IOS1 in *Arabidopsis* resistance against necrotrophic fungi such as *B. cinerea* and *A. brassicicola* was rather weak. In addition, the *ios1-1* mutant is more resistant to the biotrophic fungus *Erysiphe cruciferarum* (Hok et al., 2014). These observations suggest that IOS1 role in the *Arabidopsis* resistance against pathogens producing the MAMP chitin is not critical. As suggested for oomycete pathogens (Hok et al., 2011), fungal pathogens may produce effectors that target IOS1, and the absence of IOS1 may result in WT or enhanced *Arabidopsis* resistance levels even though IOS1 is critical for a full chitin-mediated defense response.

IOS1 Plays a Critical Role in Priming of PTI

Accumulation of positive regulators of defense such as MPK3/6 or LecRK-VI.2 prior to stress challenge is critical for priming (Beckers et al., 2009; Singh et al., 2012a). Plants over-expressing IOS1 demonstrated potentiated expression of PTI-responsive genes, primed callose deposition and increased MPK3/6 activities upon PTI elicitation. These observations further suggest that increased accumulation of positive regulators of PTI before elicitation is sufficient to prime PTI and consequently to increase resistance to pathogens. BABA-mediated accumulation of *IOS1* mRNA (Tsai et al., 2011) may thus be critical for BABA-mediated priming of PTI (Singh et al., 2012a; Po-Wen et al., 2013). We therefore investigated whether *ios1* mutants are defective in BABA-mediated priming. While *lecrk-VI.2-1* mutant is only partially defective in BABA priming (Singh et al., 2012a), *ios1-1* and *ios1-2* mutants were largely deficient in BABA-induced resistance to bacteria, priming of *FRK1* expression and callose deposition, and in BABA-mediated strengthening of stomatal innate immunity. These results suggest that IOS1 plays a predominant role during priming of PTI by the non-protein amino acid BABA. Surprisingly, BABA had no effect on flg22-mediated FLS2-BAK1 association, suggesting that the reported role of IOS1 in BABA-triggered priming involves another regulatory mechanism.

LRR-RLKs such as FLS2, EFR, and CERK1 or PEPR1/2 are receptors for MAMPs or DAMPs, respectively (Huffaker et al., 2006; Yamaguchi et al., 2006; Miya et al., 2007; Ryan et al., 2007; Wan et al., 2008; Krol et al., 2010; Yamaguchi et al., 2010). Another LRR-RLK, BAK1, functions in several PRR complexes as a co-receptor (Chinchilla et al., 2007; Heese et al., 2007; Boller and Felix, 2009; Schulze et al., 2010; Sun et al., 2013). Our data identified the malectin-like LRR-RLK IOS1 as a novel member of FLS2 and EFR PRR complexes that also associates in a ligand-independent manner with BAK1. In addition, IOS1 regulates CERK1-dependent PTI responses that are BAK1-independent (Shan et al., 2008; Kemmerling et al., 2011; Ranf et al., 2011). This work identifies a novel LRR-RLK regulating BAK1-dependent and -independent PTI responses and further reveals the intricate regulation of the PRR complex dynamics needed for transmitting and regulating PTI signaling, which requires additional components beyond the ligand-binding receptor and co-receptor.

METHODS

Biological Materials and Growth Conditions

Arabidopsis thaliana (L. Heyhn.) ecotypes Columbia (Col-0) and Landsberg erecta (*Ler*-0) were grown in commercial potting soil/perlite (3:2) at 22 °C to 24 °C day and 17 °C to 19 °C night temperature under a 9-h-light/15-h-dark photoperiod. The lighting was supplied at an intensity of $\sim 100 \mu\text{E m}^{-2} \text{ s}^{-1}$ by fluorescence tubes. The Ds transposon insertion line (*Ler*-0) *ios1-1* (GT_5_22250) and the T-DNA insertion mutants (Col-0) *ios1-2* (Salk_137388) and *ios1-3* (SAIL_343_B11) were obtained from the *Arabidopsis* Biological Resource Centre (ABRC). The mutant *bak1-4* (Salk_116202) and *bik-1* have been described elsewhere (Chinchilla et al., 2007; Lu et al., 2010a; Zhang et al., 2010). Bacterial strains *Pst* DC3000 and the *Pst* DC3000 *hrcC* mutant (CB200) were provided by B.N. Kunkel (Washington University, St. Louis, Missouri, USA), while *Psm* ES4326 was a gift from J. Glazebrook (Minnesota University, St. Paul, Minnesota, USA). All bacteria were cultivated at 28 °C and 340 rpm in King's B medium with 50 mg/L rifampicin (*Pst* DC3000), 50 mg/L rifampicin and kanamycin (CB200), or 50 mg/L streptomycin (*Psm* ES4326). The fungi *B. cinerea* and *A. brassicicola* were obtained from C.Y. Chen (National Taiwan University, Taipei, Taiwan), and grown at room temperature (18 °C ~ 25 °C) on PDA-agar plates (Zimmerli et al., 2001).

Pathogen Infection Assays

Five-week-old *Arabidopsis* were dipped in 10^6 cfu/mL *Pst* DC3000 or 5×10^5 cfu/mL *Psm* ES4326 in 10 mM MgSO_4 containing 0.01 % Silwet L-77 (Lehle Seeds) for 15 min. After inoculation, plants were kept at 100 % relative humidity, and symptoms were evaluated 3 days later. Bacterial titers were determined as previously described (Zimmerli et al., 2000). For *B. cinerea* and *A. brassicicola* infection, spores were diluted to 1×10^5 and 5×10^5 spores/mL in 1/2 PDB medium respectively. Droplets of 10 μL 1/2 PDB with *B. cinerea* or *A. brassicicola* spores were deposited on leaf surfaces of 5-week-old plants (3 leaves per plants). Leaves of same age were used for droplet-inoculation. Disease symptoms and lesion diameters were determined at 3 dpi. At least 18 lesion diameters were evaluated for each independent experiment (6 plants).

IOS1 Over-Expression Plants

The DNA plasmids (pH35GWG) expressing IOS1 protein fused with GFP at the C terminus under the control of the cauliflower mosaic virus 35S promoter were obtained from ABRC (Gou et al., 2010) (ABRC stock S1G51800HGF). *Agrobacterium tumefaciens* strain GV3101 was used for the transformation of Col-0 plants. Successful transformation were determined by screening on 0.6 % MS agar plates containing 50 mg/mL hygromycin B and raised to homozygous T3 lines. For the generation of *IOS1*-OE lines in *bak1-5* or *bik1* mutant background, mutant plants were dip-inoculated with *Agrobacterium* strains GV3101 carrying Pro35S-*IOS1*-GFP (pFAST-R05) using OLE1-TagRFP as a screenable marker (Shimada et al., 2010) and raised to T2 for analyses.

BABA and MAMP Treatments

For bacteria titer, callose deposition, and stomatal aperture evaluations, 5-week-old *Arabidopsis* were soil drenched with BABA (Fluka) at a final concentration of 225 μ M two days before bacteria inoculation or MAMP treatments. BABA was dissolved in water and controls were soil drenched with water only. For *FRK1* expression and ligand-induced FLS2-BAK1 association, seedlings grown on $\frac{1}{2}$ MS plates with or without 30 μ M BABA for 10 days were submerged in a 1 μ M or 100 nM flg22 solution, respectively for 60 min before sample collection.

The flg22 and elf26 or elf18 peptides were purchased from Biomer Technology and dissolved in 10 mM MgSO₄, MgSO₄ only or water for seedling treatment was used as control. Chitin from shrimp shells (Sigma) was dissolved in water. Water only treatments were used as controls. MAMPs were syringe-infiltrated into leaves and samples were harvest at indicated time points.

Callose Deposition

Five-week-old *Arabidopsis* leaves were syringe infiltrated with 1 μ M flg22 in 10 mM MgSO₄. Control plants were infiltrated with 10 mM MgSO₄ only. Nine leaf discs from 3 different plants were selected for analyses at the indicated time points. Callose deposition evaluation on seedlings was performed on 14-day-old *Arabidopsis* grown on $\frac{1}{2}$ MS plates that were transferred to $\frac{1}{2}$ MS liquid medium one night before treatment with 100 nM flg22, 100 nM elf18 or 0.2 mg/mL chitin for 16 h. Six seedlings were selected for analyses for each sample. Callose deposits were visualized as described (Singh et al., 2012a).

RT-PCR

For quantitative RT-PCR, *Arabidopsis* seedlings grown on ½ MS plates for 10 days were transferred to liquid ½ MS one night before treatments with 100 nM flg22 or elf26 for *ios1* mutants and with 50 nM flg22 or elf18 for the OE lines and samples were collected at the indicated time points. Total RNA isolation, complementary DNA biosynthesis and real-time PCR analyses were performed as described (Wu et al., 2010). Normalization of gene expression was conducted with At4g05320 (*UBQ10*). For RT-PCR, one microliter of cDNA was used as template and standard PCR conditions were applied as described (Singh et al., 2012a). At4g05320 (*UBQ10*) was used as a loading control. Primers used are in Supplemental Table 2 online.

MAP Kinase Assay

Twenty 10-day-old plants were incubated in ½ MS supplemented with 100 nM (*ios1* mutants) or 50 nM (OE lines) flg22 or elf18 dissolved in water, or with water only (control), 0.2 mg/mL Chitin or water (control) for 5 min before being pooled for harvest. For complementation assays, protoplasts from 5-week-old *Arabidopsis* were transfected with plasmids (pH35GWG) carrying Pro35S-*IOS1*-GFP or the vector only by polyethylene glycol (Sigma, <http://www.sigmaaldrich.com/>) and samples were collected 5 min after flg22 or water treatment. MAP kinase assays were performed as described (Singh et al., 2012a).

ROS Burst

ROS assays were performed as previously described (Huang et al., 2012). Shortly, 6 leaf discs (10 mm diameter) from three 5-week-old *Arabidopsis* (2 discs/plant) were incubated in ddH₂O in 96-well plates overnight. The following day, the water was replaced by 10 nM flg22 or elf26 in 10 mM MgSO₄ buffer or by 10 mM MgSO₄ buffer only for the mock controls containing 2 μM luminol (Sigma) and 10 μg/mL peroxidase (Sigma). The plates were analyzed every 2 min for mutants or every 2.5 min for OE lines after addition of MAMPs for a period of 30 min using a CentroLIApc LB 692 plate luminometer (Berthold Technologies, Bad Wildbad, Germany).

Stomatal Assay

Five-week-old plants were kept under light (100 μE m⁻² s⁻¹) for at least 3 h to open stomata before the beginning of the experiments. For each biological replicate, stomatal apertures were

evaluated from 12 epidermal peels from 4 plants (3 epidermal peels/plant) as described (Tsai et al., 2011).

Subcellular Localization in Protoplast

For transient expression of the GFP fusion proteins, constructs expressing *35S-IOS1-GFP* (plasmid pH35GWG, ABRC stock S1G51800HGF) or vector alone were transfected into *Arabidopsis* mesophyll protoplasts according to He et al., (2007). The GFP-fusion constructs were co-transfected with the plasma membrane marker pm-rkCD3-1007 (Nelson et al., 2007). Transfected protoplasts were visualized using a confocal laser scanning microscope (Zeiss LSM 780 Confocal, Carl Zeiss, Oberkochen, Germany) with excitation at 488 nm and emission at 490-515 nm, autofluorescence was observed at 650-700 nm. The plasma membrane marker was detected with excitation at 594 nm and emission at 595-650 nm.

BIK1 Phosphorylation

Mesophyll protoplasts were obtained as described by He et al. (2007). BIK1 phosphorylation assays on protoplasts treated with 0.75 μ M flg22 for 3.5, 7 and 10 min were performed as described (Singh et al., 2012b).

Cloning, Expression and Purification of Recombinant Proteins

In order to generate a Trx-6xHis N-terminal fusion of the IOS1 kinase domain, the sequence coding for the IOS1 cytosolic domain was amplified from the pH35GWG vector expressing the IOS1-GFP fusion using primers carrying BamHI and XhoI restriction sites (see Supplemental Table 2 online), and introduced into the polylinker of the pET-32a(+) expression vector (Novagen). To produce an inactive kinase fusion protein, a point mutation at the kinase active site (D710N) was introduced into the expression vector by primer extension (see Supplemental Table 2 online) using the Phusion polymerase (New England Biolabs), followed by DpnI (New England Biolabs) digestion according to manufacturer's instructions. The Trx-6xHis-IOS1KD and Trx-6xHis-IOS1KDM (kinase dead) fusion proteins were expressed in the *E. coli* strain Rosetta (DE3) pLysS (Novagen). After overnight induction at 16 °C with 0.4 mM IPTG, the bacteria were pelleted by centrifugation, re-suspended in 100 mL binding buffer (20 mM sodium phosphate, 0.5 M NaCl, 20 mM imidazole, 0.04% 2-mercaptoethanol) and sonicated. The soluble His-tagged proteins were affinity purified using a HisTrap FF column (GE Healthcare) according

to manufacturer's instructions. All constructs were confirmed by Sanger sequencing and the purified Trx-6xHis-*IOS1*KD and Trx-6xHis-*IOS1*KDm were analysed by LC-MS/MS.

The production of MBP-tagged *FLS2* and *EFR* kinase domain constructs was performed as described in Schwessinger et al., (2011). MBP was expressed from pMALTM-c5X (New England Biolabs). MBP and the two MBP-tagged proteins were expressed as described above, but using the *E. coli* strain BL21 (DE3) pLysS (Novagen), and purified using amylose resins (MBPTrap HP, GE Healthcare) following manufacturer's instructions. Finally, MBP and the two MBP-tagged proteins were dialysed against dialysis buffer (20 mM Tris-HCl, 100 mM NaCl, pH 7.4).

***In vitro* Pull-Down Assay**

One microgram of MBP, MBP-*FLS2*KD, or MBP-*EFR*KD was incubated with 2 µg of Trx-6xHis-*IOS1*KD in a binding buffer (20 mM Tris-HCl, 200 mM NaCl, 1 mM EDTA, pH 7.4) under agitation at 4°C. After 2 h, 50 µL of amylose resin beads (GE Healthcare) were added, and the incubation continued for another 2 h. The beads were then washed five times with washing buffer (20 mM Tris-HCl, 200 mM NaCl, 1 mM EDTA, 0.6% Triton X-100, pH 7.4). Input and pulled-down proteins were resolved by 8% SDS-PAGE and detected by Western blotting using appropriate antibodies.

BiFC Assay

Full-length coding sequences (CDS) of *FLS2*, *EFR*, *BAK1*, *CERK1*, *LTI6b* and *IOS1* without stop codon amplified from cDNA of *Arabidopsis* Col-0 were inserted into the entry vector pCR8/GW/TOPO, and subcloned into YN (pEarleyGate201-YN) or YC (pEarleyGate202-YC) vectors (Lu *et al.*, 2010c) through LR reaction (Invitrogen). The constructs were transfected into *Arabidopsis* protoplasts by polyethylene glycol (Sigma) for transient expression (Yoo *et al.*, 2007). Sixteen hours later transfected cells were treated with or without 100 nM flg22, 100 nM elf26, or 0.2 mg/mL chitin for 10 min before being imaged using a confocal laser scanning microscope (Zeiss LSM 780 Confocal, Carl Zeiss, Germany).

Transient Expression in *Arabidopsis* Protoplasts

For the co-immunoprecipitation, TOPO plasmid containing full-length coding sequences (CDS) of *FLS2*, *EFR*, *BAK1*, *CERK1*, *LTI6b* or *IOS1* without stop codon were recombined into GFP (pEarleyGate103) or HA (modified pEarleyGate100 with a AvrII-3xHA-SpeI fragment introduced

after the attR2 recombination site) vectors. Amplification of the CDS was performed using the primers described in Supplemental Table 2 online. All constructs were confirmed by DNA sequencing. The *35S-FLS2-GFP-His* and *35S-EFR-GFP-His* constructs were as described (Schwessinger et al., 2011).

Protein Extraction and Immunoprecipitation in *Arabidopsis* Protoplasts

Protein extraction and immunoprecipitation were performed as described (Yeh et al., 2015). Briefly, plasmids containing HA₃ or GFP tag constructs were co-transfected into *Arabidopsis* protoplasts by polyethylene glycol (Sigma) for transient expression (Yoo et al., 2007). Total proteins were extracted with 0.5 mL protein extraction buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 10% glycerol, 10 mM DTT, 10 mM EDTA, 1 mM NaF, 1 mM Na₂MoO₄·2H₂O, 1% [v/v] IGEPAL CA-630 [Sigma-Aldrich] and 1% [v/v] Roche protease inhibitor cocktail) and incubated with gentle shaking at 4°C for 1 h. Samples were then centrifuged at 14,000 rpm for 15 min at 4°C. Supernatants (1.5 mL) were adjusted to 2 mg/mL protein and incubated for 2 h at 4°C with 20 mL GFP Trap-A beads (Chromotek). Following incubation, beads were washed four times with TBS containing 0.5% (v/v) IGEPALCA-630. Total proteins (input) or immunoprecipitated proteins were separated by 10% SDS-PAGE and then transferred to a polyvinylidene fluoride membrane (Immobilon-P; Millipore). GFP and HA₃ fusion proteins were detected by immunoblotting with anti-GFP and anti-HA primary antibodies, respectively

Protein Extraction and Immunoprecipitation in *Arabidopsis*

The protocol for protein extraction was described in Roux et al., (2011). *Arabidopsis* seedlings grown in liquid ½ MS medium were used for immunoprecipitation assays. For immunoprecipitation of endogenous BAK1, supernatants were incubated with 25 µL true-blot anti-rabbit Ig beads (Ebioscience) and 20 µL anti-BAK1 antibody (Schulze et al., 2010) for 4 h at 4 °C. For immunoprecipitation of IOS1-GFP, supernatants were incubated with 50-200 µL of anti-GFP magnetic beads (Miltenyi Biotec) for 2 h at 4 °C (Kadota et al., 2014; Kadota et al., 2016). Following incubation, beads were washed 3-5 times with extraction buffer, before adding SDS loading buffer (Schwessinger et al., 2011).

SDS-PAGE and Immunoblotting

Eight to 10 % SDS-PAGE gels were run at 80-140 V for 2 h before electroblotting on PVDF membrane (Millipore) at 100 V for 1 h at 4 °C. Membranes were rinsed in TBS and blocked in 5 % (w/v) nonfat milk powder in TBS-Tween 0.1 % (v/v) for 2 h. Primary antibodies were diluted in TBS-Tween solution to the following ratios: mouse anti-His (Santa Cruz Biotechnology sc-8036) 1:1000; rabbit anti-MBP 1:4000 (Sigma SAB2108749); rabbit anti-GFP (Santa Cruz Biotechnology sc-9996) 1:3000; mouse anti-HA (Santa Cruz Biotechnology sc-7392) 1:3000; anti-BAK1 1:500 and anti-FLS2 1:1000 (Schwessinger et al., 2011) and incubated overnight. Membranes were washed 3 times in TBS-Tween before 1 h incubation with secondary antibodies anti-mouse-HRP (Santa Cruz Biotechnology sc-2005) 1:3000 or anti-rabbit-HRP (Santa Cruz Biotechnology sc-2004) 1:3000. Signals were visualized using an enhanced chemiluminescence system (Immobilon Western, Millipore) and a LAS-3000 (Fujifilm, Tokyo, Japan) scanner following manufacturer's instructions.

Mass Spectrometry

Proteins were separated by SDS page (Nupage precast gel system, Invitrogen) and after staining with CBB (seeblue safe stain, Invitrogen), the proteins were cut out and were digested by trypsin as described previously (Ntoukakis et al., 2009). LC-MS/MS analysis was performed using a LTQ-Orbitrap mass spectrometer (Thermo Scientific) and a nanoflow-HPLC system (nanoAcquity; Waters) as described previously (Ntoukakis et al., 2009). The entire TAIR10 (www.Arabidopsis.org) and E. coli O157 databases were searched using Mascot (with the inclusion of sequences of common contaminants, such as keratins and trypsin). Parameters were set for 65 ppm peptide mass tolerance and allowing for Met oxidation and two missed tryptic cleavages. Carbamidomethylation of Cys residues was specified as a fixed modification, and oxidized Met and phosphorylation of Ser or Thr residues were allowed as variable modifications. Scaffold (v2_06_01; Proteome Software) was used to validate MS/MS-based peptide and protein identifications.

***In vitro* Kinase Assay**

The *in vitro* kinase assay was performed as described previously (Singh et al., 2012b). Briefly, 2 µg of purified Trx-6xHis-IOS1KD and Trx-6xHis-IOS1KDM were incubated for 30 min at 28 °C in 30 µL kinase buffer (50 mM Tris-Cl pH 7.5, 50 mM KCl, 2 mM DTT, 10% v/v glycerol, 5 mM MnCl₂, 5 mM MgCl₂). Phosphorylation was initiated with the addition of 10 mM ATP and

terminated by adding 30 μ L of 2 x SDS page loading buffer. Of these, 30 μ L were separated on a 8% polyacrilamide gel and the phosphorylation level of the proteins was detected using the Pro-Q Diamond Phosphoprotein Gel Stain (Invitrogen) according to manufacturer's instructions. The fluorescent signal was imaged using a Typhoon 9400 scanner (Amersham Biosciences), and the same gel was subsequently stained for total protein with Coomassie Brilliant Blue.

Accession Numbers

Sequence data from this article can be found in the *Arabidopsis* Genome Initiative under accession number(s): *IOS1* (At1g51800), *FRK1* (At2g19190), and *UBQ10* (At4g05320).

Supplemental Data

Figure 1. The mutant *ios1-1* is a knock-out while *ios1-2* and *ios1-3* produce some *IOS1* transcripts.

Figure 2. Susceptibility phenotypes of *ios1-2* and *bak1-5* to *Pst* DC3000.

Figure 3. *ios1* mutants demonstrate increased susceptibility to *Pst* DC3000 *hrcC*.

Figure 4. Resistance of *ios1* mutants to necrotrophic fungal pathogens.

Figure 5. *IOS1* mRNA expression levels in 2 independent *IOS1* over-expression lines.

Figure 6. Stomatal innate immunity in *ios1* mutants.

Figure 7. Expression of *IOS1* is up-regulated by bacterial MAMPs.

Figure 8. Early PTI responses.

Figure 9. Bimolecular fluorescence complementation analyses of *IOS1* interactions with *EFR* and *BAK1*.

Figure 10. *FLS2-FLS2* and *IOS1-IOS1* dimerizations.

Figure 11. Complementation of defective MAPK activation in *ios1-1* and *ios1-2* mutants by *IOS1-GFP*.

Figure 12. *IOS1 in vitro* autophosphorylation.

Figure 13. *A. brassicicola*-mediated lesions in lines overexpressing *IOS1*.

Figure 14. BABA does not regulate ligand-induced *FLS2-BAK1* association.

Table 1. Identification of IOS1 Tryptic Peptides by HPLC-ESI-MS/MS Analysis of EFR Immunoprecipitates.

Table 2. Primer sequences used in this study.

ACKNOWLEDGEMENTS

We thank ABRC for providing seeds and constructs. We are grateful to B.N. Kunkel, J. Glazebrook and C.Y. Chen for the pathogens. We also thank P. He for providing the BIK1 construct and *bik1* mutant seeds, and D. Chinchilla for FLS2 antibody. We appreciate the help from the staff of Technology Commons, College of Life Science, National Taiwan University, in microscopy and for qRT-PCR equipment. We thank the Proteomics Core Laboratory sponsored by the Institute of Plant and Microbial Biology and the Agricultural Biotechnology Research Center, Academia Sinica for Mass spectrometric protein identifications and analyses. We also thank A. Jones and J. Sklenar from TSL Proteomics for their excellent service. We acknowledge Y.S. Cheng and I.F. Chang for their technical support in protein expression and analyses, B. Schulze, M. Desclos-Theveniau and members of Zimmerli's laboratory for critical comments. This work was supported by the National Science Council of Taiwan grants 99-2628-B-002-053-MY3 and 102-2628-B-002-011-MY3 (to L.Z.), the Frontier and Innovative Research grant of the National Taiwan University code number 99R70436 (to L.Z.), The European Research Council (ERC) and The Gatsby Charitable Foundation (to C.Z.). Y.K. was supported by fellowships from KAKENHI (#23580068), the Excellent Young Researcher Overseas Visit Program and the Uehara memorial foundation. M.R. was part of the John Innes Centre/The Sainsbury Laboratory PhD Rotation Program.

AUTHOR CONTRIBUTIONS

Y.H.Y., D.P., Y.K., C.Z. and L.Z. designed the research. Y.H.Y., D.P., Y.K., Y.C.H., P.Y.H., C.N.T., M.R., H.C.C., T.C.C., and P.W.C performed research. Y.H.Y., D.P., Y.K., P.Y.H., M.R.,

C.Z. and L.Z. analyzed data, and L.Z. wrote the article with contributions from Y.H.Y., D.P., Y.K., and C.Z.

This article is a resubmission after the retraction of the 2014 article “The *Arabidopsis* malectin-like leucine-rich repeat receptor-like kinase IOS1 associates with the pattern recognition receptors FLS2 and EFR and is critical for priming of pattern-triggered immunity. Chen CW, Panzeri D, Yeh YH, Kadota Y, Huang PY, Tao CN, Roux M, Chien SC, Chin TC, Chu PW, Zipfel C, Zimmerli L. *Plant Cell*. 26: 3201-3219. Retraction comments visible at [Plant Cell](https://doi.org/10.1105/PC.115.01563). (2015) 27: 1563. Shortly, experiments performed by the first author Chen CW in the 2014 article had to be redone.

For this new version of the manuscript, Y.H.Y performed the experiments for the new Figures 1A and B, 3C and D, 4A, C and D, 5A, 8A and C and Figures 9A and B. Meanwhile, Y.C.H. performed experiments for the new Figures 2C (elf18), 7E and 10C, while new Figures 2B and D and 8B were done by Y.H.Y and Y.C.H. together. Experiments for the retained Figures 1C and D were performed by Y.H.Y. and C.N.T. respectively. Retained Figure 2A was done by H.C.C. Experiments for retained Figures 2C (flg22), 3A and B were performed by Y.H.Y. Figure 4B was provided by P.Y.H., while retained Figure 5B was done by Y.K. Retained Figures 6A and C were provided by Y.K. and experiments for Figures 6 B and D were performed by D.P. Retained Figures 7A-D are from D.P. Retained Figure 10A was provided by Y.H.Y. and C.N.T. and retained Figure 10B was provided by H.C.C. and T.C.C. Finally experiments for retained Figure 10D were performed by C.N.T. The new Supplemental Figures 2, 4, 5, 7, 8C-E, 9, 10, 11, 13, and 14 were provided by Y.H.Y.. Data provided by P.W.C. were used for retained Supplemental Figure 1. Retained Supplemental Figures 3 and 6 were done by C.N.T. Retained Supplemental Figures 8A and B are from Y.H.Y. Experiments for retained Supplemental Figure 12 were performed by D.P.. Data for retained Supplemental Table 1 are from M.R.

FIGURE LEGENDS

Figure 1. A Critical Role for IOS1 in *Arabidopsis* Resistance to Hemi-Biotrophic Bacteria.

(A) Disease symptoms in *ios1* mutants. Five-week-old *Arabidopsis* were dip-inoculated in a bacterial solution of 10^6 cfu/mL *Pst* DC3000 or 5×10^5 cfu/mL *Psm* ES4326. Symptoms were evaluated at 3 dpi. These experiments were repeated at least twice with similar results.

(B) Bacterial growth in *ios1* mutants. Five-week-old *Arabidopsis* were dip-inoculated as in (A) and bacterial titers were evaluated at 2 dpi. Values are means \pm SE of 2 independent experiments each consisting of 3 plants ($n = 6$). Asterisks indicate a significant difference to the respective WT control based on a *t* test ($P < 0.01$).

(C) Growth of *Pst* DC3000 in lines over-expressing *IOS1*. Bacterial titers in 5-week-old Col-0 and *IOS1* over-expression lines OE1 and OE3 were determined at 3 dpi with 10^6 cfu/mL *Pst* DC3000. Values are means \pm SE of 3 independent biological replicates each with 3 plants ($n = 9$). Asterisks indicate a significant difference to Col-0 WT based on a *t* test (* $P < 0.01$).

(D) Stomatal innate immunity in lines over-expressing *IOS1*. Stomatal apertures in leaf epidermal peels from 5-week-old Col-0 and *IOS1* over-expression lines OE1 and OE3 were analyzed after 1.5 h or 3 h exposure to MgSO_4 (Mock) or 10^8 cfu/mL *Pst* DC3000. Values are shown as means \pm SE of 3 independent experiments each consisting of at least 60 stomata ($n > 180$). Asterisks indicate a significant difference to respective mock controls based on a *t* test analysis ($P < 0.001$).

Figure 2. Altered Late PTI Responses in *ios1* Mutants and *IOS1*-OE Lines.

(A and C) Callose deposition. Leaves of 5-week-old *ios1-1* and *ios1-2* (A) were syringe infiltrated with $1 \mu\text{M}$ flg22 or elf26 and samples were collected 9 h (flg22) or 24 h (elf26) later for aniline blue staining. For *IOS1*-OE lines (C), leaves of 5-week-old *Arabidopsis* or 10-day-old seedlings were respectively syringe infiltrated with $1 \mu\text{M}$ flg22 or treated with 100 nM elf18 and samples were collected 6 h (flg22) or 16 h (elf18) later for aniline blue staining. Mock samples were infiltrated with MgSO_4 (for flg22 and elf26) or water (for elf18). Numbers under the pictures are average \pm SD of the number of callose deposits per square millimeter from at least 2 independent experiments each consisting of 6 plants ($n = 12$). White bar = $200 \mu\text{m}$.

(B and D) PTI-responsive gene *FRK1* up-regulation. Relative *FRK1* expression levels were evaluated at 30 min post treatment (mpt) with 100 nM flg22 or elf18 in *ios1-1* and *ios1-2*

mutants (B) or at 45 mpt with 50 nM flg22 or elf18 in *IOS1*-OE lines (D). *UBQ10* was used for normalization. Relative gene expression levels were compared to WT mock (*Ler-0* or *Col-0*) (defined value of 1) by qRT-PCR analyses. The values are means \pm SD of 2 independent experiments each with 3 batches of 20 plantlets ($n = 6$). Asterisks indicate a significant difference to WT controls based on a *t* test ($P < 0.01$).

Figure 3. Early PTI Responses.

(A) ROS production in *ios1* mutants. Responsiveness of 5-week-old *Ler-0* and *Col-0* WT controls and respective mutants *ios1-1* and *ios1-2* to 10 nM flg22. *bak1-4* was used as a negative control. Production of ROS in *Arabidopsis* leaf discs is expressed as relative light units (RLU) for a period of 30 min after elicitation. Values are means \pm SE of 3 independent experiments each with 6 leaf discs ($n = 18$). Differences between *ios1* mutants and WT were not statistically significant based on a *t* test ($P < 0.01$).

(B) ROS production in *IOS1*-OE lines. Responsiveness of 5-week-old over-expression lines OE1 and OE3 and *Col-0* WT control to 10 nM flg22. Production of ROS in *Arabidopsis* leaf discs is expressed as relative light units (RLU) for a period of 30 min after elicitation. Values are means \pm SE of 3 independent experiments each with 6 leaf discs ($n = 18$). Differences between OE lines and WT were not statistically significant based on a *t* test ($P < 0.01$).

(C) MPK activation in *ios1* mutants. Ten-day-old *Ler-0* and *ios1-1* or *Col-0* and *ios1-2* were treated with 100 nM flg22 for 5 min. Immunoblot analysis using phospho-p44/42 MPK antibody is shown in top panel. Lines indicate the positions of MPK3 and MPK6. Coomassie Brilliant Blue-staining is used to estimate equal loading in each lane (bottom panel). Similar results were observed in another independent repeat.

(D) MAPK activation in *IOS1*-OE lines. Ten-day-old *Col-0* and *IOS1* over-expression lines OE1 and OE3 were treated with 50 nM flg22 for 5 min. Immunoblot analysis using phospho-p44/42 MAP kinase antibody is shown in the top panel. Lines indicate the positions of MPK3 and MPK6. Coomassie Brilliant Blue-staining is used to estimate equal loading in each lane (bottom panel). Similar results were observed in another independent repeat.

Figure 4. IOS1 Localization, Pull-Down and BiFC Analyses of IOS1 Interaction with PRRs.

(A) Subcellular localization of IOS1-GFP fusion protein in *Arabidopsis* mesophyll protoplasts. *IOS1-GFP* expression was driven by the cauliflower mosaic virus 35S promoter and transiently expressed in *Arabidopsis* mesophyll protoplasts. The images of the GFP fluorescence (GFP), the chlorophyll autofluorescence (chlorophyll), the bright-field image (bright), the plasma membrane marker (pm-rk CD3-1007)-mCherry fluorescence localization, and the combined images (merged) are shown. Similar observations were made in another independent repeat. Scale bars represent 10 μm .

(B) *In vitro* MBP pull-down assay of IOS1 interaction with FLS2 and EFR. *E. coli* expressed MBP (negative control), MBP-FLS2KD, or MBP-EFRKD were incubated with Trx-6xHis-IOS1KD and pulled down with amylose resin beads. Input and bead-bound proteins were analyzed by immunoblotting with specific antibodies. Experiments were repeated 3 times with similar results.

(C) Bimolecular fluorescence complementation analyses of IOS1 interactions with FLS2 and BAK1. *Arabidopsis* protoplasts were co-transfected with BAK1-YFP^N + FLS2-YFP^C, IOS1-YFP^N + FLS2-YFP^C and IOS1-YFP^N + BAK1-YFP^C and treated with (+) or without (-) 100 nM flg22 for 10 min. The YFP fluorescence (yellow), chlorophyll autofluorescence (red), bright field and the combined images were visualized under a confocal microscope 16 h after transfection. Images are representative of multiple protoplasts. Experiments were repeated at least twice with similar results. Scale bars represent 10 μm .

(D) Bimolecular fluorescence complementation of LTI6b and IOS1 interaction. *Arabidopsis* protoplasts were co-transfected with LTI6b-YFP^N + LTI6b-YFP^C or IOS1-YFP^N + LTI6b-YFP^C and treated with (+) or without (-) 100 nM flg22 for 10 min. The YFP fluorescence (yellow), chlorophyll autofluorescence (red), bright field and the combined images were visualized under a confocal microscope 16 h after transfection. Images are representative of multiple protoplasts. Experiments were repeated twice with similar results. Scale bars represent 10 μm .

Figure 5. IOS1 Associates with Unstimulated and Stimulated FLS2, EFR and BAK1.

(A) Co-immunoprecipitation of IOS1, FLS2, EFR and BAK1 proteins. *Arabidopsis* protoplasts expressing IOS1-GFP and FLS2-HA₃ (lane 2 and 3), IOS1-GFP and EFR-HA₃ (lane 5 and 6) or IOS1-GFP and BAK1-HA₃ (lane 8, 9 and 10) constructs were treated (+) or not (-) with 100 nM flg22 or elf18 for 10 min. LTI6b-GFP, a known plasma membrane protein was used as a control to illustrate that FLS2-HA₃, EFR-HA₃ and BAK1-HA₃ do not associate with GFP at the plasma membrane (lane 1, 4 and 7). Total proteins (input) were subjected to immunoprecipitation with GFP trap beads followed by immunoblot analysis with anti-HA antibodies to detect FLS2-HA₃, EFR-HA₃ and BAK1-HA₃. Anti-GFP antibodies detect IOS1-GFP and LTI6b-GFP. Experiments were repeated twice with similar results.

(B) Co-immunoprecipitation of FLS2, BAK1 and IOS1 proteins in *Arabidopsis*. Transgenic *Arabidopsis* seedlings over-expressing IOS1-GFP (OE3) were treated (+) or not (-) with 100 nM flg22 for 10 min. Total proteins (input) were subjected to immunoprecipitation with anti-GFP magnetic beads followed by immunoblot analysis with anti-FLS2 antibodies, anti-BAK1 antibodies or anti-GFP antibodies to detect FLS2, BAK1 and IOS1-GFP. Untransformed Col-0 *Arabidopsis* tissue was used as a control to show that FLS2 and BAK1 do not adhere non-specifically to anti-GFP magnetic beads (lane 1). LTI6b, a known plasma membrane protein was used as a control to illustrate that FLS2 and BAK1 do not associate with GFP at the plasma membrane (lane 2). This experiment is one of 2 independent replicates.

Figure 6. IOS1 Regulates Ligand-Induced FLS2/BAK1 Association.

(A and B) Ligand-dependent association of FLS2 to BAK1 is reduced in *ios1-2* mutant. Col-0 or *ios1-2* seedlings were treated (+) or not (-) with 100 nM flg22 for 10 min. Total proteins (input) were subjected to immunoprecipitation (IP) with anti-BAK1 antibodies and IgG beads followed by immunoblot analysis using anti-FLS2 and anti-BAK1 antibodies. For (A), Coomassie Brilliant Blue (CBB) is used to estimate equal loading (bottom panel). The experiment shown in (A) is one of 3 independent replicates pooled together in (B).

(C and D) Ligand-dependent association of FLS2 to BAK1 is augmented in the *IOS1*-OE3 line. Col-0 or OE3 seedlings were treated with MgSO₄ (0), 10 or 50 nM flg22 for 10 min. Total proteins (input) were subjected to immunoprecipitation (IP) with anti-BAK1 antibodies and IgG beads followed by immunoblot analysis using anti-FLS2 and anti-BAK1 antibodies. The

experiment shown in (C) is one of 3 independent replicates pooled together in (D). For both (B) and (D), signals were evaluated with the ImageJ software. Values are means \pm SD of 3 independent biological replicates ($n = 3$). Different letters denote significant difference based on a one-way ANOVA with post-hoc Tukey HSD (Honestly Significant Difference) ($P < 0.05$).

Figure 7. IOS1 Functions in a BAK1-Dependent, but BIK1-Independent Manner in the FLS2 Complex.

(A-D) Western blot analysis of BIK1 phosphorylation revealed by gel mobility shift. Non-phosphorylated (BIK1) and phosphorylated (pBIK1) BIK1 signals are indicated. Protoplasts from Col-0 leaves and *ios1-2* (A and C) or OE3 (B and D) were treated 4 h after transfection using 0.75 μ M flg22 for 3.5, 7 and 10 min. The reaction was stopped by immersion in liquid nitrogen following concentration by low speed centrifugation. Experiments were repeated at least 5 times with similar results. For (C and D), phosphorylated over non-phosphorylated BIK1 fractions were calculated by measuring digital signals with the ImageJ software. Values are means \pm SD of 5 independent biological replicates ($n = 5$). For each time point, differences between WT and the *ios1-2* mutant or the OE3 line were not statistically significant based on a *t* test ($P < 0.01$).

(E) Callose deposition upon elicitation with flg22. Fourteen-day-old Col-0 WT, *IOS1*-OE3 (OE), *bak1-5* or *bik1* mutants and *IOS1*-OE in *bak1-5* or *bik1* mutant background were treated with 100 nM flg22 and samples were collected 16 h later for aniline blue staining. Each bars are averages \pm SE of callose deposits per square millimeters from 2 independent experiments each with 6 plants ($n = 12$). For *IOS1*-OE lines in the *bak1-5* and *bik1* backgrounds, data represent 2 independent transformation events for each genotype. Different letters denote significant differences among different lines based on a one-way ANOVA with post-hoc Tukey HSD ($P < 0.01$).

Figure 8. A Role for IOS1 in the Chitin Response.

(A) MPK activation upon elicitation with chitin. Fourteen-day-old seedlings from *Ler-0* or Col-0 WT, *ios1-1* or *ios1-2* were syringe-infiltrated with 0.2 mg/mL chitin for 5 min. Immunoblot

analysis using phospho-p44/42 MPK antibody is shown in top panel. Lines indicate the positions of MPK3 and MPK6. Coomassie Brilliant Blue-staining is used to estimate equal loading in each lane (bottom panel). An independent experiment showed similar results.

(B) Callose deposition upon elicitation with chitin. Fourteen-day-old seedlings from *Ler-0* and *ios1-1* or Col-0 and *ios1-2* were treated with 0.2 mg/mL chitin and samples were collected 16 h later for aniline blue staining. Numbers are averages \pm SE of callose deposits per square millimeters from 2 independent experiments each including 6 seedlings ($n = 12$). Asterisks indicate a significant difference to WT controls based on a *t* test ($P < 0.01$).

(C) *B. cinerea*-mediated lesions. *Arabidopsis* leaves of Col-0 and *IOS1* overexpression lines were droplet-inoculated (10 μ L) with 10^5 *B. cinerea* spores/mL and lesion diameters were evaluated at 3 dpi. Data are average \pm SE of lesion diameters from 2 independent experiments each with 6 plants ($n = 12$). Asterisks indicate a significant difference to WT controls based on a *t* test ($P < 0.01$).

Figure 9. IOS1 Associates with CERK1.

(A) Co-immunoprecipitation of IOS1 with CERK1 proteins in *Arabidopsis* protoplasts.

Arabidopsis protoplasts expressing CERK1-GFP and CERK1-HA₃ (lane 1 and 2), IOS1-GFP and CERK1-HA₃ (lane 3 and 4), Empty Vector (EV)-GFP and CERK1-HA₃ (lane 5), or LTI6b-GFP and CERK1-HA₃ (lane 6) constructs were treated with (+) or without (-) 0.2 mg/mL chitin for 10 min. Total proteins (input) were subjected to immunoprecipitation (IP) with GFP trap beads followed by immunoblot analysis with anti-HA antibodies to detect CERK1-HA₃. EV-GFP and LTI6b-GFP, a known plasma membrane protein, were used as controls to illustrate that CERK1-HA₃ does not stick to GFP beads or associate with GFP at the plasma membrane, respectively. This experiment was repeated twice with similar results.

(B) Bimolecular fluorescence complementation analyses of IOS1 interactions with CERK1.

Arabidopsis protoplasts were co-transfected with CERK1-YFP^N + CERK1-YFP^C and CERK1-YFP^N + IOS1-YFP^C, and treated with (+) or without (-) 0.2 mg/mL chitin for 10 min. The YFP fluorescence (yellow), chlorophyll autofluorescence (red), bright field and the combined images were visualized under a confocal microscope 16 h after transfection. Images are representative of

multiple protoplasts. Scale bars represent 10 μm . At least 2 independent experiments were performed with similar results.

Figure 10. BABA Action is Defective in *ios1* Mutants.

(A) BABA-induced resistance. Bacterial titers in 5-week-old *Ler-0*, *ios1-1*, Col-0 and *ios1-2* were determined at 2 dpi with 10^6 cfu/mL *Pst* DC3000 or 5×10^5 cfu/mL *Psm* ES4326. Two days before bacterial inoculation, plants were soil-drenched with water as a control or 225 μM BABA. Values are means \pm SE of 3 independent experiments each with 3 plants ($n = 9$). Asterisks indicate a significant difference to respective water-treated control based on a *t* test ($P < 0.01$).

(B) BABA priming of PTI-mediated callose deposition. Leaves of water- or BABA-pretreated (225 μM) *Ler-0* and *ios1-1* or Col-0 and *ios1-2* were syringe-infiltrated with 1 μM flg22 and samples were collected 6 h later for aniline blue staining. Values are average \pm SD from 3 independent experiments each consisting of 9 plants ($n = 27$). Asterisks indicate a significant difference to water-treated respective controls based on a *t* test ($P < 0.01$).

(C) BABA priming of PTI-mediated *FRK1* expression. Ten-day-old *Ler-0* and *ios1-1* or Col-0 and *ios1-2* seedlings grown on $\frac{1}{2}$ MS medium supplemented with 30 μM BABA (BABA) or not (Water) were submerged with water (Mock) or 1 μM flg22, and *FRK1* expression levels were analyzed 60 min later by qRT-PCR. *UBQ10* was used for normalization. Relative gene expression levels were compared to respective water + mock-treated WT (defined value of 1). Values are means \pm SD of 2 independent experiments each with 3 plants ($n = 6$). Asterisks indicate a significant difference to water-treated respective controls based on a *t* test ($P < 0.01$).

(D) BABA-inhibition of bacteria-mediated stomatal re-opening. Stomatal apertures in epidermal peels from water- (W) or BABA-treated (225 μM) (B) *Ler-0* and *ios1-1* or Col-0 and *ios1-2* were analyzed after 1.5 h and 6 h exposure to MgSO_4 (Mock) or 10^8 *Pst* DC3000. Results are shown as mean \pm SE of 3 independent experiments each consisting of at least 60 stomata ($n > 180$). Asterisks indicate a significant difference to respective mock controls based on a *t* test analysis ($P < 0.001$).

REFERENCES

- Asai, T., Tena, G., Plotnikova, J., Willmann, M.R., Chiu, W.L., Gomez-Gomez, L., Boller, T., Ausubel, F.M., and Sheen, J. (2002). MAP kinase signalling cascade in *Arabidopsis* innate immunity. *Nature* 415: 977-983.
- Beckers, G.J.M., Jaskiewicz, M., Liu, Y.D., Underwood, W.R., He, S.Y., Zhang, S.Q., and Conrath, U. (2009). Mitogen-Activated Protein Kinases 3 and 6 Are Required for Full Priming of Stress Responses in *Arabidopsis thaliana*. *Plant Cell* 21: 944-953.
- Böhm, H., Albert, I., Fan, L., Reinhard, A., and Nürnberger, T. (2014). Immune receptor complexes at the plant cell surface. *Curr Opin Plant Biol.* 20: 47-54.
- Boller, T., and Felix, G. (2009). A Renaissance of Elicitors: Perception of Microbe-Associated Molecular Patterns and Danger Signals by Pattern-Recognition Receptors. *Annu. Rev. Plant Biol.* 60: 379-406.
- Boudsocq, M., Willmann, M.R., McCormack, M., Lee, H., Shan, L., He, P., Bush, J., Cheng, S.H., and Sheen, J. (2010). Differential innate immune signalling via Ca²⁺ sensor protein kinases. *Nature* 464: 418-422.
- Brooks, D.M., Hernandez-Guzman, G., Kloek, A.P., Alarcon-Chaidez, F., Sreedharan, A., Rangaswamy, V., Penaloza-Vazquez, A., Bender, C.L., and Kunkel, B.N. (2004). Identification and characterization of a well-defined series of coronatine biosynthetic mutants of *Pseudomonas syringae* pv. tomato DC3000. *Mol. Plant Microbe Interact.* 17: 162-174.
- Bücherl, C.A., van Esse, G.W., Kruis, A., Luchtenberg, J., Westphal, A.H., Aker, J., van Hoek, A., Albrecht, C., Borst, J.W., and de Vries, S.C. (2013). Visualization of BRI1 and BAK1(SERK3) membrane receptor heterooligomers during brassinosteroid signaling. *Plant Physiol.* 162: 1911-1125.
- Cao, Y., Liang, Y., Tanaka, K., Nguyen, C.T., Jedrzejczak, R.P., Joachimiak, A., and Stacey, G. (2014). The kinase LYK5 is a major chitin receptor in *Arabidopsis* and forms a chitin-induced complex with related kinase CERK1. *Elife.* doi: 10.7554/eLife.03766.

- Chinchilla, D., Zipfel, C., Robatzek, S., Kemmerling, B., Nurnberger, T., Jones, J.D.G., Felix, G., and Boller, T. (2007). A flagellin-induced complex of the receptor FLS2 and BAK1 initiates plant defence. *Nature* 448: 497-500.
- Cohen, Y.R. (2002). beta-aminobutyric acid-induced resistance against plant pathogens. *Plant Dis.* 86: 448-457.
- Conrath, U., Beckers, G.J.M., Flors, V., Garcia-Agustin, P., Jakab, G., Mauch, F., Newman, M.A., Pieterse, C.M.J., Poinssot, B., Pozo, M.J., Pugin, A., Schaffrath, U., Ton, J., Wendehenne, D., Zimmerli, L., and Mauch-Mani, B. (2006). Priming: Getting ready for battle. *Mol. Plant Microbe Interact.* 19: 1062-1071.
- Cook, D.N., Pisetsky, D.S., and Schwartz, D.A. (2004). Toll-like receptors in the pathogenesis of human disease. *Nat. Immunol.* 5: 975-979.
- Cutler, S.R., Ehrhardt, D.W., Griffitts, J.S., and Somerville, C.R. (2000). Random GFP::cDNA fusions enable visualization of subcellular structures in cells of *Arabidopsis* at a high frequency. *Proc. Natl. Acad. Sci. USA* 97: 3718-3723.
- Desclos-Theveniau, M., Arnaud, D., Huang, T.Y., Lin, G.J., Chen, W.Y., Lin, Y.C., and Zimmerli, L. (2012). The *Arabidopsis* lectin receptor kinase LecRK-V.5 represses stomatal immunity induced by *Pseudomonas syringae* pv. tomato DC3000. *PLoS Pathog.* 8: e1002513.
- Deslandes, L., and Rivas, S. (2012). Catch me if you can: bacterial effectors and plant targets. *Trends Plant Sci.* 17: 644-655.
- Durrant, W.E., and Dong, X. (2004). Systemic acquired resistance. *Annu. Rev. Phytopathol.* 42: 185-209.
- Feng, F., and Zhou, J.M. (2012). Plant-bacterial pathogen interactions mediated by type III effectors. *Curr. Opin. Plant Biol.* 15: 469-476.
- Feng, F., Yang, F., Rong, W., Wu, X.G., Zhang, J., Chen, S., He, C.Z., and Zhou, J.M. (2012). A *Xanthomonas* uridine 5'-monophosphate transferase inhibits plant immune kinases. *Nature* 485: 114-U149.

- Gassmann, W., and Bhattacharjee, S. (2012). Effector-triggered immunity signaling: from gene-for-gene pathways to protein-protein interaction networks. *Mol. Plant Microbe Interact.* 25: 862-868.
- Gimenez-Ibanez, S., Hann, D.R., Ntoukakis, V., Petutschnig, E., Lipka, V., and Rathjen, J.P. (2009). AvrPtoB targets the LysM receptor kinase CERK1 to promote bacterial virulence on plants. *Curr Biol.* 19: 423-429.
- Girardin, S.E., Sansonetti, P.J., and Philpott, D.J. (2002). Intracellular vs extracellular recognition of pathogens--common concepts in mammals and flies. *Trends Microbiol.* 10: 193-199.
- Gomez-Gomez, L., and Boller, T. (2000). FLS2: an LRR receptor-like kinase involved in the perception of the bacterial elicitor flagellin in *Arabidopsis*. *Mol. Cell* 5: 1003-1011.
- Gou, X.P., He, K., Yang, H., Yuan, T., Lin, H.H., Clouse, S.D., and Li, J. (2010). Genome-wide cloning and sequence analysis of leucine-rich repeat receptor-like protein kinase genes in *Arabidopsis thaliana*. *BMC Genomics* 11: 19.
- Halter, T., Imkampe, J., Mazzotta, S., Wierzba, M., Postel, S., Bücherl, C., Kiefer, C., Stahl, M., Chinchilla, D., Wang, X., Nürnberger, T., Zipfel, C., Clouse, S., Borst, J.W., Boeren, S., de Vries, S.C., Tax, F., and Kemmerling, B. (2014). The leucine-rich repeat receptor kinase BIR2 is a negative regulator of BAK1 in plant immunity. *Curr Biol.* 24: 134-143.
- Haweker, H., Rips, S., Koiwa, H., Salomon, S., Saijo, Y., Chinchilla, D., Robatzek, S., and von Schaewen, A. (2010). Pattern recognition receptors require N-glycosylation to mediate plant immunity. *J. Biol. Chem.* 285: 4629-4636.
- He, P., Shan, L., and Sheen, J. (2007). The use of protoplasts to study innate immune responses. *Methods Mol. Biol.* 354: 1-9.
- Heese, A., Hann, D.R., Gimenez-Ibanez, S., Jones, A.M.E., He, K., Li, J., Schroeder, J.I., Peck, S.C., and Rathjen, J.P. (2007). The receptor-like kinase SERK3/BAK1 is a central regulator of innate immunity in plants. *Proc. Natl. Acad. Sci. USA* 104: 12217-12222.
- Hodge, S., Thompson, G.A., and Powell, G. (2005). Application of DL-beta-aminobutyric acid (BABA) as a root drench to legumes inhibits the growth and reproduction of the pea aphid *Acyrtosiphon pisum* (Hemiptera: Aphididae). *Bull Entomol. Res.* 95: 449-455.

- Hok, S., Allasia, V., Andrio, E., Naessens, E., Ribes, E., Panabières, F., Attard, A., Ris, N., Clément, M., Barlet, X., Marco, Y., Grill, E., Eichmann, R., Weis, C., Hüchelhoven, R., Ammon, A., Ludwig-Müller, J., Voll, L.M., and Keller, H. (2014) The receptor kinase IMPAIRED OOMYCETE SUSCEPTIBILITY1 attenuates abscisic acid responses in *Arabidopsis*. *Plant Physiol.* 166: 1506-1518.
- Hok, S., Danchin, E.G.J., Allasia, V., Panabieres, F., Attard, A., and Keller, H. (2011). An *Arabidopsis* (malectin-like) leucine-rich repeat receptor-like kinase contributes to downy mildew disease. *Plant Cell Environ.* 34: 1944-1957.
- Huang, T.Y., Desclos-Theveniau, M., Chien, C.T., and Zimmerli, L. (2012). *Arabidopsis thaliana* transgenics overexpressing IBR3 show enhanced susceptibility to the bacterium *Pseudomonas syringae*. *Plant Biol.* 15: 832-840.
- Huang, P.Y., Yeh, Y.H., Liu, A.C., Cheng, C.P., and Zimmerli, L. (2014) The *Arabidopsis* LecRK-VI.2 associates with the pattern-recognition receptor FLS2 and primes *Nicotiana benthamiana* pattern-triggered immunity. *Plant J.* 79: 243-55.
- Huang, P.Y., and Zimmerli, L. (2014). Enhancing crop innate immunity: new promising trends. *Front Plant Sci.* 5: 624.
- Huffaker, A., Pearce, G., and Ryan, C.A. (2006). An endogenous peptide signal in *Arabidopsis* activates components of the innate immune response. *Proc. Natl. Acad. Sci. USA* 103: 10098-10103.
- Jakab, G., Cottier, V., Toquin, V., Rigoli, G., Zimmerli, L., Metraux, J.P., and Mauch-Mani, B. (2001). beta-Aminobutyric acid-induced resistance in plants. *Eur. J. Plant Pathol.* 107: 29-37.
- Jakab, G., Ton, J., Flors, V., Zimmerli, L., Metraux, J.P., and Mauch-Mani, B. (2005). Enhancing *Arabidopsis* salt and drought stress tolerance by chemical priming for its abscisic acid responses. *Plant Physiol.* 139: 267-274.
- Jaskiewicz, M., Conrath, U., and Peterhansel, C. (2011). Chromatin modification acts as a memory for systemic acquired resistance in the plant stress response. *EMBO Rep.* 12: 50-55.

- Kadota, Y., Macho, A.P., and Zipfel, C. (2016). Immunoprecipitation of Plasma Membrane Receptor-Like Kinases for Identification of Phosphorylation Sites and Associated Proteins. *Methods Mol. Biol.* 1363: 133-144.
- Kadota, Y., Shirasu, K., and Zipfel, C. (2015). Regulation of the NADPH Oxidase RBOHD During Plant Immunity. *Plant Cell Physiol.* 56: 1472-1480.
- Kadota, Y., Sklenar, J., Derbyshire, P., Stransfeld, L., Asai, S., Ntoukakis, V., Jones, J.D.G., Shirasu, K., Menke, F., Jones, A., and Zipfel, C. (2014). Direct Regulation of the NADPH Oxidase RBOHD by the PRR-Associated Kinase BIK1 during Plant Immunity. *Mol. Cell* 54: 1-13.
- Kemmerling, B., Halter, T., Mazzotta, S., Mosher, S., and Nurnberger, T. (2011). A genome-wide survey for *Arabidopsis* leucine-rich repeat receptor kinases implicated in plant immunity. *Front Plant Sci.* 2: 88.
- Kong, Q., Sun, T., Qu, N., Ma, J., Li, M., Cheng, Y.T., Zhang, Q., Wu, D., Zhang, Z., and Zhang, Y. (2016). Two redundant receptor-like cytoplasmic kinases function downstream of pattern recognition receptors to regulate activation of SA biosynthesis in *Arabidopsis*. *Plant Physiol.* Preview DOI:10.1104/pp.15.01954.
- Korasick, D.A., McMichael, C., Walker, K.A., Anderson, J.C., Bednarek, S.Y., and Heese, A. (2010). Novel Functions of Stomatal Cytokinesis-Defective 1 (SCD1) in Innate Immune Responses against Bacteria. *J. Biol. Chem.* 285: 23340-23348.
- Krol, E., Mentzel, T., Chinchilla, D., Boller, T., Felix, G., Kemmerling, B., Postel, S., Arents, M., Jeworutzki, E., Al-Rasheid, K.A., Becker, D., and Hedrich, R. (2010). Perception of the *Arabidopsis* danger signal peptide 1 involves the pattern recognition receptor AtPEPR1 and its close homologue AtPEPR2. *J. Biol. Chem.* 285: 13471-13479.
- Li, L., Li, M., Yu, L., Zhou, Z., Liang, X., Liu, Z., Cai, G., Gao, L., Zhang, X., Wang, Y., Chen, S., and Zhou, J.M. (2014). The FLS2-Associated Kinase BIK1 Directly Phosphorylates the NADPH Oxidase RbohD to Control Plant Immunity. *Cell Host Microbe* 15: 329-338.
- Liang, X., Ding, P., Lian, K., Wang, J., Ma, M., Li, L., Li, L., Li, M., Zhang, X., Chen, S., Zhang, Y., and Zhou, J.M. (2016). *Arabidopsis* heterotrimeric G proteins regulate immunity by directly coupling to the FLS2 receptor. *Elife* 2016 5. pii: e13568. Liu, T.,

- Liu, Z., Song, C., Hu, Y., Han, Z., She, J., Fan, F., Wang, J., Jin, C., Chang, J., Zhou, J.M., and Chai, J. (2012). Chitin-induced dimerization activates a plant immune receptor. *Science* 336: 1160-1164.
- Lu, D., Lin, W., Gao, X., Wu, S., Cheng, C., Avila, J., Heese, A., Devarenne, T.P., He, P., and Shan, L. (2011). Direct ubiquitination of pattern recognition receptor FLS2 attenuates plant innate immunity. *Science* 332: 1439-1442.
- Lu, D.P., Wu, S.J., Gao, X.Q., Zhang, Y.L., Shan, L.B., and He, P. (2010a). A receptor-like cytoplasmic kinase, BIK1, associates with a flagellin receptor complex to initiate plant innate immunity. *Proc. Natl. Acad. Sci. USA* 107: 496-501.
- Lu, D., Wu, S., He, P., and Shan, L. (2010b). Phosphorylation of receptor-like cytoplasmic kinases by bacterial flagellin. *Plant Signal Behav.* 5: 598-600.
- Lu, Q., Tang, X., Tian, G., Wang, F., Liu, K., Nguyen, V., Kohalmi, S.E., Keller, W.A., Tsang, E.W., Harada, J.J., Rothstein, S.J., and Cui, Y. (2010c). *Arabidopsis* homolog of the yeast TREX-2 mRNA export complex: components and anchoring nucleoporin. *Plant J.* 61: 259-270.
- Luna, E., Bruce, T.J., Roberts, M.R., Flors, V., and Ton, J. (2012). Next-generation systemic acquired resistance. *Plant Physiol.* 158: 844-853.
- Maekawa, T., Kufer, T.A., and Schulze-Lefert, P. (2011). NLR functions in plant and animal immune systems: so far and yet so close. *Nat. Immunol.* 12: 817-826.
- Melotto, M., Underwood, W., Koczan, J., Nomura, K., and He, S.Y. (2006). Plant stomata function in innate immunity against bacterial invasion. *Cell* 126: 969-980.
- Miya, A., Albert, P., Shinya, T., Desaki, Y., Ichimura, K., Shirasu, K., Narusaka, Y., Kawakami, N., Kaku, H., and Shibuya, N. (2007). CERK1, a LysM receptor kinase, is essential for chitin elicitor signaling in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* 104: 19613-19618.
- Monaghan, J., and Zipfel, C. (2012). Plant pattern recognition receptor complexes at the plasma membrane. *Curr. Opin. Plant Biol.* 15: 349-357.

- Navarova, H., Bernsdorff, F., Doring, A.C., and Zeier, J. (2012). Pipecolic acid, an endogenous mediator of defense amplification and priming, is a critical regulator of inducible plant immunity. *Plant Cell* 24: 5123-5141.
- Nelson, B.K., Cai, X., and Nebenfuhr, A. (2007). A multicolored set of in vivo organelle markers for co-localization studies in *Arabidopsis* and other plants. *Plant J.* 51: 1126-1136.
- Nicaise, V., Roux, M., and Zipfel, C. (2009). Recent advances in PAMP-triggered immunity against bacteria: pattern recognition receptors watch over and raise the alarm. *Plant Physiol.* 150: 1638-1647.
- Ntoukakis, V., Mucyn, T.S., Gimenez-Ibanez, S., Chapman, H.C., Gutierrez, J.R., Balmuth, A.L., Jones, A.M., and Rathjen, J.P. (2009). Host inhibition of a bacterial virulence effector triggers immunity to infection. *Science* 324: 784-787.
- Nuhse, T.S., Peck, S.C., Hirt, H., and Boller, T. (2000). Microbial elicitors induce activation and dual phosphorylation of the *Arabidopsis thaliana* MAPK 6. *J. Biol. Chem.* 275: 7521-7526.
- Oka, Y., Cohen, Y., and Spiegel, Y. (1999). Local and Systemic Induced Resistance to the Root-Knot Nematode in Tomato by DL-beta-Amino-n-Butyric Acid. *Phytopathology* 89: 1138-1143.
- Po-Wen, C., Singh, P., and Zimmerli, L. (2013). Priming of the *Arabidopsis* pattern-triggered immunity response upon infection by necrotrophic *Pectobacterium carotovorum* bacteria. *Mol. Plant Pathol.* 14: 58-70.
- Prins, T.W., Tudzynski, P., Tiedemann, A.V., Tudzynski, B., Have, A.T., Hansen, M.E., Tenberge, K., and van Kan, J.A.L. (2000) Infection strategies of *Botrytis cinerea* and related necrotrophic pathogens. J. Kronstad (Ed.), *Fungal Pathology*, Kluwer Academic Publishers, pp. 33-64.
- Ranf, S., Eschen-Lippold, L., Pecher, P., Lee, J., and Scheel, D. (2011). Interplay between calcium signalling and early signalling elements during defence responses to microbe- or damage-associated molecular patterns. *Plant J.* 68: 100-113.

- Rasmann, S., De Vos, M., Casteel, C.L., Tian, D., Halitschke, R., Sun, J.Y., Agrawal, A.A., Felton, G.W., and Jander, G. (2012). Herbivory in the previous generation primes plants for enhanced insect resistance. *Plant Physiol.* 158: 854-863.
- Robatzek, S., Chinchilla, D., and Boller, T. (2006). Ligand-induced endocytosis of the pattern recognition receptor FLS2 in *Arabidopsis*. *Genes Dev.* 20: 537-542.
- Roux, M., Schwessinger, B., Albrecht, C., Chinchilla, D., Jones, A., Holton, N., Malinovsky, F.G., Tor, M., de Vries, S., and Zipfel, C. (2011). The *Arabidopsis* Leucine-Rich Repeat Receptor-Like Kinases BAK1/SERK3 and BKK1/SERK4 Are Required for Innate Immunity to Hemibiotrophic and Biotrophic Pathogens. *Plant Cell* 23: 2440-2455.
- Ryan, C.A., Huffaker, A., and Yamaguchi, Y. (2007). New insights into innate immunity in *Arabidopsis*. *Cell Microbiol.* 9: 1902-1908.
- Schulze, B., Mentzel, T., Jehle, A.K., Mueller, K., Beeler, S., Boller, T., Felix, G., and Chinchilla, D. (2010). Rapid Heteromerization and Phosphorylation of Ligand-activated Plant Transmembrane Receptors and Their Associated Kinase BAK1. *J. Biol. Chem.* 285: 9444-9451.
- Schwessinger, B., Roux, M., Kadota, Y., Ntoukakis, V., Sklenar, J., Jones, A., and Zipfel, C. (2011). Phosphorylation-Dependent Differential Regulation of Plant Growth, Cell Death, and Innate Immunity by the Regulatory Receptor-Like Kinase BAK1. *Plos Genet.* 7: e1002046.
- Segonzac, C., Feike, D., Gimenez-Ibanez, S., Hann, D.R., Zipfel, C., and Rathjen, J.P. (2011). Hierarchy and roles of pathogen-associated molecular pattern-induced responses in *Nicotiana benthamiana*. *Plant Physiol.* 156: 687-699.
- Shan, L., He, P., Li, J., Heese, A., Peck, S.C., Nurnberger, T., Martin, G.B., and Sheen, J. (2008). Bacterial effectors target the common signaling partner BAK1 to disrupt multiple MAMP receptor-signaling complexes and impede plant immunity. *Cell Host Microbe* 4: 17-27.
- Shi, H., Shen, Q., Qi, Y., Yan, H., Nie, H., Chen, Y., Zhao, T., Katagiri, F., and Tang, D. (2013). BR-SIGNALING KINASE1 Physically Associates with FLAGELLIN SENSING2 and Regulates Plant Innate Immunity in *Arabidopsis*. *Plant Cell* 25: 1143-1157.

- Shimada, T.L., Shimada, T. and Hara-Nishimura, I. (2010). A rapid and non-destructive screenable marker, FAST, for identifying transformed seeds of *Arabidopsis thaliana*. *Plant J.* 61: 519-528.
- Singh, P., and Zimmerli, L. (2013). Lectin receptor kinases in plant innate immunity. *Front. Plant Sci.* 7: 124.
- Singh, P., Kuo, Y.C., Mishra, S., Tsai, C.H., Chien, C.C., Chen, C.W., Desclos-Theveniau, M., Chu, P.W., Schulze, B., Chinchilla, D., Boller, T., and Zimmerli, L. (2012a). The Lectin Receptor Kinase-VI.2 Is Required for Priming and Positively Regulates *Arabidopsis* Pattern-Triggered Immunity. *Plant Cell* 24: 1256-1270.
- Singh, P., Chien, C.C., Mishra, S., Tsai, C.H., and Zimmerli, L. (2012b). The *Arabidopsis* LECTIN RECEPTOR KINASE-VI.2 is a functional protein kinase and is dispensable for basal resistance to *Botrytis cinerea*. *Plant Signal. Behav.* 8: e22611.
- Slaughter, A., Daniel, X., Flors, V., Luna, E., Hohn, B., and Mauch-Mani, B. (2012). Descendants of Primed *Arabidopsis* Plants Exhibit Resistance to Biotic Stress. *Plant Physiol.* 158: 835-843.
- Sreekanta, S., Bethke, G., Hatsugai, N., Tsuda, K., Thao, A., Wang, L., Katagiri, F., and Glazebrook, J. (2015). The receptor-like cytoplasmic kinase PCRK1 contributes to pattern-triggered immunity against *Pseudomonas syringae* in *Arabidopsis thaliana*. *New Phytol.* 207: 78–90.
- Sun, W., Cao, Y., Jansen Labby, K., Bittel, P., Boller, T., and Bent, A.F. (2012). Probing the *Arabidopsis* flagellin receptor: FLS2-FLS2 association and the contributions of specific domains to signaling function. *Plant Cell.* 24: 1096-1113.
- Sun, Y., Li, L., Macho, A.P., Han, Z., Hu, Z., Zipfel, C., Zhou, J.M., and Chai, J. (2013). Structural basis for flg22-induced activation of the *Arabidopsis* FLS2-BAK1 immune complex. *Science* 342: 624-628.
- Tena, G., Boudsocq, M., and Sheen, J. (2011). Protein kinase signaling networks in plant innate immunity. *Curr. Opin. Plant Biol.* 14: 519-529.

- Ton, J., and Mauch-Mani, B. (2004). Beta-amino-butyric acid-induced resistance against necrotrophic pathogens is based on ABA-dependent priming for callose. *Plant J.* 38: 119-130.
- Tsai, C.H., Singh, P., Chen, C.W., Thomas, J., Weber, J., Mauch-Mani, B., and Zimmerli, L. (2011). Priming for enhanced defence responses by specific inhibition of the *Arabidopsis* response to coronatine. *Plant J.* 65: 469-479.
- Tsuda, K., and Katagiri, F. (2010). Comparing signaling mechanisms engaged in pattern-triggered and effector-triggered immunity. *Curr. Opin. Plant Biol.* 13: 459-465.
- Van Wees, S.C.M., Van der Ent, S., and Pieterse, C.M.J. (2008). Plant immune responses triggered by beneficial microbes. *Curr. Opin. Plant Biol.* 11: 443-448.
- Walter, M., Chaban, C., Schütze, K., Batistic, O., Weckermann, K., Näke, C., Blazevic, D., Grefen, C., Schumacher, K., Oecking, C., Harter, K., and Kudla, J. (2004). Visualization of protein interactions in living plant cells using bimolecular fluorescence complementation. *Plant J.* 40: 428-438.
- Wan, J., Tanaka, K., Zhang, X.C., Son, G.H., Brechenmacher, L., Nguyen, T.H., and Stacey, G. (2012). LYK4, a lysin motif receptor-like kinase, is important for chitin signaling and plant innate immunity in *Arabidopsis*. *Plant Physiol.* 160: 396-406.
- Wan, J., Zhang, X.C., Neece, D., Ramonell, K.M., Clough, S., Kim, S.Y., Stacey, M.G., and Stacey, G. (2008). A LysM receptor-like kinase plays a critical role in chitin signaling and fungal resistance in *Arabidopsis*. *Plant Cell* 20: 471-481.
- Willmann, R., Lajunen, H.M., Erbs, G., Newman, M.A., Kolb, D., Tsuda, K., Katagiri, F., Fliegmann, J., Bono, J.J., Cullimore, J.V., Jehle, A.K., Götz, F., Kulik, A., Molinaro, A., Lipka, V., Gust, A.A., and Nürnberger, T. (2011). *Arabidopsis* lysin-motif proteins LYM1 LYM3 CERK1 mediate bacterial peptidoglycan sensing and immunity to bacterial infection. *Proc. Natl. Acad. Sci. U S A.* 108: 19824-19829.
- Windram, O., Madhou, P., McHattie, S., Hill, C., Hickman, R., Cooke, E., Jenkins, D.J., Penfold, C.A., Baxter, L., Breeze, E., Kiddle, S.J., Rhodes, J., Atwell, S., Kliebenstein, D.J., Kim, Y.S., Stegle, O., Borgwardt, K., Zhang, C., Tabrett, A., Legaie, R., Moore, J., Finkenstadt, B., Wild, D.L., Mead, A., Rand, D., Beynon, J., Ott, S., Buchanan-

- Wollaston, V., and Denby, K.J. (2012). *Arabidopsis* defense against *Botrytis cinerea*: chronology and regulation deciphered by high-resolution temporal transcriptomic analysis. *Plant Cell* 24: 3530-3557.
- Wu, C.C., Singh, P., Chen, M.C., and Zimmerli, L. (2010). L-Glutamine inhibits beta-aminobutyric acid-induced stress resistance and priming in *Arabidopsis*. *J. Exp. Bot.* 61: 995-1002.
- Xiao, F., He, P., Abramovitch, R.B., Dawson, J.E., Nicholson, L.K., Sheen, J., and Martin, G.B. (2007). The N-terminal region of *Pseudomonas* type III effector AvrPtoB elicits Pto-dependent immunity and has two distinct virulence determinants. *Plant J.* 52: 595-614.
- Xu, J., Xie, J., Yan, C., Zou, X., Ren, D., and Zhang, S. (2014a). A chemical genetic approach demonstrates that MPK3/MPK6 activation and NADPH oxidase-mediated oxidative burst are two independent signaling events in plant immunity. *Plant J.* 77: 222-2234.
- Xu, P., Xu, S.L., Li, Z.J., Tang, W., Burlingame, A.L., and Wang, Z.Y. (2014b). A brassinosteroid-signaling kinase interacts with multiple receptor-like kinases in *Arabidopsis*. *Mol Plant.* 7: 441-444.
- Yamaguchi, Y., Huffaker, A., Bryan, A.C., Tax, F.E., and Ryan, C.A. (2010). PEPR2 Is a Second Receptor for the Pep1 and Pep2 Peptides and Contributes to Defense Responses in *Arabidopsis*. *Plant Cell* 22: 508-522.
- Yamaguchi, Y., Pearce, G., and Ryan, C.A. (2006). The cell surface leucine-rich repeat receptor for AtPep1, an endogenous peptide elicitor in *Arabidopsis*, is functional in transgenic tobacco cells. *Proc. Natl. Acad. Sci. USA* 103: 10104-10109.
- Yeh, Y.H., Chang, Y.H., Huang, P.Y., Huang, J.B., and Zimmerli, L. (2015). Enhanced *Arabidopsis* pattern-triggered immunity by overexpression of cysteine-rich receptor-like kinases. *Front Plant Sci.* 6: 322.
- Yoo, S.D., Cho, Y.H., and Sheen, J. (2007). *Arabidopsis* mesophyll protoplasts: a versatile cell system for transient gene expression analysis. *Nat. protoc.* 2: 1565-1572.
- Zeng, W.Q., Melotto, M., and He, S.Y. (2010). Plant stomata: a checkpoint of host immunity and pathogen virulence. *Curr. Opin. Biotech.* 21: 599-603.

- Zhang, J., and Zhou, J.M. (2010). Plant Immunity Triggered by Microbial Molecular Signatures. *Mol. Plant* 3: 783-793.
- Zhang, J., Li, W., Xiang, T.T., Liu, Z.X., Laluk, K., Ding, X.J., Zou, Y., Gao, M.H., Zhang, X.J., Chen, S., Mengiste, T., Zhang, Y.L., and Zhou, J.M. (2010). Receptor-like Cytoplasmic Kinases Integrate Signaling from Multiple Plant Immune Receptors and Are Targeted by a *Pseudomonas syringae* Effector. *Cell Host Microbe* 7: 290-301.
- Zimmerli, L., Hou, B.H., Tsai, C.H., Jakab, G., Mauch-Mani, B., and Somerville, S. (2008). The xenobiotic beta-aminobutyric acid enhances *Arabidopsis* thermotolerance. *Plant J.* 53: 144-156.
- Zimmerli, L., Jakab, C., Metraux, J.P., and Mauch-Mani, B. (2000). Potentiation of pathogen-specific defense mechanisms in *Arabidopsis* by beta-aminobutyric acid. *Proc. Natl. Acad. Sci. USA* 97: 12920-12925.
- Zimmerli, L., Metraux, J.P., and Mauch-Mani, B. (2001). beta-aminobutyric acid-induced protection of *Arabidopsis* against the necrotrophic fungus *Botrytis cinerea*. *Plant Physiol.* 126: 517-523.
- Zipfel, C. (2014). Plant pattern-recognition receptors. *Trends Immunol.* 35: 345-351.
- Zipfel, C., and Robatzek, S. (2010). Pathogen-associated molecular pattern-triggered immunity: veni, vidi...? *Plant Physiol.* 154: 551-554.
- Zipfel, C., Kunze, G., Chinchilla, D., Caniard, A., Jones, J.D.G., Boller, T., and Felix, G. (2006). Perception of the bacterial PAMP EF-Tu by the receptor EFR restricts *Agrobacterium*-mediated transformation. *Cell* 125: 749-760.