



Spatial memory deficits in a mouse model of late-onset Alzheimer's disease are caused by zinc supplementation and correlate with amyloid-beta levels

Jane M. Flinn^{1*}, P. Lorenzo Bozzelli¹, Paul A. Adlard² and Angela M. Railey¹

¹ Department of Psychology, George Mason University, Fairfax, VA, USA

² Synaptic Neurobiology Laboratory, The Florey Institute of Neuroscience and Mental Health, The University of Melbourne, Parkville, VIC, Australia

Edited by:

Rodrigo Orlando Kuljiš, Zdrav Mozak
Limitada, Chile

Reviewed by:

Jacques Micheau, University of
Bordeaux 1, France
Luc Buee, INSERM, France
José M. Delgado-García, University
Pablo de Olavide, Spain

*Correspondence:

Jane M. Flinn, Department of
Psychology, George Mason
University, 4400 University Drive,
MSN 3F5, Fairfax, VA 22030, USA
e-mail: jflinn@gmu.edu

Much of the research in Alzheimer's disease (AD) that uses mouse models focuses on the early-onset form of the disease, which accounts for less than 5% of cases. In contrast, this study used a late-onset AD model to examine the interaction between increased dietary zinc (Zn) and the apolipoprotein E (ApoE) gene. ApoE $\epsilon 4$ is overrepresented in late-onset AD and enhances Zn binding to amyloid- β (A β). This study sought to determine if elevated dietary Zn would impair spatial memory in CRND8 mice (CRND8), as well as mice who carry both the mutated human amyloid precursor protein (APP) and ApoE $\epsilon 4$ genes (CRND8/E4). Mice were provided with either lab tap water or water enhanced with 10 ppm Zn (ZnCO₃) for 4 months. At 6 months of age, spatial memory was measured by the Barnes maze. CRND8 mice exhibited significant memory deficits compared to WT mice, as shown by an increased latency to reach the escape box. For the CRND8/E4, but not the CRND8 mice, those given Zn water made significantly more errors than those on lab water. During the probe trial for the WT group, those on Zn water spent significantly less time in the target quadrant than those on lab water. These data suggest that increased dietary Zn can significantly impair spatial memory in CRND8/E4. WT mice given Zn water were also impaired on the 24-h probe trial when compared to lab water WTs. Within the CRND8/E4 group only, levels of soluble A β were significantly correlated with average primary latencies. Within the Zn-treated CRND8/E4 group, there was a significant correlation between insoluble A β and average primary errors. Levels of the zinc transporter 3, ZnT3, were negatively correlated with soluble A β ($p < 0.01$). These findings are particularly relevant because increased intake of dietary supplements, such as Zn, are common in the elderly—a population already at risk for AD. Given the effects observed in the CRND8/E4 mice, ApoE status should be taken into consideration when evaluating the efficacy of therapies targeting metals.

Keywords: apolipoprotein E, hAPP, CRND8, transgenic mice, Barnes maze, metals, ZnT3, copper

INTRODUCTION

Alzheimer's disease (AD) is the primary cause of dementia in the elderly and currently affects more than five million Americans (Alzheimer's Association, 2013). The present research focuses on how the biometals, such as zinc (Zn), copper (Cu), and iron (Fe), interact with amyloid beta (A β)—the key constituent of the plaques that are characteristic of the disease, which mediates plaque formation and consequent oxidative damage.

Zn, Cu, and Fe are found in high concentrations in and around amyloid plaques (Lovell et al., 1998; Maynard et al., 2005), consistent with the notion that A β is a metalloprotein that possesses binding sites for both Zn and Cu (Bush et al., 1993; Hesse et al., 1994). Zn has been shown to be particularly effective in promoting A β aggregation (Bush et al., 1993), and the sequestration of Zn by A β leads to an intracellular deficiency of this metal (Grabrucker et al., 2011). Cu, on the other hand, causes

a shift in the processing of the amyloid precursor protein (APP) towards the non-amyloidogenic pathway (Borchard et al., 1999) but once A β is present, it binds both Cu and Fe. Therefore, Cu and Fe binding is problematic as these metals are redox-active and A β reduces both metals, resulting in the production of hydrogen peroxide (H₂O₂; Huang et al., 1999). H₂O₂ further reacts with the reduced metal to produce hydroxyl radicals that can cause oxidative damage (Bush, 2003). Both aging and plaque-related inflammation may promote a more acidic brain environment, which is conducive to Fe and/or Cu binding (Atwood et al., 1998).

Evidence suggests that Zn is crucial for A β deposition, and the presence of Zn *in vitro* prevents the proteolytic degradation of A β _{1–42} by matrix metalloprotease 2 (Crouch et al., 2011). Zn also inhibits the activity of alpha-secretase and promotes gamma- and beta-secretase activity (Capasso et al., 2005). The zinc transporter, ZnT3, is responsible for loading zinc into the synaptic vesicles

(Palmiter et al., 1996; Cole et al., 1999; Linkous et al., 2008). Eliminating synaptic Zn by genetic ablation of zinc transporter 3 (ZnT3) in an AD mouse model resulted in a 50% reduction in amyloid (Lee et al., 2002). ZnT3 levels, however, have been shown to decrease as a function of age and AD (Adlard et al., 2010). Interestingly, Bjorklund et al. (2012) found that in brains exhibiting AD neuropathology, ZnT3 levels were lowest in those patients who received a diagnosis of AD, while those who did not receive a diagnosis had ZnT3 levels similar to those of healthy controls.

Given the complexity of biometals in AD, the behavioral outcome due to changes in metal levels is difficult to predict, but this has been studied in various mouse models. Long-term dietary Zn enhancement in transgenic (Tg) mouse models of AD (TgCRND8, Tg2576, and APP/presenilin 1 (PS1) mouse models) has been shown to result in cognitive deficits (Linkous et al., 2009; Railey et al., 2011), as well as an increase in both brain Zn levels and the number and size of Zn-positive plaques in the cortex and hippocampus (Wang et al., 2010). Other findings have found that a severe Zn deficiency (<10 ppm Zn) enlarged amyloid plaques (Stoltenberg et al., 2007); thus either an overabundance or deficiency in Zn can potentiate amyloid deposition.

The observations derived from these early-onset mouse models may be of diminished utility for the majority of AD cases that are classified as late-onset (manifest after age 65 years) (Harvey et al., 2003). The emphasis on early-onset genetics dominating AD research is due to the strong relationship between those genes and development of the disease. Genetic linkages to the late-onset form of AD are less specific, but the $\epsilon 4$ allele of the apolipoprotein E (ApoE) gene has been identified as a major risk factor (Saunders et al., 1993; Blacker et al., 1998; Huang and Mucke, 2012). The ApoE gene has three alleles ($\epsilon 2$, $\epsilon 3$, and $\epsilon 4$) resulting in three isoforms of the protein (Zannis et al., 1981), with ApoE $\epsilon 4$ overrepresented in patients with late-onset AD (Weisgraber and Mahley, 1996). While ApoE $\epsilon 4$ is considered a susceptibility factor it does not guarantee the development of disease (Weisgraber and Mahley, 1996). The exact mode of action of ApoE $\epsilon 4$ in AD remains elusive. One proposed mechanism is the interaction between ApoE $\epsilon 4$ and A β , which may increase plaques and/or impede A β clearance (Mahley et al., 2006). Amongst those with AD, ApoE $\epsilon 4$ carriers typically have a larger number of plaques than ApoE $\epsilon 3$ carriers (Schmechel et al., 1993). Similarly, ApoE $\epsilon 4$ mice that overexpress human APP have an elevation in A β plaque deposition, reduced numbers of presynaptic terminals, and an impairment in learning and memory tasks as compared to their $\epsilon 3$ counterparts (Huang, 2006).

The ApoE alleles differ at two residues: $\epsilon 2$ (Cys¹¹², Cys¹⁵⁸), $\epsilon 3$ (Cys¹¹², Arg¹⁵⁸), and $\epsilon 4$ (Arg¹¹², Arg¹⁵⁸) (Rebeck et al., 2002). Cysteine binds to Zn and Cu with high affinity and the Zn binding site on APP is located in a cysteine-rich area (Bush and Tanzi, 2002). The increased number of cysteine residues in $\epsilon 2$ and $\epsilon 3$ correlates with their ability to bind zinc in the synaptic cleft and may contribute to the protective effect of these alleles by preventing zinc binding to A β (Lee et al., 2010). ApoE $\epsilon 4$ is also implicated in the formation of neurofibrillary tangles (NFTs). It may increase the phosphorylation of the tau protein (Harris et al., 2004) and Zn may be a key modulator of this process (Craddock et al., 2012).

Through their relatively high-binding affinity for zinc, A β plaques will cause a reduction in intracellular levels of zinc that will in turn lead to a destabilization of microtubules and subsequent liberation of tau (Craddock et al., 2012). These findings suggest that Zn and ApoE $\epsilon 4$ may act together to affect multiple processes in AD through the promotion of NFT formation and increased A β deposition, which may be due to increased Zn binding in the presence of ApoE $\epsilon 4$.

In order to examine the effect of excess zinc in the more prevalent form of AD we have developed a late-onset model, examining two types of Tg mice with the (1) humanized ApoE $\epsilon 4$ and mutated hAPP cross (CRND8/E4); or (2) those only with mutated hAPP (CRND8), and investigated the effect of Zn supplementation on spatial memory in these mice. Protein levels of soluble and insoluble A β , and ZnT3 were measured to elucidate molecular mechanisms that may be associated with excess dietary zinc in AD.

MATERIALS AND METHODS

SUBJECTS

To address the primary research questions of the study, the creation of a mouse strain modeling late-onset AD was necessary. The CRND8/E4 experimental animals were obtained by breeding CRND8 males (University of Toronto) with female mice who were homozygous for the human ApoE $\epsilon 4$ knock-in and the knock-out of murine ApoE (Jackson Labs). Resultant CRND8/E4 were therefore heterozygous for human ApoE $\epsilon 4$ and murine ApoE in addition to being hemizygous for the hAPP mutation. Tg CRND8 mice carry a mutant form of APP 695 containing both the Swedish (KM670/671NL) and Indiana (V717F) mutations on a hybrid C3H/B6 genetic background and exhibit extensive amyloid deposition by 3 months of age. There is a potent increase in A β ₄₂ at around 10 weeks of age in this model making the A β ₄₂ to A β ₄₀ ratio 5:1 (Chishti et al., 2001). The Tg1HolApoE^{tm1unc}/J strain expresses human ApoE $\epsilon 4$ under the direction of human glial fibrillary acidic protein (GFAP). Experimental WT mice were those littermates who did not inherit the hAPP transgene from the cross between CRND8 males and WT females (C3H/B6). CRND8 mice were those offspring who inherited the hAPP transgene from the CRND8 males that were bred with WT females.

Mice were bred in three groups in order to facilitate behavioral testing. The offspring were group housed with same-sex littermates with 2–4 animals per cage. Each cage contained an igloo, nylabone, and running wheel (Bio-serv). The mouse colony was maintained with a 12-h light/dark cycle. Food (Harlan diet 7012) and water were provided *ad libitum*.

Behavioral group numbers are shown in **Table 1**. This study used both male and female mice, and analyses were conducted to determine possible sex differences. Zn supplementation began at 6 weeks of age and continued throughout behavioral testing. Behavioral testing began at 5 months of age and mice were sacrificed within 10 days following the conclusion of all behavioral testing (approximately 6 months of age). Mice were euthanized by CO₂ asphyxiation and brains were removed and stored in a –80° freezer for later analysis. There were no differences in body weight on the day of euthanasia when evaluated by water condition. All experiments and procedures performed on these mice were

Table 1 | Group distributions and male to female ratio for Barnes maze (BM) and Western blot (WB) analyses.

Group	BM N =	BM m:f	WB N =	WB m:f
WT + Lab Water	12	7:5	0	n/a
WT + Zn	12	6:6	0	n/a
CRND8/E4 + Lab Water	11	6:5	10	5:5
CRND8/E4 + Zn	11	5:6	11	5:6
CRND8 + Lab Water	12	8:4	10	6:4
CRND8 + Zn	11	9:2	9	7:2

approved by the George Mason University Institutional Animal Care and Use Committee.

WATER PREPARATION

The 10 ppm ZnCO₃ was supplemented to laboratory tap water (herein referred to as Zn water) and was prepared using a starting solution of 10,000 mg/L of zinc dissolved in 5% HNO₃. The final solution was buffered with Na₂CO₃ to bring it to a pH of 7.0 (Linkous et al., 2009). All waters were made and stored in separate polycarbonate carboys and dispensed to animals in 500 mL glass bottles. Water was analyzed regularly using inductively coupled plasma-optical emission spectroscopy and ion chromatography at the United States Geological Survey (USGS, Reston, VA) to confirm metal content. Laboratory tap water without any supplement, herein referred to as lab water, was used for the control group. Water consumption was measured every week and there were no significant differences when evaluated by water condition.

BARNES MAZE

The Barnes maze (BM) is designed to assess spatial memory. The maze (Med Associates) consisted of an elevated platform 122 cm in diameter (91 cm above the ground) with 40 equidistant holes located around the edges. The maze's surface color was white. An escape box was located under one of the holes and four distinctive extra-maze reference cues were placed around the maze, with one on each side (large black letters on a white background). Testing took place in a brightly illuminated room with a 300-watt light situated directly over the maze in order to make the maze surface undesirable, given the natural tendency of mice to prefer dark enclosed areas over bright/open areas. All animals started each trial from a dark enclosed start box positioned in the center of the maze. The orientation of the start chamber was varied for each trial. The maze was cleaned with 70% ethanol solution between all trials to eliminate scent trails.

Procedure

BM testing took place over seven consecutive days with the 24-h probe trial occurring on the seventh day.

Habituation: On the first day of testing all mice were given two habituation trials. The mice were placed in a start chamber in the center of the maze for 10 s after which time the start box was lifted and they were gently guided to the escape box and allowed to enter on their own. Once inside, mice stayed in the escape box for 30 s before being returned to their home cage for a 2 min inter-trial interval.

Acquisition: Days 2–6 were acquisition days. The mice received three trials per day with a 15 min inter-trial interval. The trials on the acquisition days ended when the animal entered the escape box or when the maximum trial length of 3 min was reached. Mice were allowed 30 s in the escape box after locating it. If the animal did not find the escape box in the allotted 3 min trial period they were gently guided to the escape box and allowed 30 s inside. Primary latency (amount of time elapsed prior to first reaching target hole) and primary errors (number of errors made prior to reaching target hole) were measured.

Probe Trial: On day 7 the animals received one probe trial 24 h after the last training trial. The escape box was removed from the target hole that had been previously learned and the mice were given 3 min to explore the maze. Percent of time in the correct quadrant was measured.

TISSUE PREPARATION

Frozen brain samples were minced into small pieces and homogenized in 1 mL of a mixture of PBS and protease/phosphatase inhibitor cocktail (Sigma-Aldrich). The homogenate was then centrifuged at 4°C at 24,000 RPM for 1 h. Soluble and insoluble fractions were separated. The insoluble fractions were additionally reconstituted in 1 mL of a mixture of PBS and protease/phosphatase inhibitors. Both soluble and insoluble fractions were then lyophilized for approximately 12 h until fully dehydrated.

WESTERN BLOT ANALYSES

For Western blot (WB) analysis, lyophilized insoluble fractions were reconstituted in 250 µL of PBS and protease/phosphatase inhibitor cocktail. Soluble fractions were reconstituted in 50 µL of PBS and cocktail. A BCA protein assay was done to determine protein concentration for each sample. Fifteen microliters of samples were heated at 90°C for 10 min and then spun down and loaded onto Criterion XT precast 4–12% gels. After the gels were run and transferred to a PVDF membrane they were blocked in 5% tris buffered saline with Tween20 (TBST) skim milk for 1 h. The membranes were then briefly rinsed with TBST and incubated overnight at 4°C with the following primary antibodies: WO2 (1:2000; in-house antibody prepared at The Florey Institute of Neuroscience and Mental Health); ZnT3 (1:10,000, in-house antibody prepared at The Florey Institute of Neuroscience and Mental Health); and GAPDH (1:10,000; Millipore Cat: MAB 374). After three washes of TBST for 5–10 min, secondary antibody incubation was performed (polyclonal rabbit anti-mouse IgG conjugated to horseradish peroxidase (HRP); 1:10,000; Dako P0260) at room temperature for 1 h. Three additional washes of TBST were done before enhanced chemiluminescent substrate was added for imaging using Multigauge software (Fujifilm). After imaging, the membranes were briefly rinsed with TBST and then the membranes were stripped using Re-blot Plus Strong solution (Millipore). The bands were scanned and the intensities of the bands were measured using the Fuji Reader LAS-4000. Protein levels for WO2 and ZnT3 were normalized to GAPDH (Figure 5).

STATISTICAL ANALYSES

Statistical differences among different genotypes and water conditions against measured behavioral parameters were determined by repeated-measures analysis of variance (RMANOVAs). When necessary, Greenhouse-Geisser estimates of sphericity were used to correct for degrees of freedom. Bonferroni *post-hoc* tests were used to evaluate group differences. Correlations between protein levels and measured behavioral parameters were calculated using the Pearson correlation coefficient (r). Data are expressed as mean \pm SEM. A significance level of $p < 0.05$ was used for all analyses. Trends were reported at a level of $p < 0.10$.

RESULTS

BARNES MAZE BEHAVIOR

Three genotypes (WT, CRND8/E4, CRND8) \times two water conditions (lab water and zinc water) mixed design RMANOVAs were performed for measures of primary latency, primary errors, and percent of time in the target quadrant for the 24-h probe trial. Between group variables were genotype and water type. The within subjects variable was testing day (five levels). Sex was entered as a covariate for all statistical analyses.

On days 1–5, there was a significant within-subjects effect of day for primary latency ($F_{(3,03,172.43)} = 44.29$, $p < 0.001$) indicating overall improvement across testing days for all groups. There was a significant effect on primary latency between genotypes ($F_{(2,57)} = 4.124$, $p < 0.05$). A Bonferroni *post-hoc* test revealed that primary latencies across days 1–5 were significantly shorter for WT mice as compared to CRND8 mice ($p < 0.05$), but not compared to CRND8/E4. The CRND8/E4 were also not significantly different from CRND8 mice (Figure 1A). On average CRND8/E4 animals had latencies that were shorter than CRND8 animals, but longer than WT animals. For primary latency, there were no main effects of water condition.

On days 1–5, there was a significant within-subjects effect of day for primary errors ($F_{(3,39,200.0)} = 23.32$, $p < 0.001$). For primary errors there were no significant main effects of water or genotype across all groups on days 1–5 or during probe trials (Figure 1B). Further analyses within genotypes revealed no significant differences between Zn water and lab water in WT and CRND8 mice (Figures 2A,B). However, within the CRND8/E4 genotype there was a significant main effect of water. CRND8/E4 mice on Zn water made significantly more primary errors than those CRND8/E4 on lab water on days 2–3 ($F_{(1,17)} = 5.64$, $p < 0.05$; Figure 2C).

There were no significant main effects of genotype on the 24-h probe trial. There was, however, a significant main effect of water condition within the WT group. Zn water WT mice spent significantly less time than lab water WT mice in the target (correct) quadrant (23% vs. 41% respectively) ($F_{(1,21)} = 3.28$, $p < 0.05$; Figure 3). Both CRND8/E4 and CRND8 groups raised on either Zn or lab water scored close to chance during the probe trial.

In summary, there were significant main effects of water condition within the CRND8/E4 and WT genotypes, but not in the CRND8 group. As predicted, Zn water CRND8/E4 mice had higher primary errors in the BM compared to those on lab water. Significant genotype differences did exist between CRND8






FIGURE 1 | Average primary latency and primary errors across genotypes in the Barnes maze, regardless of water condition.

(A) Average primary latencies were significantly shorter for WT mice as compared to CRND8 mice ($p < 0.05$), but not CRND8/E4s. Overall, there was a significant decrease across days in primary latency for all genotypes ($p < 0.001$). (B) Average primary errors in the Barnes maze. Primary errors were not significantly different between any of the genotypes. All groups did, however, improve significantly over time ($p < 0.001$). Points are mean \pm SEM. * denotes $p < 0.05$.

mice compared to WT mice on measures of primary latency, as was expected. Mice carrying both the ApoE $\epsilon 4$ and mutated hAPP genes were not more impaired than the CRND8 transgene. However, on the primary error measure, CRND8/E4 mice on Zn water had the highest number of primary errors. WT mice on Zn water were spent significantly less time in the target quadrant when compared to those WTs on lab water.

PROTEIN ANALYSIS

Relative levels of soluble A β , insoluble A β , and ZnT3 were obtained from WB analysis. This analysis was conducted only on Tg mice (CRND8 and CRND8/E4). The control for both Tg groups was a CRND8/E4 in the lab water group which may explain why all significant correlations with behavior were seen only in the CRND8/E4 group; all protein correlations presented here are within that genotype only. There were no significant main effects for water and genotype. There were no significant main effects on levels of ZnT3. There were also no



significant main effects for soluble A β or insoluble A β . Regardless of water condition, however, levels of soluble A β were significantly correlated with the average primary latencies for day 3 ($r = 0.657$, $p < 0.01$) and day 4 ($r = 0.463$, $p < 0.05$; **Figure 4A**). Those mice on Zn water had levels of insoluble A β that were significantly correlated with average primary errors for day 3 ($r = 0.754$, $p < 0.01$) and day 5 ($r = 0.658$, $p < 0.05$; **Figure 4B**). Those on control lab water had soluble A β levels which significantly correlated with the average primary latency on day 3 ($r = 0.712$, $p < 0.05$; **Figure 4C**). Mice on Zn water had levels of soluble A β that were negatively correlated with levels of ZnT3 ($r = -0.791$, $p < 0.01$; **Figure 4D**).

DISCUSSION


These data show that the performance of the mice modeling late-onset AD depended significantly upon the presence or absence

of Zn in the drinking water. Overall, CRND8/E4 mice did not perform worse than the CRND8 mice, as had initially been predicted. However, CRND8/E4 mice raised on Zn water made the most errors in the BM of all six groups, and also had significantly increased errors compared to those raised on lab water. On the primary latency measure, the CRND8 mice performed significantly worse than the WT mice. The CRND8/E4 group's performance was intermediate between the other groups, and did not differ significantly from either. In contrast to the effect on primary errors, there was no significant effect of Zn water on primary latency. The observation that primary errors, and not primary latencies, detected a Zn effect in the late-onset mouse model is consistent with O'Leary and Brown (2013) who have suggested that errors are a more sensitive measure of memory than latency in the Barnes maze.

The effect on errors may be due to enhanced Zn binding to A β when in the presence of ApoE $\epsilon 4$. ApoE $\epsilon 4$ has the weakest affinity for binding Zn, due to the amino acid substitution of arginine which is not a ligand for Zn (Lee et al., 2010). In contrast, the ApoE $\epsilon 2$ and ApoE $\epsilon 3$ isoforms with cysteine residues strongly bind Zn, and may prevent it from binding with A β , which could in turn prevent oligomerization and deposition (Lee et al., 2010). While there are numerous *in vitro* studies investigating the association between Zn and ApoE $\epsilon 4$, this is a novel behavioral study examining the relationship between the two in a mouse model of late-onset AD.

Zn water did not result in significant latency deficits in the CRND8 and WT groups. For the WT group, Zn significantly reduced the percent time spent in the target quadrant for the 24-h probe trial; thus the WT mice on Zn water were impaired on the most difficult memory task. Neither the CRND8/E4 or CRND8 groups performed above chance on either water condition (**Figure 4**); thus, additional impairments caused by Zn were difficult to assess. Given the difficulty of the 24-h probe task for the Tg mice, running probe tests as the last run on alternate days, as is done in the Morris water maze (MWM), could be a useful strategy.

We have previously demonstrated significant latency impairments in the MWM in both Zn water rats and Tg (CRND8 and



Tg2576) mice (Linkous et al., 2009; Railey et al., 2010, 2011). The MWM is a more stressful test than the BM (Hölscher, 1999) and this may account for the different effects of zinc on latency seen in this experiment. Overall, data from our lab indicate that Zn water has adverse behavioral effects on WT and CRND8 mice but that additional mechanisms are at work for CRND8/E4 mice. Although Zn supplementation exacerbated behavioral deficits in the Tg2576 and CRND8 mice, this led to a paradoxical decrease in amyloid deposits in the Tg2576 mice (Linkous et al., 2009), which was also observed by Harris et al. (2014); however they found no significant behavioral differences.

The deficits in spatial memory that were observed in the Barnes maze, in connection with dietary Zn supplementation, may be attributable to the effect of Zn on other metals in the brain, specifically Cu. Excess dietary Zn prevents adequate absorption of Cu through the intestinal wall (Milne et al., 2001). Therefore, Zn supplementation may potentiate a Cu deficiency that is inherent in Tg mice overexpressing mutant hAPP (Phinney et al., 2003). Brewer (2014), however, found a non-significant delay in cognitive impairment in AD patients who received Zn supplements, and suggests that AD is the result of a Zn deficiency. He also noted that these subjects had relatively low Zn serum levels and suggested that this would lead to an

overload of copper. Brewer (2014) considers zinc deficiency a risk factor for AD, due to its interaction with copper (Brewer, 2014).

In contrast, Klevay (2008) has suggested that AD is due to a Cu deficiency instead of Cu toxicity and spatial memory deficits in Zn-treated Tg mice can be remediated by the concurrent administration of small amounts of Cu (Railey et al., 2011). APP-overexpressing mice have lower levels of Cu in the brain (Bayer et al., 2003) which would support increasing dietary Cu to treat AD. This is consistent with another study where APP23 mice were given Cu supplementation which resulted in a reduction of A β and an increase in superoxide dismutase 1 (SOD1) activity (Bayer et al., 2003). These conflicting findings from different labs suggest that metal dyshomeostasis in AD can be due to either a deficiency or overabundance of metal ions, and that alterations to the levels of one metal can impact levels of another. The initial metal status of the patient should be an important factor in considering any form of dietary metal-based therapy.

In contrast to our observations, Zn supplementation in the 3xTg mouse actually improved performance on behavioral tasks, and decreased both A β and tau pathology (Corona et al., 2010), while decreased dietary Zn elevated plaque volume in APP/PS1 mice (Stoltenberg et al., 2007). Carrying mutations in PS1, or




FIGURE 5 | Representative blots are shown for WO2 and ZnT3 which were normalized to GAPDH.

in the case of the 3xTg, a mutation in tau, can alter the ability of these proteins to interact with metals. The divergent results may stem from differences in the mouse model used, chemical composition of zinc solutions, and age of onset and duration of Zn supplementation. Our mice were given a standard rodent chow which contains adequate levels of Zn, in addition to the Zn supplementation, which would be in contrast to the Zn deficiency that is reported in the elderly population. While Zn therapy may be beneficial for those with a Zn deficiency, excess Zn, and Zn in the presence of ApoE ϵ 4, may actually be harmful.

In a mouse model carrying the three different human ApoE alleles in the 5xTg mouse, the ApoE transgenes delayed pathology typically observed in the 5xTg strain (Youmans et al., 2012). Although having an expanded window of time to study the progression of AD pathology could be useful, using either the 3xTg or 5xTg strains may be contraindicated for studying the effects of Zn or other metals in late-onset AD. These multiple transgene models have mutations in PS1, which affects cellular turnover of Zn and Cu, since PS1 has Zn binding sites, which will affect SOD1 activity (Sensi et al., 2008; Greenough et al., 2011; Southon et al., 2013). Therefore, studies looking at the relationship between metals and pathobiology of late-onset AD should avoid incorporating mutated PS1 transgenes. Also, enzymes that modulate tau are Zn-dependent.

The current data did not support the predicted outcome of more behavioral impairment in CRND8/E4 mice as compared to CRND8 mice. On most measures the CRND8/E4 mice performed slightly, but not significantly, better than CRND8 mice. This is consistent with other recent behavioral data in younger Tg mice modeling late-onset AD (Moreau et al., 2013). Similar observations of intermediacy were reported in circadian rhythm and nest building behavior, as well as in markers for inflammatory cytokines (Graybeal et al., 2014). Recent studies have shown that any isoform of human ApoE delays AD pathology in early-onset Tg mice (Youmans et al., 2012; Tai et al., 2013).

One possible explanation for the greater impairment associated with the ApoE ϵ 4 allele in old age is due to its interaction with metals and their dysregulation with increased age. Zn and Cu have an interdependent relationship and they exist in a delicate

homeostasis whereby a shift towards either excess or deficiency may influence A β neurotoxicity. Lee et al. (2012) have shown that Zn accumulates around dystrophic neurons in aged Tg2576 mice and suggest that this excess Zn “crowded around amyloid plaques” and could directly cause the death of neurons. Such an effect would likely be greater in the case of ApoE ϵ 4 which does not bind Zn and Cu as efficiently as the other forms of ApoE.

Another interesting finding was that ZnT3, which is the transporter responsible for loading zinc into synaptic vesicles, was negatively correlated with soluble A β levels, which is the more toxic form of A β . Although a general decline in levels of ZnT3 has been associated with cognitive decline (Bjorklund et al., 2012), it has also been demonstrated that the regions of brain that express ZnT3 may change so that the overall level of expression remains unaltered, but the localization of expression may vary and be centered more around areas of pathology in the Tg CNS (Lee et al., 2012). This may explain why no overall differences for ZnT3 expression were detected. It is notable that within the Zn water group we observed a correlation between insoluble A β and primary errors, as this may be due to the ability of Zn to aggregate A β . The relationship between amyloid load and learning and memory remains controversial; for example, Gruart et al. (2008) found no relationship between hippocampal-dependent learning and amyloid load, suggesting that other mechanisms may be involved.

ApoE ablation reduces the amount of ZnT3 and results in significantly lower levels of synaptic Zn compared to WTs (Lee et al., 2010). Zn is required for both presynaptic and postsynaptic mossy fiber long-term potentiation, and dysregulation of synaptic Zn alters the excitatory input of the mossy fiber-CA3 synapse in the hippocampus (Pan et al., 2011). Furthermore, ApoE- ϵ 4 targeted replacement mice have altered levels of glutaminase and vesicular glutamate transporter 1 (Dumanis et al., 2013). ApoE- ϵ 4 is correlated with decreased dendritic spines (Dumanis et al., 2009), and treatment of neurons with ApoE- ϵ 4 recombinant protein reduces expression of glutamate receptors (Chen et al., 2010). Future research should examine the effects of A β , Zn supplementation, and ApoE status on glutamatergic pathways.

Since this study demonstrates that CRND8/E4 mice are sensitive to dietary Zn-enhancement, this suggests that therapies targeting metals should consider ApoE allele status in their analyses. Currently, the only FDA-approved drugs for treating AD work by alleviating symptoms instead of targeting the underlying etiology of the disease. A newer class of drugs currently under evaluation, the 8-hydroxyquinolines (8-OHQ), are selective Zn/Cu chaperones (Greenough et al., 2013). Two drugs in this class, Clioquinol (CQ) and PBT2 have a higher affinity for binding Zn and Cu when compared to A β . Instead of chelating these metals from the system, they are able to return metals back inside the cell, thereby restoring the bioavailability of metal ions necessary for cellular processes (Adlard and Bush, 2012). This restoration of intracellular metal concentrations has been shown to increase synaptic proteins, reduce A β accumulation, and improve cognitive functioning in Tg mice (Adlard et al., 2008, 2011; Crouch et al., 2011).

CONCLUSIONS

This experiment showed that Zn water caused significant impairments in a mouse model of late-onset AD, containing the ApoE ϵ 4 allele; consequently, studies on the role of metals in AD need to take into account the ApoE status of either the human subjects or the mouse models. Furthermore, due to the interaction between different metals such as Cu and Zn, care needs to be taken when conducting both *in vitro* and *in vivo* studies, where the outcomes and interpretations of studies may be confounded by this interrelationship. Future studies examining the effects of Zn and/or Cu deficiencies and excesses in the different human ApoE alleles of a late-onset model of AD is warranted.

ACKNOWLEDGMENTS

The authors would like to thank Lara Glenn for assistance with many aspects of the study, Kathy Conko of USGS for analyzing water samples, Caroline L. C. Neely for manuscript edits, the University of Toronto for donating the CRND8 mice, and the reviewers of this manuscript for their helpful comments. This research was funded in part by a Sigma Xi GIAR and an APA dissertation research award to Angela M. Railey. Paul Anthony Adlard is supported by the NHMRC and ARC. In addition, the Florey Institute of Neuroscience and Mental Health acknowledges the strong support from the Victorian Government and in particular the funding from the Operational Infrastructure Support Grant. Publication of this article was funded in part by the George Mason University Libraries Open Access Fund.

REFERENCES

- Adlard, P. A., Bica, L., White, A. R., Nurjono, M., Filiz, G., Crouch, P. J., et al. (2011). Metal ionophore treatment restores dendritic spine density and synaptic protein levels in a mouse model of Alzheimer's disease. *PLoS One* 6:e17669. doi: 10.1371/journal.pone.0017669
- Adlard, P. A., and Bush, A. I. (2012). Metal chaperones: a holistic approach to the treatment of Alzheimer's disease. *Front. Psychiatry* 3:15. doi: 10.3389/fpsy.2012.00015
- Adlard, P. A., Cherny, R. A., Finkelstein, D. I., Gautier, E., Robb, E., Cortes, M., et al. (2008). Rapid restoration of cognition in Alzheimer's transgenic mice with 8-hydroxy quinoline analogs is associated with decreased interstitial Abeta. *Neuron* 59, 43–55. doi: 10.1016/j.neuron.2008.06.018
- Adlard, P. A., Parncutt, J. M., Finkelstein, D. I., and Bush, A. I. (2010). Cognitive loss in zinc transporter-3 knock-out mice: a phenocopy for the synaptic and memory deficits of Alzheimer's disease? *J. Neurosci.* 30, 1631–1636. doi: 10.1523/JNEUROSCI.5255-09.2010
- Alzheimer's Association. (2013). Alzheimer's disease facts and figures. Available online at: http://www.alz.org/alzheimers_disease_facts_and_figures.asp. Accessed on July 18, 2013.
- Atwood, C. S., Moir, R. D., Huang, X., Bacarra, N. M., Scarpa, R. C., Romano, D. M., et al. (1998). Dramatic aggregation of Alzheimer A β by Cu(II) is induced by conditions representing physiological acidosis. *J. Biol. Chem.* 273, 12817–12826. doi: 10.1074/jbc.273.21.12817
- Bayer, T. A., Schafer, S., Simons, A., Kemmling, A., Kamer, T., Tepest, R., et al. (2003). Dietary Cu stabilizes brain superoxide dismutase 1 activity and reduces amyloid Abeta production in APP23 transgenic mice. *Proc. Natl. Acad. Sci. U S A* 100, 14187–14192. doi: 10.1073/pnas.2332818100
- Bjorklund, N. L., Sadagoparamanujam, V. M., Ghirardi, V., Woltjer, R. L., Tagliatalata, G. N.L., and Reese, L. C. (2012). Absence of amyloid β oligomers at the postsynapse and regulated synaptic Zn²⁺ in cognitively intact aged individuals with Alzheimer's disease neuropathology. *Mol. Neurodegener.* 7:23. doi: 10.1186/1750-1326-7-23
- Blacker, D., Wilcox, M. A., Laird, N. M., Rodes, L., Horvath, M., Go, R., et al. (1998). Alpha-2 macroglobulin is genetically associated with Alzheimer's disease. *Nat. Genet.* 19, 357–360. doi: 10.1038/1243
- Borchard, T., Camakaris, J., Cappai, R., Masters, C. L., Beyreuther, K., and Multhaup, G. (1999). Copper inhibits beta-amyloid production and stimulates the non-amyloidogenic pathway of amyloid-precursor-protein secretion. *Biochem. J.* 344, 461–467. doi: 10.1042/0264-6021:3440461
- Brewer, G. J. (2014). Alzheimer's disease causation by copper toxicity and treatment with zinc. *Front. Aging Neurosci.* 6:92. doi: 10.3389/fnagi.2014.00092
- Bush, A. I. (2003). The metallobiology of Alzheimer's disease. *Trends Neurosci.* 26, 207–214. doi: 10.1097/00002093-200307000-00005
- Bush, A. I., Multhaup, G., Moir, R. D., Williamson, T. G., Small, D. H., Rumble, B., et al. (1993). A novel zinc (II) binding site modulates the function of the β A4 amyloid protein precursor of Alzheimer's disease. *J. Biol. Chem.* 268, 16109–16112.
- Bush, A. I., and Tanzi, R. E. (2002). The galvanization of β -amyloid in Alzheimer's disease. *Proc. Natl. Acad. Sci. U S A* 99, 7317–7319. doi: 10.1073/pnas.122249699
- Capasso, M., Jeng, J. M., Malavolta, M., Mocchegiani, E., and Sensi, S. L. (2005). Zinc dyshomeostasis: a key modulator of neuronal injury. *J. Alzheimers Dis.* 8, 93–108; discussion 209–215.
- Chen, Y., Durakoglugil, M. S., Xian, X., and Herz, J. (2010). ApoE4 reduces glutamate receptor function and synaptic plasticity by selectively impairing ApoE receptor recycling. *Proc. Natl. Acad. Sci. U S A* 107, 12011–12016. doi: 10.1073/pnas.0914984107
- Chishti, M. A., Yang, D., Janus, C., Phinney, A. L., Horne, P., Pearson, J., et al. (2001). Early-onset amyloid deposition and cognitive deficits in transgenic mice expressing a double mutant form of amyloid precursor protein 695. *J. Biol. Chem.* 276, 21562–21570. doi: 10.1074/jbc.M100710200
- Cole, T. B., Wenzel, H. J., Kafer, K. E., Schwartzkroin, P. A., and Palmiter, R. D. (1999). Elimination of zinc from synaptic vesicles in the intact mouse brain by disruption of the ZnT3 gene. *Proc. Natl. Acad. Sci. U S A* 96, 1716–1721. doi: 10.1073/pnas.96.4.1716
- Corona, C., Masciopinto, F., Silvestri, E., Viscovo, A. D., Lattanzio, R., Sorda, R. L., et al. (2010). Dietary zinc supplementation of 3xTg-AD mice increases BDNF levels and prevents cognitive deficits as well as mitochondrial dysfunction. *Cell Death Dis.* 1:e91. doi: 10.1038/cddis.2010.73
- Craddock, T. J. A., Tuszyński, J. A., Chopra, D., Casey, N., Goldstein, L. E., Hameroff, S. R., et al. (2012). The zinc dyshomeostasis hypothesis of Alzheimer's disease. *PLoS One* 7:e33552. doi: 10.1371/journal.pone.0033552
- Crouch, P. J., Savva, M. S., Hung, L. W., Donnelly, P. S., Mot, A. I., Parker, S. J., et al. (2011). The Alzheimer's therapeutic PBT2 promotes amyloid- β degradation and GSK3 phosphorylation via a metal chaperone activity. *J. Neurochem.* 119, 220–230. doi: 10.1111/j.1471-4159.2011.07402.x
- Dumanis, S. B., DiBattista, A. M., Miessau, M., Moussa, C. E. H., and Rebeck, G. W. (2013). APOE genotype affects the pre-synaptic compartment of glutamatergic nerve terminals. *J. Neurochem.* 124, 4–14. doi: 10.1111/j.1471-4159.2012.07908.x
- Dumanis, S. B., Tesoriero, J. A., Babus, L. W., Nguyen, M. T., Trotter, J. H., Ladu, M. J., et al. (2009). ApoE4 decreases spine density and dendritic

- complexity in cortical neurons in vivo. *J. Neurosci.* 29, 15317–15322. doi: 10.1523/JNEUROSCI.4026-09.2009
- Grabrucker, A. M., Schmeisser, M. J., Udvardi, P. T., Arons, M., Schoen, M., Woodling, N. S., et al. (2011). Amyloid beta protein-induced zinc sequestration leads to synaptic loss via dysregulation of the ProSAP2/Shank3 scaffold. *Mol. Neurodegener.* 6:65. doi: 10.1186/1750-1326-6-65
- Graybeal, J. J., Bozzelli, P. L., Graybeal, L. L., Groeber, C. M., McKnight, P. E., Cox, D. N., et al. (2014). Human ApoE ϵ 4 alters circadian rhythm activity, IL-1 β , and GFAP in CRND8 mice. *J. Alzheimers Dis.* doi: 10.3233/JAD-132009. [Epub ahead of print].
- Greenough, M. A., Camakaris, J., and Bush, A. I. (2013). Metal dyshomeostasis and oxidative stress in Alzheimer's disease. *Neurochem. Int.* 62, 540–545. doi: 10.1016/j.neuint.2012.08.014
- Greenough, M. A., Volitakis, I., Li, Q., Laughton, K., Evin, G., Ho, M., et al. (2011). Presenilins promote the cellular uptake of copper and zinc and maintain Cu-chaperone of sod1-dependent Cu/Zn superoxide dismutase activity. *J. Biol. Chem.* 286, 9776–9786. doi: 10.1016/j.jalz.2011.05.514
- Gruart, A., López-Ramos, J. C., Muñoz, M. D., and Delgado-García, J. M. (2008). Aged wild-type and APP, PS1 and APP + PS1 mice present similar deficits in associative learning and synaptic plasticity independent of amyloid load. *Neurobiol. Dis.* 30, 439–450. doi: 10.1016/j.nbd.2008.03.001
- Harris, F. M., Brecht, W. J., Xu, Q., Mahley, R. W., and Huang, Y. (2004). Increased tau phosphorylation in Apolipoprotein E4 transgenic mice is associated with activation of extracellular signal-regulated kinase. *J. Biol. Chem.* 279, 44795–44801. doi: 10.1074/jbc.M408127200
- Harris, C. J., Voss, K., Murchison, C., Ralle, M., Frahler, K., Carter, R., et al. (2014). Oral zinc reduces amyloid burden in tg2576 mice. *J. Alzheimers Dis.* 41, 179–192. doi: 10.3233/JAD-131703
- Harvey, R. J., Skelton-Robinson, M., and Rosor, M. N. (2003). The prevalence and causes of dementia in people under the age of 65 years. *J. Neurol. Neurosurg. Psychiatry* 74, 1206–1209. doi: 10.1136/jnnp.74.9.1206
- Hesse, L., Behr, D., Masters, C. L., and Multhaup, G. (1994). The β A4 amyloid precursor protein binding to copper. *FEBS Lett.* 349, 109–116. doi: 10.1016/0014-5793(94)00658-X
- Hölscher, C. (1999). Stress impairs performance in spatial water maze learning tasks. *Behav. Brain Res.* 100, 225–235. doi: 10.1016/S0166-4328(98)00134-X
- Huang, Y. (2006). Apolipoprotein E and Alzheimer's disease. *Neurology* 66(Suppl. 1), S79–S85. doi: 10.1212/01.wnl.0000192102.41141.9e
- Huang, X., Atwood, C. S., Hartshorn, M. A., Multhaup, G., Goldstein, L. E., Scarpa, R. C., et al. (1999). The A β peptide of Alzheimer's disease directly produces hydrogen peroxide through metal ion reduction. *Biochemistry* 38, 7609–7616. doi: 10.1021/bi990438f
- Huang, Y., and Mucke, L. (2012). Alzheimer mechanisms and therapeutic strategies. *Cell* 148, 1204–1222. doi: 10.1016/j.cell.2012.02.040
- Klevay, L. M. (2008). Alzheimer's disease as copper deficiency. *Med. Hypotheses* 70, 802–807. doi: 10.1016/j.mehy.2007.04.051
- Lee, J. Y., Cho, E., Kim, T. Y., Kim, D. K., Palmiter, R. D., Volitakis, I., et al. (2010). Apolipoprotein E ablation decreases synaptic vesicular zinc in the brain. *Biometals* 23, 1085–1095. doi: 10.1007/s10534-010-9354-9
- Lee, J. Y., Cho, E., Seo, J. W., Hwang, J. J., and Koh, J. Y. (2012). Alteration of the cerebral zinc pool in a mouse model of Alzheimer's disease. *J. Neuropathol. Exp. Neurol.* 71, 211–222. doi: 10.1097/NEN.0b013e3182417387
- Lee, J. Y., Cole, T. B., Palmiter, R. D., Suh, S. W., and Koh, J. Y. (2002). Contribution by synaptic zinc to the gender-disparate plaque formation in human Swedish mutant APP transgenic mice. *Proc. Natl. Acad. Sci. U S A* 99, 7705–7710. doi: 10.1073/pnas.092034699
- Linkous, D. H., Adlard, P. A., Wanschura, P. B., Conko, K. M., and Flinn, J. M. (2009). The effects of enhanced zinc on spatial memory and plaque formation in transgenic mice. *J. Alzheimers Dis.* 18, 565–579. doi: 10.3233/JAD-2009-1162
- Linkous, D. H., Flinn, J. M., Koh, J. Y., Lanzirrotti, A., Bertsch, P. M., Jones, B. F., et al. (2008). Evidence that ZnT3 protein controls the total amount of elemental zinc in synaptic vesicles. *J. Histochem. Cytochem.* 56, 3–6. doi: 10.1369/jhc.6a7035.2007
- Lovell, M. A., Robertson, J. D., Teesdale, W. J., Campbell, J. L., and Markesbery, W. R. (1998). Copper, iron and zinc in Alzheimer's disease senile plaques. *J. Neurol. Sci.* 158, 47–52. doi: 10.1016/S0022-510X(98)00092-6
- Mahley, R. W., Weisgraber, K. H., and Huang, Y. (2006). Apolipoprotein E4: a causative factor and therapeutic target in neuropathology, including Alzheimer's disease. *Proc. Natl. Acad. Sci. U S A* 103, 5644–5651. doi: 10.1073/pnas.0600549103
- Maynard, C. J., Bush, A. I., Masters, C. L., Cappai, R., and Li, Q.-X. (2005). Metals and amyloid β in Alzheimer's disease. *Int. J. Exp. Pathol.* 86, 147–159. doi: 10.1111/j.0959-9673.2005.00434.x
- Milne, D. B., Davis, C. D., and Nielsen, F. H. (2001). Low dietary zinc alters indices of copper function and status in postmenopausal women. *Nutrition* 17, 701–708. doi: 10.1016/S0899-9007(01)00560-3
- Moreau, P., Bott, J., Zerbinatti, C., Renger, J. J., Kelche, C., Cassel, J., et al. (2013). ApoE4 confers better spatial memory than apoE3 in young adult hApp-Yac/apoE-Tr mice. *Behav. Brain Res.* 243, 1–5. doi: 10.1016/j.bbr.2012.12.043
- O'Leary, T. P., and Brown, R. E. (2013). Optimization of apparatus design and behavioral measures for the assessment of visuo-spatial learning and memory of mice on the Barnes maze. *Learn. Mem.* 20, 85–96. doi: 10.1101/lm.028076.112
- Palmiter, R. D., Cole, T. B., Quaife, C. J., and Findley, S. D. (1996). ZnT-3, a putative transporter of zinc into synaptic vesicles. *Proc. Natl. Acad. Sci. U S A* 93, 14934–14939. doi: 10.1073/pnas.93.25.14934
- Pan, E., Zhang, X., Huang, Z., Krezel, A., Zhao, M., Tinberg, C. E., et al. (2011). Vesicular zinc promotes presynaptic and inhibits postsynaptic long-term potentiation of mossy fiber-CA3 synapse. *Neuron* 71, 1116–1126. doi: 10.1016/j.neuron.2011.07.019
- Phinney, A. L., Drisaldi, B., Schmidt, S. D., Lugowski, S., Coronado, V., Liang, Y., et al. (2003). In vivo reduction of amyloid-beta by a mutant copper transporter. *Proc. Natl. Acad. Sci. U S A* 100, 14193–14198. doi: 10.1073/pnas.2332851100
- Railey, A. M., Groeber, C. M., and Flinn, J. M. (2011). The effect of metals on spatial memory in a transgenic mouse model of Alzheimer's disease. *J. Alzheimers Dis.* 23, 375–381. doi: 10.3233/JAD-2011-101452
- Railey, A. M., Micheli, T. L., Wanschura, P. B., and Flinn, J. M. (2010). Alterations in fear response and spatial memory in pre- and post-natal zinc supplemented rats: remediation by copper. *Physiol. Behav.* 100, 95–100. doi: 10.1016/j.physbeh.2010.01.040
- Rebeck, G. W., Kindy, M., and LaDu, M. J. (2002). Apolipoprotein E and Alzheimer's disease: the protective effects of ApoE2 and E3. *J. Alzheimers Dis.* 4, 145–154.
- Saunders, A. M., Strittmatter, W. J., Schmechel, D., Hyslop, P. H., Pericak-Vance, M. A., Joo, S. H., et al. (1993). Associate of apolipoprotein E allele epsilon-4 with late-onset familial and sporadic Alzheimer's disease. *Neurology* 43, 1467–1472. doi: 10.1212/wnl.43.8.1467
- Schmechel, D. E., Saunders, A. M., Strittmatter, W. J., Crain, B. J., Hulette, C. M., Joo, S. H., et al. (1993). Increased amyloid β -peptide deposition in cerebral cortex as a consequence of apolipoprotein E genotype in late-onset Alzheimer's disease. *Proc. Natl. Acad. Sci. U S A* 90, 9649–9653. doi: 10.1073/pnas.90.20.9649
- Sensi, S. L., Rapposelli, I. G., Frazzini, V., and Mascetra, N. (2008). Altered oxidant-mediated intraneuronal zinc mobilization in a triple transgenic mouse model of Alzheimer's disease. *Exp. Gerontol.* 43, 488–492. doi: 10.1016/j.exger.2007.10.018
- Southon, A., Greenough, M. A., Gano, G., Bush, A. I., Burke, R., and Camakaris, J. (2013). Presenilin promotes dietary copper uptake. *PLoS One* 8:e62811. doi: 10.1371/journal.pone.0062811
- Stoltenberg, M., Bush, A. I., Bach, G., Smidt, K., Larsen, A., Rungby, J., et al. (2007). Amyloid plaques arise from zinc-enriched cortical layers in APP/PS1 transgenic mice and are paradoxically enlarged with dietary zinc deficiency. *Neuroscience* 5, 357–369. doi: 10.1016/j.neuroscience.2007.09.025
- Tai, L. M., Bilousova, T., Jungbauer, L., Roeske, S. K., Youmans, K. L., Yu, C., et al. (2013). Levels of soluble apolipoprotein E/amyloid- β (A β) complex are reduced and oligomeric A β increased with APOE4 and Alzheimer disease in a transgenic mouse model and human samples. *J. Biol. Chem.* 288, 5914–5926. doi: 10.1074/jbc.M112.442103
- Wang, C. Y., Wang, T., Zheng, W., Zhao, B. L., Danscher, G., Chen, Y. H., et al. (2010). Zinc overload enhances APP cleavage and A β deposition in the Alzheimer mouse brain. *PLoS One* 5:e15349. doi: 10.1371/journal.pone.0015349

- Weisgraber, K. H., and Mahley, R. W. (1996). Human apolipoprotein E: the Alzheimer's disease connection. *FASEB J.* 10, 1485–1494.
- Youmans, K. L., Tai, L. M., Nwabuisi-Heath, E., Jungbauer, L., Kanekiyo, T., Gan, M., et al. (2012). APOE4-specific changes in A β x accumulation in a new transgenic mouse model of Alzheimer disease. *J. Biol. Chem.* 287, 41774–41786. doi: 10.1074/jbc.M112.407957
- Zannis, V. I., Just, P. W., and Breslow, J. L. (1981). Human apolipoprotein E isoprotein subclasses are genetically determined. *Am. J. Hum. Genet.* 33, 11–24.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 09 December 2013; accepted: 01 July 2014; published online: 22 October 2014.

Citation: Flinn JM, Bozzelli PL, Adlard PA and Railey AM (2014) Spatial memory deficits in a mouse model of late-onset Alzheimer's disease are caused by zinc supplementation and correlate with amyloid-beta levels. *Front. Aging Neurosci.* 6:174. doi: 10.3389/fnagi.2014.00174

This article was submitted to the journal *Frontiers in Aging Neuroscience*.

Copyright © 2014 Flinn, Bozzelli, Adlard and Railey. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.



Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Flinn, JM; Bozzelli, PL; Adlard, PA; Railey, AM

Title:

Spatial memory deficits in a mouse model of late-onset Alzheimer's disease are caused by zinc supplementation and correlate with amyloid-beta levels

Date:

2014-10-22

Citation:

Flinn, J. M., Bozzelli, P. L., Adlard, P. A. & Railey, A. M. (2014). Spatial memory deficits in a mouse model of late-onset Alzheimer's disease are caused by zinc supplementation and correlate with amyloid-beta levels. *FRONTIERS IN AGING NEUROSCIENCE*, 6 (JUL), <https://doi.org/10.3389/fnagi.2014.00174>.

Persistent Link:

<http://hdl.handle.net/11343/263645>

File Description:

Published version

License:

CC BY