

Benefits and Safety of Long-Term Fenofibrate Therapy in People With Type 2 Diabetes and Renal Impairment

The FIELD Study

RU-DEE TING, FRACP^{1,2}
 ANTHONY C. KEECH, FRACP, MSC^{1,2}
 PAUL L. DRURY, MA, MB, FRACP³
 MARK W. DONOGHOE, BSC¹
 JOHN HEDLEY, FRACP⁴
 ALICIA J. JENKINS, MD, FRACP^{1,5}
 TIMOTHY M.E. DAVIS, FRACP, DPHIL⁶

SEPPO LEHTO, MD, PHD⁷
 DAVID CELERMAJER, MSC, PHD, FRACP²
 R. JOHN SIMES, SM, FRACP, MD¹
 KUSHWIN RAJAMANI, MB, CHB^{1,2}
 KIM STANTON, FRACP⁸
 ON BEHALF OF THE FIELD STUDY
 INVESTIGATORS*

OBJECTIVE—Diabetic patients with moderate renal impairment (estimated glomerular filtration rate [eGFR] 30–59 mL/min/1.73 m²) are at particular cardiovascular risk. Fenofibrate's safety in these patients is an issue because it may elevate plasma creatinine. Furthermore, guidelines regarding fenofibrate dosing in renal impairment vary internationally. We investigated fenofibrate's effects on cardiovascular and end-stage renal disease (ESRD) events, according to eGFR, in the Fenofibrate Intervention and Event Lowering in Diabetes (FIELD) Study.

RESEARCH DESIGN AND METHODS—Type 2 diabetic patients (aged 50–75 years) with eGFR \geq 30 mL/min/1.73 m² were randomly allocated to a fixed dose of fenofibrate (200 mg daily) ($n = 4,895$) or placebo ($n = 4,900$) for 5 years. Baseline renal function (Modification of Diet in Renal Disease equation) was grouped by eGFR (30–59, 60–89, and \geq 90 mL/min/1.73 m²). The prespecified outcome was total cardiovascular events (composite of cardiovascular death, myocardial infarction, stroke, and coronary/carotid revascularization). Serious adverse events and instances of ESRD (plasma creatinine >400 μ mol/L, dialysis, renal transplant, or renal death) were recorded. Analysis was by intention to treat.

RESULTS—Overall, fenofibrate reduced total cardiovascular events, compared with placebo (hazard ratio 0.89 [95% CI 0.80–0.99]; $P = 0.035$). This benefit was not statistically different across eGFR groupings ($P = 0.2$ for interaction) (eGFR 30–59 mL/min/1.73 m²: 0.68 [0.47–0.97], $P = 0.035$; eGFR \geq 90 mL/min/1.73 m²: 0.85 [0.70–1.02], $P = 0.08$). ESRD rates were similar between treatment arms, without adverse safety signals of fenofibrate use in renal impairment.

CONCLUSIONS—Patients with type 2 diabetes and moderate renal impairment benefit from long-term fenofibrate, without excess drug-related safety concerns compared with those with no or mild renal impairment. Fenofibrate treatment should not be contraindicated in moderate renal impairment, suggesting that current guidelines may be too restrictive.

Diabetes Care 35:218–225, 2012

From the ¹National Health and Medical Research Council Clinical Trials Centre, University of Sydney, Sydney, New South Wales, Australia; the ²Department of Cardiology, Royal Prince Alfred Hospital, Sydney, New South Wales, Australia; the ³Auckland Diabetes Centre, Auckland, New Zealand; the ⁴Department of Internal Medicine, Wairau Hospital, Blenheim, New Zealand; the ⁵Department of Medicine, University of Melbourne, Melbourne, Victoria, Australia; the ⁶School of Medicine and Pharmacology, University of Western Australia, Perth, Western Australia, Australia; the ⁷Department of Medicine, Kuopio University Hospital, Kuopio, Finland; and the ⁸Department of Endocrinology and Diabetes, Royal Perth Hospital, Perth, Western Australia, Australia. Corresponding authors: Ru-Dee Ting, rudee.ting@ctc.usyd.edu.au, and Anthony C. Keech, tony@ctc.usyd.edu.au. Received 2 August 2011 and accepted 2 November 2011.

DOI: 10.2337/dc11-1109. Clinical trial reg. no. ISRCTN64783481, www.isrctn.org.

This article contains Supplementary Data online at <http://care.diabetesjournals.org/lookup/suppl/doi:10.2337/dc11-1109/-DC1>.

R.-D.T. and A.C.K. are joint first authors.

*A complete list of members of the FIELD Study can be found in the Supplementary Data online.

The study sponsors had no role in the data collection or analysis. Both the authors and the study management committee had final responsibility for the decision to submit the article for publication.

© 2012 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

Cardiovascular disease (CVD) is projected to remain the single leading cause of death (1) and is a major cause of morbidity and premature mortality in people with diabetic kidney disease, independent of other risk factors (2,3). The Fenofibrate Intervention and Event Lowering in Diabetes (FIELD) Study was a 5-year trial of fenofibrate versus placebo involving type 2 diabetic patients. Although there was no significant benefit from fenofibrate for the primary end point of coronary heart disease (CHD) events, the secondary end points of CVD events and nonfatal myocardial infarction were significantly reduced, as were hospitalizations for acute coronary syndromes and coronary and carotid revascularization procedures (4). Fenofibrate also significantly reduced the microvascular complications of type 2 diabetes (5–7), including nephropathy.

Set against its potential benefits, there have been safety concerns associated with fenofibrate administration in the setting of renal impairment, with current guidelines recommending dose reduction in patients with even mildly abnormal renal function (8,9). The rise in plasma creatinine in response to fenofibrate is well documented, but this does not reflect true renal injury or an actual fall in glomerular filtration rate (GFR) (4,7,10). Although renal impairment is an independent risk predictor of CVD and end-stage renal disease (ESRD) (2,11,12), it has been unclear whether the fenofibrate-associated rise in plasma creatinine might also confer excess risk, especially in patients with pre-treatment renal impairment. We therefore investigated the impact of fenofibrate on prespecified cardiovascular and renal end points of the FIELD Study in patients with impaired renal function at baseline.

RESEARCH DESIGN AND METHODS

The FIELD Study was a double-blinded placebo-controlled trial conducted in Australia, New Zealand, and Finland. The registered study (clinical trial no. ISRCTN64783481) had ethics committee approval in accordance with the

Declaration of Helsinki and Good Clinical Practice Guidelines. Study design and patient characteristics have been published elsewhere (4). In brief, 9,795 patients with type 2 diabetes, aged 50–75 years, were randomly allocated to receive 200 mg comiconized fenofibrate or placebo daily for an average of 5 years. There was no dose adjustment for any degree of renal impairment. Subjects had mild dyslipidemia, with no immediate indication at study entry for lipid-modifying therapy. Exclusion criteria included plasma creatinine $>130 \mu\text{mol/L}$, liver or symptomatic gallbladder disease, or a CVD event within 3 months before recruitment. All patients with National Kidney Foundation chronic kidney disease stage 5 (estimated GFR [eGFR] $<15 \text{ mL/min/1.73 m}^2$), stage 4 (eGFR $15\text{--}29 \text{ mL/min/1.73 m}^2$), and some with stage 3 (eGFR $30\text{--}59 \text{ mL/min/1.73 m}^2$) of the disease were therefore excluded (13). All patients provided informed consent and then completed a 16-week run-in period comprising 4 weeks of diet, 6 weeks of single-blinded placebo, and 6 weeks of single-blinded fenofibrate before being randomly assigned. Eligibility was confirmed during the run-in period independently of adherence or biochemical changes.

A telephone computer randomization service using dynamic balancing to stratify patients by prognostic variables was used; all investigators, except one statistician, were masked to treatment allocation. Patients were seen every 4–6 months against a background of usual care, and information concerning treatment tolerability and complications was obtained.

eGFR was calculated by the Modification of Diet in Renal Disease (MDRD) Study four-variable equation (14) and was grouped as eGFR ≥ 90 , 60–89, and 30–59 mL/min/1.73 m². Baseline eGFR was taken as the mean of two measurements before the active run-in phase (visits 1 and 3, median 72 days apart).

All major CVD events and all other deaths were adjudicated by an outcomes assessment committee using prespecified definitions and while unaware of treatment allocation. The safety- and data-monitoring committee reviewed the safety data at regular intervals every 6 months and undertook two formal interim analyses of the numbers of deaths from CHD.

The primary end point was CHD events (the first occurrence of either a nonfatal myocardial infarction or death from CHD). Secondary outcomes included total mortality, CVD deaths, total stroke, coronary

revascularization procedures, and total CVD events (CHD events, total stroke, coronary and carotid revascularization, and other CVD death combined). Prespecified tertiary end points included ESRD (plasma creatinine $>400 \mu\text{mol/L}$, dialysis, renal transplant, or death from renal disease).

Statistical analysis

Differences in baseline characteristics were assessed by comparing the 30–59 mL/min/1.73 m² group with the combined $\geq 60 \text{ mL/min/1.73 m}^2$ group using χ^2 tests for categorical variables and *t* tests or nonparametric tests, as appropriate, for continuous variables. Tests for linear trend across the three prespecified eGFR groups also were performed. Cox proportional hazards models were applied within each eGFR group to assess the effect of treatment on primary and secondary outcomes, with an interaction term between treatment and group used to test for the heterogeneity of the treatment effect. χ^2 Tests were used to test for a difference between the number of patients experiencing each type of nonfatal serious adverse event within each eGFR group and the logistic regression used to test for an interaction between treatment and eGFR groups. All analyses were performed on an intention-to-treat basis, with inferences drawn on a two-sided significance level of 0.05, unadjusted for multiple comparisons. SAS software (version 9.2; SAS Institute, Cary, NC) was used for all analyses.

RESULTS

Baseline characteristics

The placebo and fenofibrate groups were well matched for baseline characteristics (4) (Table 1). The 519 participants with eGFR $<60 \text{ mL/min/1.73 m}^2$ were older, had a longer duration of diabetes, and had higher systolic blood pressure than the others. A smaller proportion had a history of smoking. They were more likely to have elevated urinary albumin-to-creatinine ratios (ACRs), elevated total cholesterol and triglyceride levels, and a history of previous CVD events.

Lipid and blood pressure effects of fenofibrate by eGFR

The early (over ~6 weeks) lipid-modifying effects of fenofibrate (increased HDL cholesterol and lower LDL cholesterol and triglycerides) were independently significant within each category of eGFR, with

significantly greater improvement in lipid profile in the lowest eGFR group (Supplementary Table 1). However, the differences between eGFR groups in lipid effects were lost over 5 years (Supplementary Table 2). In addition, over the 6-week fenofibrate run-in period, there was a small but significantly greater reduction in systolic blood pressure in patients with eGFR $30\text{--}59 \text{ mL/min/1.73 m}^2$ (5.1 mmHg) compared with those with eGFR $\geq 90 \text{ mL/min/1.73 m}^2$ (3.4 mmHg) (Supplementary Table 3).

Cardiovascular risk by eGFR and fenofibrate treatment

Among placebo-allocated patients, all CVD outcomes (CHD events, coronary revascularization, CVD death, total CVD events, stroke, and total mortality) were more frequent in those with renal impairment compared with those without (Table 2). The number of CVD events in placebo-treated patients with eGFR $30\text{--}59 \text{ mL/min/1.73 m}^2$ was more than double that of those with eGFR $\geq 90 \text{ mL/min/1.73 m}^2$ (Fig. 1), and CVD and all-cause death were more than six- and fourfold higher, respectively (Table 2, Fig. 1).

Consistent with the highest absolute risks for CVD events being in those with the greatest renal impairment, fenofibrate offered the greatest absolute risk reductions in these individuals. Patients with a baseline eGFR of $30\text{--}59 \text{ mL/min/1.73 m}^2$ who received fenofibrate had independently significant absolute risk reductions compared with those allocated to the placebo group in coronary revascularization, total CVD events, and CVD mortality (Table 2). However, the relative risk reductions with fenofibrate were similar to those for other patients such that there were no statistically significant trends for varying treatment effects across eGFR groups ($P > 0.05$ for interaction), even after adjustment for on-study statin use. For example, fenofibrate reduced CVD risk overall by 11% (hazard ratio [HR] 0.89 [95% CI 0.80–0.99]; $P = 0.035$) irrespective of eGFR grouping (Fig. 2; $P = 0.22$ for interaction), with an overall number needed to treat of 70 (95% CI 36–1,112) to avoid one event. Among those with an eGFR of $30\text{--}59 \text{ mL/min/1.73 m}^2$, there was neither significantly greater benefit nor greater harm, with an estimated total CVD risk reduction of 32% (0.68 [0.47–0.97]; $P = 0.035$) and number needed to treat of 14 (7–834). These observed benefits on total CVD (as also was true for reduced albuminuria) were independent of

Table 1—Clinical and metabolic characteristics of the FIELD Study population at baseline, by eGFR group

| | Baseline eGFR (mL/min/1.73 m ²) | | | P for trend | P contrast* |
|--|---|---------------------|---------------------|-------------|-------------|
| | ≥90 | 60–89 | 30–59 | | |
| n | 4,058 | 5,218 | 519 | | |
| Fenofibrate allocated [n (%)] | 2,039 (50.2) | 2,561 (49.1) | 295 (56.8) | | |
| Male [n (%)]† | 2,693 (66.4) | 3,231 (61.9) | 214 (41.2) | <0.001 | <0.001 |
| Age (years) [mean (SD)]† | 60.20 (6.66) | 63.38 (6.66) | 66.51 (5.92) | <0.001 | <0.001 |
| Diabetes duration (years) | | | | | |
| [geometric mean (95% CI)]‡ | 4.81 (4.68–4.95) | 5.06 (4.93–5.19) | 6.02 (5.55–6.54) | <0.001 | <0.001 |
| Smoking status [n (%)] | | | | <0.001 | <0.001 |
| Never† | 1,462 (36.0) | 2,201 (42.2) | 266 (51.3) | | |
| Previous† | 2,120 (52.2) | 2,607 (50.0) | 217 (41.8) | | |
| Current† | 476 (11.7) | 410 (7.9) | 36 (6.9) | | |
| BMI (kg/m ²) [geometric mean (95% CI)] | | | | | |
| Women‡ | 32.2 (31.9–32.5) | 31.2 (31.0–31.5) | 31.5 (30.9–32.2) | 0.07 | 0.74 |
| Men‡ | 29.7 (29.5–29.9) | 29.1 (29.0–29.3) | 29.2 (28.6–29.8) | 0.11 | 0.56 |
| Systolic blood pressure (mmHg) | | | | | |
| [geometric mean (95% CI)]† | 138.5 (138.1–139.0) | 140.2 (139.7–140.6) | 143.4 (142.0–144.8) | <0.001 | <0.001 |
| HbA _{1c} (%) [geometric mean (95% CI)]‡ | 6.97 (6.93–7.02) | 6.91 (6.88–6.94) | 7.02 (6.91–7.13) | 0.44 | 0.15 |
| Total cholesterol (mmol/L) [mean (SD)]§ | 5.02 (0.70) | 5.04 (0.70) | 5.11 (0.71) | <0.01 | 0.02 |
| Triglycerides (mmol/L) | | | | | |
| [geometric mean (95% CI)]‡ | 1.76 (1.74–1.78) | 1.79 (1.77–1.81) | 1.91 (1.85–1.98) | <0.001 | <0.001 |
| HDL cholesterol (mmol/L) | | | | | |
| [geometric mean (95% CI)]‡ | 1.07 (1.06–1.07) | 1.07 (1.06–1.08) | 1.06 (1.03–1.08) | 0.33 | 0.24 |
| Dyslipidemia [n (%)] | 1,492 (36.8) | 1,971 (37.8) | 247 (47.6) | <0.001 | <0.001 |
| Hemoglobin (g/dL) [mean (SD)]† | 14.62 (1.19) | 14.51 (1.23) | 13.86 (1.37) | <0.001 | <0.001 |
| ACR [n (%)] | | | | <0.001 | <0.001 |
| Normal† | 3,026 (74.9) | 3,927 (75.4) | 307 (59.2) | | |
| Microalbuminuria† | 890 (22.0) | 1,068 (20.5) | 146 (28.2) | | |
| Macroalbuminuria† | 126 (3.1) | 213 (4.1) | 65 (12.5) | | |
| Use of RAAS inhibitors [n (%)]† | 1,518 (37.4) | 1,987 (38.1) | 286 (55.1) | <0.001 | <0.001 |
| Use of aspirin [n (%)]† | 1,012 (24.9) | 1,621 (31.1) | 196 (37.8) | <0.001 | <0.001 |
| Previous CVD [n (%)]† | 680 (16.8) | 1,245 (23.9) | 206 (39.7) | <0.001 | <0.001 |

Microalbuminuria: urine ACR ≥2.5 mg/mmol (male subjects) and ≥3.5 mg/mmol (female subjects). Macroalbuminuria: urine ACR >25 mg/mmol (male subjects) and >35 mg/mmol (female subjects). RAAS, renin-angiotensin-aldosterone system. *P for eGFR <60 vs. ≥60 mL/min/1.73 m². †P < 0.001 for trend. Smoking status and ACR are for overall values. ‡Analyses performed on log-transformed data. §P < 0.05 for trend.

the background use or nonuse of ACE inhibitors, the extent of glycemic control (by HbA_{1c}), and systolic blood pressure levels (all P interactions were not significant).

The CVD mortality risk in those with an eGFR of 30–59 mL/min/1.73 m² allocated to fenofibrate was significantly reduced by 49% (95% CI 7–72; P = 0.03) compared with placebo, with greater benefit among those with lower eGFR than others (P = 0.03 for interaction) (Table 2). However, this pattern was no longer significant after adjustment for other baseline variables (sex, age, diabetes duration, smoking status, systolic blood pressure, total cholesterol, HDL cholesterol, triglycerides, HbA_{1c}, hemoglobin, urine ACR, previous retinopathy, previous CVD, renin-angiotensin-aldosterone blocker use, β-blocker use, and aspirin use).

Plasma creatinine concentrations rose by ~12% soon after commencement of

fenofibrate (7), but there was no loss of CVD benefit among those with the greatest creatinine rise. In fact, the greatest absolute benefit was seen in this group (7).

Analysis by eGFR group showed a significant inverse association between baseline eGFR and the extent of fenofibrate-associated plasma creatinine rise, with the greatest rises seen in the lowest eGFR group, although the strength of this relationship was weak (r² = 0.03, P < 0.001). The average rise in plasma creatinine in the lowest eGFR group was approximately twice that in the highest eGFR group (17.9 vs. 8.4 μmol/L, P < 0.001 for trend across all three eGFR groups).

Analysis by the presence of different lipid abnormalities (low HDL cholesterol [male subjects <1.03 mmol/L, female subjects <1.29 mmol/L], high triglycerides [≥1.7 mmol/L], and dyslipidemia [low HDL cholesterol and high triglycerides]),

older age (≥65 years), previous CVD, or sex within each eGFR group did not reveal any statistically significant interactions. Among those with eGFR 30–59 mL/min/1.73 m², those with low HDL cholesterol or marked dyslipidemia (low HDL cholesterol plus triglycerides ≥2.3 mmol/L) had independently significant total CVD risk reductions (data not shown).

Safety of fenofibrate use in renal impairment: ESRD events overall and by eGFR

Overall, there were 26 compared with 21 ESRD events over 5 years in the placebo compared with fenofibrate group, respectively. The greatest rate of ESRD events occurred among those with baseline eGFR 30–59 mL/min/1.73 m² (stratified by treatment). There was no increase in ESRD events with fenofibrate use compared with placebo either overall (data previously

Table 2—Effect of treatment on primary and secondary outcomes by baseline eGFR

| Outcome* | Placebo | | Fenofibrate | | HR (95% CI) | P | Interaction |
|------------------------------------|---------|------|-------------|------|------------------|----------|-------------|
| | n | % | n | % | | | |
| Outcome* | | | | | | | |
| Coronary events | | | | | | | |
| eGFR (mL/min/1.73 m ²) | | | | | | | |
| 30–59 | 25 | 11.2 | 23 | 7.8 | 0.67 (0.38–1.18) | 0.17 | 0.60 |
| 60–89 | 168 | 6.3 | 148 | 5.8 | 0.91 (0.73–1.14) | 0.42 | |
| ≥90 | 95 | 4.7 | 85 | 4.2 | 0.89 (0.66–1.19) | 0.42 | |
| Coronary revascularization | | | | | | | |
| eGFR (mL/min/1.73 m ²) | | | | | | | |
| 30–59 | 25 | 11.2 | 15 | 5.1 | 0.43 (0.23–0.81) | 0.01 | 0.10 |
| 60–89 | 204 | 7.7 | 171 | 6.7 | 0.87 (0.71–1.07) | 0.18 | |
| ≥90 | 134 | 6.6 | 104 | 5.1 | 0.77 (0.59–0.99) | 0.04 | |
| CVD mortality | | | | | | | |
| eGFR (mL/min/1.73 m ²) | | | | | | | |
| 30–59 | 26 | 11.6 | 18 | 6.1 | 0.51 (0.28–0.93) | 0.03 | 0.03 |
| 60–89 | 65 | 2.4 | 78 | 3.0 | 1.25 (0.90–1.74) | 0.18 | |
| ≥90 | 36 | 1.8 | 44 | 2.2 | 1.22 (0.78–1.89) | 0.38 | |
| Total stroke | | | | | | | |
| eGFR (mL/min/1.73 m ²) | | | | | | | |
| 30–59 | 20 | 8.9 | 23 | 7.8 | 0.85 (0.47–1.55) | 0.60 | 0.33 |
| 60–89 | 91 | 3.4 | 89 | 3.5 | 1.02 (0.76–1.36) | 0.90 | |
| ≥90 | 64 | 3.2 | 46 | 2.3 | 0.71 (0.49–1.04) | 0.08 | |
| Total CVD events | | | | | | | |
| eGFR (mL/min/1.73 m ²) | | | | | | | |
| 30–59 | 60 | 26.8 | 57 | 19.3 | 0.68 (0.47–0.97) | 0.03 | 0.22 |
| 60–89 | 387 | 14.6 | 352 | 13.7 | 0.94 (0.81–1.08) | 0.38 | |
| ≥90 | 236 | 11.7 | 203 | 10.0 | 0.85 (0.70–1.02) | 0.08 | |
| Total mortality | | | | | | | |
| eGFR (mL/min/1.73 m ²) | | | | | | | |
| 30–59 | 41 | 18.3 | 43 | 14.6 | 0.77 (0.50–1.18) | 0.23 | 0.23 |
| 60–89 | 182 | 6.8 | 195 | 7.6 | 1.12 (0.91–1.37) | 0.27 | |
| ≥90 | 100 | 5.0 | 118 | 5.8 | 1.17 (0.90–1.53) | 0.24 | |
| Renal end points† | | | | | | | |
| Creatinine >400 μmol/L | | | | | | | |
| eGFR (mL/min/1.73 m ²) | | | | | | | |
| 30–59 | 2 | 0.9 | 2 | 0.7 | — | — | |
| 60–89 | 1 | 0.0 | 4 | 0.2 | — | — | |
| ≥90 | 0 | 0.0 | 0 | 0.0 | — | — | |
| Peritoneal or hemodialysis | | | | | | | |
| eGFR (mL/min/1.73 m ²) | | | | | | | |
| 30–59 | 3 | 1.3 | 5 | 1.7 | — | — | |
| 60–89 | 13 | 0.5 | 7 | 0.3 | — | — | |
| ≥90 | 5 | 0.2 | 4 | 0.2 | — | — | |
| Renal transplant | | | | | | | |
| eGFR (mL/min/1.73 m ²) | | | | | | | |
| 30–59 | 0 | 0.0 | 0 | 0.0 | — | — | |
| 60–89 | 0 | 0.0 | 0 | 0.0 | — | — | |
| ≥90 | 0 | 0.0 | 0 | 0.0 | — | — | |
| Death from renal disease | | | | | | | |
| eGFR (mL/min/1.73 m ²) | | | | | | | |
| 30–59 | 1 | 0.4 | 0 | 0.0 | — | — | |
| 60–89 | 1 | 0.0 | 1 | 0.0 | — | — | |
| ≥90 | 2 | 0.1 | 0 | 0.0 | — | — | |
| Total number of patients with ESRD | | | | | | | |
| eGFR (mL/min/1.73 m ²) | | | | | | | |
| 30–59 | 5 | 2.2 | 7 | 2.4 | — | — | |
| 60–89 | 14 | 0.5 | 10 | 0.4 | — | <0.0001‡ | |
| ≥90 | 7 | 0.3 | 4 | 0.2 | — | — | |

*Event categories are not mutually exclusive. †Most end points had insufficient numbers for statistical comparison. ‡P for trend in the rate of ESRD events across eGFR groups, stratified by treatment.




Figure 1—Percentage event rate of subtypes of vascular events and mortality within each category of eGFR. ■, 30–59 mL/min/1.73 m²; ■, 60–89 mL/min/1.73 m²; □, ≥90 mL/min/1.73 m².

published [7]) or within any eGFR group (Table 2; $P = 0.75$ for interaction). Within the group with eGFR 30–59 mL/min/1.73 m², the rates were 1.7 vs. 1.3% (dialysis), 0.7 vs. 0.9% (plasma creatinine >400 μmol/L), and 0.0 vs. 0.4% (death from renal disease) in the fenofibrate- versus placebo-allocated patients. The extent of the plasma creatinine rise with fenofibrate did not predict the occurrence of ESRD events ($P = 0.60$ for interaction).

Other serious adverse events

For approximately one-half of the categories of adverse events, the greater the renal impairment the higher the rates of occurrence (Supplementary Table 4). However, adverse event rates generally were no higher among those receiving fenofibrate than those on placebo, including within the eGFR 30–59 mL/min/1.73 m² group. The only category with a statistically significant interaction was nonfatal gastrointestinal events, with more events in the fenofibrate- than placebo-allocated patients with normal renal function (20.5 vs. 17.3% in the placebo group, $P = 0.01$, unadjusted for multiple comparisons) but no significant difference for patients with eGFR 30–59 or 60–89 mL/min/1.73 m² (Supplementary Table 4). Those allocated to fenofibrate had a statistically significant, but only marginally greater, overall risk of pancreatitis and pulmonary embolism than those allocated to placebo, but there was no difference in risk across groups by eGFR (P for interaction for both events not significant). Overall, there was no increased risk of myopathy (2 vs. 1 subjects) or rhabdomyolysis (3 vs. 1 subjects) attributed to fenofibrate, compared with placebo, with

too few events to assess whether treatment effect differed by baseline renal function.

CONCLUSIONS—Fenofibrate may be an especially appropriate therapy to reduce CVD risk in the setting of renal impairment because it raises apolipoprotein A1 and HDL cholesterol levels and influences HDL particle size (15,16). However, its use has been limited by concerns that the typical drug-induced rise in plasma creatinine might adversely affect CVD risk and renal outcomes. In the FIELD Study, we found no evidence that the use of a standard dose of fenofibrate in patients with moderate renal impairment had adverse effects on either CVD or renal outcomes. In fact, the greatest absolute risk reductions for CVD events with fenofibrate were seen in those with eGFR 30–59 mL/min/1.73 m², and no excess ESRD events were seen in fenofibrate-treated patients in any eGFR group. Of interest, although the number of subjects in the low-eGFR group was relatively small, there was an independently significant CVD reduction with fenofibrate in patients with low HDL cholesterol or marked dyslipidemia. The relative CVD benefits of fenofibrate were statistically equivalent across eGFR groups, except for CVD mortality, for which fenofibrate was associated with a significantly greater risk reduction in those with lower baseline renal function.

In general, higher baseline serum creatinine is associated with an increased risk of ESRD (12), with findings confirmed in the FIELD Study. However, the fenofibrate-induced increase in circulating creatinine levels does not carry the same risk of ESRD events because these

events were nonsignificantly fewer in those allocated to fenofibrate than to placebo.

The reasons behind the initial fenofibrate-associated plasma creatinine rise have yet to be fully elucidated, and there are various hypotheses apart from reduced glomerular filtration. These include increased muscle production of creatinine, changes in active tubular creatinine secretion, and altered renal plasma flow (7,10,17). Several of these mechanisms may be involved, but regardless of the underlying physiology, the early creatinine rise is fully reversible even after 5 years of fenofibrate treatment (7) and is concurrently associated with significant reductions in total CVD events and preservation of renal function.

In both the Diabetes Atherosclerosis Intervention Study (DAIS) (18) and the Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial (19), as well as in the FIELD Study, fenofibrate was associated with a reduction in albuminuria despite similar elevations in serum creatinine to those seen in the FIELD Study (7). In a recent U.S. Food and Drug Administration hearing, the ACCORD trial reported that after fenofibrate withdrawal eGFR returned to placebo-matched levels, with some patients having an overall improvement in renal function (20). The ACCORD trial (19) and other studies (21–23) also show that fenofibrate is not associated with excess kidney-related adverse events and is, overall, safe to use even in combination with statins. However, none of these previous study reports have investigated if there is any relationship between outcomes and the drug-induced creatinine rise or examined the safety and efficacy of fenofibrate use in renal impairment. To our knowledge, this article is the first to do so, looking at both cardiovascular and renal outcomes. It provides added reassurance, demonstrating that the drug is not only safe but also just as efficacious in patients with moderate renal impairment compared with those with normal function.

Clinical implications

Diabetes is a significant risk factor for CVD events and mortality, requiring intensive treatment of conventional CVD risk factors. Unfortunately, the implementation of established preventive treatment recommendations remains incomplete (24), and even with intensive multifactorial risk reduction, the CVD complication rate in type 2 diabetic patients remains unacceptably high (25). The FIELD Study confirms that type 2 diabetic patients




Figure 2—HRs (95% CIs) for total CVD events, cardiovascular mortality, and total mortality risk by treatment allocation within different categories of eGFR. ■, relative weights of the outcome event in each group; ◇, summary HR with 95% CI.

with reduced eGFR (<90 mL/min/1.73 m²) are at exceptionally high risk of CVD (>5% per year). This underscores the need for better implementation of proven treatments, especially in those with coexistent renal impairment.

These FIELD Study findings should help allay previous concerns regarding the safety of fenofibrate use in patients with renal impairment, indicating that neither moderately poor renal function nor an initial rise in plasma creatinine with fenofibrate is a contraindication to such treatment. Furthermore, treatment benefits at least as great as those seen in patients with normal renal function could be expected. The benefits reported here support the long-term use of standard doses of fenofibrate in such patients to reduce both CVD events and death from CVD. These benefits occur despite a reversible and modest

increase in plasma creatinine, with evidence of renoprotection (eGFR preservation and albuminuria reduction) that was similar across the eGFR groups (7). The benefits of fenofibrate on CVD and albuminuria are both in addition to and independent of established proven measures, such as glycemic control, antihypertensive treatment in general, and use of the renin-angiotensin system blockade in particular (7,26–33). In this context, the results would support the use of fenofibrate being in addition to, rather than an alternative to, the above current established strategies.

Fenofibrate therapy in those in the group with low eGFR produced the greatest estimated absolute reduction in CVD risk despite the greatest early plasma creatinine rise. Therefore, not only does this early drug-induced plasma creatinine rise fail to portend any adverse cardiac or renal

outcomes, but it may in fact be a surrogate marker of drug benefit. This latter finding is hypothesis generating and will require further clinical investigation.

Current guidelines for fenofibrate use in Western countries recommend a substantial dose reduction (between 30 and 69%) when renal function is impaired but with significant variation in the threshold at which this should be implemented. Dose reduction is recommended when creatinine clearance is <100 mL/min in Canada (9,34) to <80 mL/min in the U.S. (35) and <60 mL/min in the U.K. and Australia (8,36). Recommendations for dose reduction also vary by manufacturer, which may cause confusion and undue concern about fenofibrate use in patients with any level of renal impairment (34–37). Our current findings suggest that these thresholds may be too restrictive and that fenofibrate

dose reductions may not be required until eGFR falls below 30 mL/min/1.73 m².

Study limitations

Our study had limitations. The MDRD Study four-variable formula may underestimate true GFR, particularly when eGFR is ≥ 90 mL/min/1.73 m², but is more accurate at lower levels (38). However, our findings essentially were unchanged if analyzed using the Chronic Kidney Disease Epidemiology Collaboration (EPI-CKD) formula for derived GFR (7). The severity classification of eGFR is somewhat arbitrary, and within-individual variation in eGFR may have incorrectly allocated patients who were close to category cutoffs. However, baseline plasma creatinine and eGFR values in the FIELD Study were averaged over two pretreatment visits, and samples were analyzed at a central laboratory. In addition, the eGFR categories were based on standard clinical groupings to facilitate comparison with other studies. The number of patients in the group with eGFR 30–59 mL/min/1.73 m² was low compared with the other two groups, with fewer events. Despite this, there was a general pattern of greater risk reduction secondary to fenofibrate in this group compared with the other two across a wide range of individual and composite end points, although these were not statistically significant except for CVD mortality. Although there was a slight imbalance in the numbers of individuals with eGFR 30–59 mL/min/1.73 m² between treatment groups, this was minor and unlikely to have confounded the effect of fenofibrate on total CVD reduction because it did not differ consistently across groups of renal function. Patients with eGFR < 30 mL/min/1.73 m² were excluded; therefore, the study cannot address the safety issues for this group, for whom reduced clearance of fenofibrate and its metabolites has been reported (39). In a previous study, patients with severe renal impairment had a greater unbound fraction of active fenofibrate metabolites than those with normal renal function. This increase is balanced by greater renal clearance, except when creatinine clearance falls below 20 mL/min, at which point the unbound concentration increases accordingly and dose reduction becomes necessary (39).

CONCLUSIONS—In the FIELD Study, fenofibrate did not increase CVD or renal risk in the setting of mild

or moderate renal impairment in type 2 diabetes, despite causing an elevation in plasma creatinine. In those with pre-existing moderate renal impairment, the CVD benefits of fenofibrate were at least as great as in those with normal renal function, without any additional safety concerns. Therefore, fenofibrate at standard doses should be considered as an additional therapeutic option, along with conventional risk factor management, to further reduce CVD events and mortality and afford renoprotection in patients with type 2 diabetes, even in patients with moderate renal impairment.

Acknowledgments—The FIELD Study was supported by grants from Laboratoires Fournier SA, Dijon, France (now part of Abbott Pharmaceuticals), and the National Health and Medical Research Council (NHMRC) of Australia.

R.-D.T., A.C.K., T.M.E.D., R.J.S., and K.R. have received speaker's bureau fees from Solvay (now Abbott Pharmaceuticals). A.C.K., P.L.D., A.J.J., T.M.E.D., and R.J.S. have received travel and/or research grant support from Solvay (now Abbott Pharmaceuticals). No other potential conflicts of interest relevant to this article were reported.

R.-D.T. researched data, contributed to the discussion, and wrote the manuscript. A.C.K. is the study guarantor, researched data, contributed to the discussion, and wrote the manuscript. P.L.D., A.J.J., T.M.E.D., R.J.S., and K.S. researched data, contributed to the discussion, and reviewed and edited the manuscript. M.W.D. and K.R. researched data and reviewed and edited the manuscript. J.H. researched data. S.L. reviewed and edited the manuscript. D.C. contributed to the discussion and reviewed and edited the manuscript. The authors had full access to all study data.

Parts of this study were presented in abstract form at the ESC Congress 2011 of the European Society of Cardiology, Paris, France, 27–31 August 2011, and at the 59th Annual Scientific Meeting of the Cardiac Society of Australia and New Zealand, Perth, Australia, 11–14 August 2011.

The study was designed by an independent management committee and was coordinated by the NHMRC Clinical Trials Centre, University of Sydney. Two nonvoting representatives of the main sponsor attended meetings of the management committee.

The authors thank the many patients and investigators who participated in the FIELD Study.

References

- World Health Organization. Cardiovascular diseases (CVD) [fact sheet online], 2009. Available from <http://www.who.int/mediacentre/factsheets/fs317/en/index.html>. Accessed 3 March 2010
- Hemmelgarn BR, Manns BJ, Lloyd A, et al.; Alberta Kidney Disease Network. Relation between kidney function, proteinuria, and adverse outcomes. *JAMA* 2010;303:423–429
- Drury PL, Ting R, Zannino D, et al. Estimated glomerular filtration rate and albuminuria are independent predictors of cardiovascular events and death in type 2 diabetes mellitus: the Fenofibrate Intervention and Event Lowering in Diabetes (FIELD) Study. *Diabetologia* 2011;54:32–43
- Keech A, Simes RJ, Barter P, et al.; FIELD Study Investigators. Effects of long-term fenofibrate therapy on cardiovascular events in 9795 people with type 2 diabetes mellitus (the FIELD Study): randomised controlled trial. *Lancet* 2005;366:1849–1861
- Rajamani K, Colman PG, Li LP, et al.; FIELD Study Investigators. Effect of fenofibrate on amputation events in people with type 2 diabetes mellitus (FIELD Study): a prespecified analysis of a randomised controlled trial. *Lancet* 2009;373:1780–1788
- Keech AC, Mitchell P, Summanen PA, et al.; FIELD Study Investigators. Effect of fenofibrate on the need for laser treatment for diabetic retinopathy (FIELD Study): a randomised controlled trial. *Lancet* 2007;370:1687–1697
- Davis TM, Ting R, Best JD, et al.; Fenofibrate Intervention and Event Lowering in Diabetes Study Investigators. Effects of fenofibrate on renal function in patients with type 2 diabetes mellitus: the Fenofibrate Intervention and Event Lowering in Diabetes (FIELD) Study. *Diabetologia* 2011;54:280–290
- Abbott Pharmaceuticals. Fenofibrate product information, 2011. Available from www.medicines.org.au/files/smplipid.pdf. Accessed 10 December 2011
- McPherson R, Frohlich J, Fodor G, Genest J; Canadian Cardiovascular Society. Canadian Cardiovascular Society position statement: recommendations for the diagnosis and treatment of dyslipidemia and prevention of cardiovascular disease. *Can J Cardiol* 2006;22:913–927
- Ansquer J-C, Dalton RN, Caussé E, Crimet D, Le Malicot K, Foucher C. Effect of fenofibrate on kidney function: a 6-week randomized crossover trial in healthy people. *Am J Kidney Dis* 2008;51:904–913
- Ninomiya T, Perkovic V, de Galan BE, et al.; ADVANCE Collaborative Group. Albuminuria and kidney function independently predict cardiovascular and renal outcomes in diabetes. *J Am Soc Nephrol* 2009;20:1813–1821
- Shahinfar S, Dickson T, Zhang Z, Keane WF, Brenner BM; RENAAL Investigators. Baseline predictors of end-stage renal

- disease risk in patients with type 2 diabetes and nephropathy: new lessons from the RENAAL study. *Kidney Int* 2005;67 (Suppl.): S48–S51
13. Vassalotti JA, Stevens LA, Levey AS. Testing for chronic kidney disease: a position statement from the National Kidney Foundation. *Am J Kidney Dis* 2007;50: 169–180
 14. Levey AS, Greene T, Kusek J, Beck G. A simplified equation to predict glomerular filtration rate from serum creatinine (Abstract). *J Am Soc Nephrol* 2000;11:155A
 15. Ikewaki K, Tohyama J, Nakata Y, Wakikawa T, Kido T, Mochizuki S. Fenofibrate effectively reduces remnants, and small dense LDL, and increases HDL particle number in hypertriglyceridemic men: a nuclear magnetic resonance study. *J Atheroscler Thromb* 2004;11:278–285
 16. Bilz S, Wagner S, Schmitz M, Bedynek A, Keller U, Demant T. Effects of atorvastatin versus fenofibrate on apoB-100 and apoA-I kinetics in mixed hyperlipidemia. *J Lipid Res* 2004;45:174–185
 17. Hottelart C, El Esper N, Rose F, Achard J-M, Fournier A. Fenofibrate increases creatinemia by increasing metabolic production of creatinine. *Nephron* 2002; 92:536–541
 18. Ansquer J-C, Foucher C, Rattier S, Taskinen M-R, Steiner G; DAIS Investigators. Fenofibrate reduces progression to microalbuminuria over 3 years in a placebo-controlled study in type 2 diabetes: results from the Diabetes Atherosclerosis Intervention Study (DAIS). *Am J Kidney Dis* 2005;45:485–493
 19. Ginsberg HN, Elam MB, Lovato LC, et al.; ACCORD Study Group. Effects of combination lipid therapy in type 2 diabetes mellitus. *N Engl J Med* 2010;362:1563–1574
 20. U.S. Food and Drug Administration. 2011 Meeting materials, endocrinologic and metabolic drugs advisory committee [article online], 2011. Available from <http://www.fda.gov/AdvisoryCommittees/CommitteesMeetingMaterials/Drugs/EndocrinologicandMetabolicDrugsAdvisoryCommittee/ucm252891.htm>. Accessed 26 September 2011
 21. Jones PH, Goldberg AC, Knapp HR, et al. Efficacy and safety of fenofibric acid in combination with atorvastatin and ezetimibe in patients with mixed dyslipidemia. *Am Heart J* 2010;160:759–766
 22. Goldberg AC, Bays HE, Ballantyne CM, et al. Efficacy and safety of ABT-335 (fenofibric acid) in combination with atorvastatin in patients with mixed dyslipidemia. *Am J Cardiol* 2009;103:515–522
 23. Roth EM, McKenney JM, Kelly MT, et al. Efficacy and safety of rosuvastatin and fenofibric acid combination therapy versus simvastatin monotherapy in patients with hypercholesterolemia and hypertriglyceridemia: a randomized, double-blind study. *Am J Cardiovasc Drugs* 2010; 10:175–186
 24. Saydah SH, Fradkin J, Cowie CC. Poor control of risk factors for vascular disease among adults with previously diagnosed diabetes. *JAMA* 2004;291:335–342
 25. Gaede P, Lund-Andersen H, Parving H-H, Pedersen O. Effect of a multifactorial intervention on mortality in type 2 diabetes. *N Engl J Med* 2008;358:580–591
 26. Cooper ME, Jandeleit-Dahm K, Thomas MC. Targets to retard the progression of diabetic nephropathy. *Kidney Int* 2005; 68:1439–1445
 27. Gaede P, Vedel P, Parving H-H, Pedersen O. Intensified multifactorial intervention in patients with type 2 diabetes mellitus and microalbuminuria: the Steno Type 2 Randomised Study. *Lancet* 1999;353: 617–622
 28. Gerstein HC, Yusuf S, Mann JFE, et al.; Heart Outcomes Prevention Evaluation Study Investigators. Effects of ramipril on cardiovascular and microvascular outcomes in people with diabetes mellitus: results of the HOPE study and MICRO-HOPE substudy. *Lancet* 2000;355:253–259
 29. Patel A, MacMahon S, Chalmers J, et al.; ADVANCE Collaborative Group. Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. *N Engl J Med* 2008;358:2560–2572
 30. UK Prospective Diabetes Study Group. Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38. *BMJ* 1998;317:703–713
 31. Ismail-Beigi F, Craven T, Banerji MA, et al.; ACCORD Trial Group. Effect of intensive treatment of hyperglycaemia on microvascular outcomes in type 2 diabetes: an analysis of the ACCORD randomised trial. *Lancet* 2010;376:419–430
 32. Turner RC, Holman RR, Cull CA, et al.; UK Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* 1998;352:837–853
 33. Brenner BM, Cooper ME, de Zeeuw D, et al.; RENAAL Study Investigators. Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy. *N Engl J Med* 2001;345:861–869
 34. Cipher Pharmaceuticals. Fenomax product monograph, 2006. Available from webprod3.hc-sc.gc.ca/dpd-bdpp/index-eng.jsp. Accessed 10 December 2011
 35. Abbott Laboratories. Tricor product information [article online], 2010. Available from <http://www.rxabbott.com/pdf/tricorpi.pdf>. Accessed 16 December 2010
 36. Abbott Healthcare Products Limited. Lipantil product information [article online], 2009. Available from <http://www.medicines.org.uk/emc/medicine/2567>. Accessed 16 December 2010
 37. Shionogi Pharma. Triglide product information [article online], 2010. Available from <http://www.triglide.com/triglide%20pi.pdf>. Accessed 16 December 2010
 38. Chudleigh RA, Dunseath G, Evans W, et al. How reliable is estimation of glomerular filtration rate at diagnosis of type 2 diabetes? *Diabetes Care* 2007;30:300–305
 39. Laboratoires Fournier. *Fenofibrate Investigator's Brochure*. Daix, France, Laboratoires Fournier Research Centre, 2004

Minerva Access is the Institutional Repository of The University of Melbourne

Author/s:

Ting, R-D; Keech, AC; Drury, PL; Donoghoe, MW; Hedley, J; Jenkins, AJ; Davis, TME;
Lehto, S; Celermajer, D; Simes, RJ; Rajamani, K; Stanton, K

Title:

Benefits and Safety of Long-Term Fenofibrate Therapy in People With Type 2 Diabetes and Renal Impairment The FIELD Study

Date:

2012-02-01

Citation:

Ting, R. -D., Keech, A. C., Drury, P. L., Donoghoe, M. W., Hedley, J., Jenkins, A. J., Davis, T. M. E., Lehto, S., Celermajer, D., Simes, R. J., Rajamani, K. & Stanton, K. (2012). Benefits and Safety of Long-Term Fenofibrate Therapy in People With Type 2 Diabetes and Renal Impairment The FIELD Study. *DIABETES CARE*, 35 (2), pp.218-225.
<https://doi.org/10.2337/dc11-1109>.

Persistent Link:

<http://hdl.handle.net/11343/264390>

File Description:

Published version

License:

CC BY-NC-ND