

Tilburg University

Fissioned Triangular Schemes Via the Cross-Ratio

van Dam, E.R.; de Caen, D.

Publication date:
1999

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):

van Dam, E. R., & de Caen, D. (1999). *Fissioned Triangular Schemes Via the Cross-Ratio*. (FEW Research Memorandum; Vol. 782). Operations research.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Fissioned triangular schemes via the cross-ratio

D. de Caen and E.R. van Dam

Abstract

A construction of association schemes is presented; these are fission schemes of the triangular schemes $T(n)$ where $n = q + 1$ with q any prime power. The key observation is quite elementary, being that the natural action of $PGL(2, q)$ on the 2-element subsets of the projective line $PG(1, q)$ is generously transitive. Also some observations on the intersection parameters and fusion schemes of these association schemes are made.

1 The construction

This paper is a sequel to [4]. In that paper, it was observed that almost all known self-dual classical association schemes have natural fission schemes (fissioning the maximum-distance relation only); whereas in the non-self-dual case there seemed to be no analogous fission schemes. Subsequently, we found that there is at least one such non-self-dual classical association scheme that admits an interesting fission scheme, namely the triangular scheme $T(n) = J(n, 2)$ where $n = q + 1$ with q any prime power; this is the object of the present work. For terminology and background, we refer to Bannai and Ito [2] for association schemes and Hirschfeld [7] for finite geometry. Recall that the group $PGL(2, q)$ acts (as Möbius transformations) on the projective line $PG(1, q)$; this action is (sharply) 3-transitive. There is a natural induced action on the 2-element subsets of the projective line, namely $M(\{x, y\}) := \{M(x), M(y)\}$ for each M in $PGL(2, q)$. In the proof below we apply the basic fact (cf. [7], p. 135) that the cross-ratio

$$\rho(a, b, c, d) := \frac{(a - c)(b - d)}{(a - d)(b - c)}$$

is a complete invariant for ordered quadruples of distinct points on the projective line, i.e. one quadruple may be mapped to another quadruple (via a Möbius transformation) if and only if they have the same cross-ratio.

Theorem. *The action of $PGL(2, q)$ on the two-element subsets of $PG(1, q)$ is generously transitive.*

Proof. Given intersecting 2-sets $\{a, b\}$ and $\{a, c\}$, there is some M in $PGL(2, q)$ that swaps them, since the group is triply transitive. And given disjoint 2-sets $\{a, b\}$ and $\{c, d\}$, there is also some Möbius transformation that interchanges them, because the ordered quadruples (a, b, c, d) and (c, d, a, b) have the same cross-ratio. \square

Given any transitive permutation group G acting on a set Ω , the orbitals are the orbits in $\Omega \times \Omega$ under the natural action of G on pairs. If G is generously transitive, then the orbitals form the relations (associate classes) of a symmetric association scheme (cf. [2], p. 54). In our case, the relations can be described as follows. One relation, say R_1 , is the line-graph of the complete graph (i.e. one relation of the triangular scheme $T(q + 1)$ has remained unfissioned). Next, for each reciprocal pair $\{s, s^{-1}\}$ of elements in $GF(q) \setminus \{0, 1\}$, there is a relation $R_{\{s, s^{-1}\}}$ where $\{a, b\}$ and $\{c, d\}$ are in this relation

when $\rho(a, b, c, d)$ equals s or s^{-1} . Note that $\rho(b, a, c, d) = \rho(a, b, c, d)^{-1}$ so this makes sense as a definition for unordered pairs $\{a, b\}$. Henceforth we will write R_s instead of $R_{\{s, s^{-1}\}}$ for typographical reasons; note that since the field element 1 cannot occur as a cross-ratio, this notation will not conflict with that of relation R_1 above.

We now easily find that this fissioned triangular scheme, which we shall denote by $FT(q+1)$, has $\frac{1}{2}(q+1)$ associate classes if q is odd and $\frac{1}{2}q$ classes if q is even. When q is odd the field element -1 is equal to its own reciprocal; thus the relation R_{-1} has valency $\frac{1}{2}(q-1)$ which is half the valency of the other relations R_s with s in $GF(q) \setminus \{0, 1, -1\}$. The relation R_1 has valency $2(q-1)$.

We remark that for small odd q the relation R_{-1} is a familiar object: for $q = 5$ it is the line-graph of Petersen's graph; for $q = 7$ it is the Coxeter graph (this was apparently known to Coxeter himself, cf. p. 122 in [6]); for $q = 9$ it is the line-graph of Tutte's 8-cage. There seem to be some other such "sporadic isomorphisms": for example when $q = 11$ the relation $R_2 = R_{\{2, 6\}}$ is the line-graph of the point-block incidence graph of the (unique) symmetric $(11, 6, 3)$ -design; and when $q = 9$ and $\{s, s^{-1}\}$ is the pair of primitive fourth roots of unity, then R_s is the second subconstituent of the Gewirtz graph (cf. [5], page 106).

2 Intersection parameters

It is possible to give explicit formulas for the intersection parameters p_{ij}^k of the association scheme $FT(q+1)$; we now sketch the main points of the derivation. The cases q odd and q even are similar, with the latter case being slightly cleaner since the exceptional case " $\rho = -1$ " doesn't occur. So we will only present the case q even; besides, this case is the more pertinent one in the discussion of fusion schemes in Section 3.

So let $q = 2^e$ be any power of two. The scheme $FT(2^e + 1)$ has 2^{e-1} classes. The relation R_1 has valency $2(q-1)$ and each of the other relations $R_s = R_{\{s, s^{-1}\}}$ (for s in $GF(q) \setminus \{0, 1\}$) has valency $q-1$. The intersection parameters involving R_1 are easy to work out and we list them without proof: for distinct r and s (and $s \neq r^{-1}$) in $GF(q) \setminus \{0, 1\}$, $p_{11}^1 = q-1$, $p_{11}^r = 4$, $p_{1r}^1 = 2$, $p_{rr}^1 = 1$, and $p_{rs}^1 = 2$.

Now let the symbols r, s and t represent three (not necessarily distinct) elements of $GF(q) \setminus \{0, 1\}$; we aim at a formula for p_{st}^r . What one has to do is fix a pair of 2-sets

$\{a, b\}$ and $\{c, d\}$ in relation R_r , and count the number of 2-sets $\{x, y\}$ such that $\{a, b\}$ and $\{x, y\}$ are in relation R_s and $\{c, d\}$ and $\{x, y\}$ are in relation R_t . The triple transitivity of $PGL(2, q)$ is useful here, since it implies that we may take, without loss of generality, $\{a, b\} = \{\infty, 0\}$ and $\{c, d\} = \{1, r\}$. For the unknown pair $\{x, y\}$ we then get the two equations

$$s \text{ or } s^{-1} = \frac{(\infty - x)(0 - y)}{(\infty - y)(0 - x)} = \frac{y}{x} \quad (1)$$

and

$$t \text{ or } t^{-1} = \frac{(1 - x)(r - y)}{(1 - y)(r - x)} \quad (2)$$

The equations (1) and (2) together involve two essentially different cases, not four, since $\{y, x\} = \{x, y\}$; thus we may fix the left-hand side of (1) as being s , and examine the two cases for (2) in turn. In the first case we have $y = sx$ and

$$t = \frac{(1 - x)(r - y)}{(1 - y)(r - x)} = \frac{(1 - x)(r - sx)}{(1 - sx)(r - x)}$$

This leads to the following quadratic for x (after changing all minus signs to plus signs, as we may since we are in characteristic two):

$$s(t + 1)x^2 + (rst + r + s + t)x + r(t + 1) = 0 \quad (3)$$

The other case (when the left-hand side of (2) is t^{-1}) leads to the similar quadratic

$$s(t + 1)x^2 + (rs + rt + st + 1)x + r(t + 1) = 0 \quad (4)$$

Note that since r, s and t are all in $GF(q) \setminus \{0, 1\}$, the equations (3) and (4) are genuine quadratics, with non-zero quadratic and constant terms. The linear coefficient $(rst + r + s + t)$ in (3) could equal 0, in which case the unique solution for x is the square root of $\frac{r}{s}$. If $rst + r + s + t \neq 0$, then (3) has (two) solutions x if and only if

$$Tr \left[\frac{rs(t + 1)^2}{(rst + r + s + t)^2} \right] = 0 \quad (5)$$

where $Tr(z)$ is the trace map from $GF(2^e)$ onto $GF(2)$. Similarly, if $rs + rt + st + 1 \neq 0$ then (4) has (two) solutions x if and only if

$$Tr \left[\frac{rs(t+1)^2}{(rs + rt + st + 1)^2} \right] = 0 \quad (6)$$

Thus p_{st}^r has a value of anywhere from 0 to 4. A reasonably concise formula is the following: let $A = A(r, s, t)$ be the expression for the argument of the trace map in (5), and $B = B(r, s, t)$ the one for (6). Then, when $rst + r + s + t \neq 0$ and $rs + rt + st + 1 \neq 0$

$$p_{st}^r = 2 + (-1)^{Tr[A]} + (-1)^{Tr[B]} \quad (7)$$

with the obvious modifications being made in the other cases. Incidentally, it is easy to check that $(rst + r + s + t)$ and $(rs + rt + st + 1)$ cannot simultaneously equal 0.

We make one more remark concerning the form of the intersection parameters. The expressions $A(r, s, t)$ and $B(r, s, t)$ are not symmetric in s and t , hence the formula (7) for p_{st}^r appears not to be symmetric either. This may seem strange, since we know from general principles that $p_{st}^r = p_{ts}^r$. An explanation for this is the following. $A(r, s, t)$ has the same trace as $C(r, s, t) := \frac{rs+rt+st}{(rst+r+s+t)^2}$ since their sum is of the form $\frac{xy}{x^2+y^2}$ and such field elements, in characteristic two, must have trace 0 (exercise for the reader). Similarly $B(r, s, t)$ has the same trace as $D(r, s, t) := \frac{rst(r+s+t)}{(rs+rt+st+1)^2}$. Thus we may replace A by C and B by D in (7) without changing the value of the right-hand side; and C and D are both symmetric functions of the three variables r, s and t . This confirms the fact that, since the valencies n_r are the same for all r in $GF(q) \setminus \{0, 1\}$, the intersection parameter p_{st}^r is symmetric in all three variables.

It would be interesting to find explicit formulas for the entries of the eigenmatrix (character table) of $FT(q+1)$. One strategy for doing this (used by Bannai and his co-workers in several papers; see [1] for a survey) is the following. First calculate all of the intersection parameters; it is usually feasible to do this, at least in some reasonable algebraic form perhaps involving character sums. This tells us what the intersection matrices $B_i(k, j) := p_{ij}^k$ are. Secondly, from these B_i 's (at small values of q) it may be possible to guess what the eigenmatrix P should be. Once the right guess has been made it is usually straightforward to actually prove the result, using Theorem II.4.1 in [2]. Unfortunately, we have been unable so far to guess the general shape of P for our

schemes $FT(q+1)$; we generated by computer these character tables for all prime powers q less than 40, and they seem to have a very complicated form.

3 Fusion schemes

Given any association scheme, it is of interest to determine all of its fusion schemes (also called subschemes). This is in general a very hard problem that has not been worked out completely even for quite classical examples such as the Johnson schemes (cf. [8]). In the case of the schemes $FT(q+1)$, there is of course the original two-class triangular scheme $T(q+1)$. Observe also that if $q = p^e$ is a proper power of a prime p , then the Frobenius map $x \mapsto x^p$ (and its iterates) gives a fusion scheme. In other words $PGL(2, q)$ is an overgroup of $PGL(2, p)$, and the orbitals under $PGL(2, q)$ constitute a fusion scheme of $FT(q+1)$.

Limited computational evidence suggests that $FT(q+1)$ has no other nontrivial fusions, except maybe in some sporadic cases, and when $q = 4^f$ (f any integer at least 2) where there seems to be an interesting 4-class fusion scheme. We say “seems” because we are lacking a proof that this is indeed an association scheme. To describe this (putative) scheme, let the ground-set be all 2-element subsets of the projective line $PG(1, 4^f)$; the four possible relations for two distinct 2-sets $\{a, b\}$ and $\{c, d\}$ are:

- S_1 : $\{a, b\} \cap \{c, d\} \neq \emptyset$, i.e. R_1 in the earlier notation.
- S_2 : $\{a, b\} \cap \{c, d\} = \emptyset$ and the cross-ratio $\rho = \rho(a, b, c, d)$ satisfies $\rho^{2^f-1} = 1$,
i.e. ρ lies in the subfield $GF(2^f)$.
- S_3 : $\{a, b\} \cap \{c, d\} = \emptyset$ and the cross-ratio $\rho = \rho(a, b, c, d)$ satisfies $\rho^{2^f+1} = 1$.
- S_4 : The remainder.

We have been able to show by computer that these four relations do indeed form a scheme when f is less than or equal to 6. Also we can prove in general that some of the intersection parameters, such as p_{23}^3 , are well defined; but certain other parameters such as p_{33}^3 have left us baffled. An explicit knowledge of the eigenmatrix of $FT(4^f+1)$ would theoretically settle this question (cf. [8], Lemma 1), which is partly why we earlier raised the issue of computing it.

Conjecture. *The above relations S_i on the 2-subsets of $PG(1, 4^f)$ do form a 4-class association scheme for all $f \geq 2$. The corresponding eigenmatrix is given by*

$$P = \begin{bmatrix} 1 & 2(4^f - 1) & (2^{f-1} - 1)(4^f - 1) & 2^{f-1}(4^f - 1) & 2^f(2^{f-1} - 1)(4^f - 1) \\ 1 & 4^f - 3 & 2 - 2^f & -2^f & -2^f(2^f - 2) \\ 1 & -2 & 1 - 2^f & 0 & 2^f \\ 1 & -2 & (2^{f-1} - 1)(2^f - 1) & 2^{f-1}(2^f - 1) & -2^f(2^f - 2) \\ 1 & -2 & 2^{f-1}(2^f - 1) + 1 & -2^{f-1}(2^f + 1) & 2^f \end{bmatrix}$$

We note finally that, granting this conjecture, one can merge S_2 and S_3 to get a 3-class scheme, and then further merge S_1 with S_2 and S_3 to get a 2-class scheme. The resulting graph $G = S_1 \cup S_2 \cup S_3$ is strongly regular with parameters $v = 2^{2f-1}(2^{2f} + 1)$, $k = (2^f + 1)(2^{2f} - 1)$, $\lambda = (2^f - 1)(3 \cdot 2^f + 2)$, $\mu = 2^{f+1}(2^f + 1)$. Graphs with these parameters have already been constructed by Brouwer and Wilbrink (cf. [3], 7B); it was checked that in the smallest case $f = 2$ ($v = 136$) the two constructions yield isomorphic strongly regular graphs. We know nothing for larger values; but the two constructions look totally different, so that it is a reasonable guess that they are not isomorphic in general.

References

- 1 E. Bannai, Character tables of commutative association schemes, pp. 105-128 in “Finite geometries, buildings, and related topics”, Oxford Univ. Press, 1990.
- 2 E. Bannai and T. Ito, “Algebraic Combinatorics I: Association schemes”, Benjamin/Cummings Publ., 1984.
- 3 A.E. Brouwer and J.H. van Lint, Strongly regular graphs and partial geometries, pp. 85-122 in “Enumeration and Design”, Academic Press, 1984.
- 4 D. de Caen and E.R. van Dam, Fissions of classical self-dual association schemes, Journal of Combinatorial Theory Series A 88 (1999), 167-175.
- 5 P.J. Cameron and J.H. van Lint, “Designs, graphs, codes and their links”, Cambridge Univ. Press, 1991.
- 6 H.S.M. Coxeter, My graph, Proc. London Math. Soc. 46 (1983), 117-136.
- 7 J.W.P. Hirschfeld, “Projective geometry over finite fields”, Oxford Univ. Press, 2nd edition 1998.
- 8 M.E. Muzychuk, Subschemes of the Johnson scheme, European Journal of Combinatorics 13 (1992), 187-193.

Authors' addresses:

<p>D. de Caen Dept. of Mathematics & Statistics Queen's University Kingston, Canada K7L 3N6 decaen@mast.queensu.ca</p>	<p>E.R. van Dam Dept. of Econometrics Tilburg University P.O. Box 90153, 5000 LE Tilburg, The Netherlands Edwin.vanDam@kub.nl</p>
--	---