

Tilburg University

One-year follow-up results of a randomized controlled clinical trial on internet-based cognitive behavioural therapy for subthreshold depression in people over 50 years

Spek, V.R.M.; Cuijpers, P.; Nyklicek, I.; Smits, N.; Riper, H.; Keyzer, J.J.; Pop, V.J.M.

Published in:
Psychological Medicine

Publication date:
2008

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):

Spek, V. R. M., Cuijpers, P., Nyklicek, I., Smits, N., Riper, H., Keyzer, J. J., & Pop, V. J. M. (2008). One-year follow-up results of a randomized controlled clinical trial on internet-based cognitive behavioural therapy for subthreshold depression in people over 50 years. *Psychological Medicine*, 38(5), 635-639.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

One-year follow-up results of a randomized controlled clinical trial on internet-based cognitive behavioural therapy for subthreshold depression in people over 50 years

V. Spek¹*, P. Cuijpers², I. Nyklíček¹, N. Smits², H. Riper³, J. Keyzer⁴ and V. Pop¹

¹ Department of Psychology and Health, Tilburg University, The Netherlands

² Department of Clinical Psychology, Vrije Universiteit Amsterdam, The Netherlands

³ Trimbos-instituut, Netherlands Institute of Mental Health and Addiction, The Netherlands

⁴ Diagnostic Centre Eindhoven, The Netherlands

Background. Internet-based cognitive behaviour therapy (CBT) is a promising new approach for the treatment of depressive symptoms. The current study had two aims: (1) to determine whether, after 1 year, an internet-based CBT intervention was more effective than a waiting-list control group; and (2) to determine whether the effect of the internet-based CBT differed from the group CBT intervention, 1 year after the start of treatment.

Method. A total of 191 women and 110 men (mean age=55 years, s.d.=4.6) with subthreshold depression were randomized into internet-based treatment, group CBT (Lewinsohn's Coping with Depression Course), or a waiting-list control condition. The main outcome measure was treatment response after 1 year, defined as the difference in pretreatment and follow-up scores on the Beck Depression Inventory (BDI). Missing data were imputed using the multiple imputation procedure of data augmentation. Analyses were performed using multiple imputation inference.

Results. In the waiting-list control group, we found a pretreatment to follow-up improvement effect size of 0.69, which was 0.62 in the group CBT condition and 1.22 with the internet-based treatment condition. Simple contrasts showed a significant difference between the waiting-list condition and internet-based treatment ($p=0.03$) and no difference between both treatment conditions ($p=0.08$).

Conclusions. People aged over 50 years with subthreshold depression can still benefit from internet-based CBT 1 year after the start of treatment.

Received 5 July 2007; Revised 20 November 2007; Accepted 28 November 2007; First published online 21 January 2008

Key words: Cognitive behaviour therapy, depression, internet.

Introduction

A promising new approach regarding treatment of depressive symptoms is internet-based cognitive behaviour therapy (CBT). The short-term effectiveness of this type of treatment for symptoms of depression has been proven in several studies (Christensen *et al.* 2004; Andersson *et al.* 2005; Clarke *et al.* 2005). We have also reported a significant effect of internet-based therapy at short-term follow-up (Spek *et al.* 2007a). However, there has been very little research conducted on the long-term effects of internet-based treatments. The longest follow-up period so far reported has been 6 months (Andersson *et al.* 2005).

The current study evaluated the 1-year follow-up effects of internet-based CBT for subthreshold depression in people aged over 50 years. We compared the effect of internet-based therapy with a waiting-list condition and with group CBT.

Method

Participants

To be included in the study, participants had to meet the following criteria: an Edinburgh Depression Scale (EDS) score of 12 or more, but no DSM-IV diagnosis of depression, signed informed consent, age between 50 and 75 years, access to the internet and the ability to use the internet. Exclusion criteria were: suffering from any other psychiatric disorder in immediate need of treatment (assessed by means of an anamnesis

* Address for correspondence: Dr V. Spek, Tilburg University, PO Box 90153, 5000 LE Tilburg, The Netherlands.
(Email: v.r.m.spek@uvt.nl)




Fig. 1. Flow chart of participants.

during the interview) and suicidal ideation (assessed in the screening questionnaire and later in depth during the CIDI) (see Spek *et al.* 2007a).

Measures

The EDS (Cox *et al.* 1987, 1996; Murray & Carothers, 1990) was used as the screening instrument in this study. For the primary outcome measure, we used the 21-item Beck Depression Inventory – second edition (BDI-II; Beck *et al.* 1996). The World Health Organization CIDI (WHO, 1997) was used to assess depression in this study at inclusion.

Procedure

Participants with an EDS score of 12 or more were invited for a face-to-face clinical interview, demographic data were collected, and a structured interview was conducted to assess the DSM-IV criteria of depression. Eligible participants were randomized, and asked to fill in the BDI at home, after which the treatment started. Ten weeks after the start of treatment or after 10 weeks on the waiting-list, participants were asked

to complete the post-treatment BDI (Spek *et al.* 2007a). One year after the interview, participants were asked to complete another BDI.

The study protocol was approved by the Maxima Medisch Centrum (local hospital) ethics committee, which is certified by the Central Committee on Research involving Human Subjects in The Netherlands.

Interventions

The group CBT protocol was the Coping with Depression Course (Lewinsohn *et al.* 1992). It has been shown to be effective (Cuijpers, 1998; Allart-Van Dam *et al.* 2003, 2007; Haringsma *et al.* 2005) and consists of 10 weekly group sessions on psycho-education, cognitive restructuring, behaviour change, and relapse prevention. The internet-based CBT is an intervention of eight modules. It was studied as a self-help intervention, and no professional support was offered to the participants of this study. The intervention was found to be effective at post-treatment (Spek *et al.* 2007a).

Analyses

The target sample size of 300 participants was calculated to yield 78% power to detect a small effect (Cohen's $f=0.10$). The study was *a priori* powered to detect a small effect because we wanted to test whether there was a difference between the two interventions. The calculation was based on an ANOVA with an α of 0.05 (Cohen, 1988).

Preliminary analyses included checks for normality and the computation of descriptive statistics. All variables were distributed acceptably close to normal. ANOVAs, t tests and χ^2 tests were used to compare people who completed all questionnaires *versus* non-completers on baseline characteristics.

Analyses regarding the main hypotheses were performed according to the intention-to-treat approach. Missing data were imputed using the multiple imputation procedure of data augmentation in NORM (Schafer, 1999), as explained in our previous paper (Spek *et al.* 2007a).

The effects of the interventions were tested by means of contrasts. We calculated improvement effect sizes (d_{impr}) by dividing the absolute difference between the post-treatment average score ($\text{mean}_{\text{post}}$) and the pretreatment average score (mean_{pre}) by the pretreatment standard deviation (s.d._{pre}). An effect size of 0.5 thus indicates that the post-treatment average score is half a standard deviation (of the prescore) larger than the pretreatment average score. For between-group effect sizes, we calculated effect sizes by subtracting the effect size of the experimental group from the effect size of the control group.

We did not conduct the CIDI at the end of the trial. However, we wanted to estimate in how many participants a major depression might have been averted. For this, we used a cut-off score of 13 on the BDI. Scores above 13 can be interpreted as referring to subthreshold depression (see Van der Does, 2002). For our estimation of averted (subthreshold) depressions, we calculated numbers needed to treat.

Results

Follow-up measures at 12 months were completed by 58 of 102 participants (57%) in the internet group, 66 of 99 participants (67%) in the group course condition and 66 of 100 participants (66%) on the waiting-list (see Fig. 1). The number of participants who provided follow-up data was larger than the number of participants who provided post-treatment data because at 1-year follow-up there was a larger time window in which questionnaires could be returned. Intention-to-treat analyses were performed on imputed data of all 301 participants.

Table 1. Means (standard deviations) for depressive symptoms according to the Beck Depression Inventory (BDI)

	Pretreatment	Follow-up
Internet-based intervention ($n=102$)	19.07 (7.04)	10.45 (8.05)
Group intervention ($n=99$)	17.99 (9.39)	12.14 (8.76)
Waiting-list ($n=100$)	18.31 (7.88)	12.88 (10.10)

At post-treatment, we found that a mean of 98.3% of treatment sessions was completed in the group course and a mean of 78.1% of the modules was completed in the internet-based intervention. Group treatment was completed by 94.5% of the participants; the internet course was completed by 48.3% of the participants. We did not find a dose-effect relationship in the internet-based treatment condition. It is unclear if internet-based treatment participants repeated or continued treatment after the post-treatment assessment. They still had access to the intervention. However, participants in group treatment received a treatment manual that they could continue to use after the end of treatment.

Participants who did not complete follow-up measures did not differ from participants who did complete follow-up measures regarding age, gender, having a partner, educational level, employment status, assigned condition, EDS scores at screening, and BDI baseline scores (Spek *et al.* 2007a, Table 1).

For the intention-to-treat analyses, we fitted contrasts to the imputed data to test the hypotheses about the differences between conditions. The first contrast was designed to test whether internet-based treatment differed from the waiting-list control group. The second contrast was designed to test whether the internet-based treatment and the group CBT were different from each other. The first fitted contrast showed a significant difference between the internet-based treatment and the waiting list condition ($p=0.03$). We found no difference in effects of internet-based CBT and group CBT ($p=0.08$). For means and standard deviations of all conditions, see Table 1.

For improvement within the waiting-list control group, we found a large improvement effect size of 0.69. The group CBT condition also had a large improvement effect size (0.62), while an even larger improvement effect size of 1.22 was found within the internet-based treatment condition. When comparing the group treatment with the waiting-list group, we found an effect size of -0.07 . For internet-based treatment compared with the waiting-list, we found an effect size of 0.53.

Using the cut-off score of 13 on the BDI, we found that 62% of participants in the internet-based treatment scored below 13 at follow-up, which was 45% for group treatment, and 38% for the control condition. For the internet-based treatment, we found a risk difference of $0.62 - 0.38 = 0.26$. If we use this risk difference to calculate the number needed to treat, we find that $1/0.26 = 3.85$ people have to be treated for one averted case of (subthreshold) depression. For group treatment, the risk difference is $0.62 - 0.55 = 0.07$. Calculating the number needed to treat, we find that $1/0.07 = 14.29$ people have to be treated for one averted case of (subthreshold) depression.

Finally, analysis of only complete cases (instead of imputation techniques) showed almost the same results as analyses on imputed data: internet-based treatment differed significantly from the waiting-list condition ($p = 0.04$) but did not differ significantly from group treatment ($p = 0.13$).

Discussion

In this study, 1 year after the start of treatment, internet-based CBT was significantly more effective than a waiting-list condition in people over 50 years of age with subthreshold depression. There was a non-significant trend towards better outcomes for the internet intervention compared to the group treatment.

One year after the start of treatment, we found a moderate effect size of 0.53 for the internet-based CBT compared to the waiting-list condition. The improvement in effect size of 1.22 we found for internet-based CBT roughly corresponds with a 6-month improvement effect size of 1.03 found in the study by Andersson *et al.* (2005).

For group CBT we found an improvement effect size of 0.62, which does not correspond with the 1-year improvement effect size of 0.78 of the same intervention found in an earlier study (Allart-Van Dam *et al.* 2007). This might be because our participants preferred to be treated with an internet-based intervention. As only people with access to the internet were eligible for inclusion, many participants expected to participate in the internet-based treatment; some of those who were randomized to the group intervention were disappointed not to be treated by the internet-based treatment.

In this study we were faced with a large amount of missing data: 37% of our participants did not provide follow-up data. This is a common problem even in short-term trials on internet-based treatment for symptoms of depression, as was shown in a recent meta-analysis (Spek *et al.* 2007b). We dealt with missing values through the application of data augmentation; more information about this procedure

can be found in Spek *et al.* (2007a). This technique is regarded as the most conscientious and conservative way to handle missing data.

Apart from the above-mentioned drop-out, this study has several other limitations. Participants could only be included in the study if they had computer skills and access to internet. The participants in this study were more highly educated than the general population in this age group (Statistics Netherlands, 2005). Therefore, it is uncertain whether the results of this study can be generalized to the general population. In addition, as our participants suffered from subthreshold depression, we cannot draw any conclusions about the long-term effects of internet-based CBT on major depressive episodes.

Despite these limitations, our findings suggest that people over 50 years old with subthreshold depression can continue to benefit from internet-based CBT 1 year after the start of treatment. This is an important finding, especially when looking at the current problems we are faced with regarding the treatment of depression. A large number of people suffer from depression and subthreshold depression. With the current capacity and treatments, only 16% of the burden of depression is averted (Andrews *et al.* 2004). If, within our current capacity to offer treatment, we only provided evidence-based treatments, we would be able to avert 23% of the burden (Andrews *et al.* 2004). Even if we could treat everyone who is suffering from (subthreshold) depression, we could still only avert 34% of the burden of depression (Andrews *et al.* 2004). Internet-based treatment for subthreshold depression has three important assets regarding these problems: (1) internet-based treatment is an evidence-based treatment, even 1 year after the start of treatment, (2) internet-based treatment can be used by any number of people, thereby helping us to reach as many people as possible, and (3) it is a treatment that can prevent the occurrence of new depressive episodes.

Acknowledgements

This study was supported by a grant from ZON-MW, The Netherlands Organization for Health Research and Development.

Declaration of Interest

None.

References

- Allart-Van Dam E, Hosman CMH, Hoogduin CAL, Schaap CPDR (2003). The Coping with Depression course: short-term outcomes and mediating effects of a

- randomized controlled trial in the treatment of subclinical depression. *Behavior Therapy* **34**, 381–396.
- Allart-Van Dam E, Hosman CMH, Hoogduin CAL, Schaap CPDR** (2007). Prevention of depression in subclinically depressed adults: follow-up effects on the 'Coping with Depression' course. *Journal of Affective Disorders* **97**, 219–228.
- Andersson G, Bergström J, Holländare F, Carlbring P, Kaldo V, Ekselius L** (2005). Internet-based self-help for depression: randomised controlled trial. *British Journal of Psychiatry* **187**, 456–461.
- Andrews G, Issakidis C, Sanderson K, Corry J, Lapsley H** (2004). Utilising survey data to inform public policy: comparison of the cost-effectiveness of treatment for ten mental disorders. *British Journal of Psychiatry* **184**, 526–533.
- Beck AT, Steer RA, Brown GK** (1996). *Beck Depression Inventory Manual*, 2nd edn. Psychological Corporation: San Antonio, TX.
- Christensen H, Griffiths KM, Jorm AF** (2004). Delivering interventions for depression by using the Internet: randomised controlled trial. *British Medical Journal* **328**, 265–267.
- Clarke G, Eubanks D, Reid E, Kelleher C, O'Connor E, DeBarr L, Lynch F, Nunley S, Gullion C** (2005). Overcoming depression on the internet (ODIN) (2): a randomised trial of a self-help depression skills program with reminders. *Journal of Medical Internet Research* **7**, e16.
- Cohen J** (1988). *Statistical Power Analysis for the Behavioral Sciences*, 2nd edn. Lawrence Erlbaum Associates: Hillsdale, NJ.
- Cox JL, Chapman G, Murray D, Jones P** (1996). Validation of the Edinburgh Postnatal Depression Scale (EPDS) in non-postnatal women. *Journal of Affective Disorders* **39**, 185–189.
- Cox JL, Holden JM, Sagovsky R** (1987). Detection of postnatal depression: Development of the 10-item Edinburgh Postnatal Depression Scale. *British Journal of Psychiatry* **150**, 782–786.
- Cuijpers P** (1998). A psychoeducational approach to the treatment of depression: a meta-analysis of Lewinsohn's 'Coping with Depression' course. *Behavior Therapy* **29**, 521–533.
- Haringsma R, Engels GI, Cuijpers P, Spinhoven P** (2005). Effectiveness of the Coping With Depression (CWD) course for older adults provided by the community-based mental health care system in the Netherlands: a randomized controlled trial. *International Psychogeriatrics* **17**, 1–19.
- Lewinsohn PM, Munoz RF, Youngren MA, Zeiss AM** (1992). *Control Your Depression*. Fireside: New York, NY.
- Murray L, Carothers AD** (1990). The validation of the Edinburgh Post-natal Depression Scale on a community sample. *British Journal of Psychiatry* **157**, 288–290.
- Schafer JL** (1999). NORM: multiple imputation of incomplete multivariate data under a normal model. Software for S-PLUS 4.0 for Windows (www.stat.psu.edu/~jls/misoftwa.html).
- Spek V, Cuijpers I, Nyklíček I, Riper H, Keyzer J, Pop V** (2007b). Internet-based cognitive behaviour therapy for symptoms of depression and anxiety: a meta-analysis. *Psychological Medicine* **37**, 319–328.
- Spek V, Nyklíček I, Smits N, Cuijpers P, Riper H, Keyzer J, Pop, V** (2007a). Internet-based cognitive behavioral therapy for subthreshold depression in people over 50 years old: a randomized controlled trial. *Psychological Medicine* **37**, 1797–1806.
- Statistics Netherlands (CBS)** (2005). Education level in Dutch population increasing (www.cbs.nl/en-GB/menu/themas/onderwijs/publicaties/artikelen/archief/2005/2005-1745-wm.htm). Accessed 21 December 2007.
- Van der Does AJW** (2002). *Manual of the Dutch Version of the Beck Depression Inventory (BDI-II-NL)*. Ipskamp: Enschede, The Netherlands.
- WHO** (1997). *Composite International Diagnostic Interview*, version 2.1. World Health Organization: Geneva.