

## Tilburg University

### Consistent estimation of regression models with incompletely observed exogenous variables

Nijman, T.E.; Palm, F.C.

*Publication date:*  
1987

[Link to publication in Tilburg University Research Portal](#)

*Citation for published version (APA):*

Nijman, T. E., & Palm, F. C. (1987). *Consistent estimation of regression models with incompletely observed exogenous variables*. (Research memorandum / Tilburg University, Department of Economics; Vol. FEW 272). Unknown Publisher.

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.


CBM  
R

UNIVERSITEIT  
BRABANT

7626  
1987  
272


12

POSTBOX 90153  
5000 LE TILBURG  
THE NETHERLANDS



DEPARTMENT OF ECONOMICS  
RESEARCH MEMORANDUM

~~MA~~  
~~7626~~  
~~1987~~  
~~272~~



K.U.B.  
DISLOOFMEK  
TILBURG

CONSISTENT ESTIMATION OF REGRESSION  
MODELS WITH INCOMPLETELY OBSERVED  
EXOGENOUS VARIABLES

Th.E. Nijman and F.C. Palm

FEW 272

330.115

CONSISTENT ESTIMATION OF REGRESSION MODELS WITH INCOMPLETELY  
OBSERVED EXOGENOUS VARIABLES.

Th.E. Nijman \* and F.C. Palm \*\*

June, 1987

ABSTRACT

We consider consistent estimation of regression models in which the exogenous variables are incompletely observed assuming that the response mechanism is ignorable. Consistent estimates can be obtained from complete observations only. If the unobserved variables are related to observed variables through an auxiliary regression model, more efficient estimators of the parameters of interest can be obtained by using all available sample information. In the literature on imputed data and on proxy variables estimators several estimators have been proposed which are based on approximations for the missing data. We discuss conditions under which these proxy variables estimators are asymptotically more efficient than the estimator based on complete observations only and show how an optimal proxy variables estimator for which these conditions are always satisfied can be obtained. Moreover for a simple case, we derive the relative efficiency of several proxy variables estimators compared with the Gaussian maximum likelihood (ML) estimator. Finally extensions of the general results to cases where only aggregates of the exogenous variables are observed and to dynamic models are considered. Again relative efficiencies compared to ML are presented for simple examples. The findings indicate that by using the

\* Tilburg University, Department of Econometrics, P.O. Box 90153,  
5000 LE Tilburg, The Netherlands.

\*\* University of Limburg, Department of Economics, P.O. Box 616,  
6200 MD Maastricht, The Netherlands.

Information provided by the auxiliary model for the regressors, it is possible to design proxy variables estimators that are almost as efficient as the ML estimator which in the presence of missing observations is often computationally unattractive. When the normality assumption does not hold, the pseudolikelihood estimator can even become less efficient than some proxy variables estimators.

## 1. INTRODUCTION

In applied research, it is common practice to impute the missing values of variables which are incompletely observed. Imputation is applied to cross section data which suffer from partial non-response (for a survey of the literature on the analysis of models in the context of non-response in sample surveys, see e.g. Little (1982)) and to time series which are available on a high temporal aggregation level only. In a common imputation procedure, the observations on the incompletely observed variable are regressed on auxiliary variables. The missing values are then approximated by the predictions from this auxiliary regression equation. Nevertheless little attention seems to have been paid to the implications of using proxies in a subsequent statistical analysis. In this paper, we are concerned with the efficiency of consistent estimators based on an imputed data set and on the set of complete observations respectively. It is shown that a regression using imputed observations does not necessarily yield more efficient parameter estimates than a regression based on data points for which all variables are observed (in the sequel called complete observations). We discuss conditions under which an estimator based on approximations for unobserved variables is asymptotically more efficient than an estimator based on complete observations only and we show how an optimal proxy variables estimator can be obtained. We also consider the estimation of standard errors of proxy variables estimators. For a simple case, we derive the relative efficiency of several proxy variables estimators compared with the maximum likelihood estimator under the normality assumption. Finally extensions of the results to cases where only aggregates of the exogenous variables are observed and to dynamic

models are considered.

The first model we consider is

$$y_i = \sum_{k=1}^K \beta_k x_{ik} + \epsilon_i, \quad i = 1, \dots, N, \quad (1)$$

$$x_{ik} = \sum_{l=1}^L \alpha_{lk} z_{il} + v_{ik}, \quad \begin{array}{l} i = 1, \dots, N, \\ k = 1, \dots, K, \end{array} \quad (2)$$

where the regression disturbances  $\epsilon_i$  and  $v_{ik}$  are i.i.d. with mean zero and variances  $\sigma^2$  and  $\sigma_{kk}$  respectively, have finite fourth moments, are independent of the corresponding regressors and satisfy

$$E\epsilon_i v_{ik} = 0,$$

$$E v_{ik} v_{il} = \sigma_{lk}, \text{ for } i = 1, \dots, N, \text{ and } l, k = 1, \dots, K, l \neq k.$$

Assume moreover that  $\text{plim } Z'Z/N$  is finite and non-singular where the matrix  $Z$  has typical element  $z_{il}$ . We consider the case where  $y_i$  and  $z_{il}$ , ( $l = 1, \dots, L$ ), are observed for  $i = 1, \dots, N$ , whereas  $x_{ik}$  is observed if and only if the random variable  $\delta_{ik}$  takes the value 1. The random variables  $\delta_{ik}$  are assumed to be independent of  $\epsilon_j$ ,  $z_{jl}$  and  $v_{jl}$ . Note that we do not exclude that some of the regressors in (1) are completely observed and are used as regressors in (2).

If one assumes that a fraction of the observations is complete in large samples, a first consistent estimator of  $\beta' = [\beta_1, \dots, \beta_K]$  can be obtained from the regression (1) using complete observations only. Evidently if only a few of the right hand side variables in (1) are complete and these variables can be closely approximated using equation (2) an estimator based on complete observations only will not be very efficient. Alternatively the missing values can be approximated by

$$\hat{x}_{ik} = \sum_{l=1}^L z_{il} \hat{\alpha}_{lk},$$

where  $\hat{\alpha}_{lk}$  is an estimate of  $\alpha_{lk}$ . If one defines

$$\hat{x}_{ik} = x_{ik} \text{ if } \delta_{ik} = 1,$$

an estimate of  $\beta$  can subsequently be obtained by regressing  $y_i$  on  $\hat{x}_{ik}$ ,  $k = 1, \dots, K$ . This procedure is known as the first order method of Afifi and Elashoff (1966). Nijman and Palm (1986) refer to it as a proxy variables estimator. Special cases have been considered by e.g. Gouriéroux and Monfort (1981) who derived the large sample distribution of several proxy variables estimators and by Conniffe (1983a) who considered small sample properties.

The plan of this paper is as follows. In section 2 we analyze the model in (1) and (2) assuming that  $K = L = 1$ , and that  $x_{i1}$  is observed if  $i \leq N/2$  only. This special case illustrates very well the main issues related to proxy variable estimators. Numerical results on the relative efficiency of these estimators compared to the Gaussian ML estimator are presented for this model. In section 3 we consider the general case and show how the use of proxies can lead to an efficiency gain over the estimator based on complete observation only. In section 4 the analysis of proxy variables estimators is extended to observations of temporal aggregates of the exogenous variables and to dynamic models. Again numerical results on the relative efficiency of a number of estimators are presented for a simple model. Finally some concluding remarks are given in section 5. Three appendices contain the technical details.

## 2. AN EXAMPLE

In this section we analyze the model in (1) and (2) assuming that  $K = L = 1$  and that the exogenous variable in (1) is observed if  $i \leq N/2$  only. Deleting redundant subscripts, the model can be written as



$$y_i = \beta x_i + \epsilon_i \quad (3)$$

$$x_i = \alpha z_i + v_i. \quad (4)$$

The variance of  $v_i$  will be denoted by  $\sigma_v^2$ . It is useful to notice that the model (3)-(4) is a restricted version of a model analyzed by Gouriéroux and Monfort (1981) who assume that  $z_i$  is also included in the regression equation for  $y$ ,

$$y_i = \beta_1 x_i + \beta_2 z_i + \epsilon_i. \quad (3')$$

If normality of  $\epsilon_i$  and  $v_i$  is assumed, asymptotically efficient ML estimators of the parameter in (3) and (4) can be obtained by maximizing the likelihood function

$$L(\alpha, \beta, \sigma^2, \sigma_v^2) = \prod_{i=1}^N L_i(\alpha, \beta, \sigma^2, \sigma_v^2)$$

with

$$L_i(\alpha, \beta, \sigma^2, \sigma_v^2) = C \cdot (\sigma_v \tilde{\sigma})^{-1} \exp\left\{-\delta_i (y_i - \beta x_i)^2 / 2\sigma^2 - \delta_i (x_i - \alpha z_i)^2 / 2\sigma_v^2 - (1 - \delta_i)(y_i - \alpha \beta z_i)^2 / 2\tilde{\sigma}^2\right\},$$

where  $\tilde{\sigma}^2 = \sigma^2 + \beta^2 \sigma_v^2$ ,  $\delta_i = 1$  if  $x_i$  is observed ( $i \leq N/2$ ) and  $\delta_i = 0$  otherwise and  $C$  is a constant independent of the unknown parameters. Note that a computationally convenient reparametrization proposed by Gouriéroux and Monfort (1981) for the model (3') - (4), no longer applies when  $\beta_2$  is known to be zero. Following an approach similar to that of Anderson (1957), Gouriéroux and Monfort reparametrize the joint distribution for  $y_i$  and  $x_i$  given  $z_i$  as a product of the marginal distribution of  $y_i$  given  $z_i$  and the conditional distribution of  $x_i$  given  $y_i$  and  $z_i$  and they show that his reparametrization provides an immediate solution for the ML estimator. When  $\beta_2 = 0$ , the computational advantage of this approach is lost. If the normality assumption is satisfied, the ML estimator will be asymptotically efficient but in general ML estimation will be computationally cumbersome for other than simple models. If the normality does not hold, the Gaussian ML estimator is still consistent but no

longer efficient, a point to which we will return below.

Alternatively, the parameter  $\beta$  can be consistently estimated by OLS using the complete observations only

$$\hat{\beta}_C = \Sigma_C x_i y_i / \Sigma_C x_i^2, \quad (5)$$

where  $\Sigma_C$  denotes summation over complete observations ( $i \leq N/2$ ). In the sequel we will also use the notation  $\Sigma_I$  and  $\Sigma_A$  to denote summation over incomplete and all, complete and incomplete, observations respectively. Intuitively there seems to be a case for using imputed data and considering the following proxy variables estimator

$$\hat{\beta}_P = \Sigma_A \hat{x}_i y_i / \Sigma_A \hat{x}_i^2, \quad (6)$$

where  $\hat{x}_i = x_i$  if  $i \leq N/2$  and  $\hat{x}_i$  is some approximation for  $x_i$  if  $i > N/2$ . As mentioned in the introduction, a natural choice for the approximation is

$$\hat{x}_i = \hat{\alpha} z_i \text{ if } i \geq N/2, \text{ where } \hat{\alpha} = \Sigma_C z_i x_i / \Sigma_C x_i^2.$$

The condition for consistency of the resulting estimator  $\hat{\beta}_P$  is that

$$\text{plim } \Sigma_A \hat{x}_i w_i / \Sigma_A \hat{x}_i^2 = 0 \text{ with } w_i = y_i - \hat{x}_i \beta = \epsilon_i + \beta(x_i - \hat{x}_i). \quad (7)$$

In applied work it is not only important to have a consistent estimator, but also to be able to estimate its large sample variance consistently.

Substituting (7) into (6), we have for the first order method

$$\hat{\beta}_P - \beta = (\Sigma_C x_i^2 + \Sigma_I \hat{\alpha}^2 z_i^2)^{-1} \{ \Sigma_C x_i \epsilon_i + \hat{\alpha} \Sigma_I z_i \epsilon_i + \beta \hat{\alpha} \Sigma_I z_i v_i + \beta \hat{\alpha} \Sigma_I z_i^2 (\alpha - \hat{\alpha}) \}.$$

The large sample variance (avar) of  $\sqrt{N} \hat{\beta}_P$  can be derived via substitution of (7) into (6) and the use of the appropriate limiting theory

$$\text{avar}(\sqrt{N} \hat{\beta}_p) = (\sigma_{\hat{x}}' \sigma^2 + \alpha' \beta' \sigma_z^2 \sigma_v^2) \sigma_{\hat{x}}^{-4} \quad (8)$$

with  $\sigma_{\hat{x}}^2 = \alpha^2 \sigma_z^2 + \frac{1}{2} \sigma_v^2$ . Three remarks have to be made.

First, although the distinction between the case where  $\alpha$  is known and that when  $\alpha$  is estimated could be neglected in proving consistency, it is essential for the computation of the large sample variance of  $\hat{\beta}_p$ . When  $\alpha$  is known, the asymptotic variance of  $\tilde{\beta}_p$  ( $\sim$  denotes that the true value of  $\alpha$  is used) is given by

$$\text{Avar}(\sqrt{N} \tilde{\beta}_p) = (\sigma_{\hat{x}}^2 \sigma^2 + \frac{1}{2} \alpha^2 \beta^2 \sigma_z^2 \sigma_v^2) \sigma_{\hat{x}}^{-4}. \quad (9)$$

This point is often missed in the literature, but has recently been stressed in the context of using approximations for unobserved expectations by Pagan (1984) and by Murphy and Topel (1985). Second, as is obvious from a comparison of their asymptotic variances in appendix A,  $\hat{\beta}_p$  can be more efficient as well as less efficient than  $\hat{\beta}_c$ , a finding which also holds for the unrestricted model considered by Gouriéroux and Monfort (1981). (In remark 2 on p. 583 they incorrectly state that  $\hat{\beta}_p$  is more efficient than  $\hat{\beta}_c$  as noted by Griliches (1986)). Third, the formula for the standard errors in a least squares regression does not yield a consistent estimator of the asymptotic standard errors for  $\hat{\beta}_p$  and  $\hat{\beta}_c$  as

$$\text{plim} \frac{\sum_A (y_i - \hat{\beta}_p \hat{x}_i)^2}{(\sum_A \hat{x}_i^2)^{-1}} = (\sigma^2 + \frac{1}{2} \beta^2 \sigma_z^2 \sigma_v^2) \sigma_{\hat{x}}^{-2}. \quad (10)$$

A standard least squares regression of  $y_i$  on  $\hat{x}_i$  produces a consistent estimate of  $\beta$ . The resulting standard errors, however, will be incorrect. It is obvious from (8) and (10) that the order of magnitude and the sign of the bias depend on the value of  $\alpha^2 \sigma_z^2$ . Some information on the order of magnitude of the bias will be provided in table 1.

When using a proxy variable  $\hat{x}_t$  for the missing values of  $x_t$ , the disturbance  $w_t = \epsilon_t + \beta(x_t - \hat{x}_t)$  is no longer homoscedastic. It is natural therefore to consider generalized least squares (GLS) estimation, as Gouriéroux and Monfort (1981) did for an exactly identified model, which can be denoted as

$$\hat{\beta}_V = (\hat{X}'V^{-1}\hat{X})^{-1}\hat{X}'V^{-1}y, \quad (11)$$

where  $y = (y_1, y_2, \dots, y_N)'$ ,  $X = (\hat{x}_1, \hat{x}_2, \dots, \hat{x}_N)'$  and  $V$  is a weighting matrix.

Dagenais (1973) proposed to take  $V$  diagonal with  $v_{ij} = \hat{\sigma}^2$  for  $i \leq N/2$  and

$v_{ij} = \hat{\sigma}^2 + \hat{\beta}^2 \hat{\sigma}_V^2$ ,  $i > N/2$ , and " $\hat{\cdot}$ " indicating a consistent estimate of the corresponding parameter. This estimator will be referred too as  $\hat{\beta}_D$ . Although

every element of the matrix  $V$  proposed by Dagenais converges in probability to the corresponding element of the covariance matrix of  $w_t$ ,  $\Omega$ , the matrix  $N^{-1}\hat{X}'V^{-1}\hat{X}$  does not converge to the same limit as  $N^{-1}\hat{X}'\Omega^{-1}\hat{X}$ .

That the choice of the weights by Dagenais (1973) is not optimal has been pointed out by Conniffe (1983b) who proposed another weighting matrix with constant elements. That these weights are not optimal either can be seen by comparing them with the elements of  $\Omega$ . Assuming for the ease of simplicity that  $z_t$  is nonstochastic and writing

$$\begin{aligned} w_t &= \epsilon_t + \beta(x_t - \alpha z_t) + \beta z_t(\alpha - \hat{\alpha}) & \text{if } t > N/2 \\ &= \epsilon_t & \text{if } t \leq N/2. \end{aligned} \quad (12)$$

we have

$$\begin{aligned} w_{ij} &= \sigma^2 + \beta^2 \sigma_V^2 + \beta^2 z_i z_j (\sum_C z_i^2)^{-1} \sigma_V^2 & i = j & \quad i > N/2 \\ &= \beta^2 z_i z_j (\sum_C z_i^2)^{-1} \sigma_V^2 & i \neq j & \quad i, j > N/2 \\ &= \sigma^2 & i = j & \quad i \leq N/2 \\ &= 0 & \text{otherwise.} & \end{aligned} \quad (13)$$

A feasible GLS estimator  $\hat{\beta}_g$  is obtained if we substitute consistent estimates for the unknown parameters in (3) and (4) and use  $\hat{\Omega}$  instead of  $V$  in (12). The inversion of  $\hat{\Omega}$  does not seem to be very attractive at first sight as it contains non-zero off-diagonal elements. However, we can avoid direct inversion of such a matrix by using the binomial inversion theorem that will be needed in more complex cases as well. First we write

$$\hat{\Omega} = G + ZHZ', \quad (14)$$

where  $G$  is a diagonal with  $\hat{\sigma}^2$  and  $\hat{\sigma}^2 + \hat{\beta}^2 \hat{\sigma}_v^2$  in position  $i$  of the main diagonal for  $i \leq N/2$  and  $i > N/2$  respectively,  $H = \hat{\sigma}_v^2 \hat{\beta}^2 (\sum_C z_i^2)^{-1}$  is a scalar,  $Z$  is a  $N \times 1$  vector with  $i$ -th element being equal to zero and  $z_i$ , for  $i \leq N/2$  and  $i > N/2$  respectively. The inverse of  $\hat{\Omega}$  can be obtained straightforwardly as  $(G + ZHZ')^{-1} = G^{-1} - G^{-1}Z(H^{-1} + Z'G^{-1}Z)^{-1}Z'G^{-1}$ . (15)

The asymptotic variance of  $\hat{\beta}_g$  can be consistently estimated by  $(\hat{X}'\hat{\Omega}^{-1}\hat{X})^{-1}$ . The estimator  $\hat{\beta}_g$  is more efficient than the first order method in (6), the Dagenais (1973) estimator and the Conniffe (1983b) estimator.

The asymptotic variance of the estimators considered in this section will be given in appendix A. In table 1 we report the ratio of the variance of alternative consistent estimators compared with the variance of the ML estimator assuming normality of  $\epsilon_i$  and  $v_i$ . From the results in appendix A, it follows that the relative efficiency only depends on  $R_x^2 = \alpha^2 \sigma_z^2 \sigma_x^{-2}$  and  $R_y^2 = \beta^2 \sigma_x^2 (\beta^2 \sigma_x^2 + \sigma^2)^{-1}$ , where  $\sigma_x^2 = \alpha^2 \sigma_z^2 + \sigma_v^2$ . From the results in table 1, it appears that the OLS estimator using the complete observations only,  $\hat{\beta}_c$ , is roughly as efficient as the proxy variables estimators  $\hat{\beta}_d$  and  $\hat{\beta}_g$ , when  $R_x^2$  is small. When  $R_x^2$  is large,  $\hat{\beta}_c$  is inferior to all proxy variables estimators considered. This finding is plausible. When a large fraction of the variance of  $x_i$  is explained by  $z_i$ ,  $\alpha z_i$  is a fairly accurate approximation of  $x_i$  and it pays to use this information. However, when  $R_x^2$  is low compared to  $R_y^2$ ,  $\hat{\beta}_p$  can well be less efficient than  $\hat{\beta}_c$ .

Table 1 : Relative efficiency of the ML estimator compared with alternative consistent estimators for  $\beta$ .

$R_x^2$ (1)	$R_y^2$ (2)	Alternative estimators				$\frac{\hat{\sigma}_w^2}{\text{var}(\hat{\beta}_p) \Sigma \hat{x}_t^2}$
		$\hat{\beta}_p$ (6)	$\hat{\beta}_d$ (11) V:Dagenais	$\hat{\beta}_g$ (11) V:optimal	$\hat{\beta}_c$ (5)	
0.20	0.20	1.1306	1.1161	1.1129	1.2719	1.0312
0.20	0.40	1.3068	1.2251	1.2140	1.3315	1.0755
0.20	0.60	1.5430	1.2675	1.2491	1.3226	1.1429
0.20	0.80	2.0741	1.1894	1.1726	1.2043	1.2581
0.20	0.95	5.3364	1.0546	1.0489	1.0556	1.4176
0.40	0.20	1.0737	1.0617	1.0587	1.3845	0.9901
0.40	0.40	1.2076	1.1382	1.1259	1.3761	0.9767
0.40	0.60	1.4517	1.2000	1.1744	1.3421	0.9575
0.40	0.80	2.0770	1.1786	1.1471	1.2262	0.9277
0.40	0.95	5.7154	1.0612	1.0473	1.0649	0.8916
0.60	0.20	1.0320	1.0258	1.0240	1.5360	0.9767
0.60	0.40	1.1033	1.0658	1.0573	1.4710	0.9444
0.60	0.60	1.2644	1.1184	1.0962	1.3952	0.8966
0.60	0.80	1.7420	1.1481	1.1085	1.2669	0.8182
0.60	0.95	4.5344	1.0706	1.0442	1.0828	0.7164
0.80	0.20	1.0078	1.0060	1.0055	1.7368	0.9814
0.80	0.40	1.0283	1.0172	1.0142	1.6547	0.9536
0.80	0.60	1.0859	1.0393	1.0288	1.5432	0.9079
0.80	0.80	1.3055	1.0807	1.0501	1.3733	0.8182
0.80	0.95	2.7544	1.0805	1.0361	1.1325	0.6624
0.95	0.20	1.0005	1.0004	1.0003	1.9274	0.9941
0.95	0.40	1.0019	1.0011	1.0009	1.8922	0.9847
0.95	0.60	1.0068	1.0030	1.0019	1.8296	0.9668
0.95	0.80	1.0335	1.0107	1.0048	1.6866	0.9206
0.95	0.95	1.3274	1.0481	1.0125	1.3442	0.7660

In column 7 the ratios of the variances computed using the OLS formula for standard errors (10) and the correct asymptotic variance for  $\hat{\beta}_p$  in (8) is presented. In a few occasions the asymptotic bias for the standard errors involved in using the OLS formula appears to be quite important.

In order to explain the results on the relative efficiency of the four estimators considered in columns 3 to 6 of table 1 we express the proxy variables estimators as a linear combination of  $\hat{\beta}_c$  and a consistent estimator of  $\beta$ ,  $\hat{\beta}_{mj}$ , based on incomplete observations only (except for the estimate  $\hat{a}$ ),

$$\hat{\beta}_j = \hat{\lambda}_j \hat{\beta}_c + (1-\hat{\lambda}_j) \hat{\beta}_{mj} \text{ with } j \in \{p, d, g\}.$$

The expressions for  $\hat{\beta}_{mj}$  and  $\hat{\lambda}_j$  are given below

j	$\hat{\beta}_{mj}$	$\hat{\lambda}_j$
p	$(\Sigma_I \hat{x}_i)^{-2} \Sigma_I \hat{x}_i y_i$	$(\Sigma_C x_i + \Sigma_I \hat{x}_i)^{-2} \Sigma_C x_i$
d	$(\Sigma_I \hat{x}_i)^{-2} \Sigma_I \hat{x}_i y_i$	$(\hat{\sigma}^{-2} \Sigma_C x_i + \hat{\sigma}^{-2} \Sigma_I \hat{x}_i)^{-1} \hat{\sigma}^{-2} \Sigma_C x_i$
g	$(\Sigma_I w_{i1} \hat{x}_i \hat{x}_i)^{-1} \Sigma_I w_{i1} \hat{x}_i y_i$	$(\hat{\sigma}^{-2} \Sigma_C x_i + \Sigma_I w_{i1} x_i x_i)^{-1} \hat{\sigma}^{-2} \Sigma_C x_i$

where  $w_{ij}^*$  denotes the  $(i, i')$ -th element of  $\hat{\Omega}^{-1}$ . The large sample variance of  $\hat{\beta}_j$  is

$$\text{Avar}(\sqrt{N} \hat{\beta}_j) = \lambda_j^2 v_c + (1-\lambda_j)^2 v_{mj} \tag{16}$$

with  $\lambda_j = \text{plim} \hat{\lambda}_j$ ,  $v_c = \text{Avar}(\sqrt{N} \hat{\beta}_c)$  and  $v_{mj} = \text{Avar}(\sqrt{N} \hat{\beta}_{mj})$ .

It is straightforward to verify that  $\hat{\beta}_j$  is asymptotically more efficient than  $\hat{\beta}_c$  if

$$\frac{v_{mj} - v_c}{v_{mj} + v_c} < \lambda_j < 1 \quad (17)$$

and that the choice of  $\lambda_j$  which minimizes the asymptotic variance of  $\sqrt{N} \hat{\beta}_j$  is

$$\lambda_j^{opt} = v_{mj} (v_c + v_{mj})^{-1} \text{ which satisfies (17).}$$

As  $\lambda_g = \lambda_g^{opt}$ ,  $\hat{\beta}_g$  is efficient relative to  $\hat{\beta}_c$ . As  $\lambda_p \neq \lambda_p^{opt}$  and  $\lambda_d \neq \lambda_d^{opt}$ , the estimators  $\hat{\beta}_p$  and  $\hat{\beta}_d$  are more efficient than  $\hat{\beta}_c$  only if inequality (17) is satisfied. This will not be the case if  $\beta$  (or  $R_y^2$ ) is sufficiently large, as the lower bound in (17) tends to 1 if  $\beta$  increases, while  $\lambda_p$  and  $\lambda_d$  are not affected by a change of  $\beta$ . In this case, due to suboptimal weighting, the additional information on  $\beta$  contained in  $\hat{\beta}_{mj}$  leads to an efficiency loss of  $\hat{\beta}_j$  compared with  $\hat{\beta}_c$ . If on the contrary  $R_x^2$  is large so that  $\alpha z_t$  tends to be a better proxy,  $v_{mj}$  gets close to  $v_c$  and the proxy variables estimators  $\hat{\beta}_p$  and  $\hat{\beta}_d$  become more efficient than  $\hat{\beta}_c$  in large samples.

The efficiency of the ML estimator in table 1 arises from the assumption that the distributions of  $\epsilon_i$  and  $v_i$  are known to be normal. If normality is assumed but does not hold, the Gaussian ML estimator which maximizes  $L(\alpha, \beta, \sigma^2, \sigma_v^2)$  above will still be consistent (see e.g. Amemiya (1985), theorem 4.1.1) and the asymptotic distribution can be determined (see e.g. Amemiya (1985), theorem 4.1.3 and appendix A). This estimator is however not necessarily more efficient than the proxy variables estimators if  $\epsilon_i$  and  $v_i$  are not normal. In table 2 we present the relative efficiency of the Gaussian ML estimator compared with the optimal proxy variables estimator  $\hat{\beta}_g$ . The relative efficiency with respect to other proxy variables estimators can easily be derived from the results in tables 1 and 2. In table 2 we restrict ourselves to cases where

$$R_x^2 = R_y^2 \text{ and } (E\epsilon_i^4)/(E\epsilon_i^2)^2 = (Ev_i^4)/(Ev_i^2)^2.$$

Evidently the normality assumption does not have a very large effect on the relative efficiency unless only small fractions of the variances of  $y_i$  and  $x_i$  are explained by (3) and (4) and the true distributions of  $\epsilon_i$  and  $v_i$  have very fat tails. For the derivation of the results in table 2 we refer to appendix A.



Table 2 : Relative efficiency of the Gaussian ML estimator compared with the optimal proxy variables estimator  $\hat{\beta}_g$ .

$R_x^2, R_y^2$	Values of $E\epsilon_i^4 / (E\epsilon_i^2)^2 = (E\nu_i^4) / (E\nu_i^2)^2$				
	2	3	4	6	10
0.2	1.1623	1.1129	1.0675	0.9869	0.8575
0.4	1.1748	1.1259	1.0809	1.0011	0.8721
0.6	1.1305	1.0962	1.0638	1.0045	0.9036
0.8	1.0672	1.0501	1.0336	1.0019	0.9441
0.95	1.0167	1.0125	1.0083	1.0001	0.9841

### 3. THE GENERAL MODEL

In this section we consider the general model introduced in equations (1) and (2). As in the simple case considered in the previous section a consistent estimate of  $\beta$  can be computed from complete observations only. Define  $y_c$ ,  $X_c$  and  $Z_c$  as the vector and matrices obtained after deletion of rows of  $y$ ,  $X$  and  $Z$  respectively for which some variable is missing. The regression estimator based on complete observations only can be written as

$$\hat{\beta}_c = (X_c' X_c)^{-1} X_c' y_c. \quad (18)$$

In the model (1) and (2) several proxies can be thought of. A first possibility is to obtain estimates of the  $\alpha_{1k}$  from regressions using complete observations only

$$\hat{\alpha} = (Z_c' Z_c)^{-1} Z_c' X_c \quad (19)$$

and subsequently to approximate missing exogenous variables in (1) by

$$\hat{x}_{ik}^{(1)} = \sum_{l=1}^L z_{il} \hat{\alpha}_{lk} \text{ if } \delta_{ik} = 0.$$

If  $\hat{x}_{ik}^{(1)} = x_{ik}$  if  $\delta_{ik} = 1$  is defined for notational convenience,  $\beta$  can sub-

sequently be estimated from the regression model

$$\begin{bmatrix} y_c \\ y_I \end{bmatrix} = \begin{bmatrix} X_c \\ \hat{X}_I^{(1)} \end{bmatrix} \beta + \begin{bmatrix} \epsilon_c \\ w_I^{(1)} \end{bmatrix} \quad (20)$$

or

$$y = \hat{X}^{(1)}\beta + w^{(1)}, \quad (21)$$

where the subscript I in (20) refers to incomplete observations. From the example in the previous section we know that the ordinary least squares estimator

$$\hat{\beta}_p = \{X_c'X_c + \hat{X}_I^{(1)'}\hat{X}_I^{(1)}\}^{-1} \{X_c'y_c + \hat{X}_I^{(1)'}y_I\} \quad (22)$$

is not necessarily more efficient than  $\hat{\beta}_c$ . As in section 2 we have to analyze the structure of the covariance matrix of the disturbances in order to derive a generalized least squares estimator. Because

$$y_i - \sum_{k=1}^K \hat{x}_{ik}^{(1)} \beta_k = \epsilon_i + \sum_{k=1}^K \delta_{ik} \beta_k v_{ik} + \sum_{l=1}^L \sum_{k=1}^K \delta_{ik} \beta_k z_{il} (\hat{\alpha}_{lk} - \alpha_{lk}) \quad (23)$$

and  $\hat{\alpha}_{lk} - \alpha_{lk}$  is linear in the  $v_{ik}$  we have  $w^{(1)} = \epsilon + AV$  for a suitable chosen  $(N \times N)$  matrix A and the GLS estimator  $\hat{\beta}_g^{(1)}$  can straightforwardly be computed using (15). Moreover it is evident that  $\hat{\beta}_g^{(1)}$  will be more efficient than  $\hat{\beta}_c$  because  $\hat{\beta}_c$  coincides with the IV estimator of  $\beta$  from (20), with  $(Z_c', 0)$  being the matrix of instruments.

A natural question to ask next is how the efficiency of  $\hat{\beta}_g$  is affected if relevant regressors are excluded from the auxiliary regressions. Partition Z as  $Z = (Z_1, Z_2)$  where  $Z_1$  and  $Z_2$  are  $(N \times L_1)$  and  $(N \times (L - L_1))$  matrices respectively and assume that the regression model

$$x_{ik} = \sum_{l=1}^{L_1} \eta_{lk} z_{il} + v_{ik}^* \quad (24)$$

still satisfies the assumptions that were made with respect to (2). This model suggests the use of the proxies

$$\hat{x}_{1k}^{(2)} = \sum_{l=1}^{L_1} \hat{\eta}_{1k} z_{l1} \quad \text{if } \delta_{1k} = 0$$

$$= x_{1k} \quad \text{if } \delta_{1k} = 1$$

where  $\hat{\eta}_{1k}$  is the regression estimate from (24). Substitution of this proxy yields the model

$$y = \hat{X}^{(2)}\beta + w(2), \quad (25)$$

from which  $\beta$  can again be estimated e.g. by generalized least squares yielding  $\hat{\beta}_g^{(2)}$ . The following theorem will be useful in determining the effect of the choice of a proxy variable on the associated estimators  $\hat{\beta}_g^{(1)}$  and  $\hat{\beta}_g^{(2)}$ .

#### Theorem

Assume that  $y = X\beta + \epsilon$  holds with  $\text{plim } N^{-1}Z'\epsilon = 0$  and let  $\hat{X}$  and  $\tilde{X}$  be two proxies for  $X$ . Consider the estimators

$$\hat{\beta}_{GLS} = (\hat{X}'\hat{\Sigma}^{-1}\hat{X})^{-1}\hat{X}'\hat{\Sigma}^{-1}y \quad \text{and} \quad \tilde{\beta}_{IV} = (Z'\tilde{X})^{-1}Z'y.$$

Assume that

- (i)  $\sqrt{N}(\hat{\beta}_{GLS} - \beta) \stackrel{d}{=} N(0, V^{-1})$ , where  $V^{-1} = \text{plim } N(\hat{X}'\hat{\Sigma}^{-1}\hat{X})^{-1}$  is finite and positive definite;
- (ii)  $\text{plim } N^{-1}Z'\tilde{X} = Q$  is finite and positive definite;

$$(iii) \frac{1}{\sqrt{N}} \begin{bmatrix} Z'\hat{w} \\ Z'(\tilde{w} - \hat{w}) \end{bmatrix} \stackrel{d}{=} N(0, D) \quad \text{where } D = \text{plim } N^{-1} \begin{bmatrix} Z'\hat{\Sigma}Z & 0 \\ 0 & Z'SZ \end{bmatrix}$$

for some  $S$  and  $\hat{w}$  and  $\tilde{w}$  are the disturbances associated with  $\hat{X}$  and  $\tilde{X}$  respectively.

Then  $\hat{\beta}_{GLS}$  is asymptotically at least as efficient as  $\tilde{\beta}_{IV}$ .

Proof: see appendix B.

The third requirement is most crucial. If two proxies  $\hat{X}$  and  $\tilde{X}$  are available, an IV estimator based on  $\tilde{X}$  cannot be more efficient than a GLS estimator based on  $\hat{X}$  if  $Z'\hat{w}$  and  $Z'(\tilde{w}-\hat{w})$  are asymptotically orthogonal provided the regularity conditions of the theorem are met.

Returning to the analysis of the relative efficiency of

$\hat{\beta}_g^{(1)}$  and  $\hat{\beta}_g^{(2)}$  let us first consider the case where the parameters  $\alpha_{jk}$  and  $\eta_{jk}$  are known a priori. Then it is very simple to use the theorem to show that conditioning on the larger information set will yield more efficient estimates. Define  $\hat{X} = \hat{X}^{(1)}$  and  $\tilde{X} = \hat{X}^{(2)}$ . The disturbances in theorem are  $\hat{w} = \epsilon + (X - \hat{X})\beta$  and  $\tilde{w} = \epsilon + (X - \tilde{X})\beta$  respectively. As  $\hat{x}_{jk}^{(1)} = E[x_{jk} | I_1]$  and  $\hat{x}_{jk}^{(2)} = E[x_{jk} | I_2]$  with  $I_2 \subset I_1$ ,  $\hat{w} - \tilde{w} = (\hat{X} - \tilde{X})\beta$  will be orthogonal to  $\hat{w}$  by the properties of conditional expectations and the theorem immediately implies efficiency of the proxy variables estimator based on the larger conditioning set. Unfortunately this result does not hold true in general if  $\alpha_{jk}$  and  $\eta_{jk}$  have to be estimated. A counter-example in a slightly different model is presented in the next section. In appendix C we show that the result does hold for  $\hat{\beta}_g^{(1)}$  and  $\hat{\beta}_g^{(2)}$  if  $\delta_{jk}$  does not depend on  $k$  that is if all exogenous variables in (1) are missing when one of them is. We conjecture that more general results can be proved along the same lines. The relative efficiency of  $\hat{\beta}_g^{(1)}$  with respect to  $\hat{\beta}_g^{(2)}$  implies among other things that if a constant is included in (2), use of that auxiliary regression model will yield more efficient estimates of  $\beta$  than simple imputation of mean values for missing observations (which is equivalent to regression on a constant only) as is often done in practice.

The theorem above can also be used to demonstrate the effect of more efficient estimation of the  $\alpha_{jk}$  in (2). If prior restrictions on these parameters are available or if observations for which some but not all exogenous variables in (1) are observed are also used to estimate these parameters, Zellner's (1962)

SUR estimator will be more efficient than a regression on the set of complete observations only. If the SUR estimates are denoted by  $\hat{\alpha}_{1k}$  and  $\hat{x}_{1k}^{(3)}$  is defined as

$$\begin{aligned}\hat{x}_{1k}^{(3)} &= \sum_{l=1}^L \hat{\alpha}_{1k} z_{l1} & \text{if } \delta_{1k} = 1 \\ &= x_{1k} & \text{if } \delta_{1k} = 0,\end{aligned}$$

we obtain

$$\sum_{k=1}^K (\hat{x}_{1k}^{(1)} - \hat{x}_{1k}^{(3)}) \beta_k = \sum_{l=1}^L \sum_{k=1}^K \delta_{1k} \beta_k z_{l1} (\hat{\alpha}_{1k} - \hat{\alpha}_{1k}). \quad (26)$$

Using (23) with  $\hat{x}_{1k}^{(1)}$  and  $\hat{\alpha}_{1k}$  replaced by  $\hat{x}_{1k}^{(3)}$  and  $\hat{\alpha}_{1k}$

respectively, the well-known fact that  $\sqrt{N}(\hat{\alpha}_{1k} - \alpha_{1k})$  and  $\sqrt{N}(\hat{\alpha}_{1'l'} - \hat{\alpha}_{1'l'})$  ( $l, l' = 1, \dots, L$ ;  $k, k' = 1, \dots, K$ ) are asymptotically orthogonal because  $\hat{\alpha}_{1k}$  is efficient (see Hausman (1978)) implies that the requirements of the lemma are satisfied. Therefore the GLS proxy variables of  $\beta$  will in general be more efficient if the auxiliary regression coefficients are estimated by SUR rather than OLS.

#### 4. EXTENSIONS

In this section we will indicate extensions of the results in sections 2 and 3 to cases in which aggregates of  $x_1$  in (1) are observed and to a dynamic auxiliary model. For simplicity we consider two examples. First assume that  $x_t$  is a flow variable which is observed every second period only, that is observations are available on  $\bar{x}_t = x_t + x_{t-1}$  if  $t \in T_2 = \{2, 4, 6, \dots, T\}$ . Throughout this section, "-" will denote similar temporal aggregates. Because aggregates are observed more frequently for time series than for cross-sections, we change the notation for the subscripts.

Assume that the analogues of (3) and (4) hold,

$$y_t = \beta x_t + \epsilon_t \quad (27)$$

$$x_t = \alpha z_t + v_t. \quad (28)$$

Ordinary least squares applied to the aggregate data ("a" denotes that aggregates are observed) yields

$$\hat{\beta}_{ac} = (\sum_{T_2} \bar{x}_t^2)^{-1} \sum_{T_2} \bar{x}_t y_t \quad (29)$$

which is consistent. Using the proxy  $\hat{x}_t = \hat{\alpha} z_t + \frac{1}{2}(\bar{x}_t - \hat{\alpha} \bar{z}_t)$  for  $t \in T_2$  and  $\hat{x}_t = \hat{\alpha}_t z_t + \frac{1}{2}(\bar{x}_{t+1} - \hat{\alpha} \bar{z}_{t+1})$  for  $t \in T_2$ , where

$$\hat{\alpha} = (\sum_{T_2} \bar{z}_t^2)^{-1} \sum_{T_2} \bar{z}_t x_t, \quad (30)$$

we have

$$y_t = \hat{x}_t \beta + w_t \quad (31)$$

with

$$w_t = \epsilon_t + \beta(v_t - \frac{1}{2}\bar{v}_t) + \beta(z_t - \frac{1}{2}\bar{z}_t)(\alpha - \hat{\alpha}) \quad \text{if } t \in T_2$$

$$w_t = \epsilon_t + \beta(v_t - \frac{1}{2}\bar{v}_{t+1}) + \beta(z_t - \frac{1}{2}\bar{z}_{t+1})(\alpha - \hat{\alpha}) \quad \text{if } t \notin T_2. \quad (32)$$

OLS applied to (31), GLS applied to (31) with  $V$  being the covariance matrix of  $w_t$  assuming  $\hat{\alpha} = \alpha$ , and GLS with optimal weights, i.e.  $V$  being the covariance matrix of  $w_t$  in (32), yield the consistent estimators  $\hat{\beta}_{ap}$ ,  $\hat{\beta}_{ad}$  and  $\hat{\beta}_{ag}$  respectively. Expressions for the asymptotic variance of the estimators  $\hat{\beta}_{ac}$  and  $\hat{\beta}_{ad}$  and the ML estimator have been given by Palm and Nijman (1982) where  $\hat{\beta}_{ad}$  is called the GLS estimator. For the sake of completeness the formulae are given in appendix A.

A simple transformation of equation (31) yields

$$\begin{aligned} y_t + y_{t-1} &= (x_t + x_{t-1})\beta + \epsilon_t + \epsilon_{t-1} & t \in T_2 \\ y_t - y_{t-1} &= (\hat{x}_t - \hat{x}_{t-1})\beta + w_t - w_{t-1} & t \notin T_2 \end{aligned} \quad (33)$$

From the theorem in the previous section it follows that because of the inclusion of  $(\hat{x}_t - \hat{x}_{t-1})\beta$  in the regressor,  $\hat{\beta}_{ag}$  is asymptotically more efficient than  $\hat{\beta}_{ac}$ .

In table 3 some numerical results on the ratio of the asymptotic variance of alternative consistent estimators compared with the large sample variance of the ML estimator are reported. For simplicity we only consider the case where the disturbances  $\epsilon_t$  and  $v_t$  are normally distributed. In that case the relative efficiency depends on  $R_x^2$ ,  $R_y^2$  and  $\rho = \sigma_z^{-2} E z_t z_{t-1}$ . For cross-sections  $\rho = 0$ . Column 8 of table 3 contains the relative efficiency of the ML estimator for a complete sample with respect to that for the incomplete sample. In column 9 we compare the standard errors for  $\hat{\beta}_{ap}$  computed by means of  $\hat{\sigma}_w^2 (\hat{X}'\hat{X})^{-1}$  with the correct formula for the variance of  $\hat{\beta}_{ap}$ .

Table 3 : Relative efficiency of the ML estimator compared with alternative consistent estimators for  $\beta$ .

$R_x^2$ (1)	$R_y^2$ (2)	$\rho$	Alternative estimators					$\hat{\beta}_{ML}$ complete sampling	$\frac{\sigma_w^2}{\text{var}(\hat{\beta}_{ap}) \Sigma x_t^2}$
			$\hat{\beta}_{ap}$	$\hat{\beta}_{ad}$	$\hat{\beta}_{ag}$	$\hat{\beta}_{ac}$			
0.40	0.40	-0.95	5.8050	4.3690	1.2310	1.3221	0.4099	0.1210	
0.40	0.40	-0.80	2.1353	1.7254	1.1867	1.4380	0.4889	0.3925	
0.40	0.40	-0.40	1.3514	1.2090	1.1425	1.4689	0.6169	0.7826	
0.40	0.40	0.00	1.2076	1.1382	1.1259	1.3761	0.6880	0.9767	
0.40	0.40	0.40	1.1502	1.1189	1.1172	1.2643	0.7333	1.0929	
0.40	0.40	0.80	1.1200	1.1119	1.1119	1.1585	0.7646	1.1703	
0.40	0.40	0.95	1.1122	1.1103	1.1103	1.1218	0.7740	1.1930	
0.40	0.90	-0.95	61.9462	6.1713	1.1461	1.1528	0.3574	0.0305	
0.40	0.90	-0.80	16.1162	2.0562	1.1315	1.1532	0.3921	0.1286	
0.40	0.90	-0.40	5.6263	1.2280	1.1048	1.1436	0.4803	0.4512	
0.40	0.90	0.00	3.2832	1.1114	1.0881	1.1250	0.5625	0.9056	
0.40	0.90	0.40	2.1212	1.0799	1.0767	1.1023	0.6393	1.5931	
0.40	0.90	0.80	1.3645	1.0685	1.0684	1.0777	0.7113	2.7553	
0.40	0.90	0.95	1.1363	1.0659	1.0659	1.0682	0.7371	3.4287	
0.90	0.40	-0.95	1.1481	1.1399	1.0100	4.3439	0.3149	0.2984	
0.90	0.40	-0.80	1.0386	1.0317	1.0050	4.4937	0.6291	0.6588	
0.90	0.40	-0.40	1.0128	1.0080	1.0037	2.6205	0.8386	0.9006	
0.90	0.40	0.00	1.0074	1.0044	1.0035	1.8003	0.9002	0.9719	
0.90	0.40	0.40	1.0050	1.0035	1.0034	1.3671	0.9296	1.0061	
0.90	0.40	0.80	1.0037	1.0033	1.0033	1.1011	0.9469	1.0261	
0.90	0.40	0.95	1.0034	1.0033	1.0033	1.0261	0.9517	1.0317	
0.90	0.90	-0.95	4.0678	3.5825	1.1726	1.5562	0.1128	0.0423	
0.90	0.90	-0.80	2.1694	1.7660	1.0751	1.6972	0.2376	0.1672	
0.90	0.90	-0.40	1.5549	1.1848	1.0348	1.5441	0.4941	0.4850	
0.90	0.90	0.00	1.3209	1.0592	1.0251	1.3546	0.6773	0.7827	
0.90	0.90	0.40	1.1710	1.0258	1.0208	1.1981	0.8147	1.0619	
0.90	0.90	0.80	1.0620	1.0185	1.0183	1.0716	0.9216	1.3245	
0.90	0.90	0.95	1.0281	1.0177	1.0177	1.0305	0.9558	1.4190	



From table 3 we can conclude that  $\hat{\beta}_{ag}$  is fairly accurate in most instances. The estimator  $\hat{\beta}_{ac}$  seems to have a reasonable precision too. However,  $\hat{\beta}_{ap}$  becomes very inaccurate when the autocorrelation of  $z_t$  is negative. The estimator  $\hat{\beta}_{ad}$  is sometimes less accurate than  $\hat{\beta}_{ac}$ . In these cases using additional information in a suboptimal way leads to a loss of efficiency. The estimator  $\hat{\beta}_{ag}$  is of course more efficient than  $\hat{\beta}_{ac}$ . Finally the bias due to using  $\hat{\sigma}_w^2(\hat{X}'\hat{X})^{-1}$  to estimate the asymptotic variance of  $\hat{\beta}_{ap}$  can be quite important.

In the second extension, we consider a dynamic equation for the exogenous variables  $x_t$ . In dynamic models, the prediction of the missing observations will usually depend on auxiliary variables and on the observed values of the variable itself. Simple examples have been considered e.g. by Chow and Lin (1971, 1976), and by Litterman (1983). In more complex models the classical Wiener-Kolmogorov filtering theory or the Kalman filter can be used to derive the best approximations for missing observations, see e.g. Nijman and Palm (1986).

Here we restrict ourselves to a discussion of the relative efficiency of proxy variables estimators for the model

$$y_t = \beta x_t + \epsilon_t \quad (34)$$

$$x_t = \gamma x_{t-1} + \alpha z_t + v_t \quad |\gamma| < 1, \quad (35)$$

where the assumptions on  $\epsilon_t$  and  $v_t$  are as above. Assume that  $x_t$  is observed if  $t \leq T_2$  only, e.g. because the model is semi-annual but only annual data on  $x_t$  are available.

As a proxy for  $x_t$  if it is unobserved we can use

$$\hat{x}_t = (1+\hat{\gamma}^2)^{-1}[\hat{\gamma}x_{t-1} + \hat{\gamma}x_{t+1} + \hat{\alpha}z_t - \hat{\alpha}\hat{\gamma}z_{t+1}], \quad (36)$$

which is the expectation of  $x_t$  given past, present and future information on  $x_t$  and  $z_t$  where consistent estimates have been substituted for  $\alpha$  and  $\gamma$ . OLS applied to (34) after substitution of this proxy for  $x_t$  is consistent for  $\beta$  because (36) is an estimate of the conditional expectation which implies that

(7) is satisfied. Note that a regression on ad hoc interpolated values, e.g. using the method proposed by Boot, Feibes and Lisman (1967) can yield estimates which are strongly biased asymptotically as is shown by Palm and Nijman (1984).

Estimates of  $\alpha$  and  $\gamma$  cannot be obtained by direct regression because  $x_t$  and  $x_{t-1}$  are not observed simultaneously. We consider the following three consistent estimators of  $\alpha$  and  $\gamma$  :

First by ML applied to equation (35) after elimination of the unobserved values of  $x_t$  which can be written as

$$x_t = \gamma^2 x_{t-2} + \alpha z_t + \alpha \gamma z_{t-1} + (v_t + \gamma v_{t-1}), \quad t \in T_2 \quad (37)$$

with one nonlinear restriction on the parameters (M1).

Second by OLS applied to the unrestricted version of (37),

$$x_t = \psi_1 x_{t-2} + \psi_2 z_t + \psi_3 z_{t-1} + (v_t + \gamma v_{t-1}), \quad (38)$$

and  $\hat{\gamma} = \pm \hat{\psi}_1^{1/2}$ ,  $\hat{\alpha} = \hat{\psi}_2$  (M2). The sign of  $\hat{\gamma}$  is determined by  $\hat{\psi}_3 \hat{\psi}_2^{-1}$ .

Third, again using (38), as  $\hat{\gamma} = \hat{\psi}_3 \hat{\psi}_2^{-1}$  and  $\hat{\alpha} = \hat{\psi}_2$  (M3).

If  $z_t$  is a white noise independent of  $x_{t-1}$ , the expectation of  $x_t$  conditional on all observations on  $x_t$  is given by ML from (37) with  $\alpha = 0$  assuming the sign of  $\gamma$  to be known a priori, and this proxy can be substituted into (34) (M4).

As argued above, OLS applied to (34) after substitution of one of these four proxies  $\hat{x}_t$  will yield a consistent estimator of  $\beta$ . This estimator will be denoted  $\hat{\beta}_p$ . The error term  $w_t = \epsilon_t + \beta(x_t - \hat{x}_t)$  is heteroscedastic and serially correlated, so that one is again naturally led to consider GLS-estimators, such as proposed by Dagenais (1973), Gouriéroux and Monfort (1981) and Conniffe (1983b) for static regression models. The estimator using the weights suggested by Dagenais (1973) will be denoted as  $\hat{\beta}_d$ . The estimator using the optimal weights given by the inverse of the covariance matrix of  $w_t$  will be denoted by  $\hat{\beta}_g$ . Finally,  $\beta$  in (34) can be consistently estimated from the complete obser-

vations only by  $\hat{\beta}_C$ .

Numerical results on the relative asymptotic efficiency of the consistent estimators of  $\beta$  discussed above are given in table 4. In the last column of table 4, we compare the SE's of  $\hat{\beta}_p$ , with  $\alpha$  and  $\gamma$  estimated by ML (see M1) with the correct asymptotic standard errors. The parameter  $\gamma$ , the first order autocorrelation of  $z_t$ ,  $\rho$ , and the variance ratio's  $R_x^2$  and  $R_y^2$  determine the relative efficiency of the different estimators with respect to the ML-estimator. Computational details are given in appendix A.

Table 4 : Relative efficiency of the ML estimator compared with alternative consistent estimators for  $\beta$  in (34), when  $x_t$  is generated by (35) and the ratio of commonly used and true asymptotic standard errors.

$\gamma$	$\rho$	$R_x^2$	$R_y^2$	M1			M2		M3		M4		M5	SE
				$\hat{\beta}_p$	$\hat{\beta}_d$	$\hat{\beta}_g$	$\hat{\beta}_p$	$\hat{\epsilon}_g$	$\hat{\epsilon}_p$	$\hat{\beta}_g$	$\hat{\beta}_p$	$\hat{\beta}_g$	$\hat{\beta}_c$	
-0.90	-0.90	0.90	0.90	1.29	1.26	1.20	6.06	1.75	635.13	1.89			1.90	2.19
-0.40	-0.90	0.90	0.90	2.38	1.73	1.23	76.63	1.45	25.54	1.40			1.54	0.80
0.00	-0.90	0.90	0.90	1.32	1.06	1.03	INF	INF	1.98	1.12			1.35	0.97
0.40	-0.90	0.90	0.90	1.67	1.26	1.02	17.69	1.14	2.20	1.07			1.26	0.24
0.90	-0.90	0.90	0.90	1.21	1.18	1.00	1.45	1.04	1.23	1.01			1.64	0.04
-0.90	0.00	0.90	0.90	1.15	1.14	1.12	3.62	1.57	6.34	1.60			1.90	0.47
-0.40	0.00	0.90	0.90	1.24	1.07	1.04	5.65	1.10	1.30	1.05	6.66	1.51	1.47	0.77
0.00	0.00	0.90	0.90	1.32	1.06	1.03	INF	INF	1.32	1.03	1.35	1.35	1.35	0.97
0.40	0.00	0.90	0.90	1.23	1.07	1.03	1.68	1.05	1.28	1.04	2.04	1.42	1.47	0.78
0.90	0.00	0.90	0.90	1.00	1.00	1.00	1.01	1.01	1.03	1.02	1.45	1.30	1.90	0.55
-0.90	0.90	0.90	0.90	1.22	1.19	1.05	1.57	1.17	1.24	1.08			1.64	0.04
-0.40	0.90	0.90	0.90	1.77	1.31	1.02	32.26	1.10	2.28	1.07			1.26	0.23
0.00	0.90	0.90	0.90	1.32	1.06	1.03	INF	INF	1.98	1.12			1.35	0.97
0.40	0.90	0.90	0.90	1.30	1.13	1.08	1.57	1.17	1.49	1.10			1.54	1.47
0.90	0.90	0.90	0.90	1.00	1.00	1.00	1.01	1.01	1.41	1.15			1.90	2.82
-0.90	-0.90	0.40	0.40	2.23	1.71	1.33	20.36	1.58	7326.75	1.61			1.62	0.47
-0.40	-0.90	0.40	0.40	8.48	1.61	1.14	170.52	1.17	263.76	1.16			1.19	0.34
0.00	-0.90	0.40	0.40	3.28	1.11	1.09	INF	INF	3.76	1.09			1.13	1.10
0.40	-0.90	0.40	0.40	4.76	1.27	1.09	13.07	1.10	6.57	1.10			1.14	0.32
0.90	-0.90	0.40	0.40	1.06	1.46	1.01	2.06	1.02	1.91	1.01			1.50	0.08
-0.90	0.00	0.40	0.40	1.96	1.55	1.28	15.91	1.54	71.08	1.55	5.65	1.28	1.61	0.34
-0.40	0.00	0.40	0.40	4.38	1.26	1.11	61.13	1.13	8.87	1.12	3.59	1.13	1.17	0.52
0.00	0.00	0.40	0.40	3.28	1.11	1.09	INF	INF	3.28	1.09	1.13	1.13	1.13	1.10
0.40	0.00	0.40	0.40	2.37	1.10	1.08	2.45	1.09	2.41	1.08	2.26	1.53	1.17	0.95
0.90	0.00	0.40	0.40	1.07	1.01	1.01	1.08	1.02	1.13	1.02	1.23	1.13	1.61	0.62
-0.90	0.90	0.40	0.40	2.37	1.77	1.23	10.20	1.41	4.18	1.36			1.50	0.06
-0.40	0.90	0.40	0.40	5.88	1.36	1.11	11.01	1.12	10.68	1.12			1.14	0.26
0.00	0.90	0.40	0.40	3.28	1.11	1.09	INF	INF	3.76	1.09			1.13	1.10
0.40	0.90	0.40	0.40	2.02	1.08	1.08	4.89	1.14	2.27	1.08			1.19	1.44
0.90	0.90	0.40	0.40	1.06	1.01	1.01	1.07	1.02	1.78	1.06			1.62	0.98
-0.90	-0.90	0.90	0.40	1.02	1.02	1.02	1.39	1.28	49.90	1.95			1.98	2.65
-0.40	-0.90	0.90	0.40	1.12	1.11	1.09	7.82	1.62	3.21	1.48			1.08	1.32
0.00	-0.90	0.90	0.40	1.01	1.00	1.00	INF	INF	1.07	1.05			1.80	0.95
0.40	-0.90	0.90	0.40	1.01	1.01	1.00	2.67	1.36	1.07	1.04			1.93	0.35
0.90	-0.90	0.90	0.40	1.00	1.00	1.00	1.02	1.02	1.00	1.00			1.75	0.06
-0.90	0.00	0.90	0.40	1.01	1.01	1.01	1.20	1.16	1.41	1.26	1.48	1.16	1.90	0.52
-0.40	0.00	0.90	0.40	1.01	1.01	1.01	1.42	1.08	1.01	1.01	1.77	1.57	1.86	0.76
0.00	0.00	0.90	0.40	1.01	1.00	1.00	INF	INF	1.01	1.00	1.00	1.00	1.00	0.95
0.40	0.00	0.90	0.40	1.01	1.01	1.01	1.05	1.03	1.01	1.01	1.62	1.57	1.86	0.76
0.90	0.00	0.90	0.40	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.08	1.08	1.90	0.53
-0.90	0.90	0.90	0.40	1.00	1.00	1.00	1.03	1.03	1.01	1.00			1.93	0.06
-0.40	0.90	0.90	0.40	1.02	1.02	1.01	4.17	1.28	1.08	1.04			1.75	0.35
0.00	0.90	0.90	0.40	1.01	1.00	1.00	INF	INF	1.07	1.05			1.00	0.95
0.40	0.90	0.90	0.40	1.02	1.02	1.02	1.04	1.04	1.04	1.03			1.08	1.45
0.90	0.90	0.90	0.40	1.00	1.00	1.00	1.00	1.00	1.03	1.03			1.98	2.71
-0.90	-0.90	0.40	0.40	1.10	1.10	1.09	2.68	1.58	639.48	1.89			1.90	0.74
-0.40	-0.90	0.40	0.40	1.68	1.49	1.21	16.85	1.39	25.50	1.34			1.50	0.57
0.00	-0.90	0.40	0.40	1.21	1.14	1.13	INF	INF	1.25	1.14			1.38	0.80
0.40	-0.90	0.40	0.40	1.24	1.14	1.07	1.98	1.11	1.40	1.11			1.37	0.39
0.90	-0.90	0.40	0.40	1.04	1.03	1.00	1.05	1.01	1.04	1.00			1.02	0.11
-0.90	0.00	0.40	0.40	1.08	1.08	1.07	2.38	1.50	7.12	1.70	1.42	1.11	1.90	0.48
-0.40	0.00	0.40	0.40	1.29	1.20	1.13	6.61	1.22	1.71	1.16	1.41	1.26	1.48	0.58
0.00	0.00	0.40	0.40	1.01	1.14	1.13	INF	INF	1.21	1.13	1.38	1.38	1.38	0.80
0.40	0.00	0.40	0.40	1.10	1.07	1.06	1.11	1.07	1.10	1.06	1.29	1.29	1.40	0.40
0.90	0.00	0.40	0.40	1.00	1.00	1.00	1.00	1.00	1.01	1.01	1.04	1.03	1.90	0.52
-0.90	0.90	0.40	0.40	1.08	1.08	1.05	1.78	1.38	1.24	1.16			1.02	0.10
-0.40	0.90	0.40	0.40	1.34	1.22	1.11	1.79	1.18	1.76	1.17			1.37	0.36
0.00	0.90	0.40	0.40	1.21	1.14	1.13	INF	INF	1.25	1.14			1.38	0.80
0.40	0.90	0.40	0.40	1.08	1.06	1.06	1.35	1.18	1.10	1.06			1.58	0.89
0.90	0.90	0.40	0.40	1.00	1.00	1.00	1.00	1.00	1.06	1.03			1.98	0.01

From the results in table 4, it is quite obvious that all proxy variables estimators are fairly efficient when  $\gamma$  and  $\alpha$  are estimated by ML. Also, OLS applied to complete data only is reasonably efficient in most instances. The estimator  $\hat{\beta}_p$  can be more efficient as well as less efficient than  $\hat{\beta}_c$ . Notice that the asymptotic variance of  $\hat{\beta}_c$  is twice that of the ML or GLS estimate for the case where all values of  $x_t$  are observed. When a moment estimator (M2 or M3) is used for  $\gamma$  and  $\alpha$ , the relative efficiency is very sensitive to the parameter values. In particular, a negative value for  $\gamma$  combined with negative first order autocorrelation of  $z_t$  often leads to a large relative efficiency of ML compared with the proxy variables estimators based on M2 or M3. The Jacobian of the transformation of the moments to  $\gamma = \psi_1^{1/2}$  equals  $.5\gamma^{-1}$ , so that when  $\gamma = 0$  (which is ignored in the estimation), the large sample variance of these estimators cannot be evaluated. This is indicated by INF.

Evidently, more efficient estimation of  $\alpha$  and  $\gamma$  yields more efficient proxy variables estimators of  $\beta$ , in accordance with our theorem. The inclusion of the observations in  $z_t$  in the conditional expectation of  $x_t$  appears to improve the efficiency of  $\beta$  if (34) is estimated by GLS and (37) is efficiently estimated, which is not surprising given the results in the previous section. Note that it can be more efficient to use the smaller information set if moment estimators of  $\alpha$  and  $\gamma$  are used instead of efficient estimators. Finally the last column of table 4 indicates that the commonly used formula for standard errors can be severely biased when proxies are used. Sign and magnitude of the bias depend on the true parameter values.

## 5. CONCLUDING REMARKS

To summarize we considered several consistent estimators for regression models with missing exogenous variables. It is not difficult to obtain proxies for the

missing observations such that the resulting proxy variables estimators will be consistent. To assure consistency one should preferably use conditional expectations to construct the proxies. We have shown how to obtain proxy variables estimators that are more efficient than estimators based on complete observations only. The use of more information when constructing a proxy and estimating the parameters of the auxiliary equation, will usually yield more efficient estimators. The asymptotic efficiency of some proxy variables estimators is much lower than that of the Gaussian ML estimator. However, the optimal proxy variables estimator, which can be obtained by GLS, appears to be almost as efficient as the ML estimator which is computationally unattractive in larger models and it can be more efficient than ML estimation if normality does not hold. This finding should be very useful for empirical work on data sets which are not complete. Although the computational complexity of ML estimation and the possible deviation of the data from normality are strong arguments in favor of using imputed data, one should be aware of the fact that consistent estimation of the large sample variance of the estimators discussed can sometimes be tricky.

APPENDIX A

In this appendix we shall give the large sample variance for several estimators presented in the paper. Consider first the model presented in (3) and (4). The asymptotic distribution of the Gaussian ML estimator of  $\theta = (\alpha, \sigma^2, \sigma_v^2, \beta)$  is given by (see e.g. Amemiya (1985), theorem 4.1.3)

$$\sqrt{N}(\hat{\theta} - \theta) \approx N(0, A \quad BA^{-1})$$

where

$$A = \lim E \frac{1}{N} \frac{\partial^2 L}{\partial \theta \partial \theta'}$$

and

$$B = \lim E \frac{1}{N} \frac{\partial L_1}{\partial \theta} \frac{\partial L_1}{\partial \theta'}, \text{ where } L_1 \text{ has been defined in section 2.}$$

$$\text{Defining } (E\epsilon_t^4)/(E\epsilon_t^2)^2 = \eta_\epsilon, (E\nu_t^4)/(E\nu_t^2)^2 = \eta_\nu$$

and  $E(\epsilon_t + \beta\nu_t)^4 / (E(\epsilon_t + \beta\nu_t)^2)^2 = \tilde{\eta}$  it is straightforward to verify that

$$B = \begin{bmatrix}
 \frac{\beta^2 \sigma_z^2}{2\sigma^2} + \frac{\sigma_z^2}{2\sigma_v^2} & 0 & 0 & \frac{\alpha\beta\sigma_z^2}{2\sigma^2} \\
 0 & \frac{(\eta_v-1)}{8\sigma^4} + \frac{\beta^4(\eta-1)}{8\sigma^4} & \frac{\beta^2(\eta-1)}{8\sigma^4} & \frac{\beta^3\sigma_v^2(\eta-1)}{4\sigma^4} \\
 0 & \frac{\beta^2(\eta-1)}{8\sigma^4} & \frac{\eta\epsilon-1}{8\sigma^2} + \frac{\eta-1}{8\sigma^4} & \frac{\beta^2\sigma^2(\eta-1)}{4\sigma^4} \\
 \frac{\alpha\beta\sigma_z^2}{2\sigma^2} & \frac{\beta^3\sigma_v^2(\eta-1)}{4\sigma^4} & \frac{\beta^2\sigma^2(\eta-1)}{4\sigma^4} & \frac{\sigma_x^2}{2\sigma^2} + \frac{\alpha^2\sigma_z^2}{2\sigma^2} + \frac{(\eta-1)\beta^2\sigma_v^2}{2\sigma^2}
 \end{bmatrix} \quad (A.1)$$



with  $\sigma^2 = \sigma_v^2 + \beta^2 \sigma_z^2$ ,  $\sigma_z^2 = E z_t^2$  and  $\sigma_x^2 = E x_t^2$ . The matrix

A has the same structure and is obtained if one puts

$\eta_\epsilon = \eta_v = \tilde{\eta} = 3$  in (A.1). If normality holds B coincides with A and the large sample variance of  $\sqrt{N} \hat{\beta}_{ML}$  is simply the (4,4) element of  $A^{-1}$  which can be shown to be

$$\text{Var}(\sqrt{N} \hat{\beta}_{ML}) = \left\{ \begin{array}{l} \frac{\sigma_x^2}{2\sigma^2} + \frac{\alpha^2 \sigma_z^2}{2\tilde{\sigma}^2} \left( 1 - \frac{\beta^2 \sigma_v^2}{\beta^2 \sigma_v^2 + \tilde{\sigma}^2} \right)^2 + \\ + \frac{\beta^2 \sigma_v^2}{\tilde{\sigma}^4} \left( 1 - \frac{\beta^4 \sigma_v^4 + \sigma^4}{\tilde{\sigma}^4 + \sigma^4 + \beta^4 \sigma_v^4} \right)^{-1} \end{array} \right. \quad (\text{A.2})$$

Along the lines followed by Palm and Nijman (1982) for aggregate observations, one can obtain the asymptotic variance of consistent estimators of  $\beta$  for skipped observations.

For the proxy variables estimator  $\hat{\beta}_p$  in (6), we have :

$$\text{Var}(\sqrt{N} \hat{\beta}_p) = 2(\sigma_x^2 + \alpha \sigma_z^2) (\sigma \sigma_x + \sigma \alpha \sigma_z + \alpha \beta \sigma_v \sigma_z). \quad (\text{A.3})$$

Similarly for  $\hat{\beta}_d$  in (11), where the matrix of weights proposed by Dagenais is used, one gets :

$$\text{Var}(\sqrt{N} \hat{\beta}_d) = p + p^2 \beta^2 \alpha^2 \sigma_z^2 \sigma_v^2 / 2\tilde{\sigma}^4. \quad (\text{A.4})$$

with p being the large sample variance of the GLS estimator of  $\beta$  when  $\alpha$  is known

$$p = \left\{ \frac{\sigma_x^2}{2\sigma^2} + \frac{\alpha \sigma_z^2}{2\tilde{\sigma}^2} \right\}^{-1}. \quad (\text{A.5})$$

When the optimal weights in (14) are used, we get :

$$\text{Var}(\sqrt{N} \hat{\beta}_g) = \left\{ \frac{\sigma_x^2}{2\sigma^2} + \frac{\alpha^2 \sigma_z^2}{2\tilde{\sigma}^2} - \frac{\beta^2 \alpha^2 \sigma_v^2 \sigma_z^2}{2(\tilde{\sigma}^4 + \alpha \sigma_v \beta^2 \sigma_z^2)} \right\}^{-1}. \quad (\text{A.6})$$

Finally, when we apply OLS to the complete data, the variance is the double of the variance of OLS in the case that no data are missing

$$\text{Var}(\sqrt{N} \hat{\beta}_c) = 2\sigma^2 \sigma_x^{-2}. \quad (\text{A.7})$$

For the static model with observed aggregates of the exogenous variable  $x_t$ , the large sample variances of the ML estimator and of the estimator  $\sqrt{N} \hat{\beta}_{ad}$  are derived in Palm and Nijman (1982). For the sake of completeness, we give all formulae for the asymptotic variances.

For the proxy variables estimator  $\hat{\beta}_{ap}$ , we have

$$\text{Var}(\sqrt{T} \hat{\beta}_{ap}) = 4(\alpha \tilde{\sigma}_z^2 + E\bar{x}) (\tilde{\sigma}^2 \alpha \tilde{\sigma}_z^2 + \sigma E\bar{x} + \beta^2 \alpha^2 \sigma_v^2 \tilde{\sigma}_z^2 (E\bar{z}^2)^{-1}). \quad (\text{A.8})$$

When a temporal aggregate of  $x_t$  is available, the variance of the Dagenais estimator in (11) is

$$\text{Var}(\sqrt{T} \hat{\beta}_{ad}) = p + p^2 \alpha^2 \beta^2 \tilde{\sigma}_z^2 \sigma_v^2 \tilde{\sigma}^{-4} (E\bar{z})^{-2}. \quad (\text{A.9})$$

$$\text{where } p = \left\{ \frac{E\bar{x}^2}{4\sigma^2} + \frac{\alpha^2 \tilde{\sigma}_z^2}{4\tilde{\sigma}^2} \right\}^{-1} \text{ is the variance of } \hat{\beta}_{ad} \text{ given}$$

that  $\alpha$  is known and  $\tilde{\sigma}_z^2 = E(z_t - z_{t-1})^2$ . The asymptotic variance of the GLS estimator with optimal weights,  $\hat{\beta}_{ag}$ , is

$$\text{Var}(\sqrt{T} \hat{\beta}_{ag}) = \left\{ \frac{\alpha^2 \tilde{\sigma}_z^2}{4\sigma^2} + \frac{E\bar{x}^2}{4\sigma^2} - \frac{\alpha \beta \sigma_v^2 \tilde{\sigma}_z^2}{4\sigma^4 E\bar{z}^2 + 4\beta^2 \alpha^2 \sigma_v^2 \tilde{\sigma}_z^2} \right\}^{-1}. \quad (\text{A.10})$$

The variance of the OLS estimator applied to the periods for which all variables are observed,  $\hat{\beta}_{ac}$ , is

$$\text{Var}(\sqrt{T} \hat{\beta}_{AC}) = 4\sigma^2 / E\tilde{x}^2. \quad (\text{A.11})$$

It differs from the variance of the OLS estimator of  $\beta$  when no data are missing

$$\text{Var}(\sqrt{T} \hat{\beta}_{OLS}) = \sigma^2 / \sigma_x^2, \quad (\text{A.12})$$

$$\text{with } \sigma_x^2 = \alpha \sigma_z^2 + \sigma_v^2.$$

The asymptotic variance of the ML estimator  $\hat{\beta}_{aML}$ , is

$$\begin{aligned} \text{Var}(\sqrt{T} \hat{\beta}_{aML}) = & \left\{ \frac{E\tilde{x}^2}{4\tilde{\sigma}_z^2} + \frac{\alpha \tilde{\sigma}_z^2}{4\tilde{\sigma}_z^2} \left[ 1 - \frac{\beta \tilde{\sigma}_z^2 \sigma_v^2}{(\beta^2 \tilde{\sigma}_z^2 \sigma_v^2 + E\tilde{z}^2 \tilde{\sigma}_z^2)} \right] \right. \\ & \left. + \frac{\beta \sigma_v^2}{\tilde{\sigma}_z^4} \left[ 1 - \frac{(\sigma^4 + \beta^4 \sigma_v^2 + \tilde{\sigma}_z^4)}{(\sigma^4 + \beta^4 \sigma_v^2 + \tilde{\sigma}_z^4)} \right] \right\}^{-1}. \quad (\text{A.13}) \end{aligned}$$

Finally, we indicate briefly how table 4 was derived. If  $\hat{x}_t$  defined in (36) is substituted in (34) the resulting error term has matrix variance covariance matrix

$$\Omega = \Omega_1 + W \Omega_2 W', \quad (\text{A.14})$$

where  $\Omega_2$  is the covariance matrix of the estimates  $\hat{\gamma}$  and  $\hat{\alpha}$ ,  $W$  is a  $(T \times 2)$  matrix which contains  $\beta$  times the derivatives of  $\hat{x}_t$  with respect to  $\alpha$  and  $\gamma$  in the first and second column respectively and  $\Omega_1$  is a diagonal matrix with diagonal element  $\sigma^2$  in case of an observed  $x_t$  and  $\sigma^2 + \beta^2 \sigma_v^2 (1 + \gamma^2)^{-1}$  if  $x_t$  is not observed. Again equation (15) can be used to get the inverse of  $\Omega$ .

In order to derive the variance of the ML estimator for the dynamic regression model in (34) and (35) we write the model in recursive form as

$$y_t = \beta x_t + \epsilon_t$$

$$y_{t-1} = \beta(1+\gamma^2)^{-1}\{\gamma x_t + \gamma x_{t-2} - \alpha \gamma z_t + \alpha z_{t-1}\} + \epsilon_{t-1} \\ - \beta(1+\gamma^2)^{-1}(\gamma v_t - v_{t-1}) \quad (\text{A.15})$$

$$x_t = \gamma^2 x_{t-2} + \alpha z_t + \alpha \gamma z_{t-1} + v_t + \gamma v_{t-1}$$

for  $t \leq T_2$ . Notice that the disturbances in (A.15) are independent and orthogonal to the explanatory variables in the corresponding equation. The log-likelihood function  $L$  can therefore be obtained in a straightforward manner, as well as the associated information matrix.

APPENDIX B PROOF OF THE THEOREM

Using assumptions (ii) and (iii) one verifies that

$$\sqrt{N} (\tilde{\beta}_{IV} - \beta) \underset{a}{\sim} N(0, Q^{-1}(D_{11} + D_{22})Q^{-1}), \text{ where } D_{11} \text{ and } D_{22}$$

are the upper-left and the lower-right blocks of  $D$  respectively.

Furthermore, the asymptotic orthogonality of  $Z$  and  $\epsilon$  and assumption (iii) imply that

$$p\lim N^{-1}Z'(\tilde{w} - \hat{w}) = p\lim N^{-1}Z'(\hat{X} - \tilde{X})\beta = 0,$$

for all  $\beta$ , so that  $p\lim N^{-1}Z'\tilde{X} = p\lim N^{-1}Z'\hat{X} = Q$ . Using this result, one obtains that

$$\begin{aligned} Q^{-1}(D_{11}+D_{22})Q^{-1} &= p\lim N^{-1}(Q^{-1}Z' - V^{-1}\hat{X}'\hat{\Sigma}^{-1})\hat{\Sigma}^{-1}(ZQ^{-1} - \\ &\quad - \hat{\Sigma}^{-1}\hat{X}V^{-1}) + p\lim N^{-1}Q^{-1}Z'\hat{X}V^{-1} + \\ &\quad p\lim T^{-1}V^{-1}\hat{X}'ZQ^{-1} - V^{-1} + Q^{-1}D_{22}Q^{-1} \geq V^{-1} \end{aligned}$$

which proves the result.

APPENDIX C

PROOF OF THE RELATIVE EFFICIENCY OF THE PROXY VARIABLES ESTIMATOR BASED ON THE LARGER INFORMATION SET FOR A SPECIAL CASE.

Assume that (4) and (24) both hold, which read in matrix notation as

$$x = Z_1\alpha_1 + Z_2\alpha_2 + v \quad (C.1)$$

and

$$x = Z_1\eta + v^* \quad (C.2)$$

respectively. Evidently

$$v^* = Z_1(\alpha_1 - \eta) + Z_2\alpha_2 + v \quad (C.3)$$

and

$$\eta = \alpha_1 + \text{plim}(Z_1'Z_1)^{-1}Z_1'Z_2\alpha_2 \quad (C.4)$$

because  $E[x|Z_1] = Z_1\eta$ . Using (C.3) and (C.4) we find

$$N^{-\frac{1}{2}}(Z_1'v^* - Z_1'v) = N^{-\frac{1}{2}}(Z_1'Z_1(\alpha_1 - \eta) + Z_1'Z_2\alpha_2 + Z_1'v - Z_1'v) \xrightarrow{p} 0 \quad (C.5)$$

It is not difficult to check using the theorem in section 3

that if  $\hat{X} = \hat{X}^{(1)}$ ,  $\tilde{X} = \hat{X}^{(2)}$ ,  $\hat{\beta}_{GLS} = \hat{\beta}_g^{(1)}$  and  $\hat{\beta}_{IV} = \hat{\beta}_g^{(2)}$ , a sufficient condition for relative efficiency of  $\hat{\beta}_g^{(1)}$  is that  $N^{-\frac{1}{2}}Z'(\tilde{w} - \hat{w}) \xrightarrow{p} 0$  which is satisfied when

$$N^{-\frac{1}{2}}\{Z_{1I}'(V_I - V_I^*)\beta + Z_I(Z_C'Z_C)^{-1}Z_C'V_C\beta - Z_{1I}(Z_{1c}'Z_{1c})^{-1}Z_{1c}'V_{1c}^*\beta\} \xrightarrow{p} 0, \quad (C.6)$$

where the subscripts c and I refer to complete and incomplete observations as before. Condition (C.6) is satisfied because of (C.5).

REFERENCES

- Afifi, A.A., and R.W. Elashoff, 1966, Missing observations in multivariate statistics, I. Review of the literature. *Journal of the American Statistical Association*, 61, pp. 595-604.
- Anderson, T.W., 1957, Maximum likelihood estimates for a multivariate normal distribution when some observations are missing. *Journal of the American Statistical Association*, 67, pp. 628-632.
- Amemiya, T., 1985, *Advanced Econometrics*. Harvard University Press, Cambridge.
- Boot, J.C.G., Feibes, W. and J.H.C. Lisman, 1967, Further methods of derivation of quarterly figures from annual data. *Applied Statistics*, 16, pp. 65-75.
- Chow, G.C., and A. Lin, 1971, Best linear unbiased interpolation and extrapolation of time series by related series. *The Review of Economics and Statistics*, 53, pp. 372-375.
- Chow, G.C., and A. Lin, 1976, Best linear unbiased estimation of missing observations in an economic time series. *Journal of the American Statistical Association*, 71, pp. 719-721
- Conniffe, D., 1983a, Small-sample properties of estimators of regression coefficients given a common pattern of missing data. *Review of Economic Studies*, 50, pp. 111-120.
- Conniffe, D., 1983b, Comments on the weighted regression approach to missing values. *The Economic and Social Review*, 14, pp. 259-272.
- Dagenais, M.G., 1973, The use of incomplete observations in multiple regression analysis, a generalized least squares approach. *Journal of Econometrics*, 1, pp. 317-348.
- Gouriéroux, C., and A. Monfort, 1981, On the problem of missing data in linear models. *Review of Economic Studies*, 48, pp. 579-586.
- Griliches, Z., 1986, Economic data issues, in Z. Griliches and M.D. Intriligator, eds., *Handbook of Econometrics*, North-Holland Publ.Co, Amsterdam, pp. 1466-1514.
- Hausman, J., 1978, Specification tests in econometrics. *Econometrica*, 46, pp. 1251-1272.
- Litterman, R.B., 1983, A random walk, Markov model for the distribution of time series. *Journal of Business and Economic Statistics*, 1, pp. 169-173.

- Little, R.J.A., 1982, Models for non-reponse in sample surveys. *Journal of the American Statistical Association*, 77, pp. 237-250.
- Murphy, K.M. and R.H. Topel, 1985, Estimation and inference in two-step econometric models. *Journal of Business and Economic Statistics*, 3, pp. 370-379.
- Nijman, T.E., and F.C. Palm, 1986, The construction and use of approximations for missing quarterly observations : a model-based approach. *Journal of Business and Economic Statistics*, 4, pp. 47-58.
- Pagan, A., 1984, Econometric issues in the analysis of regressions with generated regressors. *International Economic Review*, 25, pp. 221-248.
- Palm, F.C., and T.E. Nijman, 1982, Linear regression using both temporally aggregated and temporally disaggregated data. *Journal of Econometrics*, 19, pp. 333-343.
- Palm, F.C., and T.E. Nijman, 1984, Missing observations in the dynamic regression model. *Econometrica*, 52, pp. 1415-1435.
- Zellner, A., 1962, An efficient method of estimating seemingly unrelated regressions and tests for aggregation bias, *Journal of the American Statistical Association*, 57, pp. 348-368.



## IN 1986 REEDS VERSCHENEN

- 202 J.H.F. Schilderink  
Interregional Structure of the European Community. Part III
- 203 Antoon van den Elzen and Dolf Talman  
A new strategy adjustment process for computing a Nash equilibrium in a noncooperative more-person game
- 204 Jan Vingerhoets  
Fabrication of copper and copper semis in developing countries. A review of evidence and opportunities
- 205 R. Heuts, J. van Lieshout, K. Baken  
An inventory model: what is the influence of the shape of the lead time demand distribution?
- 206 A. van Soest, P. Kooreman  
A Microeconomic Analysis of Vacation Behavior
- 207 F. Boekema, A. Nagelkerke  
Labour Relations, Networks, Job-creation and Regional Development. A view to the consequences of technological change
- 208 R. Alessie, A. Kapteyn  
Habit Formation and Interdependent Preferences in the Almost Ideal Demand System
- 209 T. Wansbeek, A. Kapteyn  
Estimation of the error components model with incomplete panels
- 210 A.L. Hempenius  
The relation between dividends and profits
- 211 J. Kriens, J.Th. van Lieshout  
A generalisation and some properties of Markowitz' portfolio selection method
- 212 Jack P.C. Kleijnen and Charles R. Standridge  
Experimental design and regression analysis in simulation: an FMS case study
- 213 T.M. Doup, A.H. van den Elzen and A.J.J. Talman  
Simplicial algorithms for solving the non-linear complementarity problem on the simplotope
- 214 A.J.W. van de Gevel  
The theory of wage differentials: a correction
- 215 J.P.C. Kleijnen, W. van Groenendaal  
Regression analysis of factorial designs with sequential replication
- 216 T.E. Nijman and F.C. Palm  
Consistent estimation of rational expectations models

- 217 P.M. Kort  
The firm's investment policy under a concave adjustment cost function
- 218 J.P.C. Kleijnen  
Decision Support Systems (DSS), en de kleren van de keizer ...
- 219 T.M. Doup and A.J.J. Talman  
A continuous deformation algorithm on the product space of unit simplices
- 220 T.M. Doup and A.J.J. Talman  
The 2-ray algorithm for solving equilibrium problems on the unit simplex
- 221 Th. van de Klundert, P. Peters  
Price Inertia in a Macroeconomic Model of Monopolistic Competition
- 222 Christian Mulder  
Testing Korteweg's rational expectations model for a small open economy
- 223 A.C. Meijdam, J.E.J. Plasmans  
Maximum Likelihood Estimation of Econometric Models with Rational Expectations of Current Endogenous Variables
- 224 Arie Kapteyn, Peter Kooreman, Arthur van Soest  
Non-convex budget sets, institutional constraints and imposition of concavity in a flexible household labor supply model
- 225 R.J. de Groof  
Internationale coördinatie van economische politiek in een twee-regio-twee-sectoren model
- 226 Arthur van Soest, Peter Kooreman  
Comment on 'Microeconomic Demand Systems with Binding Non-Negativity Constraints: The Dual Approach'
- 227 A.J.J. Talman and Y. Yamamoto  
A globally convergent simplicial algorithm for stationary point problems on polytopes
- 228 Jack P.C. Kleijnen, Peter C.A. Karremans, Wim K. Oortwijn, Willem J.H. van Groenendaal  
Jackknifing estimated weighted least squares
- 229 A.H. van den Elzen and G. van der Laan  
A price adjustment for an economy with a block-diagonal pattern
- 230 M.H.C. Paardekooper  
Jacobi-type algorithms for eigenvalues on vector- and parallel computer
- 231 J.P.C. Kleijnen  
Analyzing simulation experiments with common random numbers

- 232 A.B.T.M. van Schaik, R.J. Mulder  
On Superimposed Recurrent Cycles
- 233 M.H.C. Paardekooper  
Sameh's parallel eigenvalue algorithm revisited
- 234 Pieter H.M. Ruys and Ton J.A. Storcken  
Preferences revealed by the choice of friends
- 235 C.J.J. Huys en E.N. Kertzman  
Effectieve belastingtarieven en kapitaalkosten
- 236 A.M.H. Gerards  
An extension of König's theorem to graphs with no odd- $K_4$
- 237 A.M.H. Gerards and A. Schrijver  
Signed Graphs - Regular Matroids - Grafts
- 238 Rob J.M. Alessie and Arie Kapteyn  
Consumption, Savings and Demography
- 239 A.J. van Reeken  
Begrippen rondom "kwaliteit"
- 240 Th.E. Nijman and F.C. Palmer  
Efficiency gains due to using missing data. Procedures in regression models
- 241 S.C.W. Eijffinger  
The determinants of the currencies within the European Monetary System

## IN 1987 REEDS VERSCHENEN

- 242 Gerard van den Berg  
Nonstationarity in job search theory
- 243 Annie Cuyt, Brigitte Verdonk  
Block-tridiagonal linear systems and branched continued fractions
- 244 J.C. de Vos, W. Vervaat  
Local Times of Bernoulli Walk
- 245 Arie Kapteyn, Peter Kooreman, Rob Willemse  
Some methodological issues in the implementation  
of subjective poverty definitions
- 246 J.P.C. Kleijnen, J. Kriens, M.C.H.M. Lafleur, J.H.F. Pardoel  
Sampling for Quality Inspection and Correction: AOQL Performance  
Criteria
- 247 D.B.J. Schouten  
Algemene theorie van de internationale conjuncturele en structurele  
afhankelijkheden
- 248 F.C. Bussemaker, W.H. Haemers, J.J. Seidel, E. Spence  
On  $(v,k,\lambda)$  graphs and designs with trivial automorphism group
- 249 Peter M. Kort  
The Influence of a Stochastic Environment on the Firm's Optimal Dynamic  
Investment Policy
- 250 R.H.J.M. Gradus  
Preliminary version  
The reaction of the firm on governmental policy: a game-theoretical  
approach
- 251 J.G. de Gooijer, R.M.J. Heuts  
Higher order moments of bilinear time series processes with symmetrically  
distributed errors
- 252 P.H. Stevers, P.A.M. Versteijne  
Evaluatie van marketing-activiteiten
- 253 H.P.A. Mulders, A.J. van Reeken  
DATAAL - een hulpmiddel voor onderhoud van gegevensverzamelingen
- 254 P. Kooreman, A. Kapteyn  
On the identifiability of household production functions with joint  
products: A comment
- 255 B. van Riel  
Was er een profit-squeeze in de Nederlandse industrie?
- 256 R.P. Gilles  
Economies with coalitional structures and core-like equilibrium concepts

- 257 P.H.M. Ruys, G. van der Laan  
Computation of an industrial equilibrium
- 258 W.H. Haemers, A.E. Brouwer  
Association schemes
- 259 G.J.M. van den Boom  
Some modifications and applications of Rubinstein's perfect equilibrium model of bargaining
- 260 A.W.A. Boot, A.V. Thakor, G.F. Udell  
Competition, Risk Neutrality and Loan Commitments
- 261 A.W.A. Boot, A.V. Thakor, G.F. Udell  
Collateral and Borrower Risk
- 262 A. Kapteyn, I. Woittiez  
Preference Interdependence and Habit Formation in Family Labor Supply
- 263 B. Bettonvil  
A formal description of discrete event dynamic systems including perturbation analysis
- 264 Dr. Sylvester C.W. Eijffinger  
A monthly model for the monetary policy in the Netherlands
- 265 F. van der Ploeg, A.J. de Zeeuw  
Conflict over arms accumulation in market and command economies
- 266 F. van der Ploeg, A.J. de Zeeuw  
Perfect equilibrium in a model of competitive arms accumulation
- 267 Aart de Zeeuw  
Inflation and reputation: comment
- 268 A.J. de Zeeuw, F. van der Ploeg  
Difference games and policy evaluation: a conceptual framework
- 269 Frederick van der Ploeg  
Rationing in open economy and dynamic macroeconomics: a survey
- 270 G. van der Laan and A.J.J. Talman  
Computing economic equilibria by variable dimension algorithms: state of the art
- 271 C.A.J.M. Dirven and A.J.J. Talman  
A simplicial algorithm for finding equilibria in economies with linear production technologies

Bibliotheek K. U. Brabant



17 000 01059384 7