

Tilburg University

The COMPlaints After Stroke (COMPAS) study

van Rijsbergen, M.W.A.; Mark, R.E.; Kort, P.L.M.; Sitskoorn, M.M.

Published in:
BMJ Open

DOI:
[10.1136/bmjopen-2013-003599](https://doi.org/10.1136/bmjopen-2013-003599)

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):
van Rijsbergen, M. W. A., Mark, R. E., Kort, P. L. M., & Sitskoorn, M. M. (2013). The COMPlaints After Stroke (COMPAS) study: Protocol for a Dutch cohort study on poststroke subjective cognitive complaints. *BMJ Open*, 3(9), e003599. <https://doi.org/10.1136/bmjopen-2013-003599>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

BMJ Open

The COMPlaints After Stroke (COMPAS) study: protocol for a Dutch cohort study on poststroke subjective cognitive complaints

Marielle W A van Rijsbergen, Ruth E Mark, Paul L M de Kort, et al.

BMJ Open 2013 3:

doi: 10.1136/bmjopen-2013-003599

Updated information and services can be found at:

<http://bmjopen.bmj.com/content/3/9/e003599.full.html>

These include:

- | | |
|-------------------------------|--|
| References | This article cites 35 articles, 6 of which can be accessed free at:
http://bmjopen.bmj.com/content/3/9/e003599.full.html#ref-list-1 |
| Open Access | This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 3.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/3.0/ |
| Email alerting service | Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article. |
-

Topic Collections

Articles on similar topics can be found in the following collections

[Neurology](#) (115 articles)
[Patient-centred medicine](#) (87 articles)
[Rehabilitation medicine](#) (58 articles)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>

The COMPlaints After Stroke (COMPAS) study: protocol for a Dutch cohort study on poststroke subjective cognitive complaints

Marielle W A van Rijsbergen,¹ Ruth E Mark,¹ Paul L M de Kort,² Margriet M Sitskoorn¹

To cite: van Rijsbergen MWA, Mark RE, de Kort PLM, *et al.* The COMPlaints After Stroke (COMPAS) study: protocol for a Dutch cohort study on poststroke subjective cognitive complaints. *BMJ Open* 2013;**3**:e003599. doi:10.1136/bmjopen-2013-003599

► Prepublication history for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2013-003599>).

Received 15 July 2013
Accepted 9 August 2013

For numbered affiliations see end of article.

Correspondence to

Dr Ruth E Mark;
r.e.mark@tilburguniversity.edu

ABSTRACT

Background: Although many studies have assessed poststroke objective cognitive impairment, only a few have evaluated patients' subjective cognitive complaints (SCC). Although these SCC are found to be common in the early and chronic phases after stroke, knowledge about their risk factors, course over time, differences with healthy controls and their diagnostic relevance is limited. The aim of the COMPlaints After Stroke (COMPAS) study is therefore to determine the possible risk factors, prognosis, time course and predictive value of SCC in the first 2 years after stroke.

Methods and design: A prospective cohort study is conducted in which patients are compared to non-stroke controls at 3, 6, 12 and 24 months after stroke. Approximately 300 patients are recruited from the stroke units of three hospitals in the Netherlands, while 300 controls are sought among the relatives (spouses excluded) and social networks of participants. A wide range of subjective and objective variables is assessed in both groups using interviews, questionnaires and neuropsychological assessment. The primary outcomes include SCC and objective cognitive impairment, whereas secondary outcomes are quality of life, subjective recovery and daily life functioning.

Ethics and dissemination: The study is being carried out in agreement with the Declaration of Helsinki and the Medical Research Involving Human Subjects Act. The protocol has been approved by the medical ethics committees of the participating centres and all participants give written informed consent. The results will be published in peer-reviewed journals and disseminated to the medical society and general public.

Discussion: The COMPAS study is the first to systematically evaluate poststroke SCC in a prospective longitudinal design, taking a wide range of subjective and objective variables into account. The results obtained can be used to accurately inform patients and their families, as well as to develop patient-tailored intervention programmes to ultimately improve stroke patient care.

BACKGROUND

Poststroke cognitive impairment is common after stroke and can be evaluated either

ARTICLE SUMMARY

Strengths and limitations of this study

- Strengths of this study are that it is the first prospective cohort study on subjective cognitive complaints in patients with stroke in which patients and controls are evaluated at multiple assessments, while at the same time a wide range of subjectively and objectively measured variables are taken into account.
- A limitation is that the most seriously affected patients are unable to participate in the study. This may reduce the generalisability of the results to the stroke population as a whole.

objectively, using neuropsychological tests (ie, objective cognitive performances (OCP)), or subjectively, using interviews or self-report questionnaires (ie, subjective cognitive complaints (SCC)). Until now, the majority of studies on poststroke cognitive sequelae have focused on OCP without also evaluating patients' SCC. However, individuals' performances in test situations do not always correspond to those in daily life and vice versa.^{1 2} Evaluating one can therefore not be used to draw conclusions about the other.

In a recent systematic review, we found that SCC are common in the early and chronic phases after stroke and that they tend to increase over time.³ The prevalence rates vary between 28.6% and 92.0% and complaints about memory, mental speed and concentration were found to be those most commonly reported.³ Furthermore, patients and their proxies showed moderate agreement on the prevalence and severity of patients' SCC.³ However, one of the main problems in most of these studies is that there is no 'gold standard' to define and measure SCC, resulting in heterogenic findings. In our review, we suggested that it is important to differentiate between *content of*

SCC (SCC_c) and *worrying* about SCC (SCC_w), as these are two different concepts.³ The first focuses on the specific cognitive difficulties respondents say they experience, while the second indicates whether participants find them worrisome, irritating and whether they say they hinder daily life. A few studies have made this distinction so far.^{2 4 5} However, the majority of research on poststroke SCC has evaluated SSC_c and not SSC_w, probably without being aware of the difference between these aspects of SCC.³

Furthermore, we found in our review that poststroke SCC are inconsistently associated with demographic and clinical characteristics, current depressive symptoms and OCP,³ but they may predict future emotional and cognitive decline.^{5 6} However, most of the research on SCC after stroke carried out so far is limited in that: unvalidated methods for assessing SCC were used, there was no non-stroke control group included, and the focus was on a specific subsample of stroke patients (eg, home-living patients only), thereby impairing the generalisability of the results. While SCC are common among stroke patients, knowledge about the following aspects is only limited or practically non-existent: the risk profile for developing SCC; their course over time; their impact on quality of life (QOL), subjective recovery and activities in daily life (ADL) functioning; and their prognostic implications.

In the general non-stroke population, however, SCC have been more frequently evaluated, in particular memory-related SCC reported by the elderly.^{7 8} Factors found to be associated with these complaints include: demographic characteristics (higher age, women, lower education), psychological distress, somatic symptoms, personality traits (neuroticism in particular) and vascular risk factors.⁷⁻¹¹ Furthermore, they are thought to be clinically relevant in this group because of their association with increased healthcare consumption, reduced QoL, current OCP (this link is not always found) and their predictive value for future cognitive decline.^{7 8 12} Whether this also applies to poststroke SCC is unknown. More systematic research is therefore needed to gain further knowledge about SCC among stroke survivors, to be able to accurately inform patients and their relatives, to develop adequate treatment programmes and to ultimately improve poststroke care.

We therefore designed the *COMPlaints After Stroke* (COMPAS) study in which we have four main aims:

- ▶ Determine the prevalence, profile and course over time of SCC_c and SCC_w.
- ▶ Identify the risk profile for reporting SCC.
- ▶ Evaluate their predictive value for future cognitive functioning.
- ▶ Determine the effect of SCC on QoL, subjective recovery and ADL functioning.

Here we describe the design and protocol of the COMPAS study, which is the first prospective cohort study of SCC in patients with stroke, evaluating both patients and controls, while at the same time a wide range of variables is taken into account.

METHODS AND ANALYSIS

Design

A multicentre, prospective cohort study of patients with stroke and controls is performed. We started in 2009 and the final measurements will be made in 2014. Patients are evaluated five times, starting at the clinical phase (T0), followed by an assessment at 3 months (T1), 6 months (T2), 1 year (T3) and 2 years (T4) after stroke. Controls are seen at the same time intervals, starting at T1.

Study population

Patients with stroke are recruited consecutively from the stroke units of three hospitals in the Netherlands, including the St Elisabeth and TweeSteden Hospitals in Tilburg and the Maxima Medical Centre in Veldhoven. The control group consists of a sample from the non-stroke general population and is recruited among the relatives and the social networks of participants in the COMPAS study. Spouses of patients with stroke are excluded from the control group since they are at a higher risk of having physical, cognitive and psychosocial problems themselves due to the fact that their partner has suffered a stroke.^{13 14}

Inclusion criteria

- ▶ Clinical diagnosis of a first or recurrent ischaemic or haemorrhagic stroke (for patients only).
- ▶ At least 18 years old (no upper age limit).

Exclusion criteria

- ▶ Pre-existent health problems interfering with cognitive functioning, including for example:
 - Cognitive decline (as defined by a score >3.6 on the short version of the Informant Questionnaire on Cognitive Decline in the Elderly; IQCODE).¹⁵
 - A recent history of severe psychopathology (eg, suicide attempts, alcohol- or drug misuser, diagnosed personality or mood disorders).
 - Severe physical comorbidity (eg, malignant diseases, progressive neurological conditions).
- ▶ Severe communication difficulties (eg, insufficient understanding of the Dutch language, severe aphasia, blindness or deafness).

Procedure

Eligible patients receive oral and written information about the study from their treating physician during the clinical phase (T0). Demographic and clinical characteristics are documented and patients are scheduled for the first assessment 3 months after stroke (T1), during which written informed consent is obtained for inclusion to be definite. Participants acknowledge that they have the intention to complete all four assessments, and that they are allowed to end their participation at any time. For the follow-up assessments (T2–T4), patients are informed by letter and telephone and invited to participate after which an appointment is scheduled.

Potential controls receive oral and written information about the study from the researcher after which they are

Table 1 Primary and secondary outcomes in the COMPAS study

	Instrument	T0	T1	T2	T3	T4
Primary outcomes						
SCC	Cognitive Failures Questionnaire ^{16 17}		X	X	X	X
	Checklist for Cognitive and Emotional Consequences ⁵		X	X	X	X
OCP ^{20 21}						
Global cognitive functioning	Mini-Mental State Examination		X		X	X
Visual perception and construction	Rey Complex Figure Test—copy trial		X		X	X
Mental speed/attention	Stroop Colour word test—cards 1 and 2		X		X	X
	Digit Symbol-Coding		X		X	X
Episodic memory	Rivermead Behavioural Memory Test		X		X	X
	Rey Complex Figure Test—immediate and delayed recall trials		X		X	X
	Verbal Paired Associates		X		X	X
Working memory	Digit span Forward and Backward conditions		X		X	X
Language	Boston Naming Test—short version		X		X	X
Executive functioning	Controlled Oral Word Association Test—FAS		X		X	X
	Category Fluency Test: animals and occupations		X		X	X
	Stroop Colour Word Test—card 3		X		X	X
	Rule Shift Cards		X		X	X
	Zoo Map		X		X	X
Fine motor dexterity	Purdue Pegboard		X		X	X
Secondary outcomes						
Quality of Life	WHO Quality of Life Questionnaire—short form ²²		X		X	X
	WHO Quality of Life Questionnaire—Old module ²³		X		X	X
ADL functioning						
Basic ADL	Barthel index ²⁵	P	X	X	X	X
Instrumental ADL	Frenchay Activities Index ²⁶		X	X	X	X
Subjective stroke recovery	Item 9 of Stroke Impact Scale ²⁴		P		P	P

ADL, activities in daily life; C, control group only; COMPAS, COMPLAINTS After Stroke; OCP, objective cognitive performance; P, patient group only; SCC, subjective cognitive complaints; T0, clinical phase; T1, 3 months poststroke; T2, 6 months poststroke; T3, 1 year poststroke; T4, 2 years poststroke; X, instrument used in both patients and controls.

asked to participate in the study. The rest of the procedure is the same as that for the patient group.

The assessments are administered in a standardised way by trained neuropsychologists and take place at the participating hospitals or, when this is not possible, at the participant's home or residence (eg, rehabilitation centre).

Measures

Tables 1 and 2 give an overview of the variables assessed and instruments used at each time point.

Outcomes

Primary outcomes of the COMPAS study are SCC and OCP. To measure SCC, two instruments are used, namely: the Dutch version of the Cognitive Failures Questionnaire (CFQ)^{16 17} and the Checklist for Cognitive and Emotional consequences following stroke (CLCE-24).⁵ The CFQ focuses on SCCc and asks subjects to rate 25 items on the frequency of cognitive slips and errors in daily life on a 5-point Likert scale ranging from 0 (never) to 4 (very often). SCCw is evaluated by four additional general questions regarding the subjective increase of complaints over time, the degree to

which these hinder daily life, are annoying and are a source of concern. Each of these extra items is rated on a scale ranging from 1 (not at all) to 5 (extremely).

The CLCE-24 is a structured clinical interview developed to evaluate SCCc and SCCw among stroke survivors.⁵ It consists of 13 items concerning cognitive complaints and 9 items addressing emotional and behavioural complaints. Each item is scored as 0 (not present), 1 (doubtful), 2 (present, but not affecting daily life) or 3 (present and negatively affecting daily life).

OCP are evaluated using an extensive neuropsychological assessment covering multiple cognitive domains and containing traditional (eg, Rey Complex Figure Test¹⁸) and more ecologically valid tests (eg, Rivermead Behavioural Memory Test¹⁹). See table 1 for an overview of all OCP tests used. In Spreen and Straus²⁰ and Lezak *et al.*,²¹ a detailed description of each of the instruments we use is given.

Secondary outcomes include QoL, ADL functioning and subjective stroke recovery. Generic QoL is evaluated using the short version of the self-report WHO Quality of Life Questionnaire (WHOQOL-Bref)²² (26 items) and, because we expect the majority of our population

Table 2 Possible determinants in the COMPAS study

	Variable/instrument	T0	T1	T2	T3	T4
Demographic variables	Age, gender, education, marital status, living situation, residence, employment status, hand preference	P	X	X	X	X
Clinical characteristics						
Stroke-specific	Lifetime history of stroke, type, side, classification according to the Oxford Community Stroke Project, ²⁷ severity within 24 h after admission using the National Institutes of Health Stroke Scale, ²⁸ treatment, poststroke complications, length of hospital stay, discharge destination	P				
General	Vascular risk factors, comorbidity (Cumulative Illness Rating Scale ²⁹), (re-) admissions to hospital, medication use, current participation in rehabilitation therapy		X	X	X	X
Health status	12-Item Short Form Health Survey ³⁰		X		X	X
Premorbid status						
Cognitive decline	Informant Questionnaire on Cognitive Decline in the Elderly—short form ¹⁵	P	C			
IQ estimation	Dutch version National Adult Reading Test ³¹		X			
Cognitive complaints	Self-made item: “in the previous months (before your stroke), have you experienced cognitive complaints?”	P	C			
Depressive complaints	Self-made item: “in the previous months (before your stroke), have you experienced depressive complaints?”	P	C			
Anxiety complaints	Self-made item: “in the previous months (before your stroke), have you experienced anxiety complaints?”	P	C			
Current comorbid complaints						
Depressive complaints	Hospital Anxiety and Depression Scale—subscale depression ³²		X	X	X	X
Anxiety complaints	Hospital Anxiety and Depression Scale—subscale anxiety ³²		X	X	X	X
Fatigue	Fatigue Assessment Scale ³³		X	X	X	X
Stress	Perceived Stress Scale, 4-item version ³⁴		X		X	X
Personal factors			X		X	X
Coping style	Utrecht Coping List—15-item version ³⁵		X			
Personality trait—neuroticism	Eysenck Personality Questionnaire Revised Short Scale—subscale neuroticism ³⁶		X			
Personality trait—extraversion	Eysenck Personality Questionnaire Revised Short Scale—Extraversion subscale ³⁶		X			
Type D	Type D scale-14 ³⁷		X			
Participants' awareness of SCC	Cognitive Failures Questionnaire completed by proxy Checklist for Cognitive and Emotional Consequences completed by proxy		X	X	X	X
Life events	Self-made item concerning the presence and impact of a positive or negative life event: “Last year, did something happen in your life which had a major impact on you? This may be something either pleasant or sad”		X		X	X

C, control group only; P, patient group only; SCC, subjective cognitive complaints; T0, clinical phase; T1, 3 months poststroke; T2, 6 months poststroke; T3, 1 year poststroke; T4, 2 years poststroke; X, instrument used in both patients and controls.

to be elderly (>60 years), the additional OLD module (WHOQOL-OLD)²³ comprising 24 items. While the first covers overall well-being on the domains ‘physical’, ‘psychological’, ‘social relationships’ and ‘environment’, the OLD module evaluates aspects of life which are specific to the elderly, including: ‘intimacy’, ‘sensory abilities’, ‘autonomy’, ‘activities in the past, present and future’, ‘social participation’ and ‘dying’.

Subjective recovery after stroke is determined by a single item from the Stroke Impact Scale,²⁴ in which patients are asked to indicate on a scale ranging from 0 (‘no recovery’) to 100 (‘full recovery’) how much they feel they have recovered from their stroke.

ADL functioning is assessed in basic activities, including self-care and mobility, using the Barthel Index²⁵ (10 items) and more complex activities like housekeeping, hobbies and employment, using the Frenchay Activities Index²⁶ (15 items).

All instruments chosen are frequently used (inter) nationally in research and daily clinical practice dealing with stroke patients.

Possible determinants

Depending on the specific outcome considered, SCC, OCP, QoL, subjective recovery and ADL functioning are either dependent or independent variables. A wide range

of possible determinants are additionally taken into account, based on what is currently known from the literature on SCC in the general and the stroke population. These include: demographic variables, clinical characteristics (those related to stroke included) and health status; premorbid status (ie, cognitive decline, IQ, cognitive and emotional complaints); comorbid complaints about mood (ie, anxiety and depression), fatigue and stress; personal factors (ie, coping style, personality traits and SCC awareness), and the occurrence and impact of positive and/or negative life events. See [table 2](#) for the specific variables assessed and instruments used.

Planned statistical analyses

Cross-sectional analyses will be used to evaluate group differences at each of the individual time points (T1–T4) and include: χ^2 test for categorical variables, the Mann-Whitney U test for ordinal data, and the Student t test or (multivariate) analysis of variances ((M)ANOVA) for continuous dependent variables. Furthermore, differences across the time points will be analysed using multilevel analysis, which allows inclusion of all available data (ie, also those from participants with partly missing values).

The course of SCC over time (T1–T4) will subsequently be evaluated using latent class growth analysis. We will explore whether groups with different trajectories of SCC over time can be distinguished and, if so, what their characteristics are.

The predictive value of the determinants for the primary and secondary outcome measures (ie, SCC, OCP, QoL, subjective recovery and ADL functioning) at T3 and T4 will be determined using multivariate regression analysis. Potential predictors are defined as variables with at least a marginally significant association ($p < 0.10$) with the outcome. Only these variables will be included in the subsequent regression analyses to determine the most important predictors. In general, effects with a two-tailed $p < 0.05$ are considered statistically significant.

Sample size and power calculation

The sample size needed in the COMPAS study is calculated using the method for multilevel analysis according to Twisk.³⁸ Based on a high intraindividual correlation across the different time points ($r = 0.70$), an α level of 0.05 and power of 0.80, there are 180 participants per group needed to be able to detect a small difference (at least 0.2 SD) between the groups. We expect about 40% drop-outs during the 2-year follow-up period due to mortality, comorbidity or refusal to continue participation. Therefore, we aim to include 300 participants at baseline in each group in order to end up with the 180/group needed.

ETHICS AND DISSEMINATION

Ethical considerations

The COMPAS study is conducted in accordance with the 'Helsinki Declaration' (Seoul revision, 2008) and the

'medical Research Involving Human Subjects Act' (WMO). The study is non-invasive, imposes no risk on participants, and its protocol has been approved by the medical ethical committees of all participating hospitals (ie, St. Elisabeth and TweeSteden Hospitals in Tilburg, and the Maxima Medical Centre in Veldhoven), and has been registered by the Central committee on Research Involving Human Subjects (number NL31208.008.10). Furthermore, written informed consent is obtained from all participants.

Dissemination

The results obtained will be disseminated to the scientific, medical and general public by publication in national and international peer-reviewed journals, as well as by presentations in conferences and meetings with clinicians dealing with patients with stroke.

DISCUSSION

The COMPAS study is the first in which poststroke SCC are systematically evaluated over time, while a wide range of subjective and objective variables in patients and controls is taken into account. While numerous studies have measured poststroke OCP, only a few have also evaluated the patients' SCC. Also, while these symptoms are found to be common among patients with stroke, knowledge about their risk factors, their course over time, differences with the non-stroke population and their predictive value for future functioning is practically non-existent.

Strong elements of the COMPAS study are its prospective design with multiple assessments during the first two years after stroke, and the extensive evaluations of subjective and objective variables, which, based on the current literature, are potentially relevant to SCC after stroke. This gives us the opportunity to determine a detailed risk profile for experiencing poststroke SCC. Furthermore, the instruments chosen are widely accepted and frequently used in daily clinical practice dealing with patients with stroke. Traditional neuropsychological and more ecologically valid tests (eg, the Rivermead Behavioural Memory Test) are used to evaluate OCP, making it possible to determine whether the ecological validity of tests affects the association between SCC and OCP. Also, a healthy control group is assessed at the same time points as the patients and will be used as a reference group. This enables us to distinguish poststroke SCC in their prevalence, profile and time course from, for example, factors which are associated with ageing. A potential limitation of the study is that the most seriously affected patients with stroke are unable to participate, thereby reducing the possibility to generalise the results to the stroke population as a whole. However, our study differs from those carried out until now in this field in that we include a broad selection of patients with stroke, not only first-ever strokes or patients discharged home.

In conclusion, we feel that the COMPAS study has the potential to contribute to knowledge on poststroke SCC. Owing to the ageing population and healthcare improvements, the number of stroke survivors who will have to deal with poststroke impairment will increase in the future, and the social and economic burdens will rise accordingly.^{39 40} Clinicians are frequently confronted with patients having SCC after their stroke, but the meaning and relevance of these SCC have yet to be determined. We aim to elucidate the possible risk factors, prognosis and the predictive value of poststroke SCC. This information can subsequently be applied by clinicians in daily practice in order to more accurately inform patients and their proxies and to treat SCC. Our data may also prove useful in the future development of patient-tailored intervention programmes to ultimately improve individual stroke patient-centred care, which is the ultimate aim of the COMPAS study.

Author affiliations

¹Department of Cognitive Neuropsychology, Centre of Research on Psychology in Somatic diseases (CoRPS), Tilburg University, Tilburg, The Netherlands

²Department of Neurology, St Elisabeth Hospital, Tilburg, The Netherlands

Contributors RM, PK and MR conceptualised the study. MS and RM contributed to the procurement of funding. MR, RM and PK developed procedures for implementing the protocol. All authors contributed to and have checked the final manuscript.

Funding This study is financially supported by the Centre of Research on Psychology in Somatic diseases (CoRPS) from Tilburg University, The Netherlands.

Competing interests None.

Ethics approval Medical Ethics Committee St Elisabeth Hospital, TweeSteden Hospital in Tilburg, and Maxima Medical Centre in Veldhoven.

Provenance and peer review Not commissioned; internally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 3.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/3.0/>

REFERENCES

- Lamb F, Anderson J, Saling M, *et al*. Predictors of subjective cognitive complaint in post-acute older adult stroke patients. *Arch Phys Med Rehabil* 2013;94:1747–52.
- Winkens I, Van Heugten CM, Fasotti L, *et al*. Reliability and validity of two new instruments for measuring aspects of mental slowness in the daily lives of stroke patients. *Neuropsychol Rehabil* 2009;19:64–85.
- van Rijnsbergen MW, Mark RE, de Kort PL, *et al*. Subjective cognitive complaints after stroke: a systematic review. *J Stroke Cerebrovasc Dis*. Published Online First: 22 June 2013. doi:10.1016/j.jstrokecerebrovasdis.2013.05.003
- Duits A, Munnecom T, van Heugten C, *et al*. Cognitive complaints in the early phase after stroke are not indicative of cognitive impairment. *J Neurol Neurosurg Psychiatry* 2008;79:143–6.
- van Heugten C, Rasquin S, Winkens I, *et al*. Checklist for cognitive and emotional consequences following stroke (CLCE-24): development, usability and quality of the self-report version. *Clin Neurol Neurosurg* 2007;109:257–62.
- Wilz G, Barskova T. Predictors of psychological and somatic components of poststroke depression: a longitudinal study. *Top Stroke Rehabil* 2007;14:25–40.
- Jonker C, Geerlings MI, Schmand B. Are memory complaints predictive for dementia? A review of clinical and population-based studies. *Int J Geriatr Psychiatry* 2000;15:983–91.
- Reid LM, MacLulich AM. Subjective memory complaints and cognitive impairment in older people. *Dement Geriatr Cogn Disord* 2006;22:471–85.
- Comijs HC, Deeg DJ, Dik MG, *et al*. Memory complaints; the association with psycho-affective and health problems and the role of personality characteristics. A 6-year follow-up study. *J Affect Disord* 2002;72:157–65.
- Jorm AF, Butterworth P, Anstey KJ, *et al*. Memory complaints in a community sample aged 60–64 years: associations with cognitive functioning, psychiatric symptoms, medical conditions, APOE genotype, hippocampus and amygdala volumes, and white-matter hyperintensities. *Psychol Med* 2004;34:1495–506.
- Paradise MB, Glozier NS, Naismith SL, *et al*. Subjective memory complaints, vascular risk factors and psychological distress in the middle-aged: a cross-sectional study. *BMC Psychiatry* 2011; 11:108.
- Mol M, Carpay M, Ramakers I, *et al*. The effect of perceived forgetfulness on quality of life in older adults; a qualitative review. *Int J Geriatr Psychiatry* 2007;22:393–400.
- Berg A, Palomaki H, Lonnqvist J, *et al*. Depression among caregivers of stroke survivors. *Stroke* 2005;36:639–43.
- van Exel NJ, Koopmanschap MA, van den Berg B, *et al*. Burden of informal caregiving for stroke patients. Identification of caregivers at risk of adverse health effects. *Cerebrovasc Dis* 2005;19:11–17.
- de Jonghe JF, Schmand B, Ooms ME, *et al*. [Abbreviated form of the Informant Questionnaire on cognitive decline in the elderly]. *Tijdschr Gerontol Geriatr* 1997;28:224–9.
- Broadbent DE, Cooper PF, FitzGerald P, *et al*. The Cognitive Failures Questionnaire (CFQ) and its correlates. *Br J Clin Psychol* 1982;21(Pt 1):1–16.
- Ponds R, van Boxtel M, Jolles J. [The 'Cognitive Failure Questionnaire' as a measure of subjective cognitive functioning]. *Tijdschrift voor neuropsychologie—diagnostiek, behandeling en onderzoek* 2006;1:37–45.
- Osterrieth PA. [The test of copying a complex figure: a contribution to the study of perception and memory]. *Archives de Psychologie* 1944;30:206–353.
- Wilson B, Cockburn J, Baddeley A, *et al*. The development and validation of a test battery for detecting and monitoring everyday memory problems. *J Clin Exp Neuropsychol* 1989;11:855–70.
- Spreen O, Strauss E. *A compendium of neuropsychological tests administration, norms and commentary*. 2nd edn. New York: Oxford University Press, 1998.
- Lezak MD, Howieson DB, Loring DW, *et al*. *Neuropsychological assessment*. 4th edn. New York: Oxford, 2004.
- Skevington SM, Lofly M, O'Connell KA. The World Health Organization's WHOQOL-BREF quality of life assessment: psychometric properties and results of the international field trial. A report from the WHOQOL Group. *Qual Life Res* 2004; 13:299–310.
- Power M, Quinn K, Schmidt S. Development of the WHOQOL-old module. *Qual Life Res* 2005;14:2197–214.
- Williams LS, Weinberger M, Harris LE, *et al*. Development of a stroke-specific quality of life scale. *Stroke* 1999;30:1362–9.
- de Haan R, Limburg M, Schuling J, *et al*. [Clinimetric evaluation of the Barthel Index, a measure of limitations in activities of daily living]. *Ned Tijdschr Geneesk* 1993;137:917–21.
- Holbrook M, Skilbeck CE. An activities index for use with stroke patients. *Age Ageing* 1983;12:166–70.
- Bamford J, Sandercock P, Dennis M, *et al*. Classification and natural history of clinically identifiable subtypes of cerebral infarction. *Lancet* 1991;337:1521–6.
- Brott T, Adams HP Jr., Olinger CP, *et al*. Measurements of acute cerebral infarction: a clinical examination scale. *Stroke* 1989;20:864–70.
- Linn BS, Linn MW, Gurel L. Cumulative illness rating scale. *J Am Geriatr Soc* 1968;16:622–6.
- Ware J Jr., Kosinski M, Keller SD. A 12-Item Short-Form Health Survey: construction of scales and preliminary tests of reliability and validity. *Med Care* 1996;34:220–33.
- Schmand B, Bakker D, Saan R, *et al*. [The Dutch Adult Reading Test: a measure of premorbid intelligence]. *Tijdschr Gerontol Geria* 1991;22:15–19.
- Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67:361–70.
- Michielsen HJ, De Vries J, Van Heck GL. Psychometric qualities of a brief self-rated fatigue measure: The Fatigue Assessment Scale. *J Psychosom Res* 2003;54:345–52.

34. Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav* 1983;24:385–96.
35. Sanderman R, Ormel J. [The Utrecht Coping List (UCL): validity and reliability]. *Gedrag Gezond* 1992;20:32–7.
36. Sanderman R, Arrindell WA, Ranchor A, et al. [Measurement of personality traits using the Eysenck Personality Questionnaire: a manual]. Groningen: Noordelijk Centrum voor Gezondheidsvraagstukken, Rijksuniversiteit Groningen, 1995.
37. Denollet J. DS14: standard assessment of negative affectivity, social inhibition, and Type D personality. *Psychosom Med* 2005;67:89–97.
38. Twisk JWR. *Applied longitudinal data analysis for epidemiology*. New York: Cambridge University Press, 2003.
39. Donnan GA, Fisher M, Macleod M, et al. Stroke. *Lancet* 2008;371:1612–23.
40. Strong K, Mathers C, Bonita R. Preventing stroke: saving lives around the world. *Lancet Neurol* 2007;6:182–7.