

## Tilburg University

### **A personalized eHealth intervention for lifestyle changes in patients with cardiovascular disease:**

Broers, E.R.; Kop, W.J.; Denollet, J.; Widdershoven, J.W.G.M.; Wetzels, M.; Ayoola, I.; Piera-Jimenez, J.; Habibovic, M.

*Published in:*

Journal of Medical Internet Research (JMIR)

*DOI:*

[10.2196/14570](https://doi.org/10.2196/14570)

*Publication date:*

2020

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication in Tilburg University Research Portal](#)

*Citation for published version (APA):*

Broers, E. R., Kop, W. J., Denollet, J., Widdershoven, J. W. G. M., Wetzels, M., Ayoola, I., Piera-Jimenez, J., & Habibovic, M. (2020). A personalized eHealth intervention for lifestyle changes in patients with cardiovascular disease: Randomized controlled trial. *Journal of Medical Internet Research (JMIR)*, 22(5), [e14570]. <https://doi.org/10.2196/14570>

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Original Paper

# A Personalized eHealth Intervention for Lifestyle Changes in Patients With Cardiovascular Disease: Randomized Controlled Trial

Eva Rosalinde Broers<sup>1,2\*</sup>, PhD; Willem Johan Kop<sup>2\*</sup>, Prof Dr; Johan Denollet<sup>2\*</sup>, Prof Dr; Jos Widdershoven<sup>1,2\*</sup>, Prof Dr, MD; Mart Wetzels<sup>3\*</sup>, PhD; Idowu Ayoola<sup>3\*</sup>, PhD; Jordi Piera-Jimenez<sup>4\*</sup>, MSc; Mirela Habibovic<sup>1,2\*</sup>, PhD

<sup>1</sup>Department of Cardiology, Elisabeth-Tweesteden Hospital, Tilburg, Netherlands

<sup>2</sup>Department of Medical and Clinical Psychology, Tilburg University, Tilburg, Netherlands

<sup>3</sup>University of Technology Eindhoven, Eindhoven, Netherlands

<sup>4</sup>Badalona Serveis Assistencials, Badalona, Spain

\* all authors contributed equally

**Corresponding Author:**

Mirela Habibovic, PhD

Department of Medical and Clinical Psychology

Tilburg University

Tilburg

Netherlands

Phone: 31 13 466 4020

Email: [m.habibovic@tilburguniversity.edu](mailto:m.habibovic@tilburguniversity.edu)

## Abstract

**Background:** Behavior change methods involving new ambulatory technologies may improve lifestyle and cardiovascular disease outcomes.

**Objective:** This study aimed to provide proof-of-concept analyses of an intervention aiming to increase (1) behavioral flexibility, (2) lifestyle change, and (3) quality of life. The feasibility and patient acceptance of the intervention were also evaluated.

**Methods:** Patients with cardiovascular disease (N=149; mean age 63.57, SD 8.30 years; 50/149, 33.5% women) were recruited in the *Do Cardiac Health Advanced New Generation Ecosystem (Do CHANGE)* trial and randomized to the *Do CHANGE* intervention or *care as usual (CAU)*. The intervention involved a 3-month behavioral program in combination with ecological momentary assessment and intervention technologies.

**Results:** The intervention was perceived to be feasible and useful. A significant increase in lifestyle scores over time was found for both groups ( $F_{2,146,6}=9.99$ ;  $P<.001$ ), which was similar for CAU and the intervention group ( $F_{1,149,9}=0.09$ ;  $P=.77$ ). Quality of life improved more in the intervention group (mean 1.11, SD 0.11) than CAU (mean -1.47, SD 0.11) immediately following the intervention (3 months), but this benefit was not sustained at the 6-month follow-up (interaction:  $P=.02$ ). No significant treatment effects were observed for behavioral flexibility ( $F_{1,149,0}=0.48$ ;  $P=.07$ ).

**Conclusions:** The Do CHANGE 1 intervention was perceived as useful and easy to use. However, no long-term treatment effects were found on the outcome measures. More research is warranted to examine which components of behavioral interventions are effective in producing long-term behavior change.

**Trial Registration:** ClinicalTrials.gov NCT02946281; <https://www.clinicaltrials.gov/ct2/show/NCT02946281>

(*J Med Internet Res* 2020;22(5):e14570) doi: [10.2196/14570](https://doi.org/10.2196/14570)

**KEYWORDS**

cardiovascular diseases; lifestyle; habits; eHealth; mHealth

## Introduction

The elimination of modifiable behavioral risk factors for cardiovascular disease (eg, smoking and physical inactivity) in the general population could prevent 80% of adverse clinical outcomes [1]. In patients diagnosed with cardiovascular disease, a modest reduction in risk behaviors can decrease the mortality rates by approximately 50% [2]. However, recommended targets (eg, lifestyle and medication adherence) for secondary prevention are rarely reached [3]. To achieve sustained health behavior change, active interventions that go beyond patient education are needed [4].

Sustainable behavior changes can be enhanced by implementing a personalized patient-tailored approach [5,6]. New ambulatory technologies can now be used to provide personalized support in a low threshold, nonobtrusive, and ecologically valid manner. These devices can be used to provide feedback about ambulatory health behaviors (eg, physical activity levels), but they are not sufficient to produce long-term behavior change [7,8]. In the setting of cardiac rehabilitation, telemonitoring guidance for patients' physical activity levels was found to be feasible in the FIT@Home study, and this intervention resulted in higher patient satisfaction and trends toward lower health care costs, but not in better improvements in fitness or physical activity levels relative to standard center-based rehabilitation [8]. The impact of this intervention could potentially have been further improved if a more patient-tailored approach were added. Another study found initial support that an app using persuasive design techniques can improve biological and psychological factors in patients after cardiac rehabilitation [9]. It is therefore plausible that ambulatory assessments are likely to have better therapeutic effects when combined with prompts that promote health-related behaviors (ie, ecological momentary interventions). These new methodologies also require a deeper knowledge about patients' needs and preferences [10].

This trial (*Do Cardiac Health Advanced New Generation Ecosystem*, Do CHANGE) was specifically designed to examine this multidisciplinary approach to behavior change [11]. What is unique to this trial is that patients received the behavior change program, *Do Something Different* (DSD), which has been previously developed to change unhealthy habits through the increase of behavioral flexibility [11]. Behavioral flexibility is associated with a broad range of the behavioral repertoire, making people more open to experience and the adoption of new behaviors [12]. DSD has been evaluated in other patient samples and has shown promising results by producing health behavior change [13]. For this study, the program was adapted to meet the needs of patients with cardiovascular disease (coronary artery disease, CAD; heart failure, HF; and hypertension, HT). Hence, the aim of this study was to provide proof of concept for the behavioral intervention aiming to address (1) behavioral flexibility, (2) lifestyle change, and (3) quality of life. The feasibility and patient acceptance of the intervention were also evaluated.

## Methods

### Design

The Do CHANGE trial is an international (the Netherlands and Spain), multicenter, randomized controlled trial, designed to enhance lifestyle changes in patients with cardiac disease (NCT02946281). The trial findings described in this paper are the first (proof of concept and feasibility) phase of the Do CHANGE project (phase 1) and will serve as input for further development of a second phase of this randomized controlled trial (Do CHANGE, phase 2; NCT03178305). A detailed description of both phases of the Do CHANGE trial has been published previously [11]. As this trial was developed to provide information about proof of concept and feasibility, an a priori sample size calculation was not performed. For this phase, we aimed to include 150 patients across 2 countries, which is considered sufficient to give information about proof of concept and feasibility of the intervention.

### Study Sample

Patients diagnosed with CAD (having experienced a myocardial infarction, percutaneous coronary intervention, angina pectoris, or coronary artery bypass graft surgery), symptomatic HF (New York Heart Association class I-IV), and HT were included in the study. HT was defined as systolic blood pressure  $\geq 140$  mm Hg and/or diastolic blood pressure  $\geq 90$  mm Hg on two different measurements spaced 1 to 2 min apart and after 3 to 5 min in a sitting position. The values of the second measure were used. HF patients were included if they had a diagnosis of systolic or diastolic HF and the presence of HF symptoms.

Patients were recruited at Badalona Serveis Assistencials (Badalona, Spain) and Elisabeth-TweeSteden Hospital (Tilburg, the Netherlands). The study was approved by the medical ethics committees of the participating hospitals and was conducted in accordance with the Helsinki Declaration.

The inclusion criteria were as follows: (1) a primary diagnosis of CAD, HF, or HT; (2) aged 18 to 75 years; (3) having  $\geq 2$  of the following risk factors: positive family history, increased cholesterol, smoking, diabetes, sedentary lifestyle, and/or psychosocial risk factors; (4) sufficient knowledge of the country's native language; (5) access to the internet at home; and (6) having a smartphone compatible with the apps used in the study.

The exclusion criteria were as follows: (1) life expectancy  $< 1$  year, (2) life-threatening comorbidities (eg, malignancy), (3) history of psychiatric illness other than anxiety and/or depression, (4) significant cognitive impairments (eg, dementia), and (5) on the waiting list for heart transplantation.

### Procedure

Patients meeting the inclusion criteria were approached for participation by a cardiologist or cardiac nurse. If interested, patients received information about the study in writing and orally. After 10 days, patients were contacted to inquire about their participation. If the patient indicated that they wanted to participate, a face-to-face appointment was scheduled at the hospital. Patients were asked to sign an informed consent

document and were provided with the first set of questionnaires (baseline). After filling in the questionnaires, patients were randomized. Patients in the intervention group received information about the intervention and the use of associated devices (see the *Intervention* section). The following day, patients in the intervention group were contacted by telephone to check that the devices were installed correctly and that the system was functional.

Patients received the intervention for 3 months. Follow-up questionnaires were sent at 3 and 6 months. Patients returned the devices after completion of the intervention (ie, after 3 months).

### Randomization and Blinding

Patients were randomized (1:1) after completing the baseline questionnaires. Randomization sequences were computer generated and individually sealed before recruitment started. After completing the questionnaires, one sealed envelope was drawn by the research assistant containing the group allocation. Owing to the nature of the behavioral monitoring aspects of the study, blinding health care providers or participants to the treatment condition was not possible, whereas the initial analyses of the study outcomes were analyzed without knowledge of the treatment allocation.

### Intervention

#### *Do Cardiac Health Advanced New Generated Ecosystem Intervention Versus Care as Usual*

##### Behavior Change Technique

Patients randomized to the intervention group received a 3-month behavior change program, *DSD*, which was provided via text messages on patients' mobile phones. The *DSD* program that was used aims to change unhealthy habits through the increase of behavioral flexibility [12]. This is achieved by disrupting patients' daily behavioral routine for a short period (few seconds) with behavioral prompts, which are referred to as *Do's* (eg, "EXPLORE MORE DAY. Today instead of going the same old way, take a different route. Look around, spot 10 things you wouldn't see on your usual journey") and are provided through patients' mobile phones. These messages challenge patients to do something different and get out of their comfort zone. They have been developed by a multidisciplinary team, including cardiologists and psychologists, to make sure that the *Do's* apply to the patient population and are thus related to their daily behaviors/needs. Patients received a total of 32 *Do's* during the 3-month intervention period (2-3 *Do's* every week). *DSD* has been evaluated in other patient samples and has shown promising results with respect to behavior change [13]. For this trial, the program was adapted to the cardiac population with slight differences in the program depending on patients' primary diagnosis (eg, CAD, HF, HT), as the preferred health behaviors may vary depending on the diagnosis. For example, because of disease-specific symptoms, advice regarding fluid intake was taken into account within the program only for patients with HF. More details regarding the *DSD* program are provided in the previously published design paper of this project [11].

### Technological Tools

In addition to the *DSD* program, to obtain objective measures on patients' physical functioning, all patients received a blood pressure monitor, the Moves app (ProtoGeo, Helsinki; to register the GPS location), and the CarePortal (Docobo Ltd, Leatherhead; eg, a home monitoring device measuring daily symptoms and electrocardiogram). Owing to the disease-specific reasons, patients diagnosed with HF also received a weight scale, as daily weight monitoring is of importance in this subgroup. Data obtained from these devices will not be included in this analysis. This manuscript will focus on the primary outcome measures related to lifestyle parameters and patient-reported outcomes, which were derived from validated questionnaires (see the study by Habibović et al [11] for a description of primary and secondary outcomes).

### Control Group

Patients randomized to the care as usual (CAU) group received the treatment as usual and were only provided with the validated questionnaires at baseline and at 3 and 6 months. These patients did not receive devices for ambulatory monitoring measures.

### Measures

#### Questionnaires

##### Primary outcomes

##### Behavioral Flexibility

Behavioral flexibility was measured using the *DSD* questionnaire from scale items designed for this study [11]. This scale contains 30 different descriptions of behavior coupled in 15 pairs of opposites (Multimedia Appendix 1). Patients were asked at each measurement point to select the behaviors that best describe them (eg, *gentle* or *firm*). On the basis of a formula, the behavioral flexibility for each participant at each time point was calculated as outlined below.

$$100\% \times \left( \frac{\text{number of behaviors selected}}{30} + \frac{\text{number of opposite pairs selected}}{15} \right)$$

Every addition of behavior raises the score as well as when both of a pair of opposite behaviors are added. For example, definite, systematic, trusting, predictable, and unpredictable are selected. All these selected behaviors raise the flexibility score. However, because predictable and unpredictable are each other's opposites, these are added to the formula again and increase the flexibility score even more. The model interprets this seemingly contradictory behavior as evidence of flexibility: based on what a given situation demands, the person can use different reactions and thus be more flexible. The total score can range from 0 to 100. The internal consistency in this sample was considered acceptable (Cronbach alpha=.67 to .76).

##### Lifestyle

The Health-Promoting Lifestyle Profile questionnaire was administered to assess health-promoting lifestyle habits [14]. This survey evaluates whether the subjective perception of patients regarding their lifestyle is changed and consists of 52 items (eg, "Eat 6-11 servings of bread, cereal, rice, and pasta each day") in total. Each item can be answered on a 4-point Likert scale, ranging from 1 (never) to 4 (routinely). The total score can therefore range from 52 to 208, with a higher score



indicating a better lifestyle. Furthermore, the questionnaire includes 6 different subscales that each cover a health promotion lifestyle domain (ie, Physical Activity, Spiritual Growth, Health Responsibility, Interpersonal Relationships, Nutrition, and Stress Management). The internal consistency was considered as excellent in this sample (Cronbach alpha=.88 to .90).

### **Quality of Life**

To administer changes in the quality of life, participants completed the World Health Organization Quality of Life—BREF (WHOQOL-BREF) [15]. The WHOQOL-BREF is considered a reliable, generic multidimensional quality of life measure and consists of 26 items in total. Two items refer to the facet's overall quality of life and general health, whereas the abiding 24 items reflect 4 different domains (ie, physical health, psychological health, social relationships, and environment). The internal consistency in this sample was excellent (Cronbach alpha=.89 to .90).

### **Perceived Usefulness and Acceptance**

The Unified Theory of Acceptance 2 (UTAUT2) scale [16] was administered to assess the perceived usefulness and acceptance of the tools that were used in the intervention. Mean scores on 8 subscales are provided, namely, (1) Performance Expectancy, (2) Effort Expectancy, (3) Social Influence, (4) Facilitating Conditions, (5) Hedonic Motivation, (6) Habit, and (7) Behavioral Intention. The initial subscale, *Price Value*, of the UTAUT2 was not included, as the cost per individual for the intervention could not be estimated. The total score per subscale can range from 4 to 20, with a higher score indicating higher usefulness and acceptance [16]. The internal consistency in this sample was excellent (Cronbach alpha=.89).

### **Client Satisfaction Questionnaire**

To assess the satisfaction of the patients about the ecosystem, the 8-item Client Satisfaction Questionnaire [17] was used. This self-administered questionnaire is a general scale that consists of 8 Likert scale items (eg, "To what extent has our program met your needs?") ranging from 0 to 4, with response descriptors that vary. The overall score can range from 8 to 32, with a higher score indicating a higher satisfaction. The internal consistency was rated as excellent (Cronbach alpha=.92) [17].

### **Other Questionnaires Included in the Model**

#### **Type D Scale (Distress Scale-14)**

Type D personality was assessed using the Type D scale (Distress Scale-14) [18]. This 14-item questionnaire consists of 2 subscales with seven 5-point Likert scale items each, ranging from 0 (false) to 4 (true). Total scores on both subscales range from 0 to 28. The 2 subscales represent the characteristics of negative affectivity (NA; eg, the tendency to experience negative emotions across time and situations) and social inhibition (SI; eg, the tendency not to express feelings). When scoring  $\geq 10$  on both subscales, patients were classified as Type D. With a reported Cronbach alpha value of .86 and .88, respectively, the internal consistency of SI and NA are considered as satisfactory [18].

### **The 7-Item Generalized Anxiety Disorder Scale**

To gauge self-administered symptoms of anxiety, the 7-item Generalized Anxiety Disorder scale was administered [19]. The questionnaire is comprised of 7 items (eg, "Feeling afraid as if something terrible might happen") that can be answered on a 4-point Likert scale, ranging from 0 (not at all) to 3 (almost every day). To get an indication of anxiety symptom severity, the total score (range from 0 to 21) can be used. A higher score implies higher levels of anxiety. The internal consistency was considered excellent (Cronbach alpha=.92) [19].

### **Nine-Item Patient Health Questionnaire**

Depressive symptoms were administered at each time point by using the 9-item Patient Health Questionnaire (PHQ-9) [20]. This self-report questionnaire consists of 9 items in total (eg, *feeling down, depressed, or hopeless*), each evaluated on a 4-point Likert scale (ie, not at all, several days, and nearly every day). The total score ranges from 0 to 27, with a higher score as an indication of worse depression symptom severity. The internal consistency was considered excellent (Cronbach alpha=.90) [21].

### **Demographic and Clinical Data**

Demographic characteristics (eg, age, sex, marital status, working status, level of education, and smoking behavior) were obtained by patients' self-report. Clinical data (eg, comorbidities; prescribed cardiac medication; prescribed psychotropic medication; left ventricular ejection fraction; history of coronary artery bypass grafting; history of percutaneous coronary intervention; and resting heart rate and systolic and diastolic blood pressure measured at the most recent outpatient visit) were obtained from the medical record.

### **Statistical Analyses**

Categorical variables were compared using chi-square tests, and continuous variables were compared using a two-tailed *t* test for independent samples. To evaluate the treatment effectiveness, based on intention-to-treat, a univariate and multivariate Linear Mixed Model analysis was performed. Multivariable analyses were adjusted for age, sex, education, site of inclusion (Badalona Serveis Assistencials [BSA] or Elisabeth-TweeSteden Hospital [ETZ]), primary diagnosis, Charlson comorbidity index scores, Type D personality, baseline anxiety scores, and baseline depression scores. *F* values with two-sided *P* values were reported for main and interaction effects (group $\times$ time). For the estimated fixed effects, beta coefficients with two-sided *P* values were reported. Data were analyzed using the SPSS software package (version 24).

## **Results**


### **Sample**

The data collection took place between January 2017 and September 2017. In total, 286 eligible patients were approached for participation, of which 132/286 (46.1%) patients refused to participate. An additional 5/286 (1.7%) participants did not show up or declined participation. The reasons for refusal included that it would be too time consuming, they did not want to be confronted about their heart disease every day, and they

were reluctant to use technology. A total of 149 (response rate: 149/286; 52.0%) patients were enrolled. Enrollment per study site was as follows: BSA randomized a total of 74 (intervention: n=37 and CAU: n=37) patients; ETZ randomized 75 patients (intervention: n = 37 and CAU: n=38). Of the total sample, 4 participants within the CAU condition dropped out, as they did

not receive the intervention and were therefore not willing to continue. Of the patients randomized to the intervention condition, 82% (61/74) reported having completed the entire 3-month program. Overall, 97.3% (145/149) of the participants completed the follow-up assessments. **Figure 1** presents the flowchart of patient recruitment.

**Figure 1.** Flow diagram of patient recruitment.



**Baseline Characteristics**

The mean age of the total sample was 63.6 (SD 8.3) years, and 66% (99/149) were men. There were significant differences observed in mean completed education in years between the intervention group (mean 14.3, SD 6.2) and the CAU group (mean 11.8, SD 7.9;  $P=.03$ ). This means that patients in the intervention group completed more years of education than

those in the CAU group. Furthermore, a significant difference between the 2 groups was found on the mean PHQ-9 baseline scores, with a higher mean score on depressive symptoms in the CAU group (mean 3.61, SD 3.6 vs mean 5.56, SD 4.17;  $P=.003$ ). No other differences were found between the intervention and CAU groups. **Table 1** presents an overview of the baseline characteristics of this sample.

**Table 1.** Baseline patient characteristics of the total sample.

Variable	Total	Do Cardiac Health Advanced New Generation Ecosystem intervention (N=74)	Care as usual (N=75)	P value
<b>Site of allocation, n (%)</b>				
Badalona Serveis Assistencials	74 (49.7)	37 (50)	37 (50)	N/A <sup>a</sup>
Elisabeth-TweeSteden Hospital	75 (50.3)	37 (49.3)	38 (50.7)	N/A
Total	149 (100)	74 (49.7)	75 (50.3)	N/A
<b>Demographics</b>				
Age (years), mean (SD)	63.57 (8.30)	63.26 (8.35)	63.88 (8.30)	.65
Gender (male), n (%)	99 (66.4)	52 (70.3)	47 (62.7)	.42
Education (years), mean (SD)	13.03 (7.22)	14.30 (6.24)	11.79 (7.91)	.03
Marital status (partner), n (%)	118 (79.2)	61 (82.4)	57 (76.0)	.44
Working status (working), n (%)	55 (36.9)	28 (37.8)	27 (36.0)	.95
Smoking (yes), n (%)	27 (18.1)	10 (13.5)	17 (22.7)	.29
<b>Clinical</b>				
Diagnosis heart failure, n (%)	36 (24.2)	21 (28.4)	15 (20.0)	.96
Diagnosis hypertension, n (%)	73 (49.0)	38 (51.4)	35 (46.7)	>.99
Diagnosis coronary artery disease, n (%)	40 (26.8)	15 (20.3)	33 (33.3)	.07
Charlson comorbidity index, mean (SD)	1.14 (0.95)	1.01 (0.88)	1.27 (1.00)	.11
Systolic blood pressure (baseline), mean (SD)	138.02 (19.71)	135.00 (20.89)	141.00 (18.13)	.06
Diastolic blood pressure (baseline), mean (SD)	79.27 (10.01)	78.76 (10.33)	79.77 (9.87)	.54
Heart rate (rest), mean (SD)	69.41 (11.97)	69.95 (14.43)	68.88 (11.57)	.59
<b>Psychological</b>				
Patient Health Questionnaire-9, mean (SD)	4.59 (4.00)	3.61 (3.60)	5.56 (4.17)	.003
Generalized Anxiety Disorder-7, mean (SD)	4.03 (4.37)	3.35 (4.07)	4.69 (4.59)	.06
Type D personality (yes), n (%)	38 (25.5)	20 (27.0)	18 (24.0)	.67

<sup>a</sup>N/A: not applicable.

## Intervention Effects

### Behavioral Flexibility

The univariate analysis on behavioral flexibility scores (including group, time, and group×time) revealed no significant main effects for group ( $F_{1,148.93}=3.42$ ;  $P=.07$ ) or time ( $F_{2,146.82}=1.69$ ;  $P=.18$ ) and group×time interaction ( $F_{2,146.82}=1.09$ ;  $P=.34$ ). After adjusting for covariates (as previously described), main effects for time ( $F_{1,146.81}=1.74$ ;  $P=.18$ ), group ( $F_{1,149.00}=0.48$ ;  $P=.07$ ), and group×time

( $F_{2,146.81}=1.13$ ;  $P=.33$ ) remained nonsignificant (Figure 2). These findings indicate that behavioral flexibility scores did not significantly change over time, and that there were no differences between the 2 groups. With regard to covariates included in the model, the estimated fixed effects of HT ( $\beta=-6.07$ ;  $P=.01$ ) and CAD ( $\beta=-5.57$ ;  $P=.02$ ) were significantly associated with lower levels of behavioral flexibility scores (across all time points). In addition, the site of recruitment was associated with behavioral flexibility scores, with only patients from Spain showing an increase in behavioral flexibility over time ( $\beta=6.31$ ;  $P<.01$ ) when compared with those in the Netherlands (see Table 2).

Figure 2. Mean scores of intervention and care as usual group on primary outcome measures.




Figure 3. Standalone Equation 1.

$$100\% \times \left( \frac{\text{number of behaviors selected}}{30} + \frac{\text{number of opposite pairs selected}}{15} \right)$$



**Table 2.** Estimates of fixed effects from multivariable linear mixed models on the main outcome measures at baseline and at 3- and 6-month follow-up.

Multivariable linear mixed model	Behavioral flexibility			Lifestyle			Quality of life		
	Estimate	SE	P value	Estimate	SE	P value	Estimate	SE	P value
<b>Model 1: unadjusted model</b>									
Intervention group	4.70	2.07	.03	2.18	3.52	.54	0.51	1.89	.79
<b>Model 2: covariates adjusted model</b>									
Intervention group	2.91	1.99	.15	0.46	3.27	.89	-2.89	1.69	.09
Hypertension <sup>a</sup>	-6.07	2.27	.01	-3.89	3.72	.30	-3.40	2.07	.10
Coronary artery disease <sup>a</sup>	-5.57	2.29	.02	3.62	3.74	.33	-2.02	2.08	.33
Study site (Spain)	6.31	1.84	<.01	12.86	3.01	<.01	4.62	1.67	<.01
Sex (male)	2.36	1.79	.19	-1.66	2.93	.57	2.66	1.63	.11
Type D	-0.28	2.14	.90	8.33	13.47	.54	-10.85	7.48	.15
Age (years)	0.08	0.09	.42	-0.04	0.16	.80	0.07	0.09	.45
Higher Charlson comorbidity index	-1.73	1.00	.08	-1.09	1.63	.51	-1.76	0.90	.05
Higher education	-0.01	0.12	.95	0.08	0.19	.66	0.25	0.11	.02
Anxiety	-0.12	0.44	.79	-1.38	0.72	.06	-0.90	0.40	.03
Depression	-1.77	1.13	.12	-4.17	1.86	.03	-3.63	1.03	<.01

<sup>a</sup>Compared with the main diagnosis of heart failure.

### Lifestyle

A univariate analysis showed no significant effect for group ( $F_{1,434.91}=0.91$ ;  $P=.34$ ) or group $\times$ time ( $F_{2,282.73}=0.39$ ;  $P=.68$ ). These findings present that, without the addition of possible confounding variables, the intervention and CAU group did not differ. In addition, no interaction effect between allocation to group and time was found. However, a significant improvement for both groups on overall reported lifestyle behavior was found ( $F_{2,282.73}=4.28$ ;  $P=.02$ ). When adjusting for covariates in the multivariable analysis, this improvement remained significant ( $F_{2,146.63}=9.99$ ;  $P<.001$ ). As shown in [Figure 2](#), both groups reported improvements in lifestyle behavior over time. The effects of interaction ( $F_{2,147.02}=1.36$ ;  $P=.26$ ) and allocation to group ( $F_{1,149.90}=0.09$ ;  $P=.77$ ) remained nonsignificant in the adjusted models. This indicated that no effect of the intervention on healthy lifestyle behavior was found. The estimated fixed effect of depression ( $\beta=-4.17$ ;  $P=.03$ ) was negatively associated with lifestyle promoting behavior, indicating that patients who score higher on the depression scale report lower healthy lifestyle behaviors. Patients from Spain showed an increase in lifestyle behavior ( $\beta=12.86$ ;  $P<.01$ ), in comparison with those in the Netherlands (see [Table 2](#)).

### Quality of Life

The results of the univariate analysis of the quality of life total scores showed an interaction effect between time and group ( $F_{2,146.40}=4.22$ ;  $P=.02$ ). This finding indicates that the mean scores on quality of life of the intervention and CAU groups have different slopes over time: the intervention group showed a small improvement in the quality of life after 3 months, whereas the CAU group reported a small decline in the quality

of life (mean improvement 1.11, SD .11 vs mean -1.47, SD .11). Both groups stabilized to baseline level after 6 months. The interaction effect remained significant after adding the covariates in the multilevel analysis ( $F_{2,146.52}=4.29$ ;  $P=.02$ ; [Figure 2](#)), suggesting a significant, positive effect of the intervention on self-reported quality of life in the first 3 months. The estimated fixed effects of higher levels of education ( $\beta=.25$ ;  $P=.02$ ) and being recruited in Spain ( $\beta=4.62$ ;  $P=.008$ ) compared with those in the Netherlands were significantly associated with higher scores on quality of life. Lower scores were predicted by higher CCI scores ( $\beta=-1.76$ ;  $P=.05$ ) anxiety ( $\beta=-0.90$ ;  $P=.03$ ), and depression ( $\beta=-3.63$ ;  $P<.01$ ; see [Table 2](#)). Examining subscales of the WHOQOL revealed no specific subscale differences regarding response patterns to the intervention.

### Acceptability and Satisfaction

Overall, patients in the intervention group indicated being satisfied with the intervention (mean 26.22, SD 4.82). The intervention was perceived to be useful (mean 13.88, SD 3.96) and easy to use (mean 17.07, SD 2.57). Patients did not feel social pressure to use the devices from the intervention (mean 9.85, SD 3.63) and reported to be quite satisfied with the possibilities to receive support (mean 15.44, SD 2.43) and had a neutral opinion regarding the pleasure in using the devices offered in the intervention (mean 10.63, SD 2.44). Furthermore, the intervention was integrated relatively well in patients' lives (mean 11.71, SD 3.05). However, patients indicated that they were neutral regarding the intention to use the ecosystem in the future (mean 8.40, SD 3.34).

## Discussion

### Principal Findings

This study aimed to provide proof of concept for the Do CHANGE behavioral intervention targeting behavioral flexibility, lifestyle change, and quality of life in cardiac patients. No significant differences between the groups were observed on behavioral flexibility and lifestyle. However, a small increase in quality of life at 3 months was observed in the intervention group, but at 6 months, no significant difference between the groups was observed. With respect to the usefulness and feasibility of the intervention, the findings of this study revealed that the ecosystem is experienced as useful, easy to use, and integrated well into the daily life of the patients. It made the participants more aware of the fact that they must undertake activities themselves to feel better. Patients also reported feeling more *safe* because health care professionals were watching along. Nonadherence is a common issue in Web-based interventions for promoting health-related behavior, and the average study results in only 50% of participants adhering to the intended intervention [22]. However, 82.4% of the patients participating in this Do CHANGE intervention condition completed the intervention, which may further indicate that the intervention was not perceived as demanding.

The findings of the study are not completely in line with previous studies in other patient populations [13]. An explanation for this discrepancy could be the fact that this was the first study implementing the concept of behavioral flexibility and thus the core Do's of the DSD program in the cardiac population. In addition, the Do's might not have been tailored enough to the patients' needs that the timing of the Do's might not have been optimal. For example, one would want a patient to receive a distractive Do at the time when the *unwanted* behavior occurs. In this trial, patients from 2 different cultures (Spain and the Netherlands) and diagnosed with different cardiac disorders (ie, HT, HF, or CAD) were enrolled. This reflects the heterogeneity of the sample, which may have affected the results. Another important area for future research is the exploration of the mediating factors that drive the interplay between behavioral flexibility, lifestyle factors, and quality of life in the setting to electronic health (eHealth) interventions.

Enrollment in the study may have increased the general awareness of lifestyle change in both groups. This awareness could unknowingly lead to the adaptation of a lifestyle, independent of the allocation to a group. Previous research in cardiac patients affirms that there is a relationship between general knowledge about cardiac risk factors and self-reported lifestyle changes in the short term [23]. Furthermore, although lifestyle change is crucial in the treatment of cardiovascular disease, there is a lack of emphasis on lifestyle change and self-care of the patient in the current health care systems [24]. Addressing self-care and lifestyle change in clinical practice is therefore warranted. The Do CHANGE trial provides (longitudinal and momentary) knowledge that can be used in the further development of personalized interventions that will help patients reach recommended lifestyle goals.

Behavioral flexibility is an important construct on which behavioral change can possibly be initiated. However, the results of this study may indicate the need for a better measurement tool, as the questionnaire that is used might not be sensitive enough to reveal significant alteration in patients' behavioral flexibility over time.

The findings on quality of life, on the other hand, are not entirely in line with previous studies in cardiac samples, which have shown that there is a decline in quality of life and generally a slight increase in anxiety and depression scores within the 3 months postcardiac event [25]. This could be explained by patients having to adapt to new behaviors after visiting the hospital and being reminded of the fact that they have a chronic illness. In this study, the intervention group received the behavioral program, which could have contributed to first an increase in their quality of life, with a slight decrease after 3 months, sustaining their baseline quality of life. After the behavioral intervention ends, the quality of life in the intervention group also goes down as the additional *support* is no longer provided.

The findings of this study must be interpreted in light of a few limitations. At baseline, the intervention group and CAU group showed some differences in mean years of completed education and mean scores on depressive symptoms. The CAU group scored significantly higher on both variables. Another limitation of this study was that the sample was rather small, in relatively good health, and clinically heterogeneous, which may have limited the possibility to find substantial effects. Although the intervention was positively evaluated by participants, half of the approached patients did refuse participation. Therefore, it can be concluded that eHealth interventions similar to those described in this study are appealing for certain subgroups of patients. Future research should focus on eHealth interventions within the cardiac population based on a larger sample with significant power that is assessed over a prolonged follow-up duration (eg, beyond 6 months) to draw firm conclusions on sustainable behavior change. The results of this study showed that depression was associated with negative behavioral and psychological outcomes, which is in line with previous findings [26]. Depressive symptoms are common in patients with cardiovascular disease [27-29] and are related to various behavioral risk factors (eg, sedentary lifestyle, unhealthy diet, alcohol overconsumption, and smoking) [30,31]; this may explain the relation between depression and lower lifestyle behavior scores. Hence, future research is needed to examine which psychological and clinical factors contributing to health behavior change and potentially address these factors during the intervention.

For clinical practice, it is important to acknowledge that technology and eHealth solutions might be the feasible way forward in meeting patient needs and initiated health behavior change. However, the findings of this study underline the importance of a personalized approach that includes the assessment of a patient's demographic, clinical, and psychological profile.

## Conclusions

In conclusion, the Do CHANGE 1 intervention was perceived as useful and easy to use. However, no main effects were found

on behavioral flexibility, lifestyle behavior, and quality of life. More research is warranted to examine which components of behavioral interventions, and in which patients, are effective in producing long-term behavior change.

## Acknowledgments

The authors would like to thank all the patients for participating and the health care professionals in the participating hospitals for making this work possible. In addition, the authors would like to thank all the students/research assistants for their help with the patient recruitment and data management. Finally, the authors want to thank the consortium members of the Do CHANGE project. This study was funded by the European Commission's Horizon 2020 program (grant number: 463735). For this project, the Do CHANGE team received funding for research and innovation from the European Union. One startup (Onmi) and 2 small- and medium-sized enterprises (DSD, Docobo Ltd) are supported financially to develop their products.

## Conflicts of Interest

None declared.

## Multimedia Appendix 1

Consolidated Standards of Reporting Trials of Electronic and Mobile Health Applications and onLine TeleHealth checklist. [[PNG File , 73 KB-Multimedia Appendix 1](#)]

## Multimedia Appendix 2

CONSORT-EHEALTH checklist (V 1.6.1). [[PDF File \(Adobe PDF File\), 507 KB-Multimedia Appendix 2](#)]

## References

1. Piepoli M, Hoes A, Agewall S, Albus C, Brotons C, Catapano A, ESC Scientific Document Group. 2016 European Guidelines on cardiovascular disease prevention in clinical practice: The Sixth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of 10 societies and by invited experts) Developed with the special contribution of the European Association for Cardiovascular Prevention & Rehabilitation (EACPR). *Eur Heart J* 2016 Aug 1;37(29):2315-2381. [doi: [10.1093/eurheartj/ehw106](https://doi.org/10.1093/eurheartj/ehw106)] [Medline: [27222591](https://pubmed.ncbi.nlm.nih.gov/27222591/)]
2. Chow CK, Jolly S, Rao-Melacini P, Fox KA, Anand SS, Yusuf S. Association of diet, exercise, and smoking modification with risk of early cardiovascular events after acute coronary syndromes. *Circulation* 2010 Feb 16;121(6):750-758. [doi: [10.1161/CIRCULATIONAHA.109.891523](https://doi.org/10.1161/CIRCULATIONAHA.109.891523)] [Medline: [20124123](https://pubmed.ncbi.nlm.nih.gov/20124123/)]
3. Kotseva K, Wood D, de Bacquer D, de Backer G, Rydén L, Jennings C, EUROASPIRE Investigators. EUROASPIRE IV: A European Society of Cardiology survey on the lifestyle, risk factor and therapeutic management of coronary patients from 24 European countries. *Eur J Prev Cardiol* 2016 Apr;23(6):636-648. [doi: [10.1177/2047487315569401](https://doi.org/10.1177/2047487315569401)] [Medline: [25687109](https://pubmed.ncbi.nlm.nih.gov/25687109/)]
4. Charlson M, Wells M, Peterson J, Boutin-Foster C, Ogedegbe G, Mancuso C, et al. Mediators and moderators of behavior change in patients with chronic cardiopulmonary disease: the impact of positive affect and self-affirmation. *Transl Behav Med* 2014 Mar;4(1):7-17 [FREE Full text] [doi: [10.1007/s13142-013-0241-0](https://doi.org/10.1007/s13142-013-0241-0)] [Medline: [24653772](https://pubmed.ncbi.nlm.nih.gov/24653772/)]
5. Habibović M, Burg M, Pedersen S. Behavioral interventions in patients with an implantable cardioverter defibrillator: lessons learned and where to go from here? *Pacing Clin Electrophysiol* 2013 May;36(5):578-590 [FREE Full text] [doi: [10.1111/pace.12108](https://doi.org/10.1111/pace.12108)] [Medline: [23438053](https://pubmed.ncbi.nlm.nih.gov/23438053/)]
6. Minich DM, Bland JS. Personalized lifestyle medicine: relevance for nutrition and lifestyle recommendations. *ScientificWorldJournal* 2013;2013:129841 [FREE Full text] [doi: [10.1155/2013/129841](https://doi.org/10.1155/2013/129841)] [Medline: [23878520](https://pubmed.ncbi.nlm.nih.gov/23878520/)]
7. Young L, Buse J, Weaver M, Vu M, Mitchell C, Blakeney T, Monitor Trial Group. Glucose self-monitoring in non-insulin-treated patients with type 2 diabetes in primary care settings: a randomized trial. *JAMA Intern Med* 2017 Jul 1;177(7):920-929 [FREE Full text] [doi: [10.1001/jamainternmed.2017.1233](https://doi.org/10.1001/jamainternmed.2017.1233)] [Medline: [28600913](https://pubmed.ncbi.nlm.nih.gov/28600913/)]
8. Kraal JJ, van den Akker-van Marle M, Abu-Hanna A, Stut W, Peek N, Kemps HM. Clinical and cost-effectiveness of home-based cardiac rehabilitation compared to conventional, centre-based cardiac rehabilitation: Results of the FIT@Home study. *Eur J Prev Cardiol* 2017 Aug;24(12):1260-1273 [FREE Full text] [doi: [10.1177/2047487317710803](https://doi.org/10.1177/2047487317710803)] [Medline: [28534417](https://pubmed.ncbi.nlm.nih.gov/28534417/)]
9. Sankaran S, Dendale P, Coninx K. Evaluating the impact of the hearthab app on motivation, physical activity, quality of life, and risk factors of coronary artery disease patients: Multidisciplinary crossover study. *JMIR Mhealth Uhealth* 2019 Apr 4;7(4):e10874 [FREE Full text] [doi: [10.2196/10874](https://doi.org/10.2196/10874)] [Medline: [30946021](https://pubmed.ncbi.nlm.nih.gov/30946021/)]

10. Kangovi S, Asch DA. Behavioral phenotyping in health promotion: embracing or avoiding failure. *J Am Med Assoc* 2018 May 22;319(20):2075-2076 [FREE Full text] [doi: [10.1001/jama.2018.2921](https://doi.org/10.1001/jama.2018.2921)] [Medline: [29710244](https://pubmed.ncbi.nlm.nih.gov/29710244/)]
11. Habibović M, Broers E, Piera-Jimenez J, Wetzels M, Ayoola I, Denollet J, et al. Enhancing lifestyle change in cardiac patients through the do CHANGE system ('Do cardiac health: advanced new generation ecosystem'): randomized controlled trial protocol. *JMIR Res Protoc* 2018 Feb 8;7(2):e40 [FREE Full text] [doi: [10.2196/resprot.8406](https://doi.org/10.2196/resprot.8406)] [Medline: [29422454](https://pubmed.ncbi.nlm.nih.gov/29422454/)]
12. Pine K, Fletcher B. Time to shift brain channels to bring about effective changes in health behaviour. *Perspect Public Health* 2014 Jan;134(1):16-17. [doi: [10.1177/1757913913514705](https://doi.org/10.1177/1757913913514705)] [Medline: [24395840](https://pubmed.ncbi.nlm.nih.gov/24395840/)]
13. Fletcher B, Hanson J, Page N, Pine K. FIT – Do something different. A new behavioral program for sustained weight loss. *Swiss J Psychol* 2011;70(1):25-34. [doi: [10.1024/1421-0185/a000035](https://doi.org/10.1024/1421-0185/a000035)]
14. Walker SN, Sechrist KR, Pender NJ. Deep Blue. 1995. Health Promotion Model - Instruments to Measure Health Promoting Lifestyle : Health-Promoting Lifestyle Profile [HPLP II] URL: <https://deepblue.lib.umich.edu/handle/2027.42/85349> [accessed 2018-06-30]
15. -. Development of the World Health Organization WHOQOL-BREF quality of life assessment. *The WHOQOL Group. Psychol Med* 1998 May;28(3):551-558. [doi: [10.1017/s0033291798006667](https://doi.org/10.1017/s0033291798006667)] [Medline: [9626712](https://pubmed.ncbi.nlm.nih.gov/9626712/)]
16. Venkatesh V, Thong JY, Xu X. Consumer acceptance and use of information technology: extending the unified theory of acceptance and use of technology. *Manag Inf Syst Q* 2012;36(1):157-178. [doi: [10.2307/41410412](https://doi.org/10.2307/41410412)]
17. Larsen DL, Attkisson C, Hargreaves WA, Nguyen TD. Assessment of client/patient satisfaction: development of a general scale. *Eval Program Plann* 1979;2(3):197-207. [doi: [10.1016/0149-7189\(79\)90094-6](https://doi.org/10.1016/0149-7189(79)90094-6)] [Medline: [10245370](https://pubmed.ncbi.nlm.nih.gov/10245370/)]
18. Denollet J. DS14: standard assessment of negative affectivity, social inhibition, and Type D personality. *Psychosom Med* 2005;67(1):89-97. [doi: [10.1097/01.psy.0000149256.81953.49](https://doi.org/10.1097/01.psy.0000149256.81953.49)] [Medline: [15673629](https://pubmed.ncbi.nlm.nih.gov/15673629/)]
19. Spitzer RL, Kroenke K, Williams JB, Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med* 2006 May 22;166(10):1092-1097. [doi: [10.1001/archinte.166.10.1092](https://doi.org/10.1001/archinte.166.10.1092)] [Medline: [16717171](https://pubmed.ncbi.nlm.nih.gov/16717171/)]
20. Kroenke K, Spitzer R, Williams J. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001 Sep;16(9):606-613 [FREE Full text] [doi: [10.1046/j.1525-1497.2001.016009606.x](https://doi.org/10.1046/j.1525-1497.2001.016009606.x)] [Medline: [11556941](https://pubmed.ncbi.nlm.nih.gov/11556941/)]
21. Stafford L, Berk M, Jackson HJ. Validity of the Hospital Anxiety and Depression Scale and Patient Health Questionnaire-9 to screen for depression in patients with coronary artery disease. *Gen Hosp Psychiatry* 2007;29(5):417-424. [doi: [10.1016/j.genhosppsy.2007.06.005](https://doi.org/10.1016/j.genhosppsy.2007.06.005)] [Medline: [17888808](https://pubmed.ncbi.nlm.nih.gov/17888808/)]
22. Kelders SM, Kok RN, Ossebaard HC, van Gemert-Pijnen JE. Persuasive system design does matter: a systematic review of adherence to web-based interventions. *J Med Internet Res* 2012 Nov 14;14(6):e152 [FREE Full text] [doi: [10.2196/jmir.2104](https://doi.org/10.2196/jmir.2104)] [Medline: [23151820](https://pubmed.ncbi.nlm.nih.gov/23151820/)]
23. Alm-Roijer C, Fridlund B, Stagmo M, Erhardt L. Knowing your risk factors for coronary heart disease improves adherence to advice on lifestyle changes and medication. *J Cardiovasc Nurs* 2006;21(5):E24-E31. [doi: [10.1097/00005082-200609000-00015](https://doi.org/10.1097/00005082-200609000-00015)] [Medline: [16966907](https://pubmed.ncbi.nlm.nih.gov/16966907/)]
24. Riegel B, Moser D, Buck H, Dickson V, Dunbar S, Lee C, American Heart Association Council on Cardiovascular and Stroke Nursing; Council on Peripheral Vascular Disease; Council on Quality of Care and Outcomes Research. Self-care for the prevention and management of cardiovascular disease and stroke: a scientific statement for healthcare professionals from the American Heart Association. *J Am Heart Assoc* 2017 Aug 31;6(9):pii: e006997 [FREE Full text] [doi: [10.1161/JAHA.117.006997](https://doi.org/10.1161/JAHA.117.006997)] [Medline: [28860232](https://pubmed.ncbi.nlm.nih.gov/28860232/)]
25. Habibović M, Denollet J, Cuijpers P, van der Voort PH, Herrman J, Bouwels L, et al. Web-based distress management for implantable cardioverter defibrillator patients: a randomized controlled trial. *Health Psychol* 2017 Apr;36(4):392-401. [doi: [10.1037/hea0000451](https://doi.org/10.1037/hea0000451)] [Medline: [28192003](https://pubmed.ncbi.nlm.nih.gov/28192003/)]
26. Kashdan TB, Rottenberg J. Psychological flexibility as a fundamental aspect of health. *Clin Psychol Rev* 2010 Nov;30(7):865-878 [FREE Full text] [doi: [10.1016/j.cpr.2010.03.001](https://doi.org/10.1016/j.cpr.2010.03.001)] [Medline: [21151705](https://pubmed.ncbi.nlm.nih.gov/21151705/)]
27. Huffman JC, Celano CM, Beach SR, Motiwala SR, Januzzi JL. Depression and cardiac disease: epidemiology, mechanisms, and diagnosis. *Cardiovasc Psychiatry Neurol* 2013;2013:695925 [FREE Full text] [doi: [10.1155/2013/695925](https://doi.org/10.1155/2013/695925)] [Medline: [23653854](https://pubmed.ncbi.nlm.nih.gov/23653854/)]
28. Lichtman JH, Froelicher ES, Blumenthal JA, Carney RM, Doering LV, Frasure-Smith N, American Heart Association Statistics Committee of the Council on Epidemiology and Prevention and the Council on Cardiovascular and Stroke Nursing. Depression as a risk factor for poor prognosis among patients with acute coronary syndrome: systematic review and recommendations: a scientific statement from the American Heart Association. *Circulation* 2014 Mar 25;129(12):1350-1369. [doi: [10.1161/CIR.000000000000019](https://doi.org/10.1161/CIR.000000000000019)] [Medline: [24566200](https://pubmed.ncbi.nlm.nih.gov/24566200/)]
29. Freedland KE, Rich MW, Skala JA, Carney RM, Dávila-Román VG, Jaffe AS. Prevalence of depression in hospitalized patients with congestive heart failure. *Psychosom Med* 2003;65(1):119-128. [doi: [10.1097/01.psy.0000038938.67401.85](https://doi.org/10.1097/01.psy.0000038938.67401.85)] [Medline: [12554823](https://pubmed.ncbi.nlm.nih.gov/12554823/)]
30. Sin NL, Kumar AD, Gehi AK, Whooley MA. Direction of association between depressive symptoms and lifestyle behaviors in patients with coronary heart disease: the heart and soul study. *Ann Behav Med* 2016 Aug;50(4):523-532 [FREE Full text] [doi: [10.1007/s12160-016-9777-9](https://doi.org/10.1007/s12160-016-9777-9)] [Medline: [26817654](https://pubmed.ncbi.nlm.nih.gov/26817654/)]
31. Penninx BW. Depression and cardiovascular disease: Epidemiological evidence on their linking mechanisms. *Neurosci Biobehav Rev* 2017 Mar;74(Pt B):277-286. [doi: [10.1016/j.neubiorev.2016.07.003](https://doi.org/10.1016/j.neubiorev.2016.07.003)] [Medline: [27461915](https://pubmed.ncbi.nlm.nih.gov/27461915/)]



## Abbreviations

**CAD:** coronary artery disease

**CAU:** care as usual

**Do CHANGE:** Do Cardiac Health Advanced New Generation Ecosystem

**DSD:** Do Something Different

**eHealth:** electronic health

**HF:** heart failure

**HT:** hypertension

**NA:** negative affectivity

**PHQ-9:** Patient Health Questionnaire 9

**SI:** social inhibition

**UTAUT2:** Unified Theory of Acceptance 2

**WHOQOL-BREF:** World Health Organization Quality of Life—BREF

*Edited by G Eysenbach; submitted 02.05.19; peer-reviewed by A Evers, C Loum, S Sankaran; comments to author 14.09.19; revised version received 30.10.19; accepted 15.12.19; published 22.05.20*

*Please cite as:*

*Broers ER, Kop WJ, Denollet J, Widdershoven J, Wetzels M, Ayoola I, Piera-Jimenez J, Habibovic M*

*A Personalized eHealth Intervention for Lifestyle Changes in Patients With Cardiovascular Disease: Randomized Controlled Trial*  
*J Med Internet Res 2020;22(5):e14570*

*URL:* <https://www.jmir.org/2020/5/e14570>

*doi:* [10.2196/14570](https://doi.org/10.2196/14570)

*PMID:* [32441658](https://pubmed.ncbi.nlm.nih.gov/32441658/)

©Eva Rosalinde Broers, Willem Johan Kop, Johan Denollet, Jos Widdershoven, Mart Wetzels, Idowu Ayoola, Jordi Piera-Jimenez, Mirela Habibovic. Originally published in the Journal of Medical Internet Research (<http://www.jmir.org>), 22.05.2020. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in the Journal of Medical Internet Research, is properly cited. The complete bibliographic information, a link to the original publication on <http://www.jmir.org/>, as well as this copyright and license information must be included.