This is a post-peer-review, pre-copyedit version of an article published in Journal of the American Academy of Child and Adolescent Psychiatry. The final authenticated version is available online at: http://dx.doi.org/10.1016/j.jaac.2015.11.010.

Geoffroy MC, Boivin M, Arseneault L, Turecki G, Vitaro F, Brendgen M, Renaud J, Séguin JR, Tremblay RE, Côté SM. Associations Between Peer Victimization and Suicidal Ideation and Suicide Attempt During Adolescence: Results From a Prospective Population-Based Birth Cohort. J Am Acad Child Adolesc Psychiatry. 2016 Feb;55(2):99-105. Epub 2015 Nov 26. PMID: 26802776.

© 2016 This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/

Associations Between Peer Victimization and Suicidal Ideation and Suicide Attempt During Adolescence: Results From a Prospective Population-Based Birth Cohort

RH = Peer Victimization and Suicidality

Marie-Claude Geoffroy, PhD, Michel Boivin, PhD, Louise Arseneault, PhD, Gustavo Turecki, MD, PhD, Frank Vitaro, PhD, Mara Brendgen, PhD, Johanne Renaud, MD, FRCPC, Jean R. Séguin, PhD, Richard E. Tremblay, PhD, Sylvana M. Côté, PhD

Clinical guidance is available at the end of this article.

Accepted November 20, 2015

Drs. Geoffroy and Turecki are with McGill University, Montreal and the McGill Group for Suicide Studies at the Douglas Mental Health University Institute, Montreal. Dr. Geoffroy is also with the Research Unit on Children's Psychosocial Maladjustment at University of Montreal. Dr. Boivin is with Laval University, Quebec, the Research Unit on Children's Psychosocial Maladjustment at Laval University, and the Institute of Genetic, Neurobiological, and Social Foundations of Child Development at the Tomsk State University, Tomsk, Russian Federation. Dr. Arseneault is with the Institute of Psychiatry, Psychology and Neuroscience at King's College London. Drs. Vitaro and Séguin are with the University of Montreal, the Research Unit on Children's Psychosocial Maladjustment at the University of Montreal, and the Sainte-Justine Hospital Research Centre, Montreal. Dr. Brendgen is with the University of Quebec in Montreal, the Research Unit on Children's Psychosocial Maladjustment at the University of Montreal, and the Sainte-Justine Hospital Research Centre. Dr. Renaud is with McGill University, and with the McGill Group for Suicide Studies and the Manulife Centre for Breakthroughs in Teen Depression and Suicide Prevention at the Douglas Mental Health University Institute. Dr. Tremblay is with the University of Montreal, the University College Dublin, and the Institute of Genetic, Neurobiological, and Social Foundations of Child Development at the Tomsk State University. Dr. Côté is with the University of Montreal, the Institute of Genetic, Neurobiological, and Social Foundations of Child Development at the Tomsk State University, the Research Unit on Children's Psychosocial Maladjustment at the University of Montreal, and the Sainte-Justine Hospital Research Centre.

This study was conducted with help from the Canadian Institutes of Health Research (grants OGE#111395 and MOP#114984 awarded to M.B. and S.M.C. and R.E.T.) and Manulife research fund in teen depression (J.R.). The larger QLSCD study was also supported by funding from the Québec Government's Ministry of Health and Ministry of Family Affairs, The Lucie and André Chagnon Foundation, the Québec Health Research Fund (FQRS), The Québec Research Fund for Society and Culture (FQRSC), Canada's Social Science and Humanities Research Council, the Canadian Institutes for Health Research, the St-Justine Research Centre, and the Québec Statistics Institute (ISQ) (R.E.T.). S.M.C. is a senior research fellow of the FRQS. http://www.msss.gouv.qc.ca/en/http://fondationchagnon.org/en/index.aspx http://www.frqs.gouv.qc.ca/en/http://www.frqsc.gouv.qc.ca/fr/ accueil.php http://www.sshrc-crsh.gc.ca/homeaccueileng.aspx http://www.stat.gouv.qc.ca.

The authors wish to thank David Gunnell, PhD, of the University of Bristol, for his contribution to the paper and all the participants of the QLSCD.

Disclosure: Drs. Geoffroy, Boivin, Arseneault, Turecki, Vitaro, Brendgen, Renaud, Séguin, Tremblay, and Côté report no biomedical financial interests or potential conflicts of interest.

Correspondence to Marie-Claude Geoffroy, PhD, McGill Group for Suicide Studies, Douglas Mental Health University Institute, Frank B. Common Pavilion, 6875 LaSalle Boulevard, Montreal, Canada H4H 1R3; email: Marie-claude.geoffroy@mcgill.ca.

ABSTRACT

Objective: To test whether adolescents who are victimized by peers are at heightened risk for suicidal ideation and suicide attempt, using both cross-sectional and prospective investigations.

Method: Participants are from the Quebec Longitudinal Study of Child Development, a general population sample of children born in Quebec in 1997/98 and followed up until 15 years of age. Information about victimization and serious suicidal ideation and suicide attempt in the past year was obtained at ages 13 and 15 years from self-reports (N=1,168).

Results: Victims reported concurrently higher rates of suicidal ideation at age 13 (11.6 to 14.7%) and suicide attempt at age 15 (5.4 to 6.8%) compared to those who had not been victimized (2.7 to 4.1% for suicidal ideation and 1.6 to 1.9% for suicide attempt). Being victimized by peers at 13 years predicted suicidal ideation (odds ratio [OR]=2.27; 95% CI, 1.25 to 4.12) and suicide attempt (OR=3.05, 1.36 to 6.82) 2 years later, even after adjusting for baseline suicidality and mental health problems and a series of confounders (socioeconomic status, intelligence, family's functioning and structure, hostile-reactive parenting, maternal lifetime suicidal ideation/suicide attempt). Those who were victimized at both 13 and 15 years had the highest risk of suicidal ideation (OR=5.41, 2.53 to 11.53) and suicide attempt (OR=5.85, 2.12 to 16.18) at 15 years.

Conclusion: Victimization is associated with an increased risk of suicidal ideation and suicide attempt over and above concurrent suicidality and prior mental health problems. The longer the history of victimization, the greater the risk is.

Key words: suicidal ideation; suicide attempt; adolescence; peer victimization; bullying

INTRODUCTION

Suicide is among the top causes of mortality worldwide, especially in adolescents and young adults.¹⁻³ According to recent US epidemiological data, 12.1% of adolescents (13 to 18 years) "seriously thought about killing themselves" and 4.1% "tried to kill themselves" in their lifetimes.⁴ In Canada in 2008/09, 7.4% of adolescents aged 14-15 years "seriously considered attempting suicide" and 3.3% "attempted suicide" in the past year.⁵ Adolescents who have attempted suicide are at high risk of mortality and of psychological and social adjustment problems in adulthood, ⁶⁻⁸ indicating that efforts need to be directed toward preventing onset of suicide behaviors in adolescence. Over the last few years, considerable attention has been given to the possible link between peer victimization and suicidal ideation and/or suicide attempt. Victimization by peers is highly prevalent in adolescence, with rates ranging from 11% to 40% across 25 countries,⁹ and it has been associated with concurrent suicidal ideation and suicide attempt.¹⁰⁻¹² But surprisingly few prospective studies have tested whether peer victimization increases the risk that adolescents will later develop suicidal ideation or attempt suicide.¹³⁻¹⁷ In fact, most prospective studies have generally focused on self-harm,¹⁷⁻¹⁹ but adolescents who self-harm do not necessarily want to take their own lives. Previous prospective studies pertaining to suicidal ideation are inconsistent: a few studies indicate that peer victimization predicts suicidal ideation,^{13, 15-17} but at least one study did not find such an association.¹⁴ For example, in a large sample of Norwegian adolescents, peer victimization at 14 years was not predictive of suicidal ideation 1 year later after baseline suicidality was controlled for.¹⁴ Meanwhile, in the Avon Longitudinal Study of Parents and Children (ALSPAC) cohort, peer victimization assessed between 4 and 10 years across different informants was prospectively associated with suicidal ideation in preadolescents after adjusting for confounders.¹⁷ Importantly, few cross-sectional¹⁰ and only one prospective survey in early adulthood²⁰ have looked at the association between peer victimization and suicide attempt.

Understanding the predictive role of peer victimization in suicidal ideation and attempt in a general population of adolescents could provide important information about suicide risk factors and suicide prevention strategies. The purpose of the present study was to investigate among a representative cohort of individuals born in 1997/98 in the Canadian province of Quebec whether adolescents who are victimized by peers are at heightened risk for later suicidal ideation and suicide attempt. We expected peer victimization to co-occur with and to predict suicidal ideation and suicide attempt. As victims may exhibit pre-existing risk for suicide and mental health problems, we tested whether victimization increased the risk for suicidal ideation and suicide attempt above and beyond baseline suicidality, mental health problems, and other potential sources of influence.

METHOD

Sample

The Quebec Longitudinal Study of Child Development (QLSCD) is a representative cohort of 2,120 youths born in the Canadian Province of Quebec in 1997/98. Data was collected annually or biannually from 1998 through 2013.

In the present study, we included 1,168 participants for whom measures of peer victimization and suicidality were collected at both 13 and 15 years of age, for 55% of the original cohort. To adjust for attrition, we conducted analyses with and without inverse probability weights, representing participants' probabilities of being included in the study sample (N=1,168) conditional on variables related to attrition: (1) males (46% vs 57%, X^2 =25.68, p<.001) and participants with (2) non-Caucasian ethnicity (4.8% vs 13.8%, X^2 =52.36, p<.001), (3) high maternal depressive symptoms (1.48 vs 1.34, T (1968.69)=2.43, p=.015) and (4) low socioeconomic status (.11 vs - .15, T (2093)=-5.98, p<.001) were under-represented in the study sample. The results with and without weights did not differ; the latter are presented here. Additionally, we tested whether restriction of analyses to a sample of participants with assessments of both victimization and suicidality outcomes at ages 13 and 15 affected our results. Therefore, we repeated our analyses on a larger sample of participants with assessment of suicidality outcomes at 15 years only (imputed n=1,446, representing 69% of the surviving cohort); patterns of results remained unchanged.

Ethics consideration

Ethical approval was given by the Quebec Institute of Statistics and the St-Justine Hospital Research Centre. Informed written consent was obtained from all participating parents at each assessment and from adolescents at the 15-year assessment. Adolescents were assured confidentiality, which is essential to ensure reliability of self-reports.²¹ All adolescents and their parents were provided with a list of resources in case of needing help. If they rated highly on the suicidal behavior scale, a pop-up appeared on screen to encourage them to seek help.

Measures

Past year suicidal ideation and suicide attempt

Suicidal ideation was assessed with one question administered to participants at 13 and 15 years: "in the past twelve months, did you ever seriously think of attempting suicide"; answering "don't know"/ or refusal was coded as 0. If they answered affirmatively, they were asked: "in the past twelve months, how many times did you attempt suicide," dichotomized as 0 versus ≥ 1 .

Peer victimization

Peer victimization was assessed at 13 and 15 years using seven items (four of which are based on the Selfreport Victimization Scale²²). They were selected on the basis of their reflecting various physical, verbal, direct, and indirect instances of victimization, including the new forms of cyber victimization, through answering questions such as: how many times it happened (since the beginning of this school year) that another student in school had: (1) called you names/said mean things; (2) didn't let you play in his/her group; (3) pushed/hit/kicked you; (4) said bad things behind your back; (5) teased you in a mean way; (6) made you pay/give something so they would leave you alone; (7) insulted/threatened/intimidated you on the Internet and cell phone. Items were administered in the second half of the school year (February to June) and were rated on a 4-point scale (never=0; rarely=1; often=2; very often =3; don't know/refusal coded 0). This peer victimization scale is analogous to that used successfully in previous work²³ and has shown good internal consistency (Cronbach alphas were .81 at age 13 and .82 at age 15). Because of the skewed distribution, victimization was treated categorically. Participants were identified as being victimized by peers if they answered "often" or "very often" to at least one of the 7 questions in accordance of the conventional definition.

Prior Mental Health Problems

Oppositional/defiance and inattention/hyperactivity symptoms were rated by school teachers when participants were 6, 7, 8, 10, and 12 years of age. Questions were derived from the Canadian National Longitudinal Study of Children and Youth,²⁴ which incorporates items from the Child Behavior Checklist,²⁵ the Ontario Child Health Study Scales,²⁶ and the Preschool Behavior Questionnaire.²⁷ Oppositional/defiance was assessed with four items (e.g., defiant/refused to comply; punishment didn't change behavior), and inattention/hyperactivity was assessed with nine items (e.g., could not sit still; was unable to concentrate) rated on a 3-point scale (0=never or not true; 1=sometimes or somewhat true; 2=often or very true). Cronbach α ranged from .81 to .84 for opposition/defiance and from .89 to .91 for hyperactivity/inattention. Scores were averaged across ages. Scales demonstrated adequate sensitivity and external validity in this cohort.²⁸ Depression was self-reported at 12 years using 10 items from the Children Depression Inventory (CDI),^{29, 30} assessing symptoms of the past 2 weeks on a 3point scale. The CDI has shown good reliability and discriminative validity consistent with clinical assessment of depression;³¹ Cronbach α was .81. For analysis purposes, we generated three variables tapping into mental health problems, where a score of ≥90th percentile of symptoms was identified as an elevated symptom level.

Other Confounding Influences

Potential confounders at different ages were identified a priori from the literature.³ Covariates were measured at 13 and 15 years unless otherwise indicated. At 2¹/₂ years, mothers completed the Parental Depression Questionnaire.³² Mothers who reported they had been depressed for most of the time for two weeks or more in their

lifetimes were asked whether they ever thought of or attempted suicide. At 10 years, participants' verbal intelligence was estimated using the Peabody Picture Vocabulary Test, standardized for age in months at administration.^{33, 34} Other environmental factors included family socioeconomic status (SES; aggregate of annual gross income, parental education level, and occupational prestige), based on the scheme of Willms and Shields³⁵; family structure (biological parents/blended/single); family functioning, assessed with 8 items (i.e. communication, problem resolution, and control of disruptive behaviours); and hostile-reactive parenting, assessed with 4 items (i.e. corporal punishment, raising voice). All questionnaires are available online (http://www.jesuisjeserai.stat.gouv.qc.ca).

Statistical Analyses

In cross-sectional analyses, we used logistic regressions to test whether suicidal ideation and suicide attempt were associated with frequent victimization.

Using logistic regressions in longitudinal analyses, we tested whether victimization at 13 years was associated with suicidal ideation and suicide attempt 2 years later. The associations were adjusted for suicidality at baseline (suicidal ideation or suicide attempt at age 13 years), for three mental health problems by age 12 years (depression, oppositional/defiance and inattention/hyperactivity problems), and for six additional confounders (i.e., socioeconomic status, intelligence, family's functioning and structure, hostile-reactive parenting, and maternal suicidality history).

The temporality of longitudinal association was further examined in sensitivity analyses of incident cases of suicidal ideation and suicide attempt. Logistic regressions were conducted, in which participants who reported suicidal ideation and suicide attempt at baseline were excluded from the analyses. The results were similar to those presented here, but analyses were based on small number of incident cases. Finally, we examined whether the prospective association between victimization and later suicidality varied as a function of the chronicity of the victimization experience in adolescence (i.e. being victimized at both 13 and 15 years; being victimized at 13 or 15 years versus never being victimized) using logistic regression.

To minimize further data loss, missing information on covariates was imputed using multiple imputations by chained equations³⁶ method, and analyses were run across the 10 imputed data set.

RESULTS

The overall prevalence of suicidal ideation slightly increased from 4.5% at age 13 to 5.9% at age 15. The prevalence of suicidal ideation was higher and increased for females (5.3-8.6%) compared to males (3.7-3.0%)

(Table 1). The prevalence of suicide attempt increased between 13 and 15 years for females (2.7 to 4.5%) and decreased for males (2.1 to 0.9%) (Table 1). The prevalence of peer victimization was 16.7% for females and 25.8% for males at age 13 and 17.5% for females and 17.2% for males at age 15 years; 8.2% (n=95) of the sample reported being victimized at both ages. Although the prevalence of suicidality and victimization differed by sex, the associations between victimization and suicidality did not differ by sex. Hence we present results for both sexes combined, with adjustment for sex (see Table 2's footnote). Associations were examined with and without adjustment for confounders and prior mental health problems.

Cross-sectional analyses indicate that victimization was associated with suicidal ideation and suicide attempt at both 13 and 15 years (Table 2). The associations were attenuated but remained significant after adjusting for confounders and prior mental health problems. Being frequently victimized was concurrently associated with suicidal ideation at 13 years (Odds ratio [OR]=3.40; 95% CI=1.82-6.37) and 15 years (OR=3.74; 95% CI=2.19-6.38), as well as with suicide attempt at 13 years (OR=2.28; 95% CI=0.97-5.38) and 15 years (OR=3.16; 95% CI=1.19-6.69), in a fully adjusted model (i.e. adjusted for mental health and confounders).

Longitudinal models showed that the experience of frequent victimization at 13 years predicted both subsequent suicidal ideation and suicide attempt 2 years later (Table 3), even after taking into account suicidality at baseline. Associations were attenuated after adjustment for confounders and prior mental health problems: victimization at 13 years increased the risk of suicidal ideation at 15 years by 2.27 times (CI=1.25-4.12) and attempt by 3.05 times (CI=1.36-6.82).

Finally, as shown in Table 4, we found that only youths who experienced chronic victimization had an increased risk of suicidal ideation at age 15 (OR=5.41; 95% CI=2.53-11.53) and suicide attempt (OR=5.85; 95% CI=2.12-16.18), in models adjusting for prior suicidality, mental health problems and other confounders.

DISCUSSION

During adolescence, the peer group becomes increasingly important. Thus, being victimized by peers (i.e. being frequently teased/called names or excluded from social groups and activities) can represent a major source of stress, with potentially damaging consequences.^{37, 38} In this population-based sample, such victimization experiences were common, with 17 - 21% of 13 - 15-year-old adolescents reporting being often or very often victimized by their peers. Such victimized adolescents concurrently reported a higher likelihood of past 12-month suicidal ideation and suicide attempt. Furthermore, adolescents who were frequently victimized by their peers at age 13 were more likely

to develop suicidal ideation and make a suicide attempt two years later. More importantly, adolescents who reported having been frequently victimized both at 13 and 15 years had the highest risk of suicidal ideation and suicide attempt at 15 years.

This is the first study to show a predictive association between victimization and suicidal ideation and suicide attempt in mid adolescence while taking into account a variety of factors, including previous suicidality, mental health problems (by age 12 years) such as depression, opposition/defiance and inattention/hyperactivity problems, as well as family adversity. According to prior studies, peer victimization often occurs within the context of other risk factors, such as mental health problems and familial adversity,³⁹ which may contributed to suicidality.³ Thus, our study was able to distinguish the role of peer victimization from these co-occurring factors to more clearly establish its contributions as a distinct predictor of suicidality. The rates of suicidal ideation and suicide attempt typically increased in late adolescence,⁴⁰ thus our study suggested that victimization represents a risk factor for the relatively early emergence of suicidality.

Our findings are in line with those of many other cross-sectional and the few longitudinal studies on bullying/suicidality association.^{10, 11, 15-17} However, in a prior Norwegian study,¹⁴ the predictive association from peer victimization in adolescence to suicidal ideation (unconfounded by depressive symptoms) one year later became non-significant after taking into account suicidal ideation at baseline. Reasons for these discrepant findings could relate to differences in measurement of suicidal ideation and length of follow-up. Specifically, suicidal ideation included more passive and indirect thoughts (e.g., Life is not worth living; My family would be better off without me) than those used in the present study (e.g., ever seriously think of attempting suicide).

Several possible explanations exist for the observed association between peer victimization and suicidality. First, one etiological pathway may involve mental health, especially depression. In one study on self-harm, being victimized in childhood was associated directly with an increased risk of self-harm in late adolescence, but also indirectly via subsequent depression.¹⁸ In the present study, we showed that being victimized by peers in adolescence could lead to suicidality while accounting for prior levels of depression. However, we could not test the possibility that depression mediated the association of peer victimization with suicidality. Still, not all victimized adolescents will exhibit higher rates of suicidal ideation and suicide attempt. Hence, it has been suggested that peer victimization may trigger mental health problems or suicidality in interaction with genetic vulnerability to psychopathology. For instance, a recent study found that peer victimization increased the risk of depression in girls who carried two copies of the short allele at the promoter region of the serotonin transporter gene, but not in those without two copies.⁴¹ Second, the experience of peer victimization may lead to biological alteration with lasting effects on psychopathology and suicidality: (1) victimized children have shown dysregulation of the hypothalamic-pituitary-adrenal axis,⁴² and such dysregulations have been involved in the development of psychopathology⁴³ and suicidality⁴⁴; (2) victimized children were also shown to have elevated C reactive protein at mid-life, a marker for neuro-inflammation in their bloodstream,⁴⁵ and inflammation markers have been associated with increased risk of developing depression.⁴⁶

This representative birth cohort provided a unique opportunity to examine the temporal relation between victimization by peers and the relatively early onset of suicidal ideation and suicide attempt in a contemporary sample of Canadian adolescents. Indeed, the study's main strengths are its prospective design, with suicidality measured at two time points during adolescence to identify incident cases, self-reported and teacher-reported assessments of prior mental health problems, and inclusion of various familial adversity factors and intelligence. Another strength includes repeated assessments of peer victimization based on 7 items covering a variety of behaviors from verbal (such as teasing/calling names) to cyberbullying. Moreover, unlike previous studies, we focused on serious suicidal ideation and suicide attempt, which are of clinical significance.⁴³ Nevertheless, the present findings should be interpreted in light of their limitations. There were some missing data, with 56% of the original sample with assessment of bullying and suicidality at both 13 and 15 years. To avoid further loss of data, missing data were imputed using multiple imputation following current guidelines.⁴⁷ Sensitivity analysis using the larger sample of those with information at 15 years (n=1,446, representing 69% of the original sample) and using sample weights to adjust for attrition suggests that results reported here are robust. Second, as in most studies of victimization and suicide, the present study relied on self-reports. A potential limitation of self-reported victimization is that it reflects a perception potentially biased by the self-system and mental health status.^{23, 48} The use of self-report for both victimization and suicide may also have induced shared method variance and inflated effect sizes. However, in this population-based survey, no clinical interviews have been conducted, and adolescents were the only informants, as after childhood, parents are generally less aware of their offspring's personal experiences, especially suicidal ideation^{24, 49} and peer victimization.⁵⁰ Furthermore, we were not able to examine the separate influence of cyberbullying on suicidality due to the small numbers of cyberbullied adolescents. A recent meta-analysis suggested that cyberbullying is more strongly related to suicidal ideation than traditional bullying.¹¹

There was no death due to suicide in our study. Thus, the relation of peer victimization and suicidal ideation and non-fatal suicide attempt may differ for completed suicide. One observational study based on coroner records reported that only 6% of suicides in youths could be attributed to peer victimization,⁵¹ whereas two prospective studies reported increased rates of suicide mortality in adulthood among those who had been victimized by peers in early childhood.^{20, 52} In addition, unmeasured confounders might explain associations such as perpetration of bullying,⁵³ but in this study, victimizing others was not associated with subsequent suicidality. Another limitation is low statistical power, particularly for investigating sex differences in associations, due to the small number of males reporting suicidal ideation and/or suicide attempt. Also, it is important to underline that the present findings document the directionality of associations, but they do not prove causality. Finally, as the follow-up was only two years, we do not know whether the consequences of victimization on suicidality persist as participants enter adulthood.

The association between being victimized by peers and suicidal ideation and suicide attempt in adolescence, distinct from prior suicidality, mental health problems, and family adversity, has implications for practice. Universal prevention of peer victimization is paramount to reduce the rate of suicidal ideation and suicide attempt in adolescence. A recent systematic review of 44 anti-bullying programs found, on average, that school-based programs are effective in reducing victimization by 17-20%.⁵⁴ Future studies are urgently needed to test whether reducing the occurrence of bullying translates into a reduction of suicidal ideation and suicide attempt. Other strategies that do not aim directly at reducing bullying can also be beneficial. For instance, parental support has been shown to attenuate the negative effect of bullying on suicidality,⁵⁵ and actions to improve school climate⁵⁶ can potentially prevent bullying. In clinical settings, health professionals should consider screening for peer victimization. Although our study and many others suggest that peer victimization in clinical settings. Effective intervention may require a multidisciplinary effort involving parents, schoolteachers, and principals, as well as mental health professionals.⁵⁷

In sum, our findings along with other studies highlight that victimization is a form of adversity that could leave significant scars, including suicidal ideation and attempt, which need to be addressed urgently.

CLINICAL GUIDANCE:

-Being victimized is a common experience for many adolescents.

-The odds of developing suicidal ideation and attempting suicide at 15 years among youth who were chronically victimized, i.e. lasting at least 2 school years, are 5 to 6 times the odds of those who were never victimized, independently from previous suicidality and mental health problems.

-Evidenced-based prevention interventions in clinical settings need to be addressed, but effective management may require a multidisciplinary effort involving parents, school, and mental health professionals.

REFERENCES

- Skinner R, McFaull S. Suicide among children and adolescents in Canada: trends and sex differences, 1980–2008. *Can Med Assoc J.* 2012;184(9):1029-1034.
- McLoughlin AB, Gould MS, Malone KM. Global trends in teenage suicide: 2003-2014. *QJM*. 2015;108(10):765-780.
- **3.** Hawton K, Saunders KEA, O'Connor RC. Self-harm and suicide in adolescents. *Lancet*. 2012;379(9834):2373-2382.
- 4. Nock MK, Green JG, Hwang I, et al. Prevalence, correlates, and treatment of lifetime suicidal behavior among adolescents: results from the National Comorbidity Survey Replication Adolescent Supplement. JAMA psychiatry. 2013;70(3):300-310.
- 5. McMartin SE, Kingsbury M, Dykxhoorn J, Colman I. Time trends in symptoms of mental illness in children and adolescents in Canada. *Can Med Assoc J.* 2014;186(18):E672-E678.
- **6.** Mars B, Heron J, Crane C, et al. Clinical and social outcomes of adolescent self harm: population based birth cohort study. *BMJ*. 2014;349:g5954.
- 7. Finkelstein Y, Macdonald EM, Hollands S, et al. Long-term outcomes following self-poisoning in adolescents: a population-based cohort study. *Lancet Psychiatry*. 2015;2(6):532-539.
- **8.** Goldman-Mellor SJ, Caspi A, Harrington H, et al. Suicide attempt in young people: a signal for long-term health care and social needs. *JAMA psychiatry*. 2014;71(2):119-127.
- Nansel TR, Craig W, Overpeck MD, Saluja G, Ruan WJ. Cross-national consistency in the relationship between bullying behaviors and psychosocial adjustment. *Arch Pediatr Adolesc Med.* 2004;158(8):730-736.

- **10.** Holt MK, Vivolo-Kantor AM, Polanin JR, et al. Bullying and suicidal ideation and behaviors: a metaanalysis. *Pediatrics*. 2015;135(2):e496-e509.
- **11.** Van Geel M, Vedder P, Tanilon J. Relationship between peer victimization, cyberbullying, and suicide in children and adolescents: a meta-analysis. *JAMA pediatrics*. 2014;168(5):435-442.
- Herba CM, Ferdinand RF, Stijnen T, et al. Victimisation and suicide ideation in the TRAILS study: specific vulnerabilities of victims. *J Child Psychol Psychiatry*. 2008;49(8):867-876.
- **13.** Kim YS, Leventhal BL, Koh Y-J, Boyce WT. Bullying increased suicide risk: prospective study of Korean adolescents. *Archives of suicide research*. 2009;13(1):15-30.
- Undheim AM. Involvement in bullying as predictor of suicidal ideation among 12-to 15-year-old Norwegian adolescents. *Eur Child Adolesc Psychiatry*. 2013;22(6):357-365.
- 15. Bannink R, Broeren S, van de Looij–Jansen PM, de Waart FG, Raat H. Cyber and traditional bullying victimization as a risk factor for mental health problems and suicidal ideation in adolescents. *PLoS One*. 2014;9(4):e94026.
- Heikkilä H-K, Väänänen J, Helminen M, Fröjd S, Marttunen M, Kaltiala-Heino R. Involvement in bullying and suicidal ideation in middle adolescence: a 2-year follow-up study. *Eur Child Adolesc Psychiatry*. 2013;22(2):95-102.
- **17.** Winsper C, Lereya T, Zanarini M, Wolke D. Involvement in bullying and suicide-related behavior at 11 years: a prospective birth cohort study. *J Am Acad Child Adolesc Psychiatry*. 2012;51(3):271-283.
- **18.** Lereya ST, Winsper C, Heron J, et al. Being bullied during childhood and the prospective pathways to self-harm in late adolescence. *J Am Acad Child* Adolesc *Psychiatry*. 2013;52(6):608-618.
- Fisher HL, Moffitt TE, Houts RM, Belsky DW, Arseneault L, Caspi A. Bullying victimisation and risk of self harm in early adolescence: longitudinal cohort study. BMJ. 2012;344:e2683.
- Klomek AB, Sourander A, Niemelä S, et al. Childhood bullying behaviors as a risk for suicide attempts and completed suicides: A population-based birth cohort study. J Am Acad Child Adolesc Psychiatry. 2009;48(3):254-261.
- Lothen-Kline C, Howard DE, Hamburger EK, Worrell KD, Boekeloo BO. Truth and consequences: ethics, confidentiality, and disclosure in adolescent longitudinal prevention research. *J Adolesc Health*. 2003;33(5):385-394.

- **22.** Ladd GW, Kochenderfer-Ladd B. Identifying victims of peer aggression from early to middle childhood: analysis of cross-informant data for concordance, estimation of relational adjustment, prevalence of victimization, and characteristics of identified victims. *Psychol Assess.* 2002;14:74-96.
- **23.** Boivin M, Brendgen M, Vitaro F, et al. Strong genetic contribution to peer relationship difficulties at school entry: Findings from a longitudinal twin study. *Child Dev.* 2013;84(3):1098-1114.
- 24. Statistics Canada and Human Resources Development Canada. *National Longitudinal Survey of Children* and Youth: Overview of Survey Instruments for 1994–1995 Data Collection Cycle 1. Ottawa: Author; 1995.
- **25.** Achenbach TM, Edelbrock C, Howell CT. Empirically based assessment of the behavioral/emotional problems of 2- and 3- year-old children. *J Abnorm Child Psychol.* 1987;15(4):629-650.
- 26. Offord DR, Boyle MH, Racine Y. Ontario Child Health Study: correlates of disorder. *J Am Acad Child Adolesc Psychiatry*. 1989;28(6):856-860.
- 27. Tremblay RE, Desmarais-Gervais L, Gagnon C, Charlebois P. The Preschool Behaviour Questionnaire: Stability of its factor structure between cultures, sexes, ages and socioeconomic classes. *International Journal of Behavioral Development*. 1987;10(4):467-484.
- **28.** Galéra C, Pingault J-B, Michel G, et al. Clinical and social factors associated with attention-deficit hyperactivity disorder medication use: population-based longitudinal study. *Br J Psychiatry*. 2014;205(4):291-297.
- 29. Allgaier A-K, Frühe B, Pietsch K, Saravo B, Baethmann M, Schulte-Körne G. Is the Children's Depression Inventory Short version a valid screening tool in pediatric care? A comparison to its full-length version. J Psychosom Res. 2012;73(5):369-374.
- **30.** Kovacs M. The Children's Depression, Inventory (CDI). *Psychopharmacol Bull*. 1984;21(4):995-998.
- **31.** Stockings E, Degenhardt L, Lee YY, et al. Symptom screening scales for detecting major depressive disorder in children and adolescents: a systematic review and meta-analysis of reliability, validity and diagnostic utility. *J Affect Disord*. 2015;174:447-463.
- **32.** Roy CA, Zoccolillo M, Gruber R, Boivin M, Perusse D, Tremblay RE. Construct validity of an instrument to assess major depression in parents in epidemiologic studies. *Can J Psychiatry*. 2005;50(12):784-791.
- **33.** Dunn LM, Dunn LM. *Peabody Picture Vocabulary Test-Revised (PPVT-R)*. Circles Pines, MN: American Guidance Services; 1981.

- 34. Dunn LM, Theriault-Whalen CM, Dunn LM. Échelle de Vocabulaire en Images Peabody. Adaptation francaise du Peabody Picture Vocabulary Test-Revised. Manuel pour les formes A et B. Toronto, ON: Psycan; 1993.
 - 35. Willms DJ, Shields M. A measure of socioeconomic status for the National Longitudinal Study of Children. Fredericton, New Brunswick: Atlantic Center for Policy Research in Education, University of New Brunswick and Statistics Canada; 1996: 9607.
- **36.** Azur MJ, Stuart EA, Frangakis C, Leaf PJ. Multiple imputation by chained equations: what is it and how does it work? *International journal of methods in psychiatric research*. 2011;20(1):40-49.
- **37.** Vitaro F, Boivin M, Bukowski WM. The role of friendship in child and adolescent psychosocial development. In: Rubin K, Bukowski W, Laursen B, eds. *Handbook of peer interactions, relationships, and groups*. New York: The Guildford Press; 2009:568-588.
- Boivin M, Hymel S, Hodges EV. Toward a process view of peer rejection and harassment. In: Juvonen J,
 Graham S, eds. *Peer harassment in school: The plight of the vulnerable and victimized*. New York:
 Guilford Press; 2001:265-289.
- **39.** Fergusson DM, Woodward LJ, Horwood LJ. Risk factors and life processes associated with the onset of suicidal behaviour during adolescence and early adulthood. *Psychol Med.* 2000;30(1):23-39.
- **40.** Arseneault L, Bowes L, Shakoor S. Bullying victimization in youths and mental health problems: 'Much ado about nothing'? *Psychol Med.* 2010;40(05):717-729.
- **41.** Benjet C, Thompson RJ, Gotlib IH. 5-HTTLPR moderates the effect of relational peer victimization on depressive symptoms in adolescent girls. *J Child Psychol Psychiatry*. 2010;51(2):173-179.
- **42.** Ouellet-Morin I, Danese A, Bowes L, et al. A discordant monozygotic twin design shows blunted cortisol reactivity among bullied children. *J Am Acad Child Adolesc Psychiatry*. 2011;50(6):574-582.
- **43.** Geoffroy M-C, Hertzman C, Li L, Power C. Prospective association of morning salivary cortisol with depressive symptoms in mid-life: A life-course study. *PLoS One*. 2013;8(11):e77603.
- **44.** Coryell W, Schlesser M. The dexamethasone suppression test and suicide prediction. *Am J Psychiatry*. 2001;158(5):748-753.
- **45.** Takizawa R, Danese A, Maughan B, Arseneault L. Bullying victimization in childhood predicts inflammation and obesity at mid-life: a five-decade birth cohort study. *Psychol Med.* 2015:1-11.

- **46.** Howren MB, Lamkin DM, Suls J. Associations of depression with C-reactive protein, IL-1, and IL-6: a meta-analysis. *Psychosom Med.* 2009;71(2):171-186.
- **47.** Sterne JAC, Cox DR, Smith GD. Sifting the evidence—what's wrong with significance tests? Another comment on the role of statistical methods. *BMJ*. 2001;322:226-231.
- **48.** Boivin M, Vitaro F, Gagnon C. A reassessment of the Self-Perception Profile for Children: Factor structure, reliability, and convergent validity of a French version among second through sixth grade children. *International Journal of behavioral development*. 1992;15(2):275-290.
- **49.** Foley DL, Goldston DB, Costello EJ, Angold A. Proximal psychiatric risk factors for suicidality in youth: the Great Smoky Mountains Study. *Arch Gen Psychiatry*. 2006;63(9):1017-1024.
- **50.** Radford L, Corral S, Bradley C, Fisher HL. The prevalence and impact of child maltreatment and other types of victimization in the UK: Findings from a population survey of caregivers, children and young people and young adults. *Child Abuse Negl.* 2013;37(10):801-813.
- **51.** Sinyor M, Schaffer A, Cheung AH. An observational study of bullying as a contributing factor in youth suicide in Toronto. *Can J Psychiatry*. 2014;59:632-638.
- 52. Geoffroy MC, Gunnell D, Power C. Prenatal and childhood antecedents of suicide: 50-year follow-up of the 1958 British Birth Cohort study. *Psychol Med.* 2014;44(6):1245-1256.
- **53.** Copeland WE, Wolke D, Angold A, Costello EJ. Adult psychiatric outcomes of bullying and being bullied by peers in childhood and adolescence. *JAMA psychiatry*. 2013;70(4):419-426.
- **54.** Ttofi MM, Farrington DP. Effectiveness of school-based programs to reduce bullying: A systematic and meta-analytic review. *Journal of Experimental Criminology*. 2011;7(1):27-56.
- **55.** Elgar FJ, Napoletano A, Saul G, et al. Cyberbullying victimization and mental health in adolescents and the moderating role of family dinners. *JAMA pediatrics*. 2014;168(11):1015-1022.
- 56. Bradshaw CP. Translating research to practice in bullying prevention. *Am Psychol.* 2015;70:322-332.
 - 57. Lamb J, Pepler DJ, Craig W. Approach to bullying and victimization. *Can Fam Physician*. 2009;55(4):356-360.

21	
22	Table 1 : Prevalence of Suicidal Ideation and Suicide Attempt in the Last 12 Months and Peer Victimization at 13 and 15 Years ^a

23							
24		То	otal	Fem	ales	Ma	les
25	Variables; age in years	13	15	13	15	13	15
26 27	Suicidal ideation, % (<i>n</i>)	4.5 (56)	5.9 (86)	5.3 (35)	8.6 (65)	3.7 (21)	3.0 (21)
28	Suicide attempt, % (<i>n</i>)	2.4 (30)	2.8 (40)	2.7 (18)	4.5 (34)	2.1 (12)	0.9 (6)
29	Peer victimization, % (<i>n</i>)	21.0 (258)	17.4 (251)	16.7(111)	17.5(132)	25.8 (147)	17.2 (119)
30 31 32	Note: $n=1,234$ at 13 years and $n=1,446$ at 15 years. ^a Based on maximum available sample.						

1	6
Т	υ

20 Table 2 : C	Table 2 : Cross-Sectional Associations of Peer Victimization by Suicidality Outcomes (<i>n</i> =1,168).						
21	Peer vict	imization ^a		Suicidality outcomes			
23	Yes	No	Model 1	Model 2	Model 3		
24	% (<i>n</i>)	% (<i>n</i>)	OR (95% CI)	OR (95% CI)	OR (95% CI)		
25 Suicide idea	tion						
26 13 years	11.6 (30)	2.7 (26)	5.12 (2.89-9.09)	4.40 (2.44-7.93)	3.40 (1.82-6.37)		

4.28 (2.58-7.08)

4.02 (1.88-8.63)

3.84 (1.91-7.69)

4.1 (49)

1.6 (16)

1.9 (23)

Note: Model 1= adjusted for sex; Model 2= additionally adjusted for socioeconomic status (13 or 15 years); family structure (biological/blended/single; 13 or 15 years) and functioning (13 or 15 years); hostile-reactive parenting (13 or 15 years); intelligence (10y); maternal lifetime suicidal ideation/suicide attempt; Model 3= additionally adjusted for depression (12y); opposition/defiance (6-12y); inattention/hyperactivity (6-12y) problems; p for sex interaction= .337 and .532 for suicidal ideation at 13 and 15 years, respectively; .990 and .459 for suicide attempt at 13 and 15 years, respectively. OR = odds ratio

4.12 (2.44-6.94)

3.16 (1.45-6.91)

3.59 (1.74-7.38)

3.74 (2.19-6.38)

2.28 (0.97-5.38)

3.16 (1.49-6.69)

^a Based on maximum available sample; n=1,234 at 13 years and n=1,446 at 15 years.

14.7 (37)

5.4 (14)

6.8 (17)

15 years

Suicide attempt

13 years

15 years

		OR (95% CI)	
	Model 1	Model 2	Model 3
Peer victimization at 13 years			
Suicidal ideation at 15 ^a	2.60 (1.50-4.51)	2.53 (1.43-4.47)	2.27 (1.25-4.12)
Suicide attempt at 15 ^b	3.64 (1.78-7.43)	3.58 (1.69-7.56)	3.05 (1.36-6.82)

Note: Model 1= adjusted for sex; Model 2 = additionally adjusted for socioeconomic status (13 years); family structure (biological/blended/single;13 years) and functioning (13 years); parental hostile-reactive parenting (13 years); intelligence (10y); maternal lifetime suicidal ideation/suicide attempt; Model 3= additionally adjusted for depression (12y); opposition/defiance (6-

12y); inattention/hyperactivity (6-12y); OR= odds ratio.

^a Adjusted for suicidal ideation at 13y.

^b Adjusted for suicide attempt at 13y.

	OR (95% CI)						
	Model 1	Model 2	Model 3				
Duration of peer victimization (versus not victimized at 13 and 15 years)							
Was victimized at 13 or 15 years							
Suicidal ideation at 15 years ^a	2.06 (1.12-3.79)	1.94 (1.03-3.63)	1.88 (1.00-3.55)				
Suicide attempt at 15 years ^b	1.77 (0.75-4.21)	1.76 (0.73-4.25)	1.70 (0.69-4.18)				
Was victimized at 13 and 15 years							
Suicidal ideation at 15 years ^a	5.90 (2.93-11.87)	5.92 (2.88-12.14)	5.41 (2.53-11.53)				

Table 4: Longitudinal Associations of Duration of Victimization and Suicidality Outcomes at 15 Years (n=1,168)

Note: Model 1 = adjusted for sex; Model 2 = additionally adjusted for socioeconomic status (13 years); family structure

7.56 (3.18-17.98)

(biological/blended/single;13 years) and functioning (13 years); hostile-reactive parenting (13 years); intelligence (10y); maternal lifetime suicidal ideation/suicide attempt; Model 3 = additionally adjusted for depression (12y); opposition/defiance (6-12y); inattention/hyperactivity (6-12y) problems. OR = odds ratio.

7.52 (2.99-18.92)

5.85 (2.12-16.18)

^a Adjusted for suicidal ideation at 13y.

Suicide attempt at 15 years^b

^b Adjusted for suicide attempt at 13y.

MANUESPIET SUBMISSION FORM -- JOURNAL OF THE AMERICAN ACADEMY OF CAN SAND ACCURESCENT PSYCHAGES

Section 7. Lestractions a Alexpolor Bert Hele, which equations consider to the A<u>LE enther</u> on the same option between entered on a section of the same option of the

herber 2 Manuscript lyformation.

neret 2 Matagerigi reneration. Menositie Tele Silier elite militati gene i si a 221 fil 2011 i si del des terre de se de la se Statempi de constante presentation presentation de constant es presentation de la se de se de se Compute numer las constants de la secondation de la secondation de la secondation de la secondation de la second Enterna gene de la secondation de la se Enterna gene de la secondation de la secondation

Corresponding Author Name <u>14(15)</u>. Charge 2015 11 http: In the bay below, place the acknowledgements forsibiling funding for the research in this taken report and the interneyel discharges at a regime. To very long we need to be has far the places and the segment place the world, "see structure?" In the tax

See Attached

Section 3: Acknowledgments By an apply below. Inconvertige my acceptioner is asket certal center of the following information

Capproved of the Submitted Work and Asknowledgacent of Sole of Corresponding Asther 1 was periodially revealed and good CApproval of the Schmitted Work and Arknowledgarust of Stele of Corresponding Asthir, I have periodicity revealed and governments of the version advantation, and with the excipation of periodicity and the version advantation, and with the excipation of periodicity patients of the traductory and for behave being proposed without exception of periodicity and the traductory of the provided to the traductory of the traductory of the traductory of the provided traductory of the traductory of the provided traductory of the traductory of the traductory of the provided traductory of the traductory of the traductory of the provided traductory of the traductory of the traductory of the traductory of the provided traductory of the once a managering in assumation, no supporter of a manger in the content will be a insert

*Authorship Contribution. Here you append sufficiently in the work to the path, respondently for all of part of the contribution of the submitted work in the form of the contribution and design, endour or or submitted work in the submitted work in the form of the contribution and design. care makers of data. AND 12 duling the starie, solver teneng in minoli, for approach the factor of the staries in the staries and the comparison of the staries and the staries an

"Fenancial and Material Support. All hourses of grant or store function support, as well is any material account, or units in the repreparent partitions of this manual of the funding was received pidear story the

" Caaffiet of Interest, Butthereres, Financial, and Material Support To my anowichid, di of my preside contains of more can a seaso of paterial parameters, resolution, and electric support of all electricity of a my pressure section of the electric models and the section of the electric models and the electric models and the electric models and the electric models are not denoted by referring the relation of the payer that have construct the two years on that not expected in the electric models. to rescale times are tone in the appropriate architect of this constraint. Distribute activet, but is not bound at grive to resolution factors, employment, allifications, parents (in preparative, file), or granted, investions, speckers' butches, boronno, consideration meditics, shock representation or experimentation. It as autors has no condition of menorates destare, this must be used as crashed for sextight, the Xieania reports in biomidical formulal interests or potential confacts of criteria

"Transfer of Copyright. To the moment that this work you not performed by entitioners of each S. Bedetal Coversimeter, Lagree to constration of the American Academic of Clubband Adolescent Proclams upon acceptance of this submassion for public station in An internal Traditional the demonstrate publics to signific Johnsel Publication Agriculture on the bolt

Artes Name (Provide Marie Charles Cats (1894) Last our Suggebore Buy Part Spring MENSING. 2 the land page 2 ct GUSTING REAL 1221 9. 1 Later Labor Sin hat provide Inchie : immed A the Martin A-Human M Cather S. Same SEAN SEGUIN J. In 12 States and and find 08-31-20/5 MARA BRENDGEN FRANK VITARD DUNE LARIENERUL 02-09-2015 Jan ank

Associations Between Peer Victimization and Suicidal Ideation and Suicide Attempt During Adolescence: Results From a Prospective Population-based Birth Cohort

Marie-Claude Geoffroy, PhD, Michel Boivin, PhD, Louise Arseneault, PhD, Gustavo Turecki, MD, PhD, Frank Vitaro, PhD, Mara Brendgen, PhD, Johanne Renaud, MD, FRCPC, Jean R. Séguin, PhD, Richard E. Tremblay, PhD, Sylvana M. Côté, PhD

Funding: This study was conducted with help from the Canadian Institutes of Health Research (grants OGE#111395 and MOP#114984 awarded to MB and SMC and RET) and Manulife research fund in teen depression (JR). The larger QLSCD study was also supported by funding from the Québec Government's Ministry of Health and Ministry of Family Affaires, The Lucie and André Chagnon Foundation, the Québec Health Research Fund (FQRS), The Québec Research Fund for Society and Culture (FQRSC), Canada's Social Science and Humanities Research Council, the Canadian Institutes for Health Research, the St-Justine Research Centre, and the Québec Statistics Institute (ISQ) (RET). SMC is a senior research fellow of the FRQS. <u>http://www.msss.gouv.qc.ca/en/ http://fondationchagnon.org/en/index.aspx http://www.</u>frqs.gouv.qc.ca/en/ <u>http://www.stat.gouv.qc.ca/fr/</u> accueil.php <u>http://www.sshrc-crsh.gc.ca/homeaccueil-eng.aspx http://www.stat.gouv.qc.ca</u>.

Acknowledgment: The authors wish to thank David Gunnell, PhD, of the University of Bristol, for his contribution to the paper and all the participants of the QLSCD.

Financial Disclosure: Drs. Geoffroy, Boivin, Arseneault, Turecki, Vitaro, Brendgen, Renaud, Séguin, Tremblay, and Côté report no biomedical financial interests or potential conflicts of interest.