

1 This is a pre-proof corrected manuscript, as accepted for publication, of an article published by  
2 Taylore & Francis in *Journal of Sports Sciences* on 9<sup>th</sup> March 2015, available online:

3 [http://www.tandfonline.com/doi/abs/10.1080/02640414.2015.1013051?url\\_ver=Z39.88-  
2003&rfr\\_id=ori:rid:crossref.org&rfr\\_dat=cr\\_pub%3dpubmed#.Vbtks03bLcs](http://www.tandfonline.com/doi/abs/10.1080/02640414.2015.1013051?url_ver=Z39.88-<br/>4 2003&rfr_id=ori:rid:crossref.org&rfr_dat=cr_pub%3dpubmed#.Vbtks03bLcs)

5 **PLEASE REFER TO THE PUBLISHED VERSION FOR CITING PURPOSES**

6  
7  
8  
9 **Title: The impact of in-season national team soccer play on injury and player availability in a  
10 professional club**

11  
12 **Authors:** Christopher Carling<sup>1,2</sup>, Alan McCall<sup>2,3</sup>, Franck Le Gall<sup>2</sup>, Gregory Dupont<sup>2,3</sup>

13  
14 **Institution:**

15 <sup>1</sup>Institute of Coaching and Performance, University of Central Lancashire, UK

16 <sup>2</sup>LOSC Lille Métropole Football Club, Research and Development Department, Camphin-en-Pévèle,  
17 France

18 <sup>3</sup>Univ Lille Nord de France, 59000 Lille, France - UDSL, EA 4488

19  
20 **Corresponding author:**

21 Christopher Carling

22 Correspondance: LOSC Lille Metropole Football Club, Domain de Luchin, Camphin-en-Pévèle, 59780,  
23 France.

24 Phone: 00.33.6.0392 1863

25 Fax: 00.33.1.4891 0793

26 Email: christopher.carling@gmail.com

27

1 **The impact of in-season national team soccer play on injury and player availability in a**  
2 **professional club**

3  
4 **Running head:** Injury in national team soccer

5  
6 **Abstract**

7 This study investigated the impact of in-season national team duty on injury rates and player availability  
8 in a professional soccer club. Time-loss injuries and exposure time during club and national team duties  
9 were recorded prospectively over 5-seasons (2009-2014). A time-loss injury was sustained by 37.7% of  
10 squad members participating in national duty; all injuries occurring in match-play. The incidence (per  
11 1000hours exposure) for national team player match-play injuries did not differ ( $p=0.608$ ) to that for all  
12 players in club competitions: 48.0 (95% CI 20.9-75.5) vs. 41.9 (95% CI 36.5-47.4), incidence rate  
13 ratio=1.2 (CI: 0.8-2.4). The majority (58%) of national team injuries resulted in a layoff  $\leq 1$  week. Of all  
14 working days lost to injury generally, 5.2% were lost through injury on national duty. Injury incidence  
15 in the week following national duty was comparable ( $p=0.818$ ) in players participating or not: 7.8 (95%  
16 CI 3.6-12.0) vs. 7.1 (95% CI: 4.6-9.6), incidence rate ratio=1.1 (CI: 0.7-2.7). While approximately forty  
17 percent of participating players incurred a time-loss injury on national duty, no training injuries were  
18 sustained and injuries made up a negligible part of overall club working days lost to injury. Following  
19 duty, players had a similar injury risk to peers without national obligations.

20  
21 Key terms: injuries, performance, football, match congestion, fatigue

## 1 **Introduction**

2 Soccer injuries result from a complex interaction of multiple internal and external risk factors  
3 and player fatigue is perceived to be one of the most important factors related to injury risk in  
4 contemporary professional soccer (McCall et al, 2014). Indeed, clubs can play over 60 competitive  
5 matches per season with some players participating in up to 50 matches (Strudwick, 2012). In addition  
6 to their domestic League and Cup programmes and European competition, many professional players  
7 are regularly exposed to national team obligations both in-season (e.g., national team tournament  
8 qualification and friendly matches) and post-season (e.g., FIFA World Cup, UEFA European  
9 Championships). In a one-season study of injury in European professional soccer, a total of 148 (56%)  
10 out of 266 players were exposed to some form of national team play on at least one occasion (Waldén,  
11 Hägglund & Ekstrand, 2005). While injury rates in post-season national team soccer tournaments are  
12 generally higher in comparison to those reported at professional club standards (Dvorak, Junge, Derman  
13 & Schwelanus, 2011), little is known about injury rates and characteristics during in-season national  
14 team training sessions or qualification and friendly matches and their subsequent impact on player  
15 availability. Similarly, to our knowledge, no information is available on the proportion of players who  
16 actually sustain an injury while on in-season national duty (training and match-play), the characteristics  
17 of injuries incurred, or the subsequent effect of these injuries on availability for future training sessions  
18 and competition in their club.

19 In-season national team matches are often closely preceded and followed by club matches.  
20 These intensive in-season competitive schedules frequently lead to periods of match congestion thereby  
21 requiring some players to play 2 matches per week consecutively over several weeks. Research in a  
22 Champions League team (Dupont et al., 2010) reported injury rates over 6-times higher when players  
23 participated in 2 club matches per week compared to only 1 match per week (25.6 vs. 4.1 per 1000  
24 hours exposure). A study by Bengtsson and colleagues (2013) also demonstrated a strong association  
25 between muscle injury rates and the recovery time available between successive matches (according to  
26 the number of days) in 27 professional European clubs. In contrast, no significant difference was  
27 reported in the incidence of injury in match (27.2 vs. 33.8 per 1000 hours) or training sessions (4.1 v 6.2  
28 per 1000 hours) in players exposed or not to national duty over the course of a competitive season  
29 despite the national team players having played significantly more matches (42 vs. 28) (Waldén et al,  
30 2005). Up to now, no study has investigated the potential effect of competitive schedules prior to  
31 national team duties in which players eventually sustained an injury or national team participation on  
32 injury rates during club training sessions and match-play immediately following duty. This information  
33 could eventually aid medical and coaching practitioners in making informed judgements on player  
34 rotation and recovery strategies around national team duties.

35 The aims of this study were to document injury rates during in-season national team duties and  
36 investigate the effects of duty on subsequent injury risk and player availability in a professional soccer  
37 club.

## 1 **Methods**

2 In this prospective observational study, injuries in male professional soccer players belonging to the  
3 first-team squad of a French Ligue 1 Club (highest competitive standard in France) sustained on both  
4 national team soccer duty and in club competitions were investigated. While all data arose as a  
5 condition of employment in which players were routinely monitored over the course of the competitive  
6 season (Winter & Maughan, 2009), approval for the study from the present club and ethics committee  
7 clearance from the 'Comité de Protection des Personnes Nord Ouest IV' (France) were obtained.

8 Data were prospectively collected in a cumulated total of 130 players (59 individual players)  
9 who participated in one or more seasons over a five-season period (2009-10, 2010-11, 2011-12, 2012-  
10 13 and 2013-14). A cumulated total of 58 players (44.6%) out of the 130 participated in national duties  
11 in one or more seasons over the five-season period. The cumulated total of 58 players included 30  
12 individual players of which 17 (53.3%) belonged to European, 12 to African (43.3%) and 1 (3.3%) to  
13 South American national teams respectively. Altogether, 7 European, 6 African and 1 South American  
14 teams were individually represented.

15 Individual exposure time to official club competitions (domestic league and cups, and European  
16 Competition), pre-season friendly matches and training sessions was recorded for each individual player  
17 belonging to the first-team squad by the club's sports scientist. Individual exposure time to in-season  
18 national team match-play (including tournament qualification and friendly matches) was also recorded  
19 for each player. Due to the non-presence of club medical and sports science staff, player exposure time  
20 to training on national duty was estimated. After consultation with the participating national team  
21 players, an average of 1.0 hours duration per training session was used to calculate training exposure  
22 time.

23 Over the 5-season period, injuries sustained in club training and match-play were prospectively  
24 diagnosed and documented by the same sports physician. Injuries incurred during national duties were  
25 also diagnosed and documented on the player's return to the club. Essentially, diagnosis involved  
26 routine clinical assessment of players by the physician at the time of presentation. Where necessary,  
27 further examinations were ordered (e.g., sonography or magnetic resonance imaging) at the discretion  
28 of the physician to assist with diagnosis.

29 The definitions of injury, injury severity and recurrent injuries were based on those  
30 recommended by International Soccer Injury Consensus Groups (Fuller et al., 2006; Hagglund, Walden,  
31 Bahr & Ekstrand, 2005). Injury: time-loss injury resulting from playing soccer and leading to a player  
32 being unable to fully participate in future training or match play independent of whether a training  
33 session actually took place on the day following injury or the player was selected to play in the next  
34 match. The incidence of injury was reported as the number of injuries per 1000 player-hours exposure  
35 to play. The severity of the injury (layoff time) was determined according to the number of days the  
36 player was absent from and unable to take full part in training or competition: minimal=1-3 days,  
37 mild=4-7 days, moderate=8-28 days and severe=>28days. All injuries were followed until the final day

1 of rehabilitation. The player was considered injured until the club physician permitted full participation  
2 in collective training and availability for match selection. A recurrent injury (reinjury) was described as  
3 an injury of the same type and at the same site as an index injury and that occurred within 2 months  
4 after a player's return to full participation from the index injury. The cause of injuries was obtained  
5 using post-match visualisation of the match video when available or alternatively via direct questioning  
6 by the club physician of the player on the latter's return to the club.

7 All statistical analyses were conducted using SPSS for Windows Version 14.0 (SPSS Inc.,  
8 Chicago, IL, USA). Standard statistical procedures were used to calculate frequencies, means,  
9 proportions (%), standard deviations and ranges. Injury incidences (presented as injuries per 1000hours  
10 play) and incidence rate ratios are both presented with 95% confidence intervals using respective  
11 formulae presented in Dvorak et al (2011) and Knowles, Marshall & Guskiewicz, (2006). After  
12 normality of the data was tested, an unpaired t-test was used to compare injury incidences observed in  
13 national teams and at club level. A p-value <0.05 was considered statistically significant. The G\*Power  
14 3 (Faul, Erdfelder, Lang et al. 2007) software was used to determine statistical power ( $1-\beta > 0.80$  was  
15 considered large).

## 16 **Results**

17 **Results**  
18 Players participated either fully or partly in a total of 206 national team matches over the 5-year period.  
19 This resulted in a total participation time of 250.1 match hours of which 182.1 hours were spent in  
20 tournament qualifying matches and 68.0 hours in friendly matches.

21 Of the 30 individual squad members participating in national duty at some time over the 5-  
22 season period, 11 individual players (37.7%) sustained an injury resulting in time loss from play.  
23 Altogether, 372 time-loss injuries (231 in match-play & 141 in training) were reported over the 5-  
24 seasons with a total of 12 injuries sustained in national team match-play accounting for 3.2% of all  
25 injuries sustained. No time-loss injuries were sustained in national team training sessions (estimated  
26 exposure time: 679 hours). Of the 12 injuries, 3 were reported in friendly matches and 9 in qualifying  
27 matches, 8 (66.7%) were diagnosed as muscle strains and 3 (25.0%) as sprains respectively, 8 (66.7%)  
28 were sustained to the players dominant side, 10 (83.3%) were as a result of non-contact trauma and 2  
29 were considered reoccurrences (16.7%) (Table 1). The upper leg was the most common injury location  
30 (5 or 41.7% of which 3 affected the hamstrings and 2 the quadriceps muscles). In the week prior to the  
31 12 injury occurrences on national team duty, the injured and non-injured national team players had both  
32 played an identical mean number (and standard deviation) of  $0.9 \pm 0.6$  matches.

33 The incidence of injuries incurred in national team training was 0.0 per 1000hours play while at  
34 club level this was 3.0 (CI 2.8-3.3). The incidences of match injury occurring in national team soccer  
35 compared to that in all players in club match-play were 48.0 (CI 20.9-75.5) and 41.9 (CI: 36.5-47.4)  
36 respectively: incidence rate ratio of 1.2 (CI: 0.8-2.4). The t-test reported no significant difference

1 between these incidences ( $p=0.608$ ) while statistical power ( $1-\beta$ ) was 0.10. The respective incidences of  
2 injury in tournament qualifying and friendly matches were 49.4 (CI 32.9-65.9) and 44.2 (CI 18.7-69.6).

3 A total of 288 days layoff time due to injuries sustained in national team match-play was  
4 observed. The mean length of the injuries occurring in national team and club match-play respectively  
5 was  $24.0\pm 46.2$  days (range 2-169 days, 16.7% were classed as minimal, 41.7% as mild, 33.3% as  
6 moderate and 8.3% as severe) and  $8.9\pm 15.0$  days (range 1-94 days, 46.6% were classed as minimal,  
7 24.7% as mild, 22.4% as moderate and 6.4% as severe). Of the 12 injuries, 10 (83.3%) forced the player  
8 to miss the following club match. Of the total of 2856 training sessions, 4290 training hours, 638  
9 matches and 3494 working days (training and matches combined) missed through injury generally, 182  
10 sessions (6.4%), 281 hours (6.6%), 36 matches (5.6%) and 182 (5.2%) working days respectively were  
11 missed directly through injuries sustained on national team duty. When the 182 club training sessions  
12 and 281 training hours missed through injury sustained on national duty were coupled with the 1075  
13 training sessions and 1634 hours of club training missed while away on duty, this led to player absence  
14 from a total of 4.2% of all club training sessions and 4.3% of training hours.

15 In the week immediately following national team duty, players participated in 70.1% of their  
16 club's matches (115 out of 164 matches). Players completed at least 75-minutes in 67.0% (77 out of 115  
17 matches) and 90-minutes play in 55.7% (64 out of 115 matches) of matches respectively. Of the 29.9%  
18 of matches (49/164) in which players did not compete, 14.3% (7 out of 49) were directly missed  
19 through an injury sustained on national team duty. Altogether, 13 injuries (incidence per 1000hours  
20 play=7.8, CI 3.6-12.0) were sustained in match-play and training combined in the week directly  
21 following national duty compared to an incidence of 7.1 (CI: 4.6-9.6) in the players who were not  
22 selected for their national team: incidence rate ratio of 1.1 (CI: 0.7-2.7). The t-test reported no  
23 significant difference between these incidences ( $p=0.818$ ) while statistical power ( $1-\beta$ ) was 0.08. Of the  
24 13 injuries sustained, 5 (38.5%) were diagnosed as muscle strains, 3 (23.1%) as contusions and 2  
25 (15.4%) as tendinopathies while 61.5% were due to non-contact trauma. The mean layoff time per  
26 injury incurred directly after national duty equalled  $13.9\pm 25.4$  days (range 1-94 days) with the majority  
27 (76.9%) resulting in a layoff of one week or less. These injuries subsequently forced players to miss a  
28 total of 136 training sessions (206 hours), 34 matches and 163 working days.

## 30 Discussion

31 In epidemiological studies conducted at the highest standards of soccer, it is important to account for  
32 the impact of exposure to national team duty and injuries subsequently incurred (Waldén et al., 2005).  
33 The aim of the present study was to analyse the impact of national team duty on injury and player  
34 availability in players belonging to a professional soccer club over a 5-season period. Main findings  
35 were that approximately 40% of players participating in national team match-play sustained an injury  
36 accounting for ~3% of all the club's injuries reported over the 5-seasons while no injuries were reported  
37 during national team training sessions. While the incidence of match-play injury was ~13% higher in

1 national versus club competitions, incidence rate ratios with corresponding confidence intervals and  
2 inferential statistical testing indicated no significant difference between rates. The majority of injuries  
3 sustained in national team matches resulted in absence from soccer for one week and or less. Of the  
4 total number of club working days (training and matches combined) missed through injury generally,  
5 ~5% were directly missed through injuries incurred in national team competition. The injury incidences  
6 in play over the week immediately following national team participation did not differ between players  
7 who had or had not participated in national team duty.

8 In the present study, ~3% of all injuries in the club's players sustained over 5-seasons occurred  
9 in national team circumstances, a figure similar to the 4% reported in 11 top European soccer clubs  
10 (Waldén et al., 2005). Altogether, ~38% of players sustained an injury during exposure to national duty  
11 which compares to 29% in a population of 65 players monitored prior to and during the 2002 World  
12 Cup (Ekstrand, Waldén & Hägglund, 2004). Positively, no injuries were reported during national team  
13 training sessions in contrast to an incidence of 3.0 per 1000 hours (CI 2.5-3.5) in club training. It must  
14 be acknowledged though that some slight injuries (0 days layoff) could have occurred in training and  
15 might not have been reported. An explanation already put forward elsewhere for this positive finding  
16 for training injury might be differences in the type and/or intensity of training sessions performed on  
17 national team duty. As the preparation time available for national friendly and qualification matches is  
18 habitually short, then it might be speculated that sessions were less intensive, more recovery-based and  
19 tactical in nature (Hägglund, Waldén & Ekstrand, 2009; Waldén, Hägglund & Ekstrand, 2007). In  
20 future studies, detailed logging and comparison of training activities and workloads in both club and  
21 national contexts would be useful to confirm or refute this suggestion.

22 Results showed that the incidence of injury occurring in national team soccer match-play did  
23 not differ statistically compared to that observed during club competitions (48.2 versus 41.9 injuries per  
24 1000 hours,  $p=0.608$ ). However, this difference represents a non-negligible 13% difference and could  
25 nevertheless be interpreted as partly confirming the trend for general higher risk of injury in national  
26 team tournament match-play in comparison to professional club soccer (Dvorak et al., 2011; Ekstrand,  
27 Hägglund & Waldén, 2011). Whilst to our knowledge no comparative data exist on injury rates specific  
28 to in-season national team qualification and friendly matches as presented here, the present incidence is  
29 within the range previously reported for post-season national team tournaments over the last decade or  
30 so: 2002 FIFA World Cup=50.7 (Junge, Dvorak & Graf-Baumann, 2004), 2006 FIFA World Cup=45.9  
31 (Dvorak, Jungen, Grimm & Kirkendall, 2007), FIFA 2010 World Cup=40.1 (Dvorak et al., 2011),  
32 FIFA 2009 Confederations Cup=60.1 (Theron, Schweltnus, Derman & Dvorak, 2013), EURO  
33 2004=36.0 (Waldén et al., 2007), and EURO 2008=41.6 (Hägglund et al., 2009). Additional larger-  
34 scale research in other professional clubs to quantify and compare injury rates during in-season national  
35 team duty and identify the extent of the risk internationally is warranted.

36 The high injury rates observed above for post-season national team tournaments have been  
37 linked to pressure and associated mental stress (Hägglund et al., 2009), as well as accumulated fatigue

1 and the density of matches played prior to the competition (Ekstrand et al., 2004). Match congestion at  
2 club level (2 versus 1 game per week) has also been associated with a six-fold increased risk of injury  
3 (Dupont et al., 2010). However, analyses of the 7-day period prior to the in-season national team  
4 matches in which the 12 injuries were incurred showed that both the national team players who  
5 subsequently sustained an injury and those who did not, had, on average, played a single game. Thus, it  
6 would seem that exposure time to club match-play prior to national team duty was not linked to the  
7 injuries sustained in players participating in national team play.

8 We also investigated how involvement with national teams affected injury in participating  
9 players over the following week back in their club environment in comparison to peers not selected for  
10 duty. Similar rates of injury were reported (incidence per 1000 hours play: 7.8 vs. 7.1,  $p=0.818$ )  
11 suggesting that involvement in national teams had no effect on subsequent injury risk at club level. A  
12 reasonable explanation for this finding might be management of player participation in the following  
13 club game which tended to take place 3-4 days afterwards. Approximately 70% of the national team  
14 players participated in the club's next match but only about half of these completed the full 90-minutes  
15 of match-play. Out of the club's national team players habitually considered by the coaching staff as  
16 'first-team starters', ~82% participated in the club's next match following national duty but only 54%  
17 completed 90-minutes. Combined, these results imply that the present club coaching staff took into  
18 account prior national duties when selecting and especially substituting players potentially with the aim  
19 of reducing the recognised higher injury risk linked to match congestion (Bengtsson et al., 2013;  
20 Dupont et al., 2010) and towards the latter end of play (Ekstrand et al., 2011; Hawkins et al, 2001).  
21 Future work using interviews with coaching staff to determine their criteria for selection or not of  
22 players who had recent involvement at national team level is warranted.

23 The impact of an injury on a club can be considered in relation to its severity and the number of  
24 potential competitive matches missed (Hawkins et al., 2001). Here, the impact of injuries sustained  
25 during national duty in relation to all injuries sustained during club obligations on the frequency of  
26 training sessions (~6%) and hours (~7%), matches (~6%) and working days (~5%) subsequently missed  
27 can be considered minimal. In contrast, an average of 3.0 matches was subsequently missed per national  
28 team injury with ~83% of the injuries leading to absence in at least one club match. In national team  
29 versus club competition, ~42% of injuries were classed as moderate/severe ( $\geq 8$  days) compared to  
30 ~29%. This suggests that when an injury occurred in national team competition, its severity and  
31 subsequent effect on player availability was potentially greater. However, player absence and injury  
32 severity must be interpreted with in the context of the low number of injury cases occurring during  
33 national duty along with the presence of an important injury outlier. Indeed, when the outlier (169 days  
34 layoff time for a tendon injury) is removed from the analysis, the mean length of the injuries occurring  
35 in national team versus club match-play was relatively similar: 10.8 vs. 8.9 days.

36 Finally, of the 12 injuries reported in national team competition, ~80% were sustained in a non-  
37 contact situation and two-thirds (~67%) were muscle strains versus ~50% and ~40% in club match-play



1 respectively. It is difficult to formulate solid explanations for these discrepancies in injury patterns  
2 mainly again due to the small number of reported injuries and again additional research is necessary to  
3 verify this trend.

4 In light of the present findings, we feel that future research into the potential ‘knock-on’ effects  
5 of participation in tournaments such as the World Cup and Continental Championships on injury risk in  
6 professional soccer players over the course of the following season is warranted. Previous research has  
7 shown that forty-one per cent of injuries sustained in the European Championships 2006 to 2008  
8 resulted in absence after the tournament, with an average of nearly four weeks subsequently spent in  
9 rehabilitation (Hägglund et al., 2009). Research to examine whether players who participate in a large  
10 number of in-season qualification and friendly matches followed by a post-season tournament are  
11 consequently subjected to greater injury risk in the following season is warranted. We also suggest that  
12 analysis of the potential effects of travel (e.g, duration and distance, time zones) especially in players  
13 travelling to different continents on subsequent injury risk during exposure to national team play and on  
14 return to their clubs would be pertinent. Finally, a multifactorial model of sports injury etiology (Bahr  
15 & Homes, 2003) would help to explore the potential interrelationships between internal risk factors and  
16 their contribution to injury rates and patterns in national team players.

17 A limitation acknowledged at the outset of this study was that only one soccer club was  
18 investigated and the patterns observed are a reflection of this particular cohort. Sample size influences  
19 the power to detect real and significant effects and similar investigations involving a larger sample of  
20 professional clubs internationally to increase statistical power are necessary to verify the present  
21 findings. Indeed, power for statistical comparisons reported here was  $\leq 0.10$ . Despite prospective data  
22 collection from over 200 national team matches played over a 5-season period, only 12 injury cases  
23 were reported in 11 individual players which is insufficient to detect moderate to strong associations  
24 (20-50 injury cases) and small to moderate associations (~200 injured subjects) (Bahr & Holme, 2003).  
25 However, despite these limitations, this study has merit in that it is the first to investigate the impact of  
26 in-season national team duty on injury and player availability at professional soccer club level. In  
27 addition, the strength of this investigation was its five-season span and prospective nature and  
28 methodology the latter respecting internationally recommended injury recording systems (Fuller et al.,  
29 2006; Hägglund et al., 2005) thereby allowing these results to be compared with future research  
30 findings.

## 31 **Conclusion**

32 In conclusion, this study showed that while in the region of 40% of participating players sustained an  
33 injury on national team duty, no injuries were sustained in national team training sessions. National  
34 team match-play injury rates did not differ statistically compared to that in players in club competitions  
35 making up a negligible part of all injuries incurred in the club over a 5-season period. Following  
36

- 1 national duty, injury incidence in participating players was not greater than in peers who had no
- 2 national obligations.

## 1 **References**

- 2 Bahr, R., & Holme, I. (2003). Risk factors for sports injuries-a methodological approach. *British*  
3 *Journal of Sports Medicine*, 37, 384-392.
- 4 Bengtsson, H., Ekstrand, J., & Hagglund, M. (2013). Muscle injury rates in professional soccer  
5 increases with fixture congestion: an 11-year follow-up of the UEFA Champions League injury study.  
6 *British Journal of Sports Medicine*, 47, 743-747.
- 7 Dupont, G., Nedelec, M., McCall, A., McCormack, D., Berthoin, S., & Wisløff, U. (2010). Effect of 2  
8 soccer matches in a week on physical performance and injury rate. *American Journal of Sports*  
9 *Medicine*, 38, 1752-1758.
- 10 Dvorak, J., Junge, A., Derman, W., & Schwellnus, M. (2011). Injuries and illnesses of soccer players  
11 during the 2010 FIFA World Cup. *British Journal of Sports Medicine*, 45, 626-630.
- 12 Dvorak, J., Jungen A., Grimm K., & Kirkendall, D. (2007). Medical report from the 2006 FIFA World  
13 Cup Germany. *British Journal of Sports Medicine*, 41, 57-81.
14. Ekstrand, J., Waldén, M., & Hägglund, M. (2004). A congested soccer calendar and the wellbeing of  
15 players: Correlation between match exposure of European soccerers before the World Cup 2002 and  
16 their injuries and performances during that World Cup. *British Journal of Sports Medicine*, 38, 493-  
17 497.
- 18 Ekstrand, J., Hägglund, M., & Waldén, M. (2011). Injury incidence and injury patterns in professional  
19 soccer: the UEFA injury study. *British Journal of Sports Medicine*, 45, 553-558.
- 20 Faul, F., Erdfelder, E., Lang, A., & Buchner, A. (2007). G\*Power 3: a flexible statistical power analysis  
21 program for the social, behavioral, and biomedical sciences. *Behavioral Research Methods*, 39, 175-  
22 191.
- 23 Fuller, C.W., Ekstrand, J., Junge, A., Andersen, T.E., Bahr, R., Dvorak, J., Hägglund, M., McCrory, P.,  
24 & Meeuwisse, W.H. (2006). Consensus statement on injury definitions and data collection procedures  
25 in studies of soccer (soccer) injuries. *Clinical Journal of Sports Medicine*, 16, 97-106.
- 26 Hagglund, M., Walden, M., Bahr, R., & Ekstrand, J. (2005). Methods for epidemiological study of  
27 injuries to professional soccer players: developing the UEFA model. *British Journal of Sports*  
28 *Medicine*, 39, 340-346.
- 29 Hägglund, M., Waldén, M., & Ekstrand, J. (2009). UEFA injury study-an injury audit of European  
30 Championships 2006 to 2008. *British Journal of Sports Medicine*, 43, 483-489.
- 31 Hawkins, R.D., Hulse, M.A., Wilkinson, C., Hodson, A., Andersen, T., & Bahr, R. (2001). The  
32 association soccer medical research programme: an audit of injuries in professional soccer. *British*  
33 *Journal of Sports Medicine*, 35, 43-47.
- 34 Junge, A., Dvorak, J., & Graf-Baumann T. Soccer injuries during the World Cup 2002. (2004).  
35 *American Journal of Sports Medicine*, 32(1 Suppl), 23S-7S.
36. Knowles, S.B, Marshall, S.W., & Guskievicz, K.M. (2006). Issues in estimating risks and rates in  
37 sports injury research. *Journal of Athletic Training*, 41, 207-215.

12. McCall, A., Carling, C., Nedelec, M., Davison M., Le Gall, F., Berthoin, S., & Dupont, G. (2014) Risk  
2 factors, testing and preventative strategies for non-contact injuries in professional football: current  
3 perceptions and practices of 44 teams from various premier leagues. *British Journal of Sports Medicine*,  
4 48, 1352-1357.
- 5 Strudwick, T. (2012). Contemporary issues in the physical preparation of elite players. In: M. Williams  
6 ed. *Science & Soccer III*. London. Routledge, 335-356.
- 7 Theron, N., Schwellnus, M., Derman, W., & Dvorak, J. (2013). Illness and injuries in elite soccer  
8 players--a prospective cohort study during the FIFA Confederations Cup 2009. *Clinical Journal of*  
9 *Sports Medicine*, 23, 379-383.
- 10 Waldén, M., Hägglund, M., & Ekstrand, J. (2005). UEFA Champions League study: a prospective study  
11 of injuries in professional soccer during the 2001-2002 season. *British Journal of Sports Medicine*, 39,  
12 542-546.
- 13 Waldén, M., Hägglund, M., & Ekstrand, J. (2007). Soccer injuries during European Championships  
14 2004-2005. *Knee Surgery, Sports Traumatology, Arthroscopy*, 15, 1155-1156.
- 15 Winter, E.M., & Maughan, R.J. (2009). Requirements for ethics approvals. *Journal of Sports Sciences*,  
16 27, 985.
- 17

Table 1 Description of 12 injuries sustained in professional soccer players on international duty over a 5-season period

Competitive season	Playing position	Layoff time (days)	Training sessions absent		N° matches absent	Working days absent	Injury characteristics					
			N°	Hours			Type	Location	Dominant side	Reinjury	Contact	Cause
2009-10	Midfielder	13	8	14	1	7	Sprain	Toe	Yes	No	Yes	Tackle
2009-10	Forward	6	3	5	1	4	Strain	Pelvis	Yes	No	Yes	Collision
2010-11	Forward	2	2	3	0	2	Strain	Hamstring	No	No	No	Unknown
2010-11	Defender	2	2	3	1	2	Strain	Neck	Yes	No	No	Collision
2010-11	Forward	14	9	14	1	10	Strain	Hamstring	Yes	No	No	Kicking ball
2011-12	Defender	8	5	8	1	6	Strain	Quadriceps	Yes	No	No	Unknown
2012-13	Midfielder	17	5	8	1	6	Sprain	Knee	Yes	No	No	Tackled
2012-13	Defender	25	13	21	5	18	Strain	Foot	No	No	No	Change in direction
2012-13	Forward	7	9	14	1	9	Strain	Hamstring	No	Yes	No	Acceleration
2012-13	Midfielder	20	14	23	5	19	Sprain	Ankle	Yes	No	Yes	Collision
2012-13	Forward	169	108	162	19	95	Tendinopathy	Ankle	Yes	Yes	No	Change in direction
2013-14	Forward	5	4	6	0	4	Strain	Quadriceps	No	No	No	Change in direction