

Movement & Sport Sciences – Science & Motricité  
 © ACAPS, EDP Sciences, 2015  
 DOI: 10.1051/sm/2015029

# Side to side differences in hamstring muscle kinematics during maximal instep soccer kicking

Jonathan Sinclair

Division of Sport, Exercise and Nutritional Sciences, School of Sport Tourism and Outdoors, University of Central Lancashire, UK

Received 26 June 2015 – Accepted 29 September 2015

**Abstract.** Hamstring strains are a common non-contact injury in soccer. The current study investigates bilateral differences in hamstring kinematics during maximal instep kicking. Thirteen male soccer players performed maximal instep kicks with their dominant and non-dominant limbs. Muscle-tendon kinematics of the four hamstring muscles during the kick movement were quantified using OpenSim software. Differences between dominant and non-dominant limbs were examined using paired t-tests. The results revealed that the biceps femoris long head (dominant =  $165.28 \pm 62.46$  & non-dominant =  $137.65 \pm 52.17\%$ ), semimembranosus (dominant =  $220.75 \pm 43.35$  & non-dominant =  $131.23 \pm 36.74\%$ ) and semitendinosus (dominant =  $90.95 \pm 16.69\%$  and non-dominant =  $80.47 \pm 15.99\%$ ) experienced significantly greater strain when using the dominant limb. The current investigation provides key information regarding the mechanics of the hamstring group during maximal instep kicking, indicating that kicking with the dominant limb may place soccer players at increased risk from hamstring strain injury.

**Key words:** Hamstring, soccer, muscle-tendon, muscle strain

**Résumé.** Différence bilatérale dans la cinématique des ischio-jambiers lors d'une frappe au pied chez des joueurs de football masculin.

Les blessures aux muscles ischio-jambiers sont classiques au football. La présente étude analyse les différences bilatérales dans la cinématique des ischio-jambiers lors d'une frappe du pied maximale en football. Treize joueurs de football masculins ont réalisé des frappes maximales avec leurs membres dominants et non dominants. La cinématique du complexe muscle-tendon de quatre muscles des ischio-jambiers a été analysée lors du mouvement en utilisant le logiciel OpenSim. Les différences entre les membres dominants et non dominants ont été examinées à l'aide de tests *t* appariés. Les résultats ont révélé que les longs biceps fémoraux (côté dominant =  $165,28 \pm 62,46$ ; côté non dominant =  $137,65 \pm 52,17 \%$ ), les semi-membraneux (côté dominant =  $220,75 \pm 43,35$ ; côté non dominant =  $131,23 \pm 36,74 \%$ ) et les semi-tendineux (côté dominant =  $90,95 \pm 16,69$ ; côté non dominant =  $80,47 \pm 15,99 \%$ ) subissent plus de contraintes lorsque le membre dominant est utilisé. Ces données fournissent des informations relatives à la mécanique des ischio-jambiers pendant une frappe maximale du pied et indiquent qu'une frappe avec le membre dominant en football peut entraîner des risques accrus de blessures au niveau des ischio-jambiers.

**Mots clés :** Ischio-jambiers, football, muscle-tendon, blessure musculaire

## 1 Introduction

Instep kicking is a skill that is fundamental to soccer performance and represents the most commonly used kicking technique in soccer (Kellis & Katis, 2007; Lees & Nolan, 1998; Lees, Asai, Andersen, Nunome, & Sterzing, 2010). It is important to generate high ball velocities when executing instep kicks as this improves the likelihood of scoring

by reducing the amount of time that the goalkeeper has to react (Sinclair, Taylor, *et al.*, 2014).

As part of their typical training regimen, soccer players are required to develop competency in kicking with both limbs (Carey, *et al.*, 2001). Despite this, soccer players will typically demonstrate limb dominance in kicking mechanics (Dorge, Anderson, Sorensen, & Simonsen, 2002; Sinclair, Fewtrell, *et al.*, 2014). The unilateral

8	9	10	11	12	13	14	15
---	---	----	----	----	----	----	----

1 nature of soccer kicking has been proposed as a con-  
 2 tributing factor to the aetiology of injury in soccer players  
 3 (Dorge, *et al.*, 2002). In relation to most other sports soc-  
 4 cer is associated with a high rate of injury which ranges  
 5 from 3.7–29.1 injuries per 1000 hours of game and train-  
 6 ing activity (Agel, Evans, Dick, Putukian, & Marshall,  
 7 2007). Aetiological analyses investigating injury locations  
 8 in soccer have shown that 60–80% of injuries occur in  
 9 the lower extremities (Agel, *et al.*, 2007; Dick, Putukian,  
 10 Agel, Evans, & Marshall, 2007).

11 The majority of muscle injuries in soccer are non-  
 12 contact in nature (Ueblacker, Mueller-Wohlfahrt, &  
 13 Ekstrand, 2015). Hamstring strains are known to be the  
 14 most common non-contact injury in soccer (Arnason,  
 15 Andersen, Holme, Engebretsen, & Bahr, 2008; Dadebo,  
 16 White, & George, 2004; Ekstrand & Gillquist, 1982;  
 17 Ekstrand, Hagglund, & Walden, 2011; Orchard & Seward,  
 18 2002; Orchard, Wood, Seward, & Broad, 1998; Seward,  
 19 Orchard, Hazard, & Collinson, 1993). Strain injuries  
 20 to the hamstring muscles are characterized by pain  
 21 in the posterior aspect of the thigh with accompany-  
 22 ing damage to the hamstring muscle fibres (Verrall,  
 23 Slavotinek, Barnes, Fon, & Spriggins, 2001). Hamstring  
 24 strain injuries range in seriousness from grade I which  
 25 is characterized by microscopic tearing and minor loss  
 26 of muscle function through to grade III which repre-  
 27 sents a full muscle rupture with complete loss of func-  
 28 tion (Blankenbaker & Tuite, 2010). Aetiological research  
 29 has shown that hamstring strains occur at a rate of  
 30 3.0–4.1 per 1000 hours of match play and 0.4–0.5 per  
 31 1000 hours of training (Arnason, Gudmundsson, Dahl, &  
 32 Johannsson, 1996; Arnason, *et al.*, 2004).

33 Hamstring strains occur as a function of exces-  
 34 sive muscle lengthening during eccentric contractions  
 35 (Heiderscheit, Sherry, Silder, Chumanov, & Thelen 2010;  
 36 Mueller-Wohlfahrt, *et al.*, 2013; Liu, Garrett, Moorman,  
 37 & Yu, 2012). Therefore, sports motions that require  
 38 frequent hamstring muscle lengthening may serve as  
 39 a precursor for aetiology of hamstring muscle strains  
 40 (Garrett, 1990; Garrett, Safran, Seaber, Glisson, &  
 41 Ribbeck 1987; Mair, Seaber, Glisson, & Garrett, 1996).  
 42 Clinical research has shown that the extent of muscle fibre  
 43 strain and the rate of muscle fibre lengthening are pri-  
 44 mary determinants of muscle strain injuries (Liu, *et al.*,  
 45 2012). Therefore rapid eccentric hamstring actions that  
 46 are associated with maximal velocity kicking have been  
 47 linked to the aetiology of hamstring injuries in soccer  
 48 players (Orchard & Seward, 2002).

49 A small number of investigations have examined the  
 50 kinematics of the hamstring muscle group during sports  
 51 movements. Yu, *et al.* (2008) examined the mechanics of  
 52 the hamstring muscles during sprinting. Their findings  
 53 showed that the risk for hamstring muscle strain injuries  
 54 is greatest during the late stance and late swing phases  
 55 of overground sprinting. Higashihara, Nagano, Takahashi,  
 56 & Fukubayashi (2014) investigated the effects of forward  
 57 trunk lean on hamstring muscle kinematics during sprint-  
 58 ing. They showed that the strain load imposed on the

biceps femoris long head and semimembranosus mus- 59  
 cles was larger with forward trunk lean which lead to 60  
 the conclusion that injury risk in these specific muscles 61  
 may be enhanced. Similarly, Chumanov, Heiderscheit, 62  
 and Thelen (2011) studied hamstring muscle strain dur- 63  
 ing high velocity running. Their findings showed that the 64  
 greatest strain loads exist during the swing phase of run- 65  
 ning which led to the conclusion that the hamstrings are 66  
 most susceptible to injury during this phase of the gait 67  
 cycle. 68

69 There is currently a paucity of information regarding  
 70 the mechanics of the hamstring muscle group during kick-  
 71 ing movements nor is there any consideration given to the  
 72 potential bilateral differences that may exist in hamstring  
 73 kinematics. Therefore the aim of the current study was to  
 74 investigate bilateral differences in the kinematics of the  
 75 hamstring group during maximal instep kicking.

## 2 Methods 76

### 2.1 Participants 77

78 Fifteen male soccer players (age =  $18.20 \pm 1.0$  years;  
 79 height =  $1.79 \pm 0.11$  m; body mass =  $74.65 \pm 5.54$  kg)  
 80 were examined whilst performing maximal instep kicks  
 81 into a regulation goal with their right (dominant) and  
 82 left (non-dominant) foot. All participants were academy  
 83 level players contracted to a professional club in England.

### 2.2 Procedure 84

85 Kinematic information was calculated using a ten cam-  
 86 era motion capture system (Qualisys<sup>TM</sup> Medical AB,  
 87 Goteburg, Sweden) at a rate of 500 Hz. Each participant  
 88 performed maximal in-step kicks with a 5 m run up into  
 89 a regulation sized soccer goal. Five kicking trials were  
 90 obtained from each participant from the dominant and  
 91 non-dominant limbs. Dynamic calibration of the motion  
 92 analysis system was performed before each data collection  
 93 session.

94 Retroreflective markers (19 mm diameter) were placed  
 95 at the C7, T12 and xiphoid process landmarks and also  
 96 positioned bilaterally onto the acromion process, iliac  
 97 crest, anterior superior iliac spine, posterior super iliac  
 98 spine, medial and lateral malleoli, medial and lateral  
 99 femoral epicondyles and greater trochanter. This allowed  
 100 the trunk, pelvis, thighs, shanks and feet to be defined.  
 101 Carbon-fibre tracking clusters comprising of four non-  
 102 linear retroreflective markers were positioned onto the  
 103 thigh and shank segments. Static calibration trials were  
 104 obtained with the participant in the anatomical position  
 105 in order for the positions of the anatomical markers to be  
 106 referenced in relation to the tracking clusters/markers.

**Table 1.** Hip and knee joint kinematics (means, standard deviations and 95C.I's) from the dominant and non-dominant limbs.

	Dominant			Non-dominant			% Difference	Effect size ( $p\eta^2$ )
	Mean	SD	95% C.I	Mean	SD	95% C.I		
Pelvis								
Angle at footstrike ( $^\circ$ )	10.52	1.47	9.71–11.33	11.52	1.19	10.86–12.18	9.10	0.24
Angle at maximum hip flexion ( $^\circ$ )	17.63	1.68	16.69–18.57	23.48	2.57	22.06–24.90	28.47	0.25
Range of motion ( $^\circ$ )	7.11	1.99	6.01–8.22	11.96	2.55	10.55–13.38	50.85	0.40
Hip								
Angle at footstrike ( $^\circ$ )	-14.25	1.44	-15.03--13.45	-11.57	0.58	-10.98--11.06	20.76	0.60
Angle at maximum hip flexion ( $^\circ$ )	68.55	7.30	64.50–72.59	60.73	6.39	57.20–64.27	12.09	0.35
Range of motion ( $^\circ$ )	82.79	6.60	79.14–86.45	72.30	6.53	68.69–75.91	13.53	0.50
Knee								
Angle at footstrike ( $^\circ$ )	81.00	6.36	77.48–84.52	81.07	7.91	76.69–85.45	0.08	0.01
Angle at maximum hip flexion ( $^\circ$ )	39.05	1.98	21.95–44.15	33.23	2.37	27.08–40.69	16.10	0.42
Range of motion ( $^\circ$ )	67.95	6.91	64.13–71.78	61.84	6.53	58.22–65.46	9.42	0.23

1 **2.3 Data processing**

2 Dynamic trials were digitized using Qualisys Track Man-  
 3 ager in order to identify anatomical and tracking mark-  
 4 ers then exported as C3D files to Visual 3D (C-Motion,  
 5 Germantown, MD, USA). Kinematic data was smoothed  
 6 using a cut-off frequency of 15 Hz with a non-phase shift  
 7 low-pass Butterworth 4th order filter. Five kicking trials  
 8 were obtained from each participant from the dominant  
 9 and non-dominant limbs. Kicking trials were defined from  
 10 the instance of stance limb touch down to maximum hip  
 11 flexion (R). Kinematic parameters from the kicking limb  
 12 that were extracted for statistical analysis were 1) angle  
 13 at stance limb footstrike, 2) angle at maximum hip flexion  
 14 and 3) range of motion representing the angular range of  
 15 motion from footstrike to maximum hip flexion.

16 OpenSim software was used to quantify muscle-tendon  
 17 lengths during the kicking movements (Delp, *et al.*, 2007).  
 18 Muscle kinematics were quantified using the gait2392  
 19 model using Opensim v3.2. This model corresponds to  
 20 the eight segments exported from Visual 3D and fea-  
 21 tures ninety two muscles, eighty six of which are cen-  
 22 tred around the lower extremities and six are associated  
 23 with the pelvis and trunk. The muscle properties were  
 24 modelled using the Hill recommendations based on the  
 25 associations between force-velocity-length (Zajac, 1989).  
 26 These muscle properties were then scaled based on each  
 27 participant’s height and body mass based on the recom-  
 28 mendations of Delp, *et al.*, (1990). Muscle-tendon lengths  
 29 are determined by the positions of their proximal and dis-  
 30 tal muscles muscle origins. The muscle-tendon complexes  
 31 which were evaluated as part of the current research were  
 32 the biceps femoris long head (LH), biceps femoris short  
 33 head (SH), semimembranosus and semitendinosus. Mus-  
 34 cle kinematic parameters that were extracted for statisti-  
 35 cal analysis were 1) change in length throughout the  
 36 kicking movement 2) strain (representative of the change  
 37 in length divided by original length at the start of the  
 38 movement) and 3) maximum lengthening velocity.

**2.4 Statistical analyses**

Descriptive statistics (means, standard deviations  
 and 95% confidence intervals) were calculated. To com-  
 pare differences in hamstring muscle kinematics between  
 the dominant and non-dominant limbs, paired t-tests  
 were utilized with statistical significance accepted at the  
 $p \leq 0.05$  level (Sinclair, Taylor, & Hobbs, 2013). Effect  
 sizes were quantified using partial eta<sup>2</sup> ( $p\eta^2$ ). In addition  
 to this percentage differences were also calculated. The  
 Shapiro-Wilk statistic for each condition confirmed  
 that the data were normally distributed. All statistical  
 procedures were conducted using SPSS 22.0 (SPSS Inc.,  
 Chicago, IL, USA).

**3 Results**

**3.1 Angular kinematics**

The hip joint at footstrike was shown to be significantly  
 ( $p < 0.05$ ,  $p\eta^2 = 0.60$ ) more extended in the dominant  
 foot compared to non-dominant. In addition the hip was  
 also found to be significantly ( $p < 0.05$ ,  $p\eta^2 = 0.35$ )  
 more extended at the instance of maximum hip flexion  
 in the dominant limb. Finally, the hip range of motion  
 was significantly ( $p < 0.05$ ,  $p\eta^2 = 0.50$ ) larger when us-  
 ing the dominant foot compared to non-dominant (Tab. 1,  
 Fig. 1a).

The knee joint was significantly more flexed ( $p < 0.05$ ,  
 $p\eta^2 = 0.42$ ) at the instance of peak hip flexion in the non-  
 dominant limb (Tab. 1, Fig. 1c). Finally at the pelvis,  
 range of motion was significantly greater ( $p < 0.05$ ,  $p\eta^2 =$   
 $0.40$ ) when kicking with the non-dominant limb (Tab. 1,  
 Fig. 1c).

**3.2 Hamstring kinematics**

For the biceps femoris LH muscle the dominant limb was  
 associated with a significantly ( $p < 0.05$ ,  $p\eta^2 = 0.47$ )

39

40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51


52

53

54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68

69

70  
71



**Fig. 1.** Joint and segment kinematics (a = hip, b = knee and c = pelvis) from the dominant and non-dominant limbs (black = dominant and dash = non-dominant) (FL = flexion and PT = posterior tilt).

**Table 2.** Hamstring kinematics (means, standard deviations and 95th the dominant and non-dominant limbs).

	Dominant			Non-dominant			%	Effect size ( $p\eta^2$ )
	Mean	SD	95% C.I	Mean	SD	95% C.I		
Biceps femoris LH change in length (m)	0.34	0.05	0.30–0.40	0.29	0.08	0.24–0.34	15.70	0.47
Biceps femoris SH change in length (m)	0.05	0.02	0.04–0.06	0.06	0.01	0.05–0.07	18.27	0.25
Semimembranosus change in length (m)	0.36	0.04	0.34–0.38	0.27	0.04	0.25–0.29	29.88	0.71
Semitendinosus change in length (m)	0.32	0.03	0.29–3.34	0.28	0.04	0.26–0.30	10.95	0.39
Biceps femoris LH strain (%)	165.28	62.46	130.69–199.98	137.65	52.17	108.76–165.54	18.24	0.47
Biceps femoris SH strain (%)	25.76	10.68	19.85–31.67	30.40	6.88	26.59–34.21	16.52	0.24
Semimembranosus strain (%)	220.75	45.35	195.64–245.87	131.23	36.74	110.89–151.58	50.86	0.73
Semitendinosus strain (%)	90.95	16.69	81.71–100.19	80.47	15.99	71.61–89.32	12.23	0.37
Biceps femoris LH peak velocity (m/s)	1.53	0.06	1.31–1.74	1.55	0.02	1.39–1.68	1.38	0.08
Biceps femoris SH peak velocity (m/s)	1.57	0.18	1.47–1.67	1.60	0.13	1.53–1.67	1.30	0.08
Semimembranosus peak velocity (m/s)	2.69	0.11	2.58–2.78	2.72	0.10	2.60–2.83	1.13	0.07
Semitendinosus peak velocity (m/s)	3.20	0.21	3.08–3.33	3.41	0.15	3.30–3.50	6.28	0.22

1 greater change in length compared to the non-dominant  
 2 limb. In addition the findings also showed that the  
 3 strain experienced by the biceps femoris LH was sig-  
 4 nificantly ( $p < 0.05$ ,  $p\eta^2 = 0.47$ ) greater when using  
 5 the dominant limb (Tab. 2, Fig. 2a). In addition for  
 6 the semimembranosus the dominant limb was found to  
 7 have undergone a significantly ( $p < 0.05$ ,  $p\eta^2 = 0.71$ )  
 8 larger change in length. Also the strain experienced

by the semimembranosus was significantly ( $p < 0.05$ ,  
 $p\eta^2 = 0.73$ ) greater in the dominant limb compared to  
 non-dominant (Tab. 2, Fig. 2c). Finally, for the semi-  
 tendinosus the dominant limb was associated with a  
 significantly ( $p < 0.05$ ,  $p\eta^2 = 0.39$ ) larger change in  
 length. The strain experienced by the semitendinosus was  
 significantly ( $p < 0.05$ ,  $p\eta^2 = 0.37$ ) greater in the domi-  
 nant limb compared to non-dominant (Tab. 2, Fig. 2d).

9  
10  
11  
12  
13  
14  
15  
16




Fig. 2. Muscle-tendon lengths from the dominant and non-dominant limbs (black = dominant and dash = non-dominant).

#### 1 4 Discussion

2 The aim of the current study was to investigate bilateral  
 3 differences in the kinematics of the hamstring group dur-  
 4 ing maximal instep kicking. To the authors knowledge  
 5 this represents the first investigation to quantify ham-  
 6 string muscle kinematics during instep kicking. A study  
 7 of this nature may provide important information to soc-  
 8 cer clinicians regarding the aetiology of hamstring strain  
 9 injuries as a function of maximal kicking actions.

10 The first key observation is that all of the four primary  
 11 hamstring muscles tested in the current study exhibited  
 12 eccentric lengthening in an almost linear manner through-  
 13 out the kick movement. This is to be expected given the  
 14 joint observed joint/ segment kinematics during the in-  
 15 step kick movement; hamstring lengthening was required  
 16 support flexion and extension rotations of the hip and  
 17 knee joints and also the posterior tilt of the pelvic seg-  
 18 ment during the kick (Lees, *et al.*, 2010).

19 Of further importance is the finding that the dominant  
 20 limb was associated with significant increases in strain  
 21 magnitude of the biceps femoris LH, semimembranosus  
 22 and semitendinosus muscles. The strain imposed on the  
 23 hamstring muscle-tendon unit during the kick is a func-  
 24 tion of the flexion and extension patterns of at the hip  
 25 and knee joints (Opar, Williams, & Shield, 2012). Given  
 26 the proximal and distal attachment of the aforementioned  
 27 muscles to the ischial tuberosity and fibula/ tibial heads;

the increased angular range of the hip and extension of  
 the knee joint when using the dominant limb served to  
 enhance the strain imposed on the muscles.

28  
 29  
 30 Although differences in muscle strain were shown be-  
 31 tween the dominant and non-dominant limbs, the biceps  
 32 femoris LH, semimembranosus and semitendinosus mus-  
 33 cles all experienced a substantial degree of strain regard-  
 34 less of limb dominance. Given the proposed relationship  
 35 between muscle strain magnitude and the aetiology of  
 36 muscle strain injuries the current investigation provides  
 37 insight regarding the high incidence of hamstring strain  
 38 injuries in soccer (Orchard, *et al.*, 1998; Orchard &  
 39 Seward, 2002; Seward, *et al.*, 1993). Nonetheless, the  
 40 statistical analysis showed that the biceps femoris LH,  
 41 semimembranosus and semitendinosus muscles of the  
 42 dominant limb experience significantly greater strain,  
 43 leading to the conclusion that kicking with the dominant  
 44 limb may place soccer players at increased risk from ham-  
 45 string strain injury. Of further interest is the relatively  
 46 low amount of strain experienced by muscle-tendon unit  
 47 of the biceps femoris SH. It is hypothesized that this find-  
 48 ing relates to the unilateral nature of the biceps femoris  
 49 SH which attaches proximally to the lateral ridge of the  
 50 femur rather as opposed to the ischial tuberosity. There-  
 51 fore, this muscle unit is not involved to the same extent in  
 52 hip flexion or in posterior pelvic tilt and thus the extent  
 53 to which it is required to lengthen is reduced in relation  
 54 to the other hamstring muscles.  
 55




Fig. 3. Muscle-tendon velocities from the dominant and non-dominant limbs (black = dominant and dash = non-dominant).

1 There are some limitations to the current work which  
 2 should be acknowledged so that the observations can be  
 3 appropriately contextualized. Firstly the current investiga-  
 4 tion utilized an all-male sample which may limit its  
 5 generalizability. Barfield, *et al.* (2002) documented gender  
 6 differences in kicking kinematics during maximal instep  
 7 kicking. In addition to this clinical research investigating  
 8 the prevalence of sports injuries has shown that there are  
 9 gender differences in hamstring injury risk (Ristolainen,  
 10 *et al.*, 2010; Sallis, Jones, Sunshine, Smith, & Simon,  
 11 2001; Satterthwaite, Larmer, Gardiner, & Norton, 1996).  
 12 It is therefore recommended that the current investiga-  
 13 tion be repeated using a sample of female soccer players.

14 In addition whilst, musculoskeletal simulations have  
 15 the potential to improve our understanding of muscles be-  
 16 haviour during movement, there are some limitations to  
 17 this technique that should be recognised. Musculoskeletal  
 18 simulations utilize a generic model with a number of  
 19 mechanical assumptions such as constrained rotational  
 20 degrees of freedom, fiber pennation angles, joint articula-  
 21 tions and the origins and insertions of the muscle-tendons  
 22 may lead to incorrectly predicted muscle kinematics.  
 23 It is also important to recognise that muscle-tendon  
 24 lengthening is not necessarily linearly related to muscle  
 25 fiber strain because of the interactions between tendon  
 26 elasticity and muscle contraction states during movement  
 27 (Zajac, 1989).

In conclusion, although the mechanics of instep kicking have been examined extensively, the current knowledge regarding the mechanics of the hamstring muscles during this movement is limited. The present investigation therefore adds to the current knowledge by providing a comprehensive evaluation of hamstring kinematics during maximal instep kicking when using the dominant and non-dominant limbs. Importantly the current study showed that the amount of muscle strain in the biceps femoris LH, semimembranosus and semitendinosus muscles was significantly larger when kicking with the dominant limb. The current investigation therefore provides key information regarding the mechanics of the hamstring group during maximal instep kicking, which shows that when kicking maximally with the dominant limb soccer players may be at greater risk from hamstring strain injury.

## Bibliography

Agel, J., Evans, T.A., Dick, R., Putukian, M., & Marshall, S.W. (2007). Descriptive epidemiology of collegiate men’s soccer injuries: National Collegiate Athletic Association Injury Surveillance System, 1988–1989 through 2002–2003. *Journal of Athletic Training*, 42, 270–277.

28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51

1 Arnason, A., Gudmundsson, A., Dahl, H.A., & Johannsson, E.  
 2 (1996). Soccer injuries in Iceland. *Scandinavian Journal of*  
 3 *Medicine & Science in Sports*, 6, 40–45.

4 Arnason, A., Sigurdsson, S.B., Gudmundsson, A., Holme, I.,  
 5 Engebretsen, L., & Bahr, R. (2004). Risk factors for in-  
 6 juries in football. *American Journal of Sports Medicine*,  
 7 32, 5–16.

8 Arnason, A., Andersen, T.E., Holme, I., Engebretsen, L., &  
 9 Bahr, R. (2008). Prevention of hamstring strains in elite  
 10 soccer: an intervention study. *Scandinavian Journal of*  
 11 *Medicine & Science in Sports*, 18, 40–48.

12 Barfield, W.R., Kirkendall D.T. (2002). Kinematic instep kick-  
 13 ing differences between elite female and male soccer play-  
 14 ers. *Journal of Sports Science & Medicine*, 1, 272–279.

15 Blankenbaker, D.G., & Tuite, M.J. (2010). Temporal changes  
 16 of muscle injury. *Seminars in Musculoskeletal Radiology*,  
 17 14, 176–193.

18 Carey, D.P., Smith, G., Smith, D.T., Shepherd, J.W., Skriver,  
 19 J., Ord, L., & Rutland, A. (2001). Footedness in world soc-  
 20 cer: an analysis of France’98. *Journal of Sports Sciences*,  
 21 19, 855–864.

22 Chumanov, E.S., Heiderscheit, B.C., & Thelen, D.G. (2011).  
 23 Hamstring musculotendon dynamics during stance and  
 24 swing phases of high speed running. *Medicine & Science*  
 25 *in Sports & Exercise*, 43, 525–532.

26 Dadebo, B., White, J., & George, K.P. (2004). A survey of flex-  
 27 ibility training protocols and hamstring strains in profes-  
 28 sional football clubs in England. *British Journal of Sports*  
 29 *Medicine*, 38, 388–394.

30 Delp, S.L., Loan, J.P., Hoy, M.G., Zajac, F.E., Topp, E.L., &  
 31 Rosen J.M. (1990). An interactive graphics-based model  
 32 of the lower extremity to study orthopaedic surgical pro-  
 33 cedures. *IEEE Transactions Biomedical Engineering*, 37,  
 34 757–767.

35 Delp, S.L., Anderson, F.C., Arnold, A.S., Loan, P., Habib,  
 36 A., John, C.T., & Thelen, D.G. (2007). OpenSim: open-  
 37 source software to create and analyze dynamic simula-  
 38 tions of movement. *IEEE Trans Biomedical Engineering*,  
 39 54, 1940–1950.

40 Dick, R., Putukian, M., Agel, J., Evans, T.A., &  
 41 Marshall, S.W. (2007). Descriptive epidemiology of col-  
 42 legiate women’s soccer injuries: National Collegiate  
 43 Athletic Association Injury Surveillance System, 1988–  
 44 1989 through 2002–2003. *Journal of Athletic Training*, 42,  
 45 278–285.

46 Dorge, H.C., Anderson, T.B., Sorensen, H., & Simonsen, E.B.  
 47 (2002). Biomechanical differences in soccer kicking with  
 48 the preferred and the non-preferred leg. *Journal of Sports*  
 49 *Sciences*, 20, 293–299.

50 Ekstrand, J., & Gillquist, J. (1982). The frequency of muscle  
 51 tightness and injuries in soccer players. *American Journal*  
 52 *of Sports Medicine*, 10, 75–78.

53 Ekstrand, J., Hagglund, M., & Walden, M. (2011).  
 54 Epidemiology of muscle injuries in professional foot-  
 55 ball (soccer). *American Journal of Sports Medicine*, 39,  
 56 1226–1232.

Garrett, W.E., Safran, M.R., Seaber, A.V., Glisson, R.R., &  
 Ribbeck, B.M. (1987). Biomechanical comparison of stim-  
 ulated and non-stimulated skeletal muscle pulled to fail-  
 ure. *American Journal of Sports Medicine*, 15, 448–454.

Garrett, W.E. (1990). Muscle strain injuries: clinical and ba-  
 sic aspects. *Medicine & Science in Sports & Exercise*, 22,  
 436–443.

Heiderscheit, B.C., Sherry, M.A., Silder, A., Chumanov, E.S., &  
 Thelen, D.G. (2010). Hamstring strain injuries: recommen-  
 dations for diagnosis, rehabilitation, and injury preven-  
 tion. *Journal of Orthopaedic & Sports Physical Therapy*,  
 40, 67–81.

Higashihara, A., Nagano, Y., Takahashi, K., & Fukubayashi,  
 T. (2014). Effects of forward trunk lean on hamstring  
 muscle kinematics during sprinting. *Journal of Sports*  
*Sciences*, 33, 1366–1375.

Kellis, E., & Katis, A. (2007). Biomechanical characteristics  
 and determinants of instep soccer kick. *Journal of Sports*  
*Science and Medicine*, 6, 154–165.

Lees, A., & Nolan, L. (1998). The biomechanics of soccer: A  
 review. *Journal of Sports Sciences*, 16, 21–234.

Lees, A., Asai, T., Andersen, T.B., Nunome, H., & Sterzing,  
 T. (2010). The biomechanics of kicking in soccer: A review.  
*Journal of sports sciences*, 28, 805–817.

Liu, H., Garrett, W.E., Moorman, C.T., & Yu, B. (2012).  
 Injury rate, mechanism, and risk factors of hamstring  
 strain injuries in sports: A review of the literature. *Journal*  
*of Sport and Health Science*, 1, 92–101.

Mair, S.D., Seaber, A.V., Glisson, R.R., & Garrett, W.E.  
 (1996). The role of fatigue in susceptibility to acute mus-  
 cle strain injury. *American Journal of Sports Medicine*, 24,  
 137–143.

Mueller-Wohlfahrt, H.W., Haensel, L., Mithoefer, K.,  
 Ekstrand, J., English, B., McNally, S., & Ueblacker, P.  
 (2013). Terminology and classification of muscle injuries  
 in sport: a consensus statement. *British Journal of Sports*  
*Medicine*, 47, 6, 342–350.

Opar, M.D.A., Williams, M.D., & Shield, A.J. (2012).  
*Hamstring strain injuries. Sports Medicine*, 42, 209–226.

Orchard, J., Wood, T., Seward, H., & Broad, A. (1998).  
 Comparison of injuries in elite senior and junior Australian  
 football. *Journal of Science & Medicine in Sport*, 1, 83–88.

Orchard, J., & Seward, H. (2002). Epidemiology of injuries  
 in the Australian Football League, seasons 1997–2000.  
*British Journal of Sports Medicine*, 36, 39–44.

Ristolainen, L., Heinonen, A., Turunen, H., Mannström, H.,  
 Waller, B., Kettunen, J. A., & Kujala, U.M. (2010). Type  
 of sport is related to injury profile: A study on cross  
 country skiers, swimmers, long-distance runners and soc-  
 cer players. A retrospective 12-month study. *Scandinavian*  
*Journal of Medicine & Science in Sports*, 20, 384–393.

Sallis, R.E., Jones, K., Sunshine, S., Smith, G., & Simon,  
 L. (2001) Comparing sports injuries in men and women.  
*International Journal of Sports Medicine* 22, 420–423.

- 1 Satterthwaite, P., Larmer, P., Gardiner, J., & Norton, R. 22  
 2 (1996). Incidence of injuries and other health problems in 23  
 3 the Auckland Citibank marathon, 1993. *British Journal of* 24  
 4 *Sports Medicine*, 30, 324–326. 25
- 5 Seward, H., Orchard, J., Hazard, H., & Collinson, D. (1993). 26  
 6 Football injuries in Australia at the elite level. *The Medical* 27  
 7 *Journal of Australia*, 159, 298–301. 28
- 8 Sinclair, J., Taylor, P.J., & Hobbs, S.J. (2013). Alpha level 29  
 9 adjustments for multiple dependent variable analyses 30  
 10 and their applicability—a review. *International Journal of* 31  
 11 *Sports Science & Engineering*, 7, 17–20. 32
- 12 Sinclair, J., Fewtrell, D., Taylor, P.J., Atkins, S., Bottoms, L., 33  
 13 & Hobbs, S.J. (2014). Three-dimensional kinematic dif- 34  
 14 ferences between the preferred and non-preferred limbs 35  
 15 during maximal instep soccer kicking. *Journal of Sports* 36  
 16 *Sciences*, 32, 1914–1923. 37
- 17 Sinclair, J., Taylor, P.J., Atkins, S., Bullen, J., Smith, A., & 38  
 18 Hobbs, S.J. (2014). The influence of lower extremity kine- 39  
 19 matics on ball release velocity during in-step place kick- 40  
 20 ing in rugby union. *International Journal of Performance* 41  
 21 *Analysis in Sport*, 14, 64–72.
- Ueblacker, P., Mueller-Wohlfahrt, H. W., & Ekstrand, J. 22  
 (2015). Epidemiological and clinical outcome comparison 23  
 of indirect (‘strain’) versus direct (‘contusion’) anterior 24  
 and posterior thigh muscle injuries in male elite foot- 25  
 ball players: UEFA Elite League study of 2287 thigh in- 26  
 juries (2001–2013). *British Journal of Sports Medicine*, 27  
 doi:10.1136/bjsports-2014-094285. 28
- Verrall, G.M., Slavotinek, J.P., Barnes, P.G., Fon, G.T., & 29  
 Spriggins, A.J. (2001). Clinical risk factors for hamstring 30  
 muscle strain injury: a prospective study with correlation 31  
 of injury by magnetic resonance imaging. *British Journal* 32  
*of Sports Medicine*, 35, 435–439. 33
- Yu, B., Queen, R.M., Abbey, A.N., Liu, Y., Moorman, C.T., 34  
 & Garrett, W.E. (2008). Hamstring muscle kinematics 35  
 and activation during overground sprinting. *Journal of* 36  
*Biomechanics*, 41, 3121–3126. 37
- Zajac, F.E. (1989). Muscle and tendon: properties, mod- 38  
 els, scaling, and application to biomechanics and motor 39  
 control. *Critical Reviews in Biomedical Engineering*, 17, 40  
 359–411. 41