

Article

Post discharge medicines use review service for older patients: recruitment issues in a feasibility stud

Rutter, Paul, Ramsbottom, Helen Frances and Fitzpatrick, Ray

Available at <http://clock.uclan.ac.uk/14218/>

Rutter, Paul, Ramsbottom, Helen Frances and Fitzpatrick, Ray (2016) Post discharge medicines use review service for older patients: recruitment issues in a feasibility stud. International Journal of Clinical Pharmacy . pp. 1-5. ISSN 2210-7703

It is advisable to refer to the publisher's version if you intend to cite from the work.

<http://dx.doi.org/10.1007/s11096-015-0243-8>

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

International Journal of Clinical Pharmacy

Post Discharge Medicines Use Review (dMUR) Service for Older Patients: Recruitment Issues in a Feasibility Study --Manuscript Draft--

Manuscript Number:	IJCP-D-15-00507R2
Full Title:	Post Discharge Medicines Use Review (dMUR) Service for Older Patients: Recruitment Issues in a Feasibility Study
Article Type:	Short Research Report
Keywords:	hospital discharge; older people; Recruitment; pharmacist; Medication Review
Corresponding Author:	Helen Frances Ramsbottom, MPharm Southport and Ormskirk Hospitals NHS Trust Southport, UNITED KINGDOM
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Southport and Ormskirk Hospitals NHS Trust
Corresponding Author's Secondary Institution:	
First Author:	Helen Frances Ramsbottom, MPharm
First Author Secondary Information:	
Order of Authors:	Helen Frances Ramsbottom, MPharm Ray Fitzpatrick, BSc (Pharm), PhD Paul Rutter, BPharm, PhD
Order of Authors Secondary Information:	
Funding Information:	
Abstract:	<p>Abstract</p> <p>Background</p> <p>The community pharmacy Medicines Use Review (MUR) service in England has been identified as a way of providing support with medication to recently discharged patients; however initial uptake of post-discharge Medicines Use Reviews has been low.</p> <p>Objective</p> <p>To identify barriers to recruitment into a randomised controlled feasibility study of a hospital referral system to older patients' regular community pharmacists.</p> <p>Method</p> <p>Ward pharmacists at Southport District General Hospital identified patients aged over 65 to be approached by a researcher to assess eligibility and discuss involvement in the trial. Participants were randomised to referral for a post discharge MUR with their regular community pharmacist, or to standard discharge care. Reasons for patients not participating were collected.</p> <p>Results</p> <p>Over a 9-month period 337 potential participants were identified by ward pharmacists. Of these, 132 were eligible and 60 were recruited.</p> <p>Barriers to recruitment included competing priorities among ward pharmacists, and national restrictions placed on MURs e.g. housebound patients and those requiring carer support with medication. Lack of expected benefit resulted in a high proportion of patient refusals.</p> <p>Conclusions</p> <p>The current provisions for post discharge MURs exclude many older people from participation, including those possibly in greatest need.</p> <p>Unfamiliarity with the role of the pharmacist in transitional care may have affected patients' perceived 'cost-benefit' of taking part in this study.</p>

	<p>Impact of Findings on Clinical Pharmacy Practice</p> <p>*Engaging older patients in the hospital setting with the concept of a post discharge Medicines Use Review in community pharmacy is challenging.</p> <p>*Further support for hospital teams to identify and refer suitable patients for post discharge Medicines Use Reviews is necessary.</p> <p>*The current provisions for post discharge medicines use reviews in England appear to cater poorly for patients who are housebound and those who rely on others to support them with their medicines at home.</p>
<p>Response to Reviewers:</p>	<p>Reviewer 1: Comments and Responses</p> <p>1. In the abstract and on various other occasions throughout the manuscript, the participants are referred to as the 'elderly'. I would advise that this terminology be replaced with 'older adults' or 'older people', which would be regarded as more appropriate terms to use when referring to this population. Done</p> <p>2. In Figure 1, please change '10682 Medical Patients' to '10682 medical patients' and '337 Patients' to '337 patients', (lowercase), in keeping with the word formatting in the other boxes within the same figure. Done</p> <p>3. Line 68: 'Feedback from ward pharmacists via a focus group...' I would advise that you add (n=x) or similar, to indicate how many ward pharmacists provided feedback in the focus group. Done (n=7)</p> <p>4. Whilst this article predominantly focusses on recruitment being problematic due to it being a study attempting to recruit older adults, an important finding described in this research report was that, overall, the pharmacists only referred 3.2% of all patients admitted, which is a significantly lower proportion than one would expect given the wider patient population admitted to hospitals. As such, this clearly represented a major barrier to the recruitment process. As the author(s) states that recruitment of participants to the study was 'often forgotten about' by ward pharmacists (line 70), it is perhaps unclear to the reader what steps (if any) were taken to ensure that pharmacists were aware of their role in the recruitment process. This point is also pertinent as the author(s) also report that of the 337 patients referred to the researcher by the pharmacists, 105 were ineligible for inclusion as they did not meet the specified study inclusion criteria. As this represents nearly one third of all those who were referred, this would lead this reviewer to assume that the ward pharmacists involved in recruitment were not aware of the study's inclusion/exclusion criteria (as suggested in lines 35-36). To address this issue, I think this article would benefit from a comment in the discussion section to clarify exactly how and to what extent the pharmacists involved in the recruitment were informed of the recruitment process.</p> <p>The following comment has now been inserted into the discussion: The low referral rate was despite steps taken by the researcher to raise awareness of the study among ward pharmacists, including a 'briefing' session prior to its commencement and posters displayed around the pharmacy department detailing types of patients who might be suitable. A study 'pack' was also given to each pharmacist, containing background information about the study and its eligibility criteria. Additionally, email reminders were sent regularly to all ward pharmacists throughout the recruitment period</p> <p>In addition, this article would be improved if the author(s) could provide a short comment on whether they believe the approach taken with involving pharmacists in the recruitment process could have been improved, and if so, how?</p> <p>The following comment has now been inserted into the discussion: Further investigation of these barriers and possible solutions (for example, increased pharmacist time on wards to allow for patient screening or use of information technology to make referral easier) is ongoing.</p> <p>5. On line 98, please change 'pharmacist led' to 'pharmacist-led'. Done</p> <p>6. On lines 81-83, the author(s) argue that patients who have carers managing their medicines at home are 'frailer and more susceptible to adverse drug events'; this report would benefit from the inclusion of a reference here for this assertion, particularly as the author(s) also use this argument again to form part of their conclusion (line 128). If no specific reference exists to support the association, I would suggest that this sentence is re-worded to reflect that the proposed association is a (valid) assumption. No specific reference exists; first two paragraphs under subheading 'Patient exclusions' (lines 75-83 in original manuscript) have now been</p>

combined into a single paragraph to summarise the issue addressed (lines 91-96 in revised manuscript)

7. On lines 116-117, please add (1.5%) after 'only 2 patients...' as previous findings are presented in this way in the text (i.e. number followed by percentage in brackets). Similarly, on line 118: '11 patients...', please add the percentage for information. Done

8. On lines 75-78 the author(s) state that polypharmacy is common among older adults who have 'managed care at home'. The relevance of this statement, in the context of patients' ineligibility, is not clear to this reviewer. I would ask the author(s) to review and either remove or rephrase this point in order to clarify their intended meaning. The intended argument here is: the fact that patients who had carers managing their medicines had to be excluded is a cause for concern and needs addressing in terms of future service development. The combination of first two paragraphs under subheading 'Patient exclusions' (lines 75-83 in original manuscript) attempts to clarify this point: In fact, patients requiring carers to manage their medication may be more in need of a medication support service, as polypharmacy, including the use of inappropriate medication, is common among older populations who have 'managed care at home' [8].

9. In Table 1 please correct 'Participants community pharmacy...' to 'Participant's community pharmacy...' Done

10. Finally, I would draw the author(s) attention to the reference section of the research report. Please review and correct the formatting of references as per the requirements of the journal. The use of semi-colons, commas, spaces, full-stops, capital letters and page numbering varies throughout the reference list. Also, some journals are abbreviated and others are not. Also, in reference number 8, there is a '1' after 'Fialova D'. Reference list re-visited and re-formatted

Reviewer 2: Comments and Responses

Abstract:

1. In the methods section, please briefly state potential patients were visited by a researcher to assess eligibility and discuss involvement in the trial, and reasons for patients not participating were collected, or similar. Done
2. In the results section, please state that of the 337 potential participants, 132 were eligible and 60 were recruited. Done
3. Suggest first two sentences of conclusion are amended to summarise most common reasons and placed in the results section of the abstract. Done: following sentences now inserted in results: Barriers to recruitment included competing priorities among ward pharmacists, and national restrictions placed on Medicines Use Reviews, for example housebound patients and those requiring carer support with medication. Lack of expected benefit resulted in a high proportion of patient refusals.

Introduction:

4. If word count allows, it may assist international readers if the description of the MUR program was expanded. From memory the program is free for consumers - if so, this suggest this is stated. Lines 9-13 now read: In the UK, a national service known as Medicines Use Reviews (MUR) is offered by many community pharmacists which is free to patients and paid for centrally by the NHS. The MUR is a face-to-face consultation between a patient and pharmacist, designed to identify any problems or information needs patients have with their medicines and offer solutions [4].

5. Final sentence of introduction seems out of place (line 25) - suggest deleting or moving. Deleted, (along with preceding sentence in order to reduce word count)

6. The section of the aim which states "to investigate the logistics of setting up the study" could be a little more specific. Aim now reads: To identify barriers to recruitment into a randomised controlled feasibility study of a hospital referral system to older patients' regular community pharmacists, and identify areas for improvement of the service or study design.

Methods:

7. Line 43 (Exclusion criteria) suggest restarting numbering at 1. Done

8. Can the authors please describe the process after eligibility was assessed ie did the researcher talk to the patients, what information was provided (ie brochure) etc
The following text has been inserted (lines 44-47 of revised manuscript): The researcher approached eligible patients to discuss involvement in the study and provide an information leaflet. Patients were given up to 24 hours (or longer if requested) to consider participation. Reasons for refusal were collected via verbal discussion, entered into an Excel spreadsheet and categorised by the researcher.

9. Please describe the method used to collect reasons for refusal ie simple discussion with patient or was a survey completed etc.
See response to point 8 above

Editor comments and Responses

- The results section in the abstract is not in line with the described Objective of the study.
Objective now reads: To identify barriers to recruitment into a randomised controlled feasibility study of a hospital referral system to older patients' regular community pharmacists. Results section also amended as per reviewer 2, comment 3.
- Please insert 3-5 Statements on the impact of your findings on practice, patients or professionals, and add them at the end of the abstract in Editorial Manager. Done
- The references to the reports need more detail, see the Vancouver style formatting instructions.
All references checked and now in the authors' opinion are correct
- Replace the titles of some journals by the appropriate abbreviation (ref 9 and 10).
Done

Miss Helen Ramsbottom
Pharmacy Department, Southport and Ormskirk NHS Trust
Town Lane
Southport
PR8 6PN

School of Pharmacy
University of Wolverhampton
Wulfruna St
Wolverhampton
WV1 1SB

14th December 2015

Dear Editor,

Please accept this revised submission of a short research report:
Post Discharge Medicines Use Review (dMUR) Service for Older Patients: Recruitment Issues in a Feasibility Study.

The paper reports on the recruitment phase of a randomised controlled feasibility study of a post-discharge medicines support service for patients aged 65 and older. The study took place in England and involved referral of eligible patients from hospital to community pharmacy for post discharge Medicines Use Review (dMUR), a nationally commissioned service which has had low take-up in practice. Participants were randomised to referral for a dMUR, or to standard discharge care.

The authors feel this article sits well within the scope of the International Journal of Clinical Pharmacy as it describes the challenges faced by pharmacy practice researchers attempting to set up a controlled study of a medicines management service for one of the patient groups most vulnerable to adverse medication events; ie elderly patients who are being transferred from one care setting to another. This research is therefore of national importance within England but also of national importance in a world with an aging population who are taking more and more medicines.

Submission Statement

This article is not in submission elsewhere in any form. All authors have contributed significantly to the publication and are in agreement with this submission.

The authors are:

Miss Helen Ramsbottom, Clinical Pharmacist, Southport and Ormskirk Hospital NHS Trust and PhD researcher, University of Wolverhampton (Addresses as above)

Prof Ray Fitzpatrick, Professor of Clinical Pharmacy, School of Pharmacy, Keele University, Staffordshire, UK

Prof Paul Rutter, Professor of Pharmacy Practice, School of Pharmacy, Wolverhampton, UK

Main article length: 1530 words

Abstract length: 215 words

Number of references: 10; word count: 307

Yours sincerely

Helen Ramsbottom

1 Word count 1530

2 **Post Discharge Medicines Use Review (dMUR) Service for Older Patients: Recruitment**

3 **Issues in a Feasibility Study**

4 **Introduction**

5 The likelihood that an older medical patient will be discharged from hospital on the same medication
6 they were admitted on is less than 10%, and this extensive medication regimen modification may
7 predispose to an increased risk of mortality [1].

8 Improved discharge information and patient education are core principles for reducing unintentional
9 medication changes, patient harm and hospital re-admissions [2,3]. In the UK, a national service
10 known as Medicines Use Reviews (MUR) is offered by many community pharmacists which is free to
11 patients and paid for centrally by the NHS. The MUR is a face-to-face consultation between a
12 patient and pharmacist, designed to identify any problems or information needs patients have with
13 their medicines and offer solutions [4].

14 Since 2011, the MUR service has targeted specific patient groups, identified as being at particular
15 risk of medication related problems or poor adherence, including patients recently discharged from
16 hospital [4]. However, early evidence suggests uptake of post discharge MURs (dMURs) has been
17 poor; pilot studies have shown that less than 3% of patients 'signposted' to the service by hospital
18 pharmacy staff receive a dMUR [5].

19 In 2012, a Department of Health steering group recommended that formal communication channels
20 between hospital and community pharmacy should be established so that dMURs become an integral
21 part of the medicines pathway [2]. However, at present, most NHS hospital Trusts do not appear to
22 have such formalised systems in place [6].

23 **Aim**

24 To identify barriers to recruitment into a randomised controlled feasibility study of a hospital referral
25 system to older patients' regular community pharmacists, and identify areas for improvement of the
26 service or study design.

27 **Ethical Approval**

28 Ethics approval for the study was obtained from the Northwest Research Ethics Committee (Ref
29 13/NW/0779).

30 **Method**

31 All pharmacists working on medical wards at Southport and Ormskirk NHS Trust, England were asked
32 to identify patients over the age of 65 years who in their professional opinion might benefit from
33 receiving a dMUR. These patients were referred on to the researcher (HR), who visited each patient
34 on the wards to assess eligibility.

35 *Inclusion criteria (All must be met for eligibility)*

- 36 1. Taking at least one long-term medication prior to admission
- 37 2. Taking five or more medicines OR one or more changes to medication during hospital stay
- 38 3. Patient's community pharmacy signed up to received referrals from the hospital pharmacy.

39 *Exclusion Criteria*

- 40 1. Patient will have a relative or carer managing their medication once back at home
- 41 2. Patient unable to name a regular community pharmacy
- 42 3. Patient unable to visit their community pharmacy for an MUR (unless community pharmacy offers
43 telephone or domiciliary MURs).

44 The researcher approached eligible patients to discuss involvement in the study and provide an
45 information leaflet. Patients were given up to 24 hours (or longer if requested) to consider

46 participation. Reasons for refusal were collected via verbal discussion, entered into an Excel
47 spreadsheet and categorised by the researcher.

48 Consenting patients were randomised to receive a dMUR at their regular community pharmacy or
49 standard hospital discharge care (no formal communication with the patient's community pharmacy,
50 unless the patient received their medication in a weekly compliance aid (monitored dosage system)
51 in which case a discharge list of medication was faxed to the community pharmacy, usually within 24
52 hours of discharge).

53 Prior to discharge each patient was told which group they had been assigned to, so they knew
54 whether or not to expect their community pharmacist to contact them to arrange a dMUR.

55 **Results**

56 The study took place between 7th April 2014 and 6th January 2015. Over this 9-month period 337
57 potential participants were identified by ward pharmacists, which ultimately resulted in 60 patients
58 recruited to the study, as shown in Figure 1.

59 **'Insert Figure 1 here'**

60 The reasons for exclusion and refusal are displayed in Table 1.

61 **'Insert Table 1 here'**

62 **Discussion**

63 This study is the first to describe formalising secondary care discharge of older patients into a
64 nationally contracted community pharmacy service (dMURs) as advocated by UK health officials and
65 bodies representing pharmacy.

66 Over the study period over 10,000 patients over 65 years of age were admitted to the hospital;
67 clearly a very small proportion (3.2%) of these patients were identified by ward pharmacists and
68 referred on to the researcher. This highlights the pivotal role ward staff play in such studies as

69 identified by McMurdo et al in their review of recruitment of older people to research [7]. The low
70 referral rate was despite steps taken by the researcher to raise awareness of the study among ward
71 pharmacists, including a 'briefing' session prior to its commencement and posters displayed around
72 the pharmacy department detailing types of patients who might be suitable. A study 'pack' was also
73 given to each pharmacist, containing background information about the study and its eligibility
74 criteria. Additionally, email reminders were sent regularly to all ward pharmacists throughout the
75 recruitment period. Feedback from ward pharmacists (n=7) via a focus group highlighted that
76 identification of patients for inclusion into any type of trial was not standard practice, thus not part
77 of their daily routine and was therefore often forgotten about. Other barriers identified by the
78 pharmacists were time constraints and concerns over patients' poor health status. Further
79 investigation of these barriers and possible solutions (for example, increased pharmacist time on
80 wards to allow for patient screening or use of information technology to make referral easier) is
81 ongoing.

82 *Patient Exclusions*

83 The most frequent reason (30.8%) for ineligibility was patients' not self-medicating at home. This
84 has implications for the English service specification for dMURs, which stipulates they must be
85 conducted with the patient and not a care giver, unless the patient specifically consents for the carer
86 to be present. In fact, patients requiring carers to manage their medication may be more in need of
87 a medication support service, as polypharmacy, including the use of inappropriate medication, is
88 common among older populations who have 'managed care at home' [8].

89 The nature of the nationally agreed service specification for community pharmacists also accounted
90 for a relatively high proportion of patients (n=22, 24.2% of exclusions) being excluded from the study
91 due to being housebound and unable to attend a community pharmacy for a dMUR. Only around
92 one in 10 community pharmacies in this study were able to provide domiciliary visits. Mobility
93 issues are obviously a key consideration in this population; reduced mobility due to ill health or

94 availability of transport has been cited by others as a challenge to the recruitment of older patients
95 [7,9].

96 Twenty patients (22% of exclusions), were not well enough to engage in discussion with the
97 researcher at the time of contact, either because they were too unwell (n=9) or had cognitive
98 impairment (n=11). Delirium or other conditions may temporarily impair older patients' capacity,
99 but they may recover by the point of discharge [7]. Therefore some patients may require more than
100 one visit to properly engage with them regarding research.

101 *Patient Refusals*

102 Of those patients fully screened, 72 (32%) refused. This is within the range (8-54%) found by
103 McMurdo, and similar to a study of a pharmacist-led medication review clinic for older patients in
104 primary care (36%) [7,10].

105 The key reason for non-participation was patients saw no benefit in having a dMUR with their local
106 community pharmacist (n=20, 27.8%). A further six patients (8.3%) stated specifically that they
107 would rather see their GP to discuss the changes to their medication. This indicates that the concept
108 of the dMUR was either not fully understood by these patients, or not viewed as a useful service for
109 them.

110 Fourteen (19.4%) patients did not want to be involved in 'research'. Distrust of researchers and a
111 reluctance to change their normal routine in order to take part in research have been cited
112 elsewhere as a challenge in the recruitment of frail older patients [10].

113 Eleven patients (15.3%) expressed concern over time commitment that the study imposed. Beyond
114 the dMUR itself, the study included two subsequent follow-up postal questionnaires, up to 6 months
115 after the initial discharge. Patients may have viewed this commitment too burdensome, although
116 comments from these patients (such as '*I've got enough appointments*') also suggested that even
117 the initial referral for a dMUR was too onerous. Again, an unfamiliarity with the role of the

118 pharmacist in provision of transitional care may have affected the perceived 'cost-benefit' of taking
119 part in this study.

120 However, it appears that old age, per se, may not have been a barrier to participation (only 2
121 patients (1.5%) said this was the reason for not taking part), but rather the frailty and co-morbidity
122 that may accompany it (11 patients (15.3%) declined due to 'health reasons'). This is consistent with
123 research done by others [7,10]. This potentially biases the study sample toward those patients who
124 are more 'well' and thus may be less in need of a medication review.

125

126 **Conclusion**

127 This study has demonstrated that recruitment of older people to a study of a post discharge
128 medicines support service provided by community pharmacists is particularly difficult for a number
129 of reasons. Patient unwillingness to participate, either due to ill-health or perceived lack of benefit
130 were noted. Additionally, pertinent to this study, the current provisions in England for post
131 discharge MURs excluded many people from participation, including those who were possibly in
132 greatest need.

133

134 **Conflicts of Interest:** None

135 **References**

- 136 1. Mansur N, Weiss A, Beloosesky Y. Relationship of in-hospital medication modifications of elderly
137 patients to postdischarge medications, adherence, and mortality. *Ann Pharmacother.* 2008 June
138 01;42(6):783-789.
- 139 2. Steering Group on Improving the Use of Medicines (for better outcomes and reduced waste).
140 Improving the use of medicines for better outcomes and reduced waste: An action plan. Dept. of
141 Health; 2012.
142 [https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/212837/Improvin](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/212837/Improving-the-use-of-medicines-for-better-outcomes-and-reduced-waste-An-action-plan.pdf)
143 [g-the-use-of-medicines-for-better-outcomes-and-reduced-waste-An-action-plan.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/212837/Improving-the-use-of-medicines-for-better-outcomes-and-reduced-waste-An-action-plan.pdf). Accessed
144 14 December 2015.
- 145 3. Royal Pharmaceutical Society. Keeping patients safe when they transfer between care providers
146 – getting the medicines right. Final Report. RPS; 2012. [http://www.rpharms.com/current-](http://www.rpharms.com/current-campaigns-pdfs/rps-transfer-of-care-final-report.pdf)
147 [campaigns-pdfs/rps-transfer-of-care-final-report.pdf](http://www.rpharms.com/current-campaigns-pdfs/rps-transfer-of-care-final-report.pdf). Accessed 14 December 2015.
- 148 4. Pharmaceutical Services Negotiating Committee/NHS Employers. Medicines use review and
149 prescription intervention service specification. PSNC and NHS Employers; 2013.
150 [http://psnc.org.uk/wp-content/uploads/2013/06/MUR-service-spec-Aug-2013-](http://psnc.org.uk/wp-content/uploads/2013/06/MUR-service-spec-Aug-2013-changes_FINAL.pdf)
151 [changes FINAL.pdf](http://psnc.org.uk/wp-content/uploads/2013/06/MUR-service-spec-Aug-2013-changes_FINAL.pdf). Accessed 14 December 2015.
- 152 5. Royal Pharmaceutical Society. Keeping patients safe when they transfer between care providers –
153 getting the medicines right. Individual reports from the Early Adopter Sites. RPS; 2012.
154 <http://www.rpharms.com/current-campaigns-pdfs/eas-finalreports.pdf>. Accessed 14 December
155 2015.
- 156 6. Bhatti N, Blenkinsopp A, Devlin L, Farooq H, Kazi I, Mulla I et al. Community pharmacists’
157 experiences of managing patients’ medicines after discharge from hospital: a preliminary study
158 of discharge medicines use reviews. *Int J Pharm Prac.* 2013;21(Supp2):86-87
- 159 7. McMurdo MET, Roberts H, Parker S, Wyatt N, May H, Goodman C, et al. Improving recruitment
160 of older people to research through good practice. *Age Ageing.* 2011;40(6):659-665.
- 161 8. Fialová D, Topinková E, Gambassi G, Finne-Soveri H, Jonsson PV, Carpenter I et al. Potentially
162 inappropriate medication use among elderly home care patients in Europe. *JAMA.*
163 2005;293(11):1348–1358
- 164 9. Baqir W, Desai N, Harker N, Vyas A, Copeland R, Burdon M, et al. Can targeted medicines use
165 reviews support safe transfer of care from hospital to community? *Int J Pharm Prac.*
166 2012;20(Supp2):82-83
- 167 10. Petty DR, Zermansky AG, Raynor DK, Vail A, Lowe CJ, Freemantle N et al. “No thank you”: why
168 elderly patients declined to participate in a research study. *Pharm World Sci.* 2001;23:22-27


[Click here to view linked References](#)**Figure 1: Summary of Recruitment Phase of Study**

Table 1: Reasons for Ineligibility (n=91) and Refusal (n=72)

Ineligibility	Number (% of patients excluded)
Protocol exclusions	
Patient not self-medicating at home	28 (30.8%)
Unable to access dMUR from chosen pharmacy	22 (24.2%)
Being discharged out of study area	11 (12.1%)
Participant's community pharmacy not taking part in the study	2 (2.2%)
Patient does not use a regular pharmacy	1 (1.1%)
Patient exclusions	
Lack of capacity to consent	11 (12.1%)
Too unwell to approach	9 (9.9%)
MUR not needed	7 (7.7%)
Reason for Refusal*	
Number (% of patients refusing)	
Patient sees no benefit to MUR	20 (27.8%)
Patient already has good links with community pharmacy	15 (20.8%)
Unwilling to engage with research	14 (19.4%)
Health reasons	11 (15.3%)
Time constraints	11 (15.3%)
Would rather see GP	6 (4.5%)
Hospital counselling adequate	4 (3%)
Patient does not want telephone MUR (home visit not available and patient cannot attend pharmacy)	3 (2.3%)
Patient does not agree with changes to medicines	2 (1.5%)
"Too old"	2 (1.5%)
Belief that medicines information will automatically be transferred to community pharmacy	1 (0.8%)

*The number of reasons for refusal is greater than the number of patients refusing, as some patients gave more than one reason.

Peer review of manuscript entitled: *Post Discharge Medicines Use Review (dMUR) Service for Older Patients: Recruitment Issues in a Feasibility Study*

30/10/15

Reviewer 1: Comments and Responses

1. In the abstract and on various other occasions throughout the manuscript, the participants are referred to as the 'elderly'. I would advise that this terminology be replaced with 'older adults' or 'older people', which would be regarded as more appropriate terms to use when referring to this population. **Done**
2. In Figure 1, please change '10682 Medical Patients' to '10682 medical patients' and '337 Patients' to '337 patients', (lowercase), in keeping with the word formatting in the other boxes within the same figure. **Done**
3. Line 68: 'Feedback from ward pharmacists via a focus group...' I would advise that you add (n=x) or similar, to indicate how many ward pharmacists provided feedback in the focus group. **Done (n=7)**
4. Whilst this article predominantly focusses on recruitment being problematic due to it being a study attempting to recruit older adults, an important finding described in this research report was that, overall, the pharmacists only referred 3.2% of all patients admitted, which is a significantly lower proportion than one would expect given the wider patient population admitted to hospitals. As such, this clearly represented a major barrier to the recruitment process. As the author(s) states that recruitment of participants to the study was 'often forgotten about' by ward pharmacists (line 70), it is perhaps unclear to the reader what steps (if any) were taken to ensure that pharmacists were aware of their role in the recruitment process. This point is also pertinent as the author(s) also report that of the 337 patients referred to the researcher by the pharmacists, 105 were ineligible for inclusion as they did not meet the specified study inclusion criteria. As this represents nearly one third of all those who were referred, this would lead this reviewer to assume that the ward

30/10/15

pharmacists involved in recruitment were not aware of the study's inclusion/exclusion criteria (as suggested in lines 35-36). To address this issue, I think this article would benefit from a comment in the discussion section to clarify exactly how and to what extent the pharmacists involved in the recruitment were informed of the recruitment process.

The following comment has now been inserted into the discussion: *The low referral rate was despite steps taken by the researcher to raise awareness of the study among ward pharmacists, including a 'briefing' session prior to its commencement and posters displayed around the pharmacy department detailing types of patients who might be suitable. A study 'pack' was also given to each pharmacist, containing background information about the study and its eligibility criteria. Additionally, email reminders were sent regularly to all ward pharmacists throughout the recruitment period*

In addition, this article would be improved if the author(s) could provide a short comment on whether they believe the approach taken with involving pharmacists in the recruitment process could have been improved, and if so, how?

The following comment has now been inserted into the discussion: *Further investigation of these barriers and possible solutions (for example, increased pharmacist time on wards to allow for patient screening or use of information technology to make referral easier) is ongoing.*

5. On line 98, please change 'pharmacist led' to 'pharmacist-led'. **Done**

6. On lines 81-83, the author(s) argue that patients who have carers managing their medicines at home are *'frailer and more susceptible to adverse drug events'*; this report would benefit from the inclusion of a reference here for this assertion, particularly as the author(s) also use this argument again to form part of their conclusion (line 128). If no specific reference exists to support the association, I would suggest that this sentence is re-worded to reflect that the proposed association is a (valid) assumption. **No specific reference exists; first two paragraphs under subheading 'Patient**

30/10/15

exclusions' (lines 75-83 in original manuscript) have now been combined into a single paragraph to summarise the issue addressed (lines 91-96 in revised manuscript)

7. On lines 116-117, please add (1.5%) after '*only 2 patients...*' as previous findings are presented in this way in the text (i.e. number followed by percentage in brackets). Similarly, on line 118: '*11 patients...*', please add the percentage for information. **Done**

8. On lines 75-78 the author(s) state that polypharmacy is common among older adults who have '*managed care at home*'. The relevance of this statement, in the context of patients' ineligibility, is not clear to this reviewer. I would ask the author(s) to review and either remove or rephrase this point in order to clarify their intended meaning. **The intended argument here is: the fact that patients who had carers managing their medicines had to be excluded is a cause for concern and needs addressing in terms of future service development. The combination of first two paragraphs under subheading '*Patient exclusions*' (lines 75-83 in original manuscript) attempts to clarify this point: *In fact, patients requiring carers to manage their medication may be more in need of a medication support service, as polypharmacy, including the use of inappropriate medication, is common among older populations who have 'managed care at home' [8].***

9. In Table 1 please correct '*Participants community pharmacy...*' to '*Participant's community pharmacy...*' **Done**

10. Finally, I would draw the author(s) attention to the reference section of the research report. Please review and correct the formatting of references as per the requirements of the journal. The use of semi-colons, commas, spaces, full-stops, capital letters and page numbering varies throughout

30/10/15

the reference list. Also, some journals are abbreviated and others are not. Also, in reference number 8, there is a '1' after 'Fialova D'. **Reference list re-visited and re-formatted**

Reviewer 2: Comments and Responses

Abstract:

1. In the methods section, please briefly state potential patients were visited by a researcher to assess eligibility and discuss involvement in the trial, and reasons for patients not participating were collected, or similar. **Done**

2. In the results section, please state that of the 337 potential participants, 132 were eligible and 60 were recruited. **Done**

3. Suggest first two sentences of conclusion are amended to summarise most common reasons and placed in the results section of the abstract. **Done: following sentences now inserted in results:**

Barriers to recruitment included competing priorities among ward pharmacists, and national restrictions placed on Medicines Use Reviews, for example housebound patients and those requiring carer support with medication. Lack of expected benefit resulted in a high proportion of patient refusals.

Introduction:

4. If word count allows, it may assist international readers if the description of the MUR program was expanded. From memory the program is free for consumers - if so, this suggest this is stated.

Lines 9-13 now read: In the UK, a national service known as Medicines Use Reviews (MUR) is offered by many community pharmacists which is free to patients and paid for centrally by the NHS. The MUR is a face-to-face consultation between a patient and pharmacist, designed to identify any problems or information needs patients have with their medicines and offer solutions [4].

30/10/15

5. Final sentence of introduction seems out of place (line 25) - suggest deleting or moving. Deleted, (along with preceding sentence in order to reduce word count)

6. The section of the aim which states "to investigate the logistics of setting up the study" could be a little more specific. Aim now reads: *To identify barriers to recruitment into a randomised controlled feasibility study of a hospital referral system to older patients' regular community pharmacists, and identify areas for improvement of the service or study design.*

Methods:

7. Line 43 (Exclusion criteria) suggest restarting numbering at 1. Done

8. Can the authors please describe the process after eligibility was assessed ie did the researcher talk to the patients, what information was provided (ie brochure) etc

The following text has been inserted (lines 44-47 of revised manuscript): *The researcher approached eligible patients to discuss involvement in the study and provide an information leaflet. Patients were given up to 24 hours (or longer if requested) to consider participation. Reasons for refusal were collected via verbal discussion, entered into an Excel spreadsheet and categorised by the researcher.*

9. Please describe the method used to collect reasons for refusal ie simple discussion with patient or was a survey completed etc.

See response to point 8 above

Peer review of manuscript entitled: *Post Discharge Medicines Use Review (dMUR) Service for Older Patients: Recruitment Issues in a Feasibility Study*

30/10/15

Editor comments and Responses

- The results section in the abstract is not in line with the described Objective of the study.

Objective now reads: To identify barriers to recruitment into a randomised controlled feasibility study of a hospital referral system to older patients' regular community pharmacists. Results section also amended as per reviewer 2, comment 3.

- Please insert 3-5 Statements on the impact of your findings on practice, patients or professionals, and add them at the end of the abstract in Editorial Manager. **Done**

- The references to the reports need more detail, see the Vancouver style formatting instructions.

All references checked and now in the authors' opinion are correct

- Replace the titles of some journals by the appropriate abbreviation (ref 9 and 10). **Done**