

Evaluating the potential of high pressure high temperature and thermal processing on volatile compounds, nutritional and structural properties of orange and yellow carrots

Article

Accepted Version

Grauwet, T., Kebede, B. T., Delgado, R., Lemmens, L., Manzoni, F., Vervoort, L., Hendrickx, M., Elmore, J. S. and Van Loey, A. (2015) Evaluating the potential of high pressure high temperature and thermal processing on volatile compounds, nutritional and structural properties of orange and yellow carrots. *European Food Research and Technology*, 240 (1). pp. 183-198. ISSN 1438-2377 doi: <https://doi.org/10.1007/s00217-014-2319-4> Available at <http://centaur.reading.ac.uk/39052/>

It is advisable to refer to the publisher's version if you intend to cite from the work.

To link to this article DOI: <http://dx.doi.org/10.1007/s00217-014-2319-4>

Publisher: Springer Verlag

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 **Evaluating the potential of high pressure high temperature and thermal processing**
2 **on volatile compounds, nutritional and structural properties of orange and yellow**
3 **carrots**

4 Tara Grauwet^{a*}, Biniam T. Kebede^a, Rosa M. Delgado^b, Lien Lemmens^a, Filipa
5 Manzoni^b, Liesbeth Vervoort^a, Marc Hendrickx^a, J. Stephen Elmore^b, Ann Van Loey^a

6 ^a *Laboratory of Food Technology and Leuven Food Science and Nutrition Research*
7 *Centre (LFoRCe), Department of Microbial and Molecular Systems (M²S), Katholieke*
8 *Universiteit Leuven, Kasteelpark Arenberg 22, 3001 Leuven, Belgium.*

9 ^b *University of Reading, Department of Food & Nutritional Sciences, Whiteknights,*
10 *Reading, RG6 6AP, Berks, United Kingdom.*

11

12 **Keywords:**

13 **Headspace GC-MS fingerprinting; High pressure high temperature processing; Thermal**
14 **processing; Carrot puree; Volatile compounds; β -carotene**

15

16 **Journal: *European Food Research and Technology***

17 *Corresponding author during submission process

18 Fax: +32 16 32 19 60

19 Telephone: +32 16 32 19 47

20 E-mail: tara.grauwet@biw.kuleuven.be

21 <http://www.biw.kuleuven.be/m2s/clmt/lmt/>

22

23 **ABSTRACT**

24 The present study compares the impact of thermal and high pressure high temperature
25 (HPHT) processing on volatile profile (via a non-targeted headspace fingerprinting) and
26 structural and nutritional quality parameter (via targeted approaches) of orange and
27 yellow carrot purees. The effect of oil enrichment was also considered. Since oil
28 enrichment affects compounds volatility, the effect of oil was not studied when
29 comparing the volatile fraction. For the targeted part, as yellow carrot purees were
30 shown to contain a very low amount of carotenoids, focus was given to orange carrot
31 purees. The results of the non-targeted approach demonstrated HPHT processing exerts
32 a distinct effect on the volatile fractions compared to thermal processing. In addition,
33 different colored carrot varieties are characterized by distinct headspace fingerprints.
34 From a structural point of view, limited or no difference could be observed between
35 orange carrot purees treated with HPHT or HT processes, both for samples without and
36 with oil. From nutritional point of view, only in samples with oil, significant
37 isomerisation of all-*trans*- β -carotene occurred due to both processing. Overall, for this
38 type of product and for the selected conditions, HPHT processing seems to have a
39 different impact on the volatile profile but rather similar impact on the structural and
40 nutritional attributes compared to thermal processing.

41

42 **1. Introduction**

43 ***1.1 High pressure high temperature (HPHT) treatment as a potential*** 44 ***alternative for thermal sterilization***

45 Sterilization treatments aim to inactivate both vegetative cells and microbial spores,
46 resulting in food products which are shelf-stable. In order to obtain this stability in low-
47 acid products with high water content, thermal processing is commonly applied. The
48 high thermal load which especially needs to be applied to products with slow heat
49 transfer results in considerable quality changes compared to the untreated product [1].
50 Improving the heat transfer is one of the strategies which have been followed in
51 optimizing thermal processes. Since the year 2000, high pressure (e.g. 600 MPa)
52 combined with high temperature (e.g. 121 °C) has been discussed in literature as an
53 alternative for thermal processing [2-5]. Some authors addressed this technique as a
54 ‘high pressure-assisted thermal process’ describing the high pressure high temperature
55 (HPHT) treatment as a more optimal thermal treatment in which pressure is used to
56 quickly increase the temperature of the product due to compression heating [2, 6].
57 Others have been studying the effect of pressure at high temperature and reported a
58 clear pressure effect on quality changes at that particular high temperature [3, 7-9]. Both
59 increasing and decreasing effects on the reaction rate constants of high pressure at high
60 temperature have been described [10].

61 In the end, the impact of a HPHT process on the food product will be the integrated
62 impact of pressure, temperature and time. In order to compare the process impact of this
63 novel technique to its traditional thermal counterpart, process conditions should be
64 selected on a fair basis, for example equivalent microbial inactivation resulting in
65 products with a similar microbial shelf-life [11-13]. However, to obtain insight in the

66 individual effect of the process variables, the importance of kinetic experiments in
67 which all process variables are accurately monitored should not be forgotten [10].

68 ***1.2 Carrots (*Daucus carota*) are one of the most popular root*** 69 ***vegetables***

70 Carrots can be considered as primary vegetables in many countries. In recent decades,
71 carrots gained popularity due to the awareness of their nutritional value and health-
72 related benefits [14]. Carrots exist in different varieties, such as orange, yellow, red,
73 white and purple. This genetic variation combined with cultivation conditions and
74 exposure to ethylene affect the sensorial (e.g. volatiles and polyacetylenes) and
75 nutritional (e.g. carotenoids, vitamins and minerals) quality parameters [15-17]. It has
76 been described in literature that food processing techniques affect
77 (negatively/positively) these quality parameters. On the one hand, as a result of
78 conversion of carotenoids (*trans* to (poly-)*cis*) and degradation reactions, the beneficial
79 biological activity of the carotenoids is altered (e.g. reduced antioxidant and provitamin
80 A capacity) [18-21]. On the other hand, processing has been shown to positively affect
81 carotenoid bioaccessibility and bioavailability in most cases [22].

82

83 Comparing the process impact on carrot quality parameters between conventional
84 thermal and HPHT processes has been the topic of interest of other researchers before.
85 However, research studies in which the comparison was performed based on the
86 principle of equivalence are scarcely found in literature. Recently, fair comparisons
87 have been reported in studies of Knockaert et al. [8] (targeted approach, carrot pieces),
88 Vervoort et al. [24] (targeted approach, orange juice), Timmermans et al. [11] (targeted
89 approach, orange juice), Vervoort et al. [13] (targeted approach, carrot pieces), Vervoort
90 et al. [25] (untargeted approach, carrot pieces) and Kebede et al. [26-28] (untargeted

91 approach, wide range of vegetables). It is thus clear that additional investigations in this
92 context are needed in order to evaluate using science-based evidence the potential of
93 HPHT processing. Besides, since the quality parameters differ from variety to variety,
94 more than one carrot cultivar should be taken into account.

95

96 In this perspective, the main objective of the present work was to compare the effect of
97 thermal and HPHT processing on a range of important quality attributes. Aiming for a
98 fair comparison, processing conditions resulting in equivalent processes in terms of
99 microbial inactivation were selected. Taking into account the possible difference in
100 chemical composition among varieties, the focus is given to carrots with different
101 colour varieties, i.e. orange and yellow carrots. Given the fact that oil might have an
102 effect on the stability of lipophilic nutrients, the enrichment of oil to the carrot purees
103 was considered as well. The novelty of the present work is that, the comparison in the
104 impact of conventional and novel sterilization techniques was performed from both non-
105 targeted and targeted approaches. In the former case, the impact on volatile fractions of
106 differently processed carrot puree was compared using an untargeted fingerprinting
107 approach (integration of headspace (HS) solid-phase micro-extraction (SPME) GC-MS
108 method and multivariate statistical data analysis). In the latter case, a targeted approach
109 to analyze specific quality related parameters (color, carotenoid profile, particle size
110 distribution, microscopy) was performed. A schematic overview of the research plan of
111 this work can be found in **Fig. 1**.

112 **2 Materials and methods**

113 **2.1 Sample preparation**

114 Single batches of fresh orange and yellow carrots (*Daucus carota* cv. Nerac and cv.
115 Yellow mellow, respectively) were bought at a local market and stored at 4 °C. Carrots

116 were peeled, cut into slices as homogeneously as possible, packed in plastic bags and
117 blanched at 95 °C for 8 minutes. At all times orange and yellow carrots were kept
118 separately. Blanched carrot bags were frozen in liquid nitrogen and stored at -40 °C
119 until use. Singular carrot purees were prepared by blending blanched carrots with water
120 (1:1) for 1 min in a kitchen blender. In the case of the carrot purees enriched with oil,
121 they were stirred with 5% (w/w) extra virgin olive oil for 15 min at room temperature.

122 The carrot puree was homogenised at 10 MPa for 1 cycle using a high-pressure
123 homogeniser (Panda 2K; Gea Niro Soavi, Mechelen, Belgium). Since the product
124 temperature usually increases after homogenisation, the sample inlet and outlet were
125 thermostated at 4 °C and the pressure was controlled on a digital display.

126 **2.2 Processing**

127 Sterilization treatments were performed using both traditional technologies (thermal
128 treatments) and novel technologies (HPHT treatment). Aiming fair comparison of the
129 process impact, an equivalent industrially relevant process value $F_0 = 3$ min was put
130 forward for both processing targeting inactivation of spores of *Clostridium botulinum*.
131 For both treatment types a particular reactor holding temperature (T_h) of 117 °C was
132 selected. Each treatment was repeated six times for valid statistical data analysis
133 afterwards. Due to the lack of reliable kinetic data as a result of incomplete
134 understanding of the combined effect of pressure and temperature on *Clostridium*
135 *botulinum* spore inactivation [10], in the present work, the HPHT was considered as
136 pressure assisted thermal processing. Due to their inert nature, glass jars for the thermal
137 and Teflon sample holders for the HPHT processing were selected. A comparison of the
138 monitored profiles of the process variables during both treatments, can be found in **Fig.**
139 **2.**

140 **2.2.1 Thermal processing**

141 For the thermal treatments, glass jars (100 mL volume, 95 mm height and 45 mm
142 diameter) were filled with 85 ± 0.5 g carrot puree and treated in a static steriflow pilot
143 retort (Barriquand, Paris, France). Temperature profiles of the retort and at the coldest
144 point in the glass jars (1 cm above the bottom) were controlled by type-T thermocouples
145 (Ellab, Hillerod, Denmark). The total process time (t_p) was on average 60 minutes (**Fig.**
146 **2**).

147 **2.2.2 High pressure high temperature processing**

148 High pressure sterilization was carried out using laboratory-scale 6-vessel high pressure
149 equipment (custom-made, Resato, Roden, the Netherlands) with propylene glycol fluid
150 (PG fluid, Resato, The Netherlands) as the pressure medium. The HPHT equipment
151 allows computer controlled pressure build-up to 800 MPa, temperature control up to
152 120 °C and data logging of both sample pressure and temperature (**Fig. 2**). For pressure
153 increase, the equipment consists of a pressure prefill pump which builds up the pressure
154 to 150 MPa with a single piston displacement after which a high pressure intensifier can
155 further built up the pressure at a particular selected pressure build-up rate. The high
156 pressure sterilization processes were performed at 600 MPa combined with a process
157 temperature of 117 °C. Teflon cylindrical tubes (12 mm internal diameter, 4 mm
158 thickness, 85 mm length) were filled with carrot puree, closed with a movable cap,
159 vacuum sealed with double plastic bags, equilibrated at 10 °C and placed in the pressure
160 vessels equilibrated at the process temperature. The pressure build-up was started when
161 the temperature registered in the vessels by type-J thermocouples (Ellab) reached 75 °C
162 (i.e. initial temperature). Pressure was immediately increased to 150 MPa after which
163 pressure was further built up to 600 MPa at a rate of 10 MPa/s. Assuming no effect of
164 pressure on spore inactivation under HPHT conditions, HPHT processes were at least

165 thermally equivalent to the thermal treatment aiming a F_0 -value of 3 min. The product
166 temperature was recorded online and the holding time was adjusted to achieve the
167 targeted F_0 value. The vessels were decompressed after the required holding time. On
168 average, the total process time (t_p) was 20 min (**Fig. 2**).

169 **2.2.3 Post treatment sample handling**

170 Following treatments, samples were immediately transferred to ice water to stop any
171 process-induced reaction. Consequently, treated samples were emptied in a cooling
172 room and transferred to a small volume (10 ml) polyethylene terephthalate tubes with a
173 polyethylene cap. Hereafter, the tubes were frozen in liquid nitrogen and stored at
174 $-40\text{ }^\circ\text{C}$ until analysis.

175 **2.3 Analysis of the volatile profile by headspace fingerprinting**

176 **2.3.1 Headspace SPME-GC-MS analysis**

177 Targeting detection of a wide range of volatiles in a particular food extract, a headspace
178 (HS) fingerprinting SPME-GC-MS method of analysis was optimized beforehand. The
179 method includes incubation, extraction using an appropriate type of fiber coating and
180 GC-MS parameters. Carrot puree (3 g) was weighed into 20 mL headspace vials
181 (Supelco, Belfonte, USA). All headspace analyses were conducted on an Agilent 7890A
182 gas chromatograph (GC) coupled to a 5975C mass selective detector (MS) (Agilent
183 Technologies, Santa Clara, CA, USA) and equipped with a CombiPAL autosampler
184 (CTC Analytics, Zwingen, Switzerland). Vials were equilibrated in the incubator at 40
185 $^\circ\text{C}$ for 10 min under agitation at 500 rpm. The carrot puree volatile compounds in the
186 headspace of the vial were sampled for 30 min by means of a solid phase
187 microextraction (SPME) fiber with 50/60 μm DVB/CAR/PDMS
188 (divinylbenzene/carboxen/polydimethylsiloxane) sorptive coating (StableFlex; Supelco,

189 Bellefonte, PA). The fiber was preconditioned according to the manufacturer's
190 guidelines before its first use. Desorption was carried out for 10 min at 250 °C. The
191 volatiles were injected in splitless mode and subsequently separated on a capillary
192 column (30 m × 0.25 mm; 1.0 µm film thickness, HP-5MS; Agilent Technologies,
193 Santa Clara, CA), using helium as carrier gas at a constant flow rate of 1.2 mL/min. The
194 column oven was programmed at a starting temperature of 40 °C, which was retained
195 for 5 min, after which it was elevated to 170 °C at a rate of 5 °C/min and retained for 5
196 min, followed by a second ramp to 230 °C at 15 °C/min. After 5 min at the final
197 temperature, the oven was cooled again to the initial temperature. Mass spectra were
198 obtained by electron ionisation (EI; 70 eV), with a scanning range of m/z 29–250. MS
199 ion source and quadrupole temperatures were 250 and 150 °C, respectively.

200 **2.3.2 Data preprocessing and multivariate data analysis**

201 As commonly observed in GC-MS analysis, co-eluting compounds were present in the
202 obtained chromatograms. Therefore, all chromatograms were analyzed with Automated
203 Mass Spectral Deconvolution and Identification System (AMDIS) (Version 2.66, 2008,
204 National Institute of Standards and Technology, Gaithersburg, MD, USA) to extract
205 “pure” component spectra from complex chromatograms. The deconvoluted spectra
206 were then analyzed with Mass Profiler Professional (MPP) (Version 12.0, 2012, Agilent
207 Technologies, Diegem, Belgium) aiming filtering and peak alignment. The MPP
208 obtained a spreadsheet containing peak areas, which was used as an input for the
209 statistics. The multivariate data were analyzed with a multivariate statistical data
210 analysis (MVDA) which was carried out in Solo (Version 6.5, 2011, Eigenvector
211 Research, Wenatchee, WA, USA). As a preprocessing step, all data were mean-centered
212 and the variables were weighed by their standard deviation to give them equal variance.
213 In a first approach, principal component analysis (PCA) was performed to evaluate each

214 data set and to detect potential outliers. In this work, the MVDA was performed at two
215 parts. In a first part, the comparison of the impact of the different sterilization treatments
216 per carrot puree type (orange versus yellow) was carried out applying principal least
217 squares discriminant analysis (PLS-DA) to the data set, considering the volatiles as *X*-
218 variables and the blanched, thermal treatment and HPHT treatment as the three
219 categorical *Y*-variables. In a second part, PLS-DA was performed to study the influence
220 of the carrot variety (orange versus yellow) per treatment type. In this case, the volatiles
221 were considered as *X*-variables and the carrot varieties as two *Y*-variables. Determining
222 the complexity of the model, the lowest number of latent variables (LVs) resulting in a
223 class separation were used. In PLS-DA, to qualitatively investigate impact differences
224 among the classes, bi-plots were plotted. To quantitatively select discriminant
225 headspace components, Variable IDentification (VID) coefficients were calculated [20].
226 These values correspond to the correlation coefficient between each original *X*-variable
227 and *Y*-variable (s). The volatile identification was carried out for compounds which had
228 a VID coefficient higher than 0.800 in absolute value. Identification was performed by
229 comparing the compound's mass spectra from the NIST spectral library (NIST08,
230 version 2.0, National Institute of Standards and technology, Gaithersburg, MD, USA).
231 A visual inspection of the spectral matching was carried out accepting a threshold match
232 and reverse match of 80%. Retention time indices were calculated. All plots were made
233 using OriginPro 8 (Origin Lab Corporation, Northampton, MA, USA). A Tukey's
234 multiple comparison was used to test for significant differences between the mean peak
235 areas ($p < 0.05$) of the discriminant headspace components.

236 **2.4 Analysis of nutritional and structural characteristics**

237 **2.4.1 Determination of the carotenoid profile and content**

238 In order to determine the carotenoid profile and content of the carrot purees, an
239 extraction step was performed based on the procedure of Sadler et al. [29]. Briefly,
240 carrot puree was mixed with CaCl₂ (ratio 1:1) and 50 ml extraction solution (containing
241 50 % hexane, 25 % ethanol, 25 % acetone and 0.1 % BHT) and the mixture was stirred
242 for 20 minutes at 4 °C. After adding 15 ml reagent grade water, the mixture was stirred
243 for another 10 minutes at 4 °C. The organic layer, which could be separated from the
244 aqueous layer and which contains the carotenoids, was filtered (Chromafil PET filters,
245 0.20 µm pore size, 25 mm diameter; Macherey-nagel, Duren, Germany) and stored in
246 dark vials before further analysis. The extraction procedure was carried out under
247 dimmed red light.

248 The carotenoids in the extract were separated and quantified using an HPLC system
249 (Agilent Technologies 1200 Series, Diegem, Belgium), equipped with a C₃₀ column (5
250 µm x 250 mm x 4.6 mm, YMC Europe, Dinslaken, Germany) and a diode array
251 detector. During the analysis, the temperatures of the autosampler and the column were
252 kept at 4 °C and 25 °C, respectively. To separate the different carotenoid isomers, linear
253 gradient elution was used. The gradient was built up in 20 min from 81% methanol,
254 15% methyl-t-butyl-ether and 4% reagent grade water to 41% methanol, 55% methyl-t-
255 butyl-ether and 4% reagent grade water at a flow rate of 1 ml/min [30]. Identification
256 and quantification of the carotenoids was performed at 450 nm. Calibration curves for
257 all-*trans*-β-carotene, 15-*cis*-β-carotene, 13-*cis*-β-carotene, 9-*cis*-β-carotene and lutein
258 (CaroteNature, Lupsingen, Switzerland) were used to quantify the β-carotene and lutein
259 content of the orange and yellow carrot purees, respectively.

260 **2.4.2 Objective colour measurement**

261 Colour measurements (CIE $L^*a^*b^*$ values) were conducted using a Hunterlab
262 ColourQuest colorimeter (45°/0° geometry, Illuminant D65, Reston, VA, USA). The
263 instrument was calibrated daily with a black and green ceramic tile. At a 10° angle, the
264 CIE colour space coordinates were determined in triplicate, whereby L^* is indicating the
265 lightness (varying from 0, black, to 100, white), a^* is a measure for the redness (varying
266 from -60, green, to +60, red) and b^* is a measure for the yellowness (varying from -60,
267 blue, to +60, yellow).

268 **2.4.3 Determination of the particle size distribution**

269 The particle size distribution of the carrot purees was measured by laser diffraction
270 using a Malvern Mastersizer S long bench diffractor (Malvern Instruments Ltd., Great
271 Malvern, UK). Laser light (HeNe Laser, wavelength 633 nm, diameter 18 mm) was sent
272 through a suspension of carrot puree (± 10 g) in water. The light that was scattered by
273 particles between 0.06 and 880 μm was measured by a series of photodetectors (42
274 element composite solid state detector array). From the intensity distribution of the
275 scattered light, the particle size distribution of the sample was calculated by the
276 instrument software using the Mie theory. The parameter $d(v, 0.5)$ was calculated
277 which indicates the median diameter or the value of the particle diameter at 50 % in the
278 cumulative distribution.

279 **2.4.4 Analysis of the microstructure**

280 The microstructure of the different carrot purees was visualised using light microscopy.
281 To 1 mL of carrot puree, 4 mL water and 5 mL 0.01% toluidine blue solution were
282 added and the mixture was incubated at room temperature for 10 min. The mixture was
283 analysed using an Olympus BX-41 light microscope (Olympus, Optical Co. Ltd.,

284 Tokyo, Japan) at a magnification level of 10x. Micrographs were taken using image
285 analysis software (AnalySIS pro 5.0 Soft Imaging System GmbH, Bensheim,
286 Germany).

287 **3 Results and discussion**

288 In the following sections, the results will be discussed starting from impact comparison
289 on the volatile profile of each carrot variety using a headspace fingerprinting method
290 (untargeted approach, section 3.1) to impact on nutritional and structural quality aspects
291 (targeted approach, section 3.2).

292 Oil enrichment of the puree directly affects the volatilizable food extract leading to a
293 decrease in volatility of particular compounds at selected incubation and extraction
294 conditions during the HS-SPME analysis. Consequently, comparing fingerprints of a
295 particular carrot puree (e.g. yellow carrot) enriched with oil or not might lead to biased
296 results not necessarily explaining the effect of oil in particular food reactions but more
297 probably revealing the clear effect of oil on the volatility of particular compounds.
298 Consequently, when comparing the volatile fraction with the headspace fingerprinting
299 (section 3.1), the effect of oil was not studied.

300 ***3.1 Comparing impact of thermal and HPHT processes on volatile*** 301 ***profile by headspace fingerprinting***

302 In the present work, prior to treatment, vegetables were blanched (section 2.1).
303 Therefore, enzymatic activities were not expected to have a significant impact on the
304 formation of volatiles and consequently changes will be related to non-enzymatic
305 process-induced chemical reactions. A representative total ion chromatogram of the
306 headspace profile of both the blanched/reference orange and yellow carrot samples is
307 depicted in **Fig. 3**. Over 100 distinct headspace components were detected in carrot
308 purees, terpenes and aldehydes being the most abundant. As explained in section 2.3.2

309 and as schematically shown in **Fig. 1**, deconvoluted spectra were analyzed with Mass
310 Profiler Professional (MPP) aiming filtering and peak alignment. The MPP yielded a
311 spreadsheet containing peak areas per detected compound, which was used as an input
312 for the multivariate data analysis (MVDA). In this work, the selected discriminative
313 markers are particularly discussed for their consequences on flavour.

314 After using PCA as an exploratory technique, PLS-DA was applied as a supervised
315 method to find differences among the three treated sample classes (blanched, HT,
316 HPHT) taking into account the available knowledge on sample class. Considering the
317 volatiles as *X*-variables and the different treatments as categorical *Y*-variables, biplots of
318 scores and correlation loadings were constructed. This type of plot offers a tool to
319 graphically summarise the analytical data to reveal relationships between samples and
320 to determine volatiles characterising a certain group of samples (section 3.1.1). Variable
321 IDentification coefficients (VID's) were calculated in a following step as a more
322 quantitative tool to select discriminant markers (i.e. compounds which detected amount
323 was different in one class compared to the other classes) (section 3.1.2).

324 **3.1.1 Visualisation of impact differences**

325 In **Fig. 4**, biplots, based on PLS-DA, are representing the process impact differences for
326 orange and yellow carrot purees. From the figures, both the similarity of samples within
327 one group (e.g. repetition of treatments) as well as the differences among the samples of
328 different groups (e.g. thermal versus HPHT processing) can be derived. Fingerprints of
329 samples from the same group clustered and showed clear resemblance. In general, a
330 clear effect of sterilization on the headspace fraction could be observed: large distance
331 was observed between samples from the blanched class and the thermal and HPHT
332 treated sample class. In addition, equivalent sterilization treatments showed different
333 volatile profiles. The separation between the three classes (blanched, HT, HPHT) in

334 orange carrot purees can be quantified by a variance in the *Y*-variables of 93% described
335 by the first two latent variables. Similar value (88%) was determined in yellow carrot
336 purees. Constructing ellipses representing correlation coefficients of 70 % and 100 %
337 some idea about the importance of volatiles for a specific group/class can be obtained:
338 all volatiles placed between the two ellipses explained the first two latent variables in
339 more than 70 % of its variability. Graphically, if those volatiles are projected between
340 those two ellipses close to a particular group, it means that they are important and are
341 characterized by a higher concentration within that group, compared to the others.
342 However, it is a challenge to deduce from the biplots information about variables which
343 concentration is clearly different in a particular group compared to another group (i.e.
344 discriminant markers), since these are the first interesting compounds to zoom further
345 into in order to understand the observed difference in process impact.

346 **3.1.2 Selection of discriminative markers**

347 Variable IDentification (VID) coefficients as defined by Ooms (1996), serve a
348 quantitative measure to select discriminative markers from the headspace fingerprints
349 (section 2.3.2). In other words, it ranks the components based on their importance for a
350 particular class compared to the other classes: high positive values demonstrating high
351 concentration of a certain compound for that particular class compared to the other
352 classes and low negative values demonstrating the opposite. In this work, for both
353 orange and yellow carrot, volatiles with an absolute value of VID more than 0.800 were
354 considered relevant, identified and further zoomed into (**Table 1**). These discriminant
355 volatiles were plotted individually as a function of processing. To clearly show the most
356 important of the selected discriminant components, those with VID's higher than 0.900
357 are represented in **Fig. 5** (for orange carrot) and to **Fig. 6** (for yellow carrot). In these
358 plots, the mean areas and the standard errors calculated from the six replicates were

359 depicted. From these figures, the concentration of those compounds compared to the
360 other groups can be deduced. Several trends could be observed. Terpenes showed
361 relevant VID coefficients in both orange and yellow carrot purees. However, their
362 identity was different depending on the matrix. As indicated in the introduction,
363 terpenes are typical, naturally present flavour compounds. On the one hand,
364 monoterpenes such as sabinene's concentration was the highest in the blanched class.
365 Sterilization might have degraded this terpene. On the other hand, higher concentrations
366 of p-cymene and α -ionone were found after applying both sterilization treatments.
367 The changes detected in degradation products from carotenoids such as α -ionone are in
368 agreement with previous studies [32, 33], which showed that sterilization processes
369 affected the total β -carotene concentration as well as its isomerization and bio-
370 accessibility. Degradation products of limonene such as terpinolene and γ -terpinene,
371 where significantly less detected after HPHT process compared to the other groups. The
372 same effect was observed in the case of other monoterpenes such as o-cymene,
373 sabinene, sesquithujene. Sesquithujene and β -caryophyllene epoxide were even not
374 detected after high pressure treatments (**Fig. 5**). Published reports demonstrated that
375 terpinolene and caryophyllene contribute significantly to carrot flavour intensity [34].
376 These conclusions are in agreement with Trejo Araya et al. [7] who determined the
377 headspace volatiles of carrot sticks pasteurized by high pressure processing (600 MPa, 2
378 min) and thermal treatments (90°C, 5 min). They showed that all monoterpenes and
379 terpinolenes were still present after treatment, and, in some cases, even increased. Since
380 correlations between carrot volatile changes after treatments (either 600 MPa, 2min or
381 90°C, 5min) and sensorial changes was observed before [7], sensorial studies of these
382 purees could give interesting information about the relevance of the flavour profile

383 modification. In future, further research should be done to investigate a possible
384 relationship between the flavour changes and carotenoid profile in carrot purees.

385 The selected aldehydes as markers in carrot purees were characterized by a negative
386 VID in blanched purees and by a positive VID in the sterilized purees (**Table 1**). In
387 other words, their formation was clearly enhanced by processing and their presence
388 discriminated the blanched from the sterilized samples. This fact could be explained
389 because aliphatic aldehydes can be formed from unsaturated fatty acids due to thermal
390 oxidation [35]. In this study, aldehydes from the degradation of oleic, linoleic and
391 linolenic acid were identified as markers. Thus, aldehydes from oxidized oleic acid such
392 as heptanal, octanal, decanal and 2-decenal were detected in higher concentration after
393 the treatment in orange carrot purees (**Fig. 4**). This trend was also observed in previous
394 studies in carrot pieces where heptanal, octanal and *trans*-2-decenal were selected as
395 markers in the study of the impact of thermal and high pressure processing technologies
396 [25]. Other aldehydes which showed a relevant VID in these purees such as hexanal and
397 2-octenal could be formed by oxidation of linoleic acid [34]. In the present study, the
398 carrot variety had an influence on the determination of markers being only heptanal and
399 octanal markers in the case of yellow carrot purees (**Table 1**). Both aldehydes showed a
400 negative VID in blanched purees indicating that they were formed after treatment in
401 yellow carrot purees. Although both sterilization conditions were established targeting a
402 particular processing value, the temperature histories of the treatments were not the
403 same (**Fig. 2**): the temperature history of the HPHT treatment coming more close to the
404 High-Temperature-Short-Time principle. Taking this knowledge into account, particular
405 reactions (e.g. thermal oxidation of fatty acid) would be estimated to be less pronounced
406 after HPHT treatment (possibly explaining the higher concentration of heptanal after
407 conventional thermal treatment compared to HPHT treatment). However, particular

408 aldehydes such as 2-nonenal and 2-octenal were significantly more detected after HPHT
409 treatment compared to its thermal equivalent (**Fig. 5**). This observation can be explained
410 by the effect of pressure on the oxygen solubility. This is in line with reports in which
411 oxidative chemical reactions were enhanced under increased pressure [26-28].

412 **3.2 Comparing impact of thermal and HPHT processes on** 413 ***nutritional and structural quality aspects***

414 **3.2.1 Characterisation of blanched carrot purees**

415 The blanched orange and yellow carrot purees were characterised in terms of
416 microscopy and particle size (structural characteristics), carotenoid content and
417 isomerisation (nutritional characteristics) and colour values as this might be related to
418 changes in carotenoids. Both samples without and with the addition of olive oil were
419 evaluated. An overview is given in this section. Based on the results, relevant samples
420 for comparing the impact of thermal and HPHT process were selected.

421 In **Fig. 7**, typical light micrographs of blanched orange carrot purees (A = without oil
422 addition; B = with oil addition) and blanched yellow carrot purees (C = without oil
423 addition; D = with oil addition) are presented. It can be observed that due to mixing and
424 high pressure homogenisation, the carrot tissue was broken down to cell fragments,
425 individual cells and cell clusters. In **Fig. 7B** and **Fig. 7D**, the emulsified oil droplets can
426 clearly be visualised. Comparing micrographs of orange and yellow carrot purees, it is
427 hard to observe clear differences in cell shape and size. Based on the results of the
428 particle size distribution measurements (**Fig. 8**), it can however be seen that in general
429 the particles in the yellow carrot purees are somewhat larger than the particles in the
430 orange carrot purees. Nevertheless, the differences are very limited. Additionally, it can
431 be observed in **Fig. 8** that the particle size distribution curves for samples with and
432 without oil addition are coinciding. This means that the addition of olive oil to the

433 samples does not have an impact on the carrot tissue particle size after homogenisation.
434 For the samples where oil was added, the particle size distribution curves show a higher
435 volume percentage at small particle size (around 10 μm), compared to the particle size
436 distribution curves of the samples without oil addition. These small particles are a
437 representation of the emulsified oil droplets.

438 Overall, from a structural point of view, it can be concluded that the blanched yellow
439 and orange carrot purees, which were used as starting material for thermal and HPHT
440 processes, are quite similar with regard to their microstructure and their particle size.

441 **Table 2** summarises the characterisation of the yellow and orange carrot puree in terms
442 of their carotenoid and carotenoid-isomer content and colour values. In orange carrot
443 purees, all-*trans*- β -carotene and its *cis*-isomers were identified to be the main
444 carotenoids. This is in agreement with previous studies on carrots (e.g. [36-38]). In the
445 samples where oil was added, a slightly higher amount of all-*trans*- β -carotene was
446 detected. This could be due to a slightly higher extraction yield as a result of the oil. In
447 general, for all orange carrot purees, the *trans*-isomer accounts for at least 75 %. In the
448 yellow carrot purees, lutein was shown to be the main carotenoid [14], although some
449 smaller unidentified peaks were present in the chromatogram. From the quantitative
450 results in **Table 2**, it can be observed that the concentration of lutein in the blanched
451 yellow carrot purees was very low. Compared to the carotenoid content in the orange
452 carrot puree, the lutein content in the yellow carrot puree was a factor 10 lower. Based
453 on this observation, it was decided not to include the yellow carrot puree in the
454 comparison study (thermal versus HPHT processing) as changes in the lutein content as
455 a result of processing would be difficult to perceive. With regard to the colour values of
456 the purees, the following observations could be made. The samples where oil was added
457 were in both cases lighter (higher L^* value) and less red (lower a^* value) compared to

458 their counterparts without oil addition. For the orange carrot purees, the samples with
459 oil were more yellow (higher b^* value), whereas the opposite was observed for the
460 yellow carrot purees. When putting the yellow carrot purees next to the orange carrot
461 purees, the yellow carrot purees were shown to be lighter and less red. In general, the
462 trends in the experimentally determined L^* , a^* and b^* values were a good reflection of
463 the visual observations of the carrot purees.

464 As explained above, the remaining part of this work, i.e. the actual comparison of the
465 impact of thermal and HPHT processes on nutritional and structural quality aspects, has
466 been focussed on the orange carrot purees.

467 **3.2.2 Impact on structural quality aspects**

468 The sterilised orange carrot purees were analysed for their particle size distribution and
469 their microstructure. With regard to the particle size, no changes could be observed as a
470 consequence of processing (data not shown). This was independent on the technology
471 that was applied, i.e. thermal or HPHT processing and on the presence or absence of oil.
472 This result implies that the samples were stable during processing in terms of particle
473 size. On the micrographs, cell separation and cell wall swelling can be observed as a
474 result of the sterilisation processes. This has mainly been attributed to β -eliminative
475 pectin depolymerisation and consecutively pectin solubilisation, which are known to
476 occur during processes of high thermal intensity [39, 40]. However, when comparing
477 samples from thermal and HPHT processes, it was hard to differentiate between the two
478 groups of samples based on the degree of pectin solubilisation (data not shown). In the
479 present study, no clear statement can thus be made on the differential effect of thermal
480 and HPHT processes on the microstructure of the carrot samples.

481 **3.2.3 Impact on nutritional quality aspects and colour**

482 **Fig. 9** shows the results for the carotenoids in the sterilised orange carrot purees. It
483 should be noted that the results are presented as contributions instead of absolute
484 concentrations. Carotenoid contributions are defined as the proportion of a particular
485 carotenoid relative to the total carotenoid content. This transformation (from absolute
486 carotenoid concentration to carotenoid contribution) has been performed, since as a
487 result of processing, the carotenoid extractability might change [38] which can bias the
488 results and conclusions. By expressing the data as contributions, the changing
489 extractability effect is filtered out.

490 From **Fig. 9A**, it can be observed that for all plain orange carrot purees (without oil
491 addition), whether or not processed, the contribution of all-*trans*- β -carotene was around
492 70 – 80 %, implying a total *cis*-isomer contribution of 20 – 30 %. The three different
493 *cis*-isomers were present in similar amounts, however the contribution of 13-*cis*- β -
494 carotene was always shown to be lower than the contribution of 9-*cis*- and 15-*cis*- β -
495 carotene. The results show that the sterilisation processes applied in this study
496 ($F_0 = 3$ min) did only result in a limited or no effect on the carotenoids for carrot
497 samples where no oil was present. A differentiation between thermal and HPHT
498 processing in terms of carotenoids can hardly be made for these samples.

499 For the carrot purees where oil was added during the preparation (**Fig. 9B**), a clear
500 effect of the sterilisation processes on the carotenoid content can be observed. The all-
501 *trans*- β -carotene contribution was decreased to around 50 %, indicating that in these
502 samples, isomerisation took place during processing. Compared to the samples without
503 oil, 13-*cis*- β -carotene was more important in this case. Difference might be explained
504 by the presence of the oil. Comparing samples enriched with oil but processed with
505 different technologies (**Fig. 9B**), a slightly higher degree of 13-*cis*- β -carotene was
506 detected after traditional thermal processing compared to HPHT processing. The fact

507 that at temperatures above 100 °C, increased processing time results in a higher degree
508 of isomerization was already observed by Knockaert et al. [13] studying β -carotene
509 isomerization in an oil/carrot emulsion.

510 As a final point of comparison, the colour values of the orange carrot purees were
511 evaluated (**Table 3**). Similar to the observations for the carotenoids, orange carrot
512 purees where no oil was added showed no changes in L^* , a^* and b^* values upon the
513 different sterilisation processes. It can thus be deduced that there is a clear correlation
514 between colour and carotenoids and this confirms that the colour of the orange carrot
515 purees is largely determined by the carotenoids. Also in the study of Vervoort et al.
516 [13], correlations between the colour values and specific carotenoids have been
517 revealed. For the orange carrot purees where oil was added during preparation, some
518 changes can be observed as a result of sterilisation. **Table 3** indicates that the sterilised
519 samples had a similar lightness, but they were less red than the blanched sample,
520 independent on the processing technology that was applied. This observation could be
521 related to decrease in all-*trans*- β -carotene which has been noted (**Fig. 9B**). Interestingly,
522 the b^* value of the sterilised carrot purees did not change for HPHT treated samples
523 compared to blanched samples, whereas for HT treated sample, a limited increase of the
524 b^* value could be seen. A similar behavior has been reported by Vervoort et al. [13] in
525 their comparative study on carrot pieces.

526 In literature, other studies on carrots have been conducted comparing specific quality
527 attributes of high pressure and thermally processed carrots. However, in most cases, no
528 fair comparison between both technologies could be made, since the processing
529 conditions that were applied did not result in an equivalent microbial inactivation (e.g.
530 [41, 42]). More recently, this fact was taken into consideration in studies of Knockaert
531 et al. [8] and Vervoort et al. [13] on carrot pieces. By applying specific processing

532 conditions for both the thermal and high pressure process, the comparison between both
533 technologies could occur on a fair basis. In the former study, focus was given to
534 carotenoid concentration and bioaccessibility and carrot tissue microstructure, in the
535 latter study, a whole range of quality attributes was considered, including enzymes,
536 sugars, vitamins, carotenoids and colour. In common with our study is the analysis of
537 the carotenoids. Some main differences and similarities are listed up. At first, the
538 contribution of all-*trans*- β -carotene is higher in the studies performed on carrot pieces.
539 In the present work, more isomerisation has probably been induced during the mixing
540 and homogenisation process. In the studies on carrot pieces (no oil present), HPHT
541 processing has been shown to result in a better retention of all-*trans*- β -carotene
542 compared to thermal processing (a significantly lower all-*trans*- β -carotene contribution
543 was detected in thermally sterilised carrot pieces). In the present work on orange carrot
544 puree, thermal and HPHT processing affected the carotenoid contribution in a similar
545 way, i.e. a limited or no effect in case of plain orange carrot puree and a distinct
546 decrease in all-*trans*- β -carotene contribution in case of carrot puree with oil. The
547 presence of oil clearly increased the sensitivity of β -carotene towards isomerisation,
548 which is in line with literature (e.g. [43]). Overall, it thus seems that the state of the
549 matrix (homogenised puree versus pieces) plays an important role in determining
550 whether there is a different effect of thermal and high pressure high temperature
551 sterilisation processes.

552 **4 Conclusion**

553 In this case study on orange and yellow carrots, the effect of thermal and high pressure
554 high temperature (HPHT) processing on a range of important quality attributes was
555 compared. The impact on the quality parameters was investigated from non-targeted and
556 targeted approach. In the first part, a HS-SPME-GC-MS fingerprinting technique was

557 used to compare the volatile fraction of differently processed carrot puree. In a second
558 part, a targeted approach to analyze specific quality related parameters (color,
559 carotenoid profile, particle size distribution, microscopy) was performed.

560 From the first part, the research outputs clearly showed the potential headspace
561 fingerprinting approach to zoom into discriminative markers which were clearly
562 detected in other concentrations depending on the processing technology used. In this
563 context, fingerprinting can be seen as a fast, data-based, hypothesis-free comparative
564 starting point, but it is not a result on its own. In the end, the identity of the fingerprint
565 marker should be studied in more detail, for example in the suggestion of the
566 consequence of the difference detected for the consumer. In this work, the identity of
567 the markers was specifically linked to flavour considerations. Given the fact that
568 different sterilization technologies and also different colored carrot varieties seem to
569 generate distinct headspace fingerprints suggests the potential that by particular mixing
570 and selection of the sterilized purees designing targeted carrot flavor profile could
571 become within reach. However, to establish models enabling process and product
572 design, more quantitative insight will be indispensable. In this context, kinetic studies in
573 which exact concentrations of selected markers are evaluated within a range of
574 processing variables should be the first step to be taken [44].

575 For the second part, as yellow carrot purees were shown to contain a very low amount
576 of carotenoids, focus was given to orange carrot purees for the actual comparative
577 study. The addition of olive oil to the orange carrot purees was considered as well.
578 HPHT processing and HT processing were shown to affect the particle size and the
579 microstructure of the orange carrot purees in a similar way. From a nutritional point of
580 view, in plain orange carrot puree, there was no or a limited effect of sterilisation
581 processes on the β -carotene content and isomerisation, independent on the technology

582 that was applied. In case oil was added to the orange carrot purees, significant
583 isomerisation of all-*trans*- β -carotene occurred, both during HPHT processing and
584 during HT processing. Overall, the colour values were shown to be a good
585 representation of the changes in carotenoids as a result of processing. Overall, for this
586 type of product and for the selected conditions and the selected nutritional and structural
587 quality parameters, it can be stated that no clear distinction can be made between HPHT
588 and HT sterilisation.

589 In general, when comparing the impact of HPHT and thermal processing, HPHT
590 processing resulted in a clear different effect on the volatile profile of orange and yellow
591 carrots, whereas no clear distinction could be with respect to the effect on structural and
592 nutritional attributes. In future, further research should also be done to investigate a
593 flavour consequence of the change in the headspace fraction.

594 **5 Acknowledgements**

595 This research was financially supported by the European Commission, Seventh
596 Framework Programme (FP7), Marie Curie Actions ‘HST Food Train’ and the KU
597 Leuven Research Fund. Tara Grauwet is a Postdoctoral Researcher funded by the
598 Research Foundation Flanders (FWO).

599 The authors thank Agilent technologies, Diegem, Belgium for providing us the Mass
600 Profiler professional (MPP) software.

601

602 6 References

- 603 1. Richardson P (2004) Improving the thermal processing of foods. Publishing Limited 507p
- 604 2. Hoogland H, De Heij W, Van Schepdael L (2001). High pressure sterilisation: novel technology,
605 new products, new opportunities. *New food* 3:21-26
- 606 3. Krebbers B, Matser A, Koets M, Bartels P, Van den Berg R (2002) High pressure-temperature
607 processing as an alternative for preserving basil. *High Pressure Research* 22:711-714
- 608 4. Matser AM, Krebbers B, van den Berg RW, Bartels PV (2004) Advantages of high pressure
609 sterilisation on quality of food products. *Trends Food Sci Technol* 15:79-85
- 610 5. Wilson DR, Dabrowski IZ, Stringer S, Moezelaar R, Brocklehurst TF (2008) High pressure in
611 combination with elevated temperature as a method for the sterilisation of food. *Trends Food Sci*
612 *Technol* 19:289-299
- 613 6. Sizer CE, Balasubramaniam VM, Ting EY (2002) Validating high pressure processes for low-
614 acid foods. *Food Technol* 56:36-42
- 615 7. Trejo Araya XI, Smale N, Zabarás D, Winley E, Forde C, Stewart CM, Mawson AJ (2009)
616 Sensory perception and quality attributes of high pressure processed carrots in comparison to
617 raw, sous-vide and cooked carrots. *Innov Food Sci Emerg Technol* 10:420-433
- 618 8. Knockaert G, De Roeck A, Lemmens L, Van Buggenhout S, Hendrickx M, Van Loey A (2011)
619 Effect of thermal and high pressure processes on structural and health-related properties of
620 carrots (*Daucus carota*). *Food Chem* 125:903-912
- 621 9. Verbeyst L, Bogaerts R, Van der Plancken I, Hendrickx M, Van Loey A (2012) Modelling of
622 vitamin C degradation during thermal and high pressure treatments of red fruit. *Food Bioprocess*
623 *Technol* 6:1015-1023
- 624 10. Van der Plancken I, Verbeyst L, De Vleeschouwer K, Grauwet T, Heiniö R-L, Husband FA,
625 Lille M, Mackie AR, van Loey A, Viljanen K, Hendrickx M (2012) (Bio)chemical reactions
626 during high pressure/high temperature processing affect safety and quality of plant-based Foods.
627 *Trends Food Sci Technol* 23:28-38
- 628 11. Timmermans RAH, Mastwijk HC, Knol JJ, Quataert MC, Vervoort L, van der Plancken I,
629 Hendrickx M, Matser AM (2011) Comparing equivalent thermal, high pressure and pulsed
630 electric field processes for mild pasteurization of orange juice. Part I: Impact on overall quality
631 attributes. *Innov Food Sci Emerg Technol* 12:235-243
- 632 12. Vervoort L, Van der Plancken I, Grauwet T, Timmermans RAH, Mastwijk HC, Matser AM,
633 Hendrickx M, Van Loey A (2011) Comparing equivalent thermal, high pressure and pulsed
634 electric field processes for mild pasteurization of orange juice. Part II: Impact on specific
635 chemical and biochemical quality parameters. *Innov Food Sci Emerg Technol* 12:466-477
- 636 13. Vervoort L, Van der Plancken I, Grauwet T, Verlinde P, Matser A, Hendrickx M, Van Loey A
637 (2012b) Thermal versus high pressure processing of carrots: A comparative pilot-scale study on
638 equivalent basis. *Innov Food Sci Emerg Technol* 15:1-3
- 639 14. Arscott SA, Tanumihardjo SA (2010) Carrots of many colors provide basic nutrition and
640 bioavailable phytochemicals acting as a functional food. *Compr Rev Food Sci F* 9:223-239
- 641 15. Yoo KS, Pike LM, Hamilton BKA (1997) A direct headspace sampling technique to rapidly
642 measure low boiling point volatile terpenoids in carrots. *HortScience* 32:714-716
- 643 16. Kjeldsen F, Christensen LP, Edelenbos M (2001) Quantitative analysis of aroma compounds in
644 carrot (*Daucus carota* L.) cultivars by capillary gas chromatography using large volume injection
645 technique. *J Agr Food Chem* 49:4342-4348
- 646 17. Ulrich D, Nothnagel T, Hoberg E, Straka P (2003) Analysis of aroma types by rapid GC methods
647 in carrot breeding research. In: *Flavour Research at the Dawn of the Twenty-First Century.*
648 *Proceedings of the 10th Weurman Flavour Research Symposium* (edited by J.L. Le Quéré & P.X.
649 Étievant). Pp. 440-444. Intercept Scientific Technical Medical Publisher
- 650 18. Fraser PD, Bramley PM (2004) The biosynthesis and nutritional uses of carotenoids. *Prog Lipid*
651 *Res* 43:228-265
- 652 19. Rodriguez-Amaya DB, Kimura M (2004) *HarvestPlus Handbook for Carotenoid Analysis.*
653 *Washington: International Food Policy Research Institute and International Center for Tropical*
654 *Agriculture*
- 655 20. Schieber A, Carle R (2005) Occurrence of carotenoid cis-isomers in food: Technological,
656 analytical, and nutritional implications. *Trends Food Sci Technol* 16:416-422

- 657 21. Böhm V, Puspitasari-Nienaber NL, Ferruzzi MG, Schwartz SJ (2002) Trolox equivalent
658 antioxidant capacity of different geometrical isomers of alpha-carotene, beta-carotene, lycopene, and
659 zeaxanthin. *J Agr Food Chem* 50:221-226
- 660 22. Van Buggenhout S, Alminger M, Lemmens L, Colle I, Knockaert G, Moelants K, Van Loey A,
661 Hendrickx M (2010) *In vitro* approaches to estimate the effect of food processing on carotenoid
662 bioavailability need thorough understanding of processing induced microstructural changes.
663 *Trends Food Sci Technol* 21:607-618
- 664 23. Patras A, Bruntona N, Da Pieve S, Butler F, Downey G (2009) Effect of thermal and high
665 pressure processing on antioxidant activity and instrumental colour of tomato and carrot purees.
666 *Innov Food Sci Emerg Technol* 10:16-22
- 667 24. Vervoort L, Grauwet T, Kebede BT, Van der Plancken I, Timmermans R, Hendrickx M, Van
668 Loey A (2012a) Headspace fingerprinting as an untargeted approach to compare novel and
669 traditional processing technologies: A case-study on orange juice pasteurisation. *Food Chem*
670 134:2303-2312
- 671 25. Vervoort L, Grauwet T, Njoroge DN, Van der Plancken I, Matser A, Hendrickx M, Van Loey A
672 (2013) Comparing thermal and high pressure processing of carrots at different processing
673 intensities by headspace fingerprinting. *Innov Food Sci Emerg Technol* 18:31-42
- 674 26. Kebede, B. T., Grauwet, T., Mutsokoti, L., Palmers, S., Vervoort, L., Hendrickx, M., Van Loey,
675 A (2014) Comparing the impact of high pressure high temperature and thermal sterilization on
676 the volatile fingerprint of onion, potato, pumpkin and red beet. *Food Res Int* 56:218-225.
- 677 27. Kebede, B. T., Grauwet, T., Palmers, S., Vervoort, L., Carle, R., Hendrickx, M., Van Loey, A
678 (2014) Effect of high pressure high temperature processing on the volatile fraction of differently
679 colored carrots. *Food Chemistry* 153: 340-352
- 680 28. Kebede, B. T., Grauwet, T., Tabilo-Munizaga, G., Palmers, S., Vervoort, L., Hendrickx, M., Van
681 Loey, A (2013) Headspace components that discriminate between thermal and high pressure
682 high temperature treated green vegetables: identification and linkage to possible process induced
683 chemical changes. *Food Chemistry* 141:1603-1613
- 684 29. Sadler G, Davis J, Dezman D (1990) Rapid extraction of lycopene and beta-carotene from
685 reconstituted tomato paste and pink grapefruit homogenates. *J Food Sci* 55:1460-1461
- 686 30. Lemmens L, Van Buggenhout S, Oey I, Van Loey A, Hendrickx M (2009) Towards a better
687 understanding of the relationship between the beta-carotene in vitro bio-accessibility and pectin
688 structural changes: A case study on carrots. *Food Res Int* 42:1323-1330
- 689 31. Ooms K (1996) Identification of potentially causal regressors in PLS models. Dissertation.
690 Katholieke Universiteit Leuven, Belgium
- 691 32. Knockaert G, Lemmens L, Van Buggenhout S, Hendrickx M, Van Loey A (2012) Changes in beta-
692 carotene bioaccessibility and concentration during processing of carrot puree. *Food Chem*
693 133:60-67
- 694 33. Knockaert G, Pulissery SK, Lemmens L, Van Buggenhout S, Hendrickx M, Van Loey A (2013)
695 Isomerization of carrot beta-carotene in presence of oil during thermal and combined thermal/high
696 pressure processing. *Food Chem* 138:1515-1520
- 697 34. Simon PW, Lindsay RC (1983) Effects of processing upon objective and sensory variables of
698 carrots. *J Amer Soc Hort Sci* 108:928-931
- 699 35. Frankel E.N (2005) In: *Lipid Oxidation*, 2nd edn. The Oily Press, Bridgwater, England
- 700 36. Marx, M, Schieber, A, Carle, R (2000) Quantitative determination of carotene stereoisomers in
701 carrot juices and vitamin supplemented (ATBC) drinks. *Food Chem* 70:403-408
- 702 37. Imsic M, Winkler S, Tomkins B, Jones R (2010) Effect of storage and cooking on beta-carotene
703 isomers in carrots (*Daucus carota* L. cv. 'Stefano'). *J Agr Food Chem* 58:5109-5113
- 704 38. Lemmens L, De Vleeschouwer K, Moelants K, Colle I, Van Loey A, Hendrickx M (2010) beta-
705 Carotene isomerization kinetics during thermal treatments of carrot puree. *J Agr Food Chem*
706 58:6816-6824
- 707 39. Sila DN, Smout C, Elliot F, Van Loey A, Hendrickx M (2006) Non-enzymatic depolymerisation
708 of carrot pectin: toward a better understanding of carrot texture during thermal processing. *J*
709 *Food Sci* 71:E1-E9
- 710 40. De Roeck A, Duvetter T, Fraeye I, Van der Plancken I, Sila DN, Van Loey A, Hendrickx M
711 (2009) Effect of high pressure/high temperature processing on chemical pectin conversions in
712 relation to fruit and vegetable texture. *Food Chem* 115:207-213
- 713 41. Trejo Araya XI, Smale N, Zabarás D, Winley E, Forde C, Stewart CM, Mawson AJ (2009)
714 Sensory perception and quality attributes of high pressure processed carrots in comparison to
715 raw sous-vide and cooked carrots. *Innov Food Sci Emerg Technol* 10:420-433

- 716
717
718
719
720
721
722
723
724
42. Nguyen LT, Tay A, Balasubramaniam VM, Legan JD, Turek EJ, Gupta R (2010) Evaluating the impact of thermal and pressure treatment in preserving textural quality of selected foods. *LWT – Food Sci Technol* 43:525-534
 43. Knockaert G, Sudheer KP, Lemmens L, Van Buggenhout S, Hendrickx, M, Van Loey A (2012) Carrot β -carotene degradation and isomerization kinetics during thermal processing in the presence of oil. *J Agr Food Chem* 60:10312-10319
 44. Grauwet T, Vervoort L, Colle I, Van Loey A, Hendrickx M (2014) From fingerprinting to kinetics in evaluating food quality changes. *Trends in Biotechnol* 32:125-131.




Fig. 1. Schematic overview of the general objective of this work. T_p: holding process temperature; t_h: holding time at process temperature; t_p: total process time including heating and cooling of the sample. T_i: initial product temperature at which the pressure build-up was initiated.




Fig 2. The profiles of thermal treatment with product temperature (thin grey solid line) and HPHT treatment with product temperature (dashed thick grey line) and pressure (thick dark solid line).




Fig 3. Total ion chromatogram of the headspace of blanched carrot purees obtained by SPME-GC-MS: (A) orange carrot; (B) yellow carrot.










Fig. 4. PLS-DA biplots of the effect of the treatment on carrot headspace fraction in blanched purees (▲) and after applying thermal treatment (HT, ●) or high pressure high temperature treatment (HP/HT, ★). (A) Orange carrot purees; (B) yellow carrot purees. Different volatiles are represented by small, open squares. Volatiles with VID higher than 0.800 in absolute value are named and marked in bold (small filled squares). Vectors indicate the correlation loadings for the categorical Y-variables. The percentages of the variances in X and Y explained by each latent variable (LV1 and LV2) are indicated on the respective axes.






Fig. 5. Discriminative headspace components for comparison of treatment impact on orange carrot purees. Volatiles with VID higher than 0.900 in absolute value in **Table 1** are represented. The Y-axis indicates the peak area and error bars represent the standard error of the analysis (n = 6).




Fig. 6. Discriminative headspace components for comparison of treatment impact on yellow carrot purees. Volatiles with VID higher than 0.900 in absolute value in **Table 2** are represented. The Y-axis indicate the peak area and error bars represent the standard error of the analysis (n = 6).




Fig. 7: Light micrographs of blanched orange carrot purees without oil addition (A) and with oil addition (B) and of blanched yellow carrot purees without oil addition (C) and with oil addition (D).




Fig. 8: Cumulative particle size distribution curves of blanched orange carrot purees (black curves) and of blanched yellow carrot purees (grey curves) without oil addition (full curves) and with oil addition (dashed curves).




Fig. 9: β -Carotene contribution (mean \pm standard error) in blanched orange carrot purees and in sterilized orange carrot purees using traditional thermal processing (HT) and high pressure high temperature processing (HPHT) without the addition of oil (A) and with the addition of oil (B). (■) all-*trans*- β -carotene contribution; (■) 9-*cis*- β -carotene contribution; (■) 13-*cis*- β -carotene contribution; (□) 15-*cis*- β -carotene contribution.

Table 1. Discriminative headspace components for each treatment in orange and yellow carrot purees, selected through the VID procedure (higher than 0.800 in absolute value), and listed in increasing order of VID coefficient. Positive/negative VID coefficients indicate an increase/decrease of its concentration respectively, after the corresponding treatment in comparison with the other treatments. The corresponding PLS-DA model contained two latent variables, explaining more than 93 and 88 % of Y-variance in orange and yellow carrot, respectively. The retention time index (RTI) per compound is listed.

	Blanched			HT			HPHT		
	Compound	RTI	VID	Compound	RTI	VID	Compound	RTI	VID
Orange carrot puree	Octanal	1153	-0.972	γ -Terpinene	1211	0.804	Sesquithujene	1537	-0.907
	α -Ionone	1558	-0.954	α -Terpinene	1171	0.806	β -Caryophyllene epoxide	1833	-0.899
	Heptanal	1048	-0.942	α -Pinene	1087	0.821	Cedrene	1587	-0.898
	2-Nonenal	1307	-0.933	β -Caryophyllene	1568	0.825	β -Elemene	1531	-0.852
	p-Cymenene	1244	-0.912	α -Caryophyllene	1604	0.842	β -Caryophyllene	1568	-0.814
	Hexanal	933	-0.876	β -Elemene	1531	0.861	α,β -Dihydropseudoionone	1574	0.802
	Decanal	1349	-0.815	Terpinolene	1240	0.945	2,4-Decadienal	1468	0.803
						2-Decenal	1401	0.827	
						2-Octenal	1207	0.906	
Yellow carrot puree	Heptanal	1048	-0.819	γ -Terpinene	1211	0.804	Terpinolene	1240	-0.953
	Octanal	1153	-0.816	p-Cymenene	1244	0.808	γ -Terpinene	1211	-0.934
	p-Cymenene	1244	-0.814	Heptanal	1048	0.828	α -Pinene	1087	-0.867
	α -Thujene	1077	0.885	Benzeneacetaldehyde	1195	0.936	β -Pinene	1134	-0.824
	Sabinene	1127	0.937	o-Cymene	1178	0.959	α -Caryophyllene	1604	-0.818

Table 2: Characterisation of the lipophilic extract of the blanched orange and yellow carrot purees without or with oil addition in terms of carotenoid content and colour values (L*, a*, b*).

orange carrot puree			yellow carrot puree		
	without oil	with oil		without oil	with oil
all-<i>trans</i>-β-carotene ($\mu\text{g/g}$ carrot puree)	25.4 \pm 2.4	33.7 \pm 2.0	lutein ($\mu\text{g/g}$ carrot puree)	3.1 \pm 0.7	2.5 \pm 0.2
9-<i>cis</i>-β-carotene ($\mu\text{g/g}$ carrot puree)	2.9 \pm 0.1	2.6 \pm 0.3			
13-<i>cis</i>-β-carotene ($\mu\text{g/g}$ carrot puree)	1.5 \pm 0.1	1.8 \pm 0.1			
15-<i>cis</i>-carotene ($\mu\text{g/g}$ carrot puree)	2.8 \pm 0.4	1.8 \pm 0.2			
L*	47.5 \pm 2.5	64.1 \pm 0.1	L*	58.3 \pm 0.1	75.1 \pm 0.1
a*	31.7 \pm 2.7	24.6 \pm 0.1	a*	5.8 \pm 0.1	1.5 \pm 0.1
b*	43.3 \pm 4.2	68.8 \pm 0.3	b*	61.0 \pm 0.4	51.3 \pm 0.2

Table 3: Colour values (L^* , a^* , b^*) of the lipophilic extract of blanched orange carrot purees and of sterilised orange carrot purees using thermal (HT) and high pressure high temperature (HPHT) processing, without and with the addition of oil.

orange carrot puree without oil			
	Blanched	HT	HPHT
L*	47.5 ± 2.5	46.1 ± 0.1	48.0 ± 0.5
a*	31.7 ± 2.7	31.6 ± 0.2	31.9 ± 1.0
b*	43.3 ± 4.2	46.5 ± 0.2	39.5 ± 2.0

orange carrot puree with oil			
	Blanched	HT	HPHT
L*	64.1 ± 0.1	65.9 ± 0.3	66.2 ± 0.4
a*	24.6 ± 0.1	18.6 ± 0.2	18.8 ± 0.6
b*	68.8 ± 0.3	74.1 ± 1.2	66.7 ± 2.9