



Effect of milk type and processing on iodine concentration of organic and conventional winter milk at retail: implications for nutrition

Article

Accepted Version

Payling, L. M., Juniper, D. T., Drake, C., Rymer, C. and Givens, D. I. (2015) Effect of milk type and processing on iodine concentration of organic and conventional winter milk at retail: implications for nutrition. *Food Chemistry*, 178. pp. 327-330. ISSN 0308-8146 doi: <https://doi.org/10.1016/j.foodchem.2015.01.091> Available at <http://centaur.reading.ac.uk/39199/>

It is advisable to refer to the publisher's version if you intend to cite from the work.

Published version at: <http://www.sciencedirect.com/science/article/pii/S0308814615001053>

To link to this article DOI: <http://dx.doi.org/10.1016/j.foodchem.2015.01.091>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1

2 **Effect of milk type and processing on iodine concentration of organic and conventional**
3 **winter milk at retail: implications for nutrition**

4

5 Laura M. Payling, Darren T. Juniper, Chris Drake, Caroline Rymer and D. Ian Givens¹

6 Food Production and Quality Division, Faculty of Life Sciences, University of Reading,

7 Reading RG6 6AR, UK

8

9 **Short title:** Iodine in organic and normal winter milk

10

11 **Keywords:** Iodine: Milk: Organic: UHT, Fat class

¹ Corresponding author, Tel: +44 118 378 8491; email: d.i.givens@reading.ac.uk

12 **Abstract**

13

14 Milk is the largest source of iodine in UK diets and an earlier study showed that organic
15 summer milk had significantly lower iodine concentration than conventional milk. There are
16 no comparable studies with winter milk or the effect of milk fat class or heat processing
17 method. Two retail studies with winter milk are reported. Study 1 showed no effect of fat
18 class but organic milk was 32.2% lower in iodine than conventional milk (404 vs. 595 $\mu\text{g/L}$;
19 $P<0.001$). Study 2 found no difference between conventional and Channel Island milk but
20 organic milk contained 35.5% less iodine than conventional milk (474 vs. 306 $\mu\text{g/L}$;
21 $P<0.001$). UHT and branded organic milk also had lower iodine concentrations than
22 conventional milk (331 $\mu\text{g/L}$; $P<0.001$ and 268 $\mu\text{g/L}$; $P<0.0001$ respectively). The results
23 indicate that replacement of conventional milk by organic or UHT milk will increase the risk
24 of sub-optimal iodine status especially for pregnant/lactating women.

25

26 **Keywords: Iodine: Milk: Organic: UHT, Fat class**

27

28 **1. Introduction**

29

30 Iodine is a key component of the hormones produced by the thyroid gland which are
31 particularly important during pregnancy for foetal brain development (Zimmermann, 2009).
32 The UK Reference Nutrient Intakes (RNI) for iodine are 130 and 140 $\mu\text{g/d}$ for children aged
33 11-14 years and adults respectively with no increase during pregnancy or lactation
34 (Department of Health, 1991), compared with 150 μg iodine/d for adults by the US Institute of
35 Medicine (IOM) and the World Health Organisation (WHO). IOM also proposes an increase
36 of 47 and 93% during pregnancy and lactation respectively whereas WHO advises an

37 increase of 67% for both (Zimmermann, 2009). In many countries provision of iodised salt
38 has reduced the incidences of iodine deficiency, but in countries such as the UK widespread
39 interventions have not been enforced (Phillips, 1997). This has led to a sustained level of
40 mild iodine deficiency in many populations, notably about 44% of children and adults in
41 Europe (Zimmermann & Andersson, 2011).

42

43 For some time the UK population has been assumed to be of adequate iodine status
44 (Wenlock, Buss, Moxon & Bunton, 1982). However, a study in UK schoolgirls recorded mild
45 iodine deficiency in 51% of the participants based on urinary iodine concentrations
46 (Vanderpump et al., 2011) and the recently published UK National Diet and Nutrition Survey
47 (Bates et al., 2014) reports that on average, young females aged 11 to 18 years consume only
48 81% of the RNI for iodine and that 22% of young females have iodine intakes below the
49 Lower RNI (70 µg iodine/d). In addition, mild-to-moderate iodine deficiency has been
50 demonstrated in a large UK cohort of pregnant women (Bath, Walter, Taylor, Wright &
51 Rayman, 2014) together with evidence of an association between low maternal iodine status
52 in early pregnancy and poorer verbal IQ, reading accuracy and reading comprehension in the
53 children (Bath, Rayman, Steer, Goulding & Emmett, 2013).

54

55 In the UK, milk and dairy foods are the largest dietary source of iodine providing 40 and 39
56 % of the daily intake of iodine for 11-18 year old males and females respectively (Bates et
57 al., 2014). Both genders obtain the majority (~43%) of the dairy food contribution from semi-
58 skimmed milk (Bates et al., 2014). Survey studies on milk iodine concentrations carried out
59 in the last 14 years (Ministry of Agriculture, Fisheries and Food, 2000; Food Standards
60 Agency, 2008) are not suggestive of any overall reduction in UK milk iodine concentration
61 but they do show that iodine concentration is highly variable and that milk produced in the

62 summer has on average, a 50% lower iodine concentration than winter milk. Moreover, the
63 studies of The Food Standards Agency (2008) and Bath, Button & Rayman (2012) both
64 showed that summer milk from organic dairy systems had significantly lower (~ -40%)
65 iodine concentrations than from conventional systems. These findings clearly have
66 implications for human iodine intake and status but there is little evidence on conventional
67 vs. organic milk produced in winter, the effect of milk fat class (whole, semi-skimmed,
68 skimmed) or the effect of milk processing method. Accordingly, two UK retail studies were
69 therefore undertaken with the objectives of comparing the iodine content of 1) organic and
70 conventional winter milk, 2) whole, semi-skimmed and skimmed milk and 3) pasteurised and
71 ultra-high temperature (UHT) treated milk.

72

73 **2. Materials and methods**

74

75 *2.1 Milk samples*

76

77 In Study 1, the intention was to purchase two samples of pasteurised conventionally and
78 organically produced whole (full fat), semi-skimmed and skimmed milk from two leading
79 supermarkets in the Reading area in late January 2014. However, Supermarket 1 did not stock
80 organic skimmed milk when visited, leading to a total of 22 samples. In Study 2, five
81 different milk product types (conventional semi-skimmed, organic semi-skimmed, branded
82 organic semi-skimmed, UHT semi-skimmed and conventional whole milk from Channel
83 Island breeds of cow) were purchased from four leading supermarkets in the Reading area in
84 each of three consecutive weeks, beginning in the first week of February 2014 giving a total
85 of 60 samples. Except for the UHT milk, all other types were pasteurised. In both studies all

86 milks were supermarket own brand except for the branded organic milk in Study 2. All milk
87 samples were stored at -20°C pending analysis.

88

89 *2.2 Sample analysis*

90

91 Milk samples were allowed to defrost overnight and thoroughly mixed by vortexing before
92 analysis. Samples from Study 1 had fat, protein and lactose concentrations measured by
93 methods of International Organisation for Standardisation (2008), Davis & Macdonald (1953)
94 and Sanchez-Manzanera, Fernandez-Villacaas, Marin-Iniesta & Laencina (1993)
95 respectively. Iodine concentration was measured in all samples by alkali extraction followed
96 by inductively coupled plasma mass spectrometry (ICP-MS) based on the method of Fecher,
97 Goldman & Nagengast (1998). In brief 100 µl of milk were diluted to 10 ml with 2%
98 tetramethyl ammonium hydroxide in ultra-pure water and containing 5 µg/L of rhodium as an
99 internal standard. Analysis was then undertaken by ICP-MS (iCAP Q, Thermo Scientific
100 Scientific Inc. Waltham, MA, USA). Certified standard solutions (Romil Ltd, Cambridge,
101 UK) based on ammonium iodide were treated as for milk samples leading to final
102 concentrations of 0, 2, 4, 6, 8 and 10 µg/L iodine.

103

104 *2.3 Statistical analysis*

105

106 The effect of milk product type (conventional production, Channel Island, organic
107 production, UHT, branded organic), milk fat class (whole, semi-skimmed, skimmed) and
108 supermarket of origin were determined as appropriate to each study, by fixed effect analysis
109 of variance using a general linear model (Mintab version 16; Minitab Inc., State College, PA,

110 USA). Tukey's pairwise multiple comparison test was then used to identify which treatments
111 were significantly different from each other when the significance was $P < 0.05$.

112

113 **3. Results**

114

115 *3.1 Study 1*

116

117 There was no significant effect of conventional vs. organic production system on fat, protein
118 or lactose contents of the milks. Fat content was significantly ($P < 0.001$) affected by milk fat
119 class with mean values of 3.55, 1.42 and 0.01 g/100g for whole, semi-skimmed and skimmed
120 milk respectively. The iodine concentrations of the milks are shown in Table 1. There was no
121 effect of milk fat class and overall, no fat class x production system interaction although such
122 an interaction was seen for milk from Supermarket 2 which was related to some small fat
123 class effects ($P < 0.05$) in conventional but not organic milk. Overall, there was a significant
124 ($P < 0.001$) effect of production system with organic milk having consistently lower iodine
125 concentrations than conventional. Mean iodine concentration in organic milk was 32.2%
126 lower than of conventional.

127

128 *3.2 Study 2*

129

130 The iodine concentrations of the milk types are shown in Table 2. Overall, milk iodine
131 concentration was not affected by supermarket but was significantly ($P < 0.001$) influenced by
132 milk production system with organic and branded organic milk having lower iodine
133 concentrations than conventional. Branded organic milk had the lowest mean iodine
134 concentration and this had a tendency to be lower than own-brand organic ($P < 0.059$).

135 Overall, the iodine concentration of organic (including branded organic) milk was 35.5%
136 lower than conventional. A production system x supermarket interaction ($P<0.05$) was seen
137 due to Supermarket 2 having unusually high iodine concentrations in its milk labelled as
138 organic. If data from this supermarket are excluded the iodine concentration of organic
139 (including branded organic) milk was 43.9% lower than conventional. Interestingly,
140 conventionally produced UHT milk had a lower ($P<0.05$) iodine concentration than
141 conventional pasteurised milk with a mean value not significantly different to organic milk.
142 Iodine in Channel Island milk was not significantly different to conventional milk.

143

144 **4. Discussion**

145

146 To our knowledge, this is the first balanced comparison of the iodine concentration of
147 conventional and organic UK retail milk produced in the winter. The Food Standards Agency
148 (2008) reported a study on milk purchased between January and March 2007 but that study
149 involved only whole milk and was not balanced, involving 62 conventional and 14 organic
150 milks. Despite this, the mean values of the current study agree quite well with those of the
151 Food Standards Agency (2008) which reported values of 441 and 391 $\mu\text{g/L}$ for conventional
152 and organic milk respectively, though no statistical comparison was reported. The key finding
153 of the current study that winter produced organic milk has a significantly lower iodine
154 concentration (-32.2, -35.5%; Study1, Study 2) than winter produced conventional milk,
155 agrees with an earlier UK study (Bath et al., 2012) and two in Scandinavia (Rasmussen,
156 Larsen & Ovesen, 2000; Dahl, Opsahl, Meltzer & Julshamn, 2003) comparing conventional
157 with organic summer milk. However both the conventional and organic milk in the present
158 study had higher iodine concentrations than the respective values in the UK summer milk
159 study (Bath et al., 2012). It has been recognised for some time that milk produced in the

160 winter has higher iodine concentrations than that produced in summer (Flachowsky, Franke,
161 Meyer, Leiterer, & Schöne, 2014). This is because iodine intake by the dairy cow is a key
162 determinant of milk iodine concentration and housed cows in winter will receive more iodine
163 supplementation via concentrate feed than cows grazing outdoors in summer (Flachowsky et
164 al., 2014). Indeed since diets for dairy cows on an organic system are mandated to rely more
165 heavily on forages than in conventional systems (European Union, 2008), this is also likely to
166 be the main reason for organic milk having lower iodine concentrations although additional
167 reasons have been proposed (Flachowsky et al., 2014). The finding that branded organic milk
168 had a tendency ($P<0.059$) to have a lower iodine content than supermarket own brand organic
169 milk agrees with the findings of Bath et al. (2012) who showed that branded organic summer
170 milk (mean 118.3 $\mu\text{g/L}$) was significantly ($P<0.001$) lower than supermarket own-brand
171 organic (mean 159.8 $\mu\text{g/L}$). The reasons for this are not known but may relate to less iodine
172 supplementation to the dairy cows and perhaps less use of iodine containing animal health
173 products (e.g. teat dips) than for supermarket own-brand organic milk.

174

175 One report from a large Spanish milk study in 2008 (Soriguer et al., 2011) showed that the
176 iodine concentration was greater ($P<0.001$) in skimmed milk (mean 273 $\mu\text{g/L}$; $n=107$) than in
177 semi-skimmed milk (mean 254 $\mu\text{g/L}$; $n=121$) or whole milk (mean 251 $\mu\text{g/L}$; $n=134$). Whilst
178 the logic for this might relate to iodine dilution as fat content is increased, no such effect was
179 seen in the present study and as Soriguer et al. (2011) indicate, the variation due to fat content
180 is very small compared with the large variation in iodine concentration seen in most studies.
181 There are also few reports on the effect of milk processing on iodine concentration.
182 Flachowsky et al. (2014) summarised the results of two Iranian studies (Norouzian,
183 Valizadeh, Azizi, Hedayati, Naserian, & Eftekhari Shahroodi, 2009; Norouzian, 2011) and
184 concluded that between 18 and 53% of iodine in raw milk was lost during pasteurisation

185 ($P<0.01$) with the loss being inversely related to the iodine content of the raw milk. It was
186 proposed that the losses were due to the sublimation characteristic of iodine as >90% of milk
187 iodine is in the inorganic form (Flachowsky et al., 2014) although how this occurs in an
188 essentially sealed system is unclear. In the present study all the milk was pasteurised
189 (typically 72°C for 15 seconds) except for the UHT milk which would have been processed
190 typically at 140°C for 3 to 5 seconds. It seems possible therefore that the significantly lower
191 iodine concentration in the UHT milk compared with conventional may be a result of the
192 high temperature treatment. Whilst in the UK, UHT milk accounts for only 5-6% of the liquid
193 milk market (Dairy Co, 2014), in many neighbouring EU Member States UHT milk
194 represents most of the milk consumed with for example France, The Netherlands and
195 Germany where UHT accounts for about 90, 80 and 65% respectively (Department for
196 Environment, Food and Rural Affairs, 2007). The impact of UHT processing should therefore
197 be explored in greater depth.

198

199 For adults the UK the RNI for iodine is 140 µg/d with no increase during pregnancy or
200 lactation (Department of Health, 1991) despite evidence that potassium iodide
201 supplementation (providing 153 µg I/d) during lactation significantly increases breast milk
202 iodine concentration and iodine status of premature infants (González-Iglesias et al., 2012).
203 Assuming the mean intake of milk by UK women (aged 19-64 years) is 125 ml/d (Bates et
204 al., 2014) the results in Table 2 would predict iodine intakes from milk of 59 and 43 µg/d
205 from conventional and organic/UHT/branded organic winter milk respectively, representing
206 42 and 31% of the RNI. Milk consumption in the UK has declined by about 23 % over the
207 last 20 years (Department for Environment, Food and Rural Affairs, 2013) with a greater
208 reduction by young women than men, and whilst this is likely to be a major contributory
209 factor to the low iodine status seen in pregnant women (Bath et al., 2014), a key conclusion

210 of the present study is that choosing to consume winter organic or UHT milk will increase the
211 risk of sub-optimal iodine status, particularly during pregnancy and lactation, in agreement
212 with the conclusions of Bath et al. (2012) for summer milk.

213

214 The present study has weaknesses. The sample number was not large and fewer samples than
215 planned were available for Study 1. Nevertheless, the findings regarding the relative iodine
216 concentrations in winter milk from differing production systems and processing methods are
217 important issues in understanding the effects of diet on iodine status. The present study was
218 also limited to supermarkets in the Reading area although the retail study of Bath et al. (2012)
219 found no significant effect of area of milk purchase, arguing that retail milk is relatively
220 homogenous as result of collections from wide geographical locations being taken for central
221 processing and distribution to supermarkets over a wide area. There were however
222 supermarket x production system interactions seen in both current studies which were related
223 to inconsistencies between supermarkets in the effect of production system on iodine content.
224 The reasons for this are unknown but possibly point to differences between supermarkets'
225 milk producers or to labelling errors.

226

227 **5. Conclusions**

228

229 Both studies showed that organic winter milk was significantly lower (~34%) in iodine
230 concentration than conventional milk in agreement with an earlier study with summer milk.
231 No effect of fat class was seen but UHT and branded organic milk had lower iodine
232 concentrations than conventional milk. The results for winter milk agree with conclusions in
233 an earlier study with summer milk that replacement of conventional milk by organic milk will
234 increase the risk of sub-optimal iodine status especially for young women in periods of

235 increased iodine demand such as pregnancy and lactation. It seems that conventionally
236 produced UHT milk will have the same effect as organic milk although more work on this is
237 warranted.

238

239 **Acknowledgements**

240

241 We thank Dr Yan Gao for her expert contribution to the milk analysis and the University of
242 Reading for funding the studies. All authors contributed to the writing and reviewing of the
243 paper and approved the final document. None of the authors had any conflict of interest to
244 declare.

245

246 **References**

247

248 Bates, B., Lennox, A., Prentice, A., Bates, C., Page, P., Nicholson, S., & Swan, G. (2014).
249 *National Diet and Nutrition Survey, Results from Years 1-4 (combined) of the Rolling*
250 *Programme (2008/2009-2011/12)*. A survey carried out on behalf of Public Health England
251 and the Food Standards Agency.

252 Bath, S. C., Button, S., & Rayman, M. P. (2012). Iodine concentration of organic and
253 conventional milk: implications for iodine intake. *British Journal of Nutrition*, 107, 935-940.

254 Bath, S. C., Rayman, M. P., Steer, C. D., Golding, J., & Emmett, P. (2013). Effect of
255 inadequate iodine status in UK pregnant women on cognitive outcomes in their children:
256 results from the Avon Longitudinal Study of Parents and Children (ALSPAC). *The Lancet*,
257 382 (9889), pp. 331-337.

258 Bath, S. C., Walter, A., Taylor, A., Wright, J., & Rayman, M. P. (2014). Iodine deficiency in
259 pregnant women living in the South East of the UK: the influence of diet and nutritional
260 supplements on iodine status. *British Journal of Nutrition*, 111, 1622 – 1631,

261 Dahl, L., Opsahl, J. A., Meltzer, H. M., & Julshamn, K. (2003). Iodine concentration in
262 Norwegian milk and dairy products. *British Journal of Nutrition*, 90, 679–685.

263 Dairy Co (2014). *Market Information*. [http://www.dairyco.org.uk/market-information/dairy-](http://www.dairyco.org.uk/market-information/dairy-sales-consumption/liquid-milk-market/liquid-milk-market/)
264 [sales-consumption/liquid-milk-market/liquid-milk-market/](http://www.dairyco.org.uk/market-information/dairy-sales-consumption/liquid-milk-market/liquid-milk-market/) Accessed 01.05.14.

265 Davis, J. G., & Macdonald, F. J. (1953). Total protein content from acidity and pH
266 determination. In *Richmond's Dairy Chemistry Fifth Edition*. (p 362). London: Charles
267 Griffin & Co. Ltd.

268 Department for Environment, Food and Rural Affairs (2013). *Family Food 2012, Report*
269 *Tables and Charts*. <https://www.gov.uk/government/collections/family-food-statistics>
270 (accessed 13 May 2014).

271 Department for Environment, Food and Rural Affairs (DEFRA) (2007). *The Potential for*
272 *GB-European Trade in Liquid Milk*. (71 pp). London: DEFRA,

273 Department of Health (1991). *Report of the panel on dietary reference values of the*
274 *Committee on Medical Aspects of Food Policy (COMA)*. In *Dietary Reference Values for*
275 *Food Energy and Nutrients for the United Kingdom*. London: The Stationery Office.

276 European Union (2008). Commission Regulation (EC) no. 889/2008 of 5 September 2008
277 laying down detailed rules for the implementation of Council Regulation (EC) no. 834/2007
278 on organic production and labelling of organic products with regard to organic production,
279 labelling and control. *Official Journal of the European Union* 51, L250.

280 Fecher, P. A., Goldmann, I., & Nagengast, A. (1998). Determination of iodine in food
281 samples by inductively coupled plasma mass spectrometry after alkaline extraction. *Journal*
282 *of Analytical Atomic Spectrometry*, 13, 977–982.

283 Flachowsky, G., Franke, K., Meyer, U., Leiterer, M., & Schöne, F. (2014). Influencing
284 factors on iodine content of cow milk. *European Journal of Nutrition*, 53, 351-365.

285 Food Standards Agency (2008). *Retail Survey of Iodine in UK Produced Dairy Foods, FSIS*
286 *02/08*. London: Food Standards Agency.

287 González-Iglesias, H., de la Flor St Remy, R.R., López-Sastre, J., Fernández-Colomer, B.,

288 Ibáñez-Fernández, A., Solís, G., Sanz-Medel, A., & Fernández-Sánchez, M.L. (2012).
289 Efficiency of iodine supplementation, as potassium iodide, during lactation: A study in
290 neonates and their mothers. *Food Chemistry*, 133, 859–865.

291 International Organisation for Standardisation (2008). *Milk: Determination of fat content*.
292 ISO 2446:2008. Geneva: International Organisation for Standardisation.
293 http://www.iso.org/iso/home/store/catalogue_tc/catalogue_detail.htm?csnumber=51019
294 Accessed 02.02.14.

295 Ministry of Agriculture, Fisheries and Food (2000). *Iodine in Milk, Food Surveillance*
296 *Information Sheet 198*. London: MAFF Joint Food Safety and Standards Group.

297 Norouzian, M. A. (2011). Iodine in raw and pasteurized milk of dairy cows fed different
298 amounts of potassium iodide. *Biological Trace Element Research*, 139, 160–167.

299 Norouzian, M. A., Valizadeh, R., Aziz, F., Hedayati, M., Naserian, A. A., & Eftekhari
300 Shahroodi, F. (2009). The effect of feeding different levels of potassium iodide on
301 performance, T-3 and T-4 concentrations and iodine excretion in Holstein dairy cows.
302 *Journal of Animal and Veterinary Advances*, 8, 111–114.

303 Phillips, D. I. (1997). Iodine, milk, and the elimination of endemic goitre in Britain: the story
304 of an accidental public health triumph. *Journal of Epidemiology and Community Health*, 51,
305 391–393.

306 Rasmussen, L. B., Larsen, E. H., & Ovesen, L. (2000). Iodine content in drinking water and
307 other beverages in Denmark. *European Journal of Clinical Nutrition*, 54, 57–60.

308 Sanchez-Manzanera, J. A., Fernandez-Villacaas, M. R., Marin-Iniesta, F., & Laencina, J.
309 (1993). Determination of lactose by an enzymatic method. *Food Chemistry*, 46, 425-427.

310 Soriguer, F., Gutierrez-Repiso, C., Gonzalez-Romero, S., Olveira, G., Garriga, M. J.,
311 Velasco, I., Santiago, P., de Escobar, G., & Garcia-Fuentes, E. (2011). Iodine concentration
312 in cow's milk and its relation with urinary iodine concentrations in the population. *Clinical*
313 *Nutrition*, 30, 44-48.

314 Vanderpump, M. P. J., Lazarus, J. H., Smyth, P. P., Laurberg, P., Holder, R. L., Boelaert, K.,
315 & Franklyn, J. A. (2011). Iodine status of UK schoolgirls: a cross-sectional survey. *The*
316 *Lancet*, 377 (9782), pp.2007-2012.

317 Wenlock, R. W., Buss, D. H., Moxon, R. E., & Bunton, N. G. (1982). Trace nutrients. 4.
318 Iodine in British food. *British Journal of Nutrition*, 47, 381-390.

319 Zimmermann, M. B. (2009). Iodine deficiency. *Endocrine Reviews*, 30, 376-408.

320 Zimmermann, M. B., & Andersson, M. (2011). Prevalence of iodine deficiency in Europe in
321 2010. *Annales d'Endocrinologie*, 72, 164-166.

322

323

Table 1. Study 1: Least square mean (\pm SE) iodine concentrations of retail milk as influenced by milk fat class (FC), production system (conventional vs. organic; PS) and supermarket (SM)

Super-market	Mean iodine concentration ($\mu\text{g/L}$)						SED	P-value for			
	Conventional			Organic				FC	PS	FC x PS	SM x PS
	Whole n=4	Semi-skimmed n=4	Skimmed n=4	Whole n=4	Semi-skimmed n=4	Skimmed n=2					
1	486.0 \pm 7.0	560.0 \pm 20.0	502.5 \pm 3.5	482.5 \pm 7.0	454.0 \pm 8.0	nd	nd	NS	NS	nd	-
2	669.5 \pm 3.5 ^{ab}	585.5 \pm 21.5 ^b	768.5 \pm 24.5 ^a	316.0 \pm 21.0 ^c	419.0 \pm 41.0 ^c	325.5 \pm 12.5 ^c	33.43	NS	<0.001	0.003	-
Overall	577.8 \pm 53.1 ^a	572.8 \pm 14.1 ^a	635.5 \pm 77.4 ^a	399.2 \pm 48.9 ^b	436.5 \pm 19.8 ^b	325.5 \pm 12.5 ^b	40.67	NS	<0.001	NS	<0.001

^{a,b}Means within a row with no superscripts or those with a common superscript are not significantly different ($P>0.05$); NS, not significant ($P>0.05$); nd, no or insufficient data due to missing samples; SED, standard error of the difference

Table 2. Study 2: Least square mean (\pm SE) iodine concentrations of retail milk as influenced by type (T) and supermarket (SM).

Supermarket	Mean iodine concentration ($\mu\text{g/L}$)					SED	<i>P</i> -value for		
	Conventional n=12	Channel Island n=12	Organic n=12	UHT n=12	Branded organic n=12		T	SM	T x SM
1	433.9 \pm 27.8 ^{ab}	436.8 \pm 59.9 ^a	327.5 \pm 15.1 ^{ab}	336.9 \pm 6.85 ^{ab}	274.5 \pm 36.6 ^b	48.86	0.029	-	-
2	411.1 \pm 62.8	482.1 \pm 17.4	478.7 \pm 74.0	280.7 \pm 33.8	301.5 \pm 43.3	71.38	0.046	-	-
3	490.9 \pm 23.6 ^a	445.3 \pm 56.9 ^a	279.4 \pm 15.5 ^b	341.6 \pm 33.6 ^{ab}	245.8 \pm 11.9 ^b	47.17	0.002	-	-
4	559.7 \pm 39.2 ^a	398.8 \pm 35.5 ^b	287.8 \pm 57.2 ^b	366.4 \pm 14.2 ^b	249.2 \pm 28.2 ^b	48.53	<0.001	-	-
Overall	473.9 \pm 24.9 ^a	440.8 \pm 21.5 ^a	343.4 \pm 30.3 ^b	331.4 \pm 14.3 ^b	267.7 \pm 15.3 ^b	27.10	<0.001	NS	0.013

^{a,b}Means within a row with no superscripts or those with a common superscript are not significantly different ($P>0.05$); NS, not significant ($P>0.05$); SED, standard error of the difference