



Methane emissions from cattle: estimates from short-term measurements using a Green Feed system compared with measurements obtained using respiration chambers or sulphur hexafluoride tracer

Article

Accepted Version

Hammond, K., Humphries, D., Crompton, L., Green, C. and Reynolds, C. (2015) Methane emissions from cattle: estimates from short-term measurements using a Green Feed system compared with measurements obtained using respiration chambers or sulphur hexafluoride tracer. *Animal Feed Science and Technology*, 203. pp. 41-52. ISSN 0377-8401 doi: <https://doi.org/10.1016/j.anifeedsci.2015.02.008> Available at <http://centaur.reading.ac.uk/39710/>

It is advisable to refer to the publisher's version if you intend to cite from the work.

Published version at: <http://www.sciencedirect.com/science/article/pii/S0377840115000590>

To link to this article DOI: <http://dx.doi.org/10.1016/j.anifeedsci.2015.02.008>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law,

including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24

Methane emissions from cattle: estimates from short-term measurements using a GreenFeed system compared with measurements obtained using respiration chambers or sulphur hexafluoride tracer

K.J. Hammond, D.J. Humphries, L.A. Crompton, C. Green, and C.K. Reynolds*

School of Agriculture, Policy and Development, Centre for Dairy Research, University of Reading, PO Box 237, Earley Gate, Reading RG 6AR, United Kingdom

*** Corresponding author. Tel: +44 118 3784684; Fax +44 1183786595**

E-mail address: c.k.reynolds@reading.ac.uk (C.K. Reynolds)

25 Abstract

26 The GreenFeed (GF) system (C-Lock Inc., Rapid City, USA) is used to estimate total
27 daily methane emissions of individual cattle using short-term measurements obtained over
28 several days. Our objective was to compare measurements of methane emission by growing
29 cattle obtained using the GF system with measurements using respiration chambers (RC) or
30 sulphur hexafluoride tracer (SF_6). It was hypothesised that estimates of methane emission for
31 individual animals and treatments would be similar for GF compared to RC or SF_6
32 techniques. In experiment 1, maize or grass silage-based diets were fed to four growing
33 Holstein heifers, whilst for experiment 2, four different heifers were fed four haylage
34 treatments. Both experiments were a 4 x 4 Latin square design with 33 d periods. GreenFeed
35 measurements of methane emission were obtained over 7 d (days 22-28) and compared to
36 subsequent RC measurements over 4 d (days 29-33). For experiment 3, 12 growing heifers
37 rotationally grazed three swards for 26 d, with simultaneous GF and SF_6 measurements over
38 two 4 d measurement periods (days 15-19 and days 22-26). Overall methane emissions (g/d
39 and g/kg dry matter intake [DMI]) measured using GF in experiments 1 (198 and 26.6,
40 respectively) and 2 (208 and 27.8, respectively) were similar to averages obtained using RC
41 (218 and 28.3, respectively for experiment 1; and 209 and 27.7, respectively, for experiment
42 2); but there was poor concordance between the two methods (0.1043 for experiments 1 and 2
43 combined). Overall, methane emissions measured using SF_6 were higher ($P < 0.001$) than GF
44 during grazing (186 vs. 164 g/d), but there was significant ($P < 0.01$) concordance between
45 the two methods (0.6017). There were fewer methane measurements by GF under grazing
46 conditions in experiment 3 (1.60/d) compared to indoor measurements in experiments 1
47 (2.11/d) and 2 (2.34/d). Significant treatment effects on methane emission measured using
48 RC and SF_6 were not evident for GF measurements, and the ranking for treatments and
49 individual animals differed using the GF system. We conclude that under our conditions of

50 use the GF system was unable to detect significant treatment and individual animal
51 differences in methane emissions that were identified using both RC and SF₆ techniques, in
52 part due to limited numbers and timing of measurements obtained. Our data suggest that
53 successful use of the GF system is reliant on the number and timing of measurements
54 obtained relative to diurnal patterns of methane emission.

55

56 *Keywords:* Dairy cattle, methane, respiration chamber, SF₆, GreenFeed

57

58 *Abbreviations:* CI, confidence interval; DM(I), dry matter (intake); GF, GreenFeed; LW, live
59 weight; LWG, LW gain; NDIR, non-dispersive infrared; RC, respiration chambers; RFID,
60 radio frequency identification; SF₆, sulphur hexafluoride tracer

61 **1. Introduction**

62 Accurate and robust measurements of methane emissions from individual animals are
63 required for national inventories and assessment of mitigation strategies. There are a number
64 of methods for determining methane emissions from ruminants, including respiration
65 chambers (RC) and sulphur hexafluoride tracer (SF_6) techniques. Precise measurements of
66 methane emission can be obtained by housing animals in RC, which allow direct
67 measurement of total methane emission. However, RC are relatively expensive, have a
68 limited throughput, and are disruptive to normal behaviour as animal by environment
69 interactions that occur within grassland ecosystems are prevented. Respiration chambers are
70 impractical for simulating grazing applications, and if the diet offered in the RC is fresh
71 forage, then diet selection is limited, and eating patterns are likely to be determined by the
72 feeding regime.

73 The SF_6 technique (Zimmerman, 1993; Johnson et al., 1994) can be used to make
74 estimations of eructated and expired methane emissions from animals which can select their
75 diet in a manner representative of farmed livestock (*e.g.* grazing). However, evaluations have
76 challenged the precision of the SF_6 technique for estimating methane emissions (Vlaming et
77 al., 2007; Pinares-Patiño and Clark, 2008; Pinares-Patiño et al., 2011), with greater between-
78 animal variation compared to RC (Hammond et al., 2009). The SF_6 technique has also
79 provided variable estimates of methane emission from animals on different herbage that
80 have not been corroborated by RC measurements (*e.g.* Hammond et al., 2011; Waghorn et al.,
81 2002; Sun et al., 2011 and 2012). Halter and collection canisters placed on the animal for
82 methane estimates can interfere with grazing (Pinares-Patiño et al., 2008), especially with
83 young animals, and a lower than predicted feed dry matter intake (DMI) will overestimate
84 methane yields (g/kg DMI). Rumen SF_6 boluses must also be administered, and frequent

85 animal handling is needed, all of which can be disruptive to normal behaviour, and is
86 relatively labour intensive.

87 In 2010, the commercial GreenFeed (GF) system (C-Lock Inc., Rapid City, South
88 Dakota, USA) was introduced as a static short-term measurement device that measures
89 methane emission from individual cattle, and uses head position sensors in combination with
90 decision rules to assess the validity of measures obtained. The animal is free to move about
91 and voluntarily enters a hood where a feed supplement (*i.e.* a reward for visiting the GF unit)
92 is delivered. Measurements of methane emission by GF are typically over short (3-7 min)
93 periods, several times within a day, over several days. The system is programmed using C-
94 Lock Inc. software to control timing of feed availability and thus, encourage animals to
95 distribute their voluntary GF visitation and hence methane measurements over a 24 h period
96 so that ultimately a 24 h individual methane emission profile can be extrapolated from several
97 days of short-term measurements. Cattle are typically not handled during GF operation and
98 one GF unit can be used for numerous animals, with manufacturer recommendations of 15-20
99 animals/unit when grazing and 20-25 animals/unit if housed in free stalls. Because the GF
100 system is relatively new, little is known about its operation, precision, accuracy, and the
101 extent to which animal interaction with GF affects methane measurements.

102 The objectives of the present study were addressed across three experiments that
103 included measurements of methane emission from individual growing dairy cattle using a
104 single GF unit. Our objectives were to compare measurements of methane emission by
105 growing dairy cattle obtained using GF with measurements using RC (experiments 1 and 2)
106 and SF₆ (experiment 3). It was hypothesised that estimates of methane emission for
107 individual animals and treatments would be similar for GF and RC or SF₆ techniques.

108

109 **2. Materials and methods**

110 Three experiments were used in this study whereby growing dairy cattle were fed a
111 variety of diets and methane emission was measured using GF, RC and/or SF₆ techniques.
112 Measurements in all experiments were individual DMI and methane production (g/d),
113 calculated methane yield (g/kg DMI), and frequency of GF visitation (*i.e.* methane
114 measurement frequency). All procedures used were approved and monitored under the UK
115 Home Office Animals (Scientific Procedures) Act 1986.

116

117 2.1 Experiments

118 2.1.1 Experiment 1

119 Four Holstein Friesian dairy heifers aged 14 months with an initial live weight (LW)
120 of 317 ± 20 kg were fed once daily (10:00 h) either maize or grass silage diets supplemented
121 with or without an extruded linseed product (Lintec; 26% fat) at 6% of ration DM ($n = 4$
122 animals/treatment).

123 Experiment 1 was a 4 x 4 Latin square design with each period 33 d in duration,
124 commencing with 21 d adaptation where access to GF was allowed, and GF data used for
125 analysis was obtained during 7 d (days 22-28), after which animals were confined to RC for
126 measurement of methane emission over 4 d (days 29-33).

127 Feed was offered to achieve target daily LW gains (LWG) of 0.75 kg. Feed intakes
128 were measured on a daily basis using an electronic Calan Broadbent individual feeding
129 system (American Calan Inc., Northwood, New Hampshire, USA) with feed refusals
130 collected once daily before morning feeding. Animals were loose-housed and bedded on
131 wood shavings with rubber mats and had *ad libitum* access to water.

132 2.1.2 Experiment 2

133 Four Holstein Friesian dairy heifers, aged 14 months with an initial LW of 339 ± 16
134 kg were fed twice daily (10:00 and 16:00 h in equal amounts), one of four conserved forage

135 (haylage) treatments of ryegrass, clover, trefoil and flowers ($n = 4$ animals/treatment). Further
136 details of these treatments are given in Hammond et al. (2014). Similar to experiment 1,
137 experiment 2 was a 4 x 4 Latin Square design with 33 d periods, with animals fed and housed
138 in a similar manner as detailed for experiment 1.

139 *2.1.3 Experiment 3*

140 Twelve Holstein Friesian dairy heifers aged eight months, with a starting LW of 230
141 ± 6 kg, grazed the same treatments used to make haylage in experiment 2 (ryegrass, clover
142 and flowers; $n = 12$ animals/treatment; Hammond et al., 2014). Heifers rotationally strip
143 grazed each sward treatment for 26 d in a sequence of flowers, clover, then ryegrass. Each 26
144 d period commenced with 14 d adaptation where GF access was allowed, with simultaneous
145 GF and SF₆ data obtained over two 4-d measurement periods (days 15-19 and days 22-26).

146 Dry matter intake was estimated using a rising plate meter (Farmworks Precision
147 Farming Systems, Feilding, New Zealand) by taking 20 sward height readings before and
148 after each days grazing period. Sward DM yield estimations were calibrated every second day
149 by taking 5 x 0.5 m² quadrat cuts of the sward at a target post-grazing height of 6 cm and
150 oven drying (100°C) the sample to give sward DM yield per m² which was applied to each
151 sward height measurement.

152

153 *2.2 Methane measurements*

154 *2.2.1 GreenFeed*

155 The GF system measured methane emission using sensors that identified the animal
156 and its head position within a sampling hood, air flow, and methane and carbon dioxide
157 concentrations in exhaust air. GreenFeed operation was initiated when the animal placed its
158 head inside the hood. A radio frequency identification (RFID) reader identified the animal's
159 ear tag and GF sampling was activated when the animals head (located by an infrared sensor)

160 was in the correct location within the hood, and it was deemed that sufficient time had
161 elapsed since the previous methane measurement for that animal.

162 Animal head position was critical for successful measurements as the animal is free to
163 move its head in and out of the hood and thus only data captured with uninterrupted
164 measurements was retained for statistical analysis. Position of the animals head within the
165 hood was monitored using sensors to ensure complete breath collection. Adequate animal
166 head position resulted in the dispensing of feed pellets which were used for enticement and
167 encouraged the animal to maintain a suitable head position for accurate measurements.
168 Pellets were dispensed from a hopper above the GF using a computer controlled rotating cup
169 dispenser.

170 Animals were able to use the GF unit at any time, provided it was not in use by
171 another animal, however, this did not necessarily generate a measurement of methane. A
172 'visit' is defined as a visit that results in a methane measurement. Thus, a 'visit' is only
173 considered when a certain time has elapsed between visits (as dictated by the user) and a food
174 reward is dropped, generating a methane measurement for that animal.

175 The concentration of the gas emitted by the animal was calculated using background
176 gas concentration, the differential concentration of gas during the animal's time in the GF
177 hood, and the calibration coefficient for concentration. The calibration coefficient was based
178 on nitrogen, carbon dioxide and methane gases used to calculate the response of the sensors.
179 The GF analysers were zeroed and calibrated weekly using zero baseline gas (oxygen-free
180 nitrogen) and a span gas mixture nitrogen containing 5000 ppm carbon dioxide and 1000 ppm
181 methane (BOC Ltd., Manchester, UK). This was to account for any drift in the calibration of
182 the analysers, which was found to be negligible. A known amount of propane or carbon
183 dioxide was released near where the animal's nose would be when feeding to check recovery

184 of expired gases when the physical location of the GF unit changed. There was no recovery
185 correction required in the current study.

186 To measure gas production (mass per unit of time) an extractor fan was used to draw
187 air past the animal's head into an exhaust pipe and airflow rate was measured. Airflow rate
188 was multiplied by the increase in gas concentration when the animals head was in the hood.
189 The duration the animals head was in the GF hood was recorded thus giving the time interval
190 for calculation of mass per unit of time. The concentrations of methane and carbon dioxide
191 gases were measured by non-dispersive infrared (NDIR) sensors, and an air filter was used to
192 filter and remove any fine particulate material from the air that was subsampled to the sensors
193 to prevent damage. The air filter was changed every two weeks. Data from GF was available
194 real-time using mobile phone communication through a web-based data management system
195 provided by C-Lock Inc.

196 For all experiments, the GF was programmed using C-Lock Inc. software to deliver a
197 maximum of five rotations of a feed dispensing cup, delivering approximately 55 g of pellet
198 (as fed) per rotation, with intervals of 45 sec between each rotation, so that 275 g of pellet
199 was delivered during each visit. A maximum of four visits per day (24 h) was allowed, with a
200 minimum of 4 h required between visits. Therefore, if an animal attempted to use the GF less
201 than 4 h from the previous visit pellets would not be dispensed. Commercial calf pellets
202 (Rearer18 Nuts, Wynnstay Group PLC, UK) were used for GF enticement and had a
203 chemical composition (g/kg DM) of ash, 85.1; oil, 46.5; acid detergent fibre, 174; neutral
204 detergent fibre, 289; starch, 259; water soluble carbohydrate, 91.3; nitrogen, 27.3, crude
205 protein, 171; and gross energy (MJ/kg), 18.1. In all experiments total daily feed allocations
206 included 1 kg of expected pellet DM provided by the GF unit.

207 The GF unit was set up indoors for experiments 1 and 2 at one end of animal housing,
208 with gates positioned to restrict access to one animal at a time. Barn ventilation was used to

209 maintain ambient concentrations of methane in background air. For experiment 3, the GF unit
210 was located outdoors under an awning at a point central to the experimental paddocks. The
211 GF was located next to the only available water trough to encourage visitation, and fences
212 and tracks were established to provide continuous access from grazed paddocks.

213 2.2.2 *Respiration chambers*

214 Details of the RC and measurements of methane emission are given by Reynolds et al.
215 (2001) and Cammell et al. (1986). For measurements of gaseous emissions, two open-circuit
216 RC were used (internal volume approximately 21 m³), with air-locks enabling access for
217 faecal and urine collection (Cammell et al., 1986). An integrative sample of ambient and RC
218 exhaust air was analysed at 4-min intervals, and every 4 h there was a switch to calibration
219 gases (oxygen-free nitrogen and nitrogen carrier with 20.5%, 3000 ppm, and 200 ppm
220 oxygen, carbon dioxide, and methane, respectively) to provide gas analyses with variation
221 coefficients of 5% or less.

222 2.2.3 *Sulphur hexafluoride*

223 Experiment 3 used the SF₆ technique, as detailed previously by Hammond et al.
224 (2014). Two weeks prior to experiment 3 commencing, heifers were each dosed by mouth
225 with a SF₆ permeation tube (supplied by AFBI, Hillsborough, Northern Ireland, UK) into the
226 rumen. The SF₆ gas release rates from the permeation tubes (5.176 ± 0.248 mg/d) were
227 measured prior to dosing by oven incubation at 39°C and weighing twice weekly for six
228 weeks. Daily methane emissions from heifers were estimated from analysis of air collected
229 from around the nose and mouth over a 24 h period into a pre-evacuated PVC canister which
230 was suspended under the neck. Based on recommendations given by Berndt et al. (2014), air
231 was sampled using a crimped stainless steel capillary 0.004" ID 10 cm tube, with a flow rate
232 between 0.45 to 0.55 ml/min. Canisters had a volume of approximately 2.3 L and a pre-
233 collection vacuum of 90 kPa. Canisters were changed once daily at the same time each

234 morning and were rejected if vacuum post-collection was > 75 or < 50 kPa. A background air
235 sample was also obtained daily from the paddock adjacent to that being grazed. Samples from
236 canisters were analysed daily in our laboratory using gas chromatography to determine
237 methane and SF₆ concentrations as described by Muñoz et al. (2012).

238

239 *2.3 Data and statistical analyses*

240 Data from GF and RC during periods 1 and 2 of experiment 1 were excluded from the
241 analyses because the methane concentration of the calibration gas used for the GF unit was
242 too low. Thus, comparisons for experiment 1 were restricted to periods 3 and 4 ($n = 8$) which
243 meant treatment effects were not tested due to the limited observations obtained with the
244 Latin Square design experiment. Each animal and period emissions data generated by GF was
245 averaged over 7 d, whereas RC data was averaged over 4 d, with data expressed on a per min
246 basis over 24 h and as a daily average (g/h and g/d).

247 For experiment 2, data from all four animals and periods were analysed statistically (n
248 = 16) using the Mixed Models Procedures of SAS (2011) for random effects of animal and
249 fixed effects of period and treatment. Like experiment 1, each animal and period emissions
250 data generated by GF was averaged over 7 d, whereas RC data was averaged over 4 d in
251 experiment 2, with data expressed on a per min basis over 24 h and as a daily average (g/h
252 and g/d).

253 Experiment 3 provided methane data for 12 heifers grazing three fresh forage
254 treatments for two 4-d methane measurement periods and two treatment periods (May to July
255 and August to October; Hammond et al., 2014). Analysis of methane emission data were
256 undertaken on daily averages across 4 d of measurements for both GF and SF₆ techniques
257 (obtained simultaneously) for individual animals during each measurement period. Twelve
258 heifers were used for the first grazing rotation of ryegrass, clover, and flowers, and also for

259 the second rotation of flowers. However, for the second rotation on ryegrass and clover, two
260 animals were removed because there was insufficient sward cover. Therefore, a total of 136
261 GF and SF₆ individual animal average emission rate observations were analysed using Mixed
262 Models Procedure of SAS (2011) for effects of forage treatment and treatment period (1 or
263 2), with 4-d measurement period within forage treatment period as a repeated effect within
264 heifers (Hammond et al., 2014). When significant effects occurred, means of forage mixtures
265 (clover and flowers) were differentiated from ryegrass control using Dunnett's adjusted mean
266 comparisons.

267 Differences in methane emission (g/h, g/d and g/kg DMI) between techniques (GF vs.
268 RC and GF vs. SF₆) across all experiments were tested using Lin's Concordance Correlation
269 Coefficient analysis (Lin et al., 1998) in GenStat (2010) and the Univariate Procedure of SAS
270 (2011) to determine if the difference between the two methods for each experiment was
271 different from zero.

272 Within each experiment, the Least Squares Mean option of the GLM procedure (SAS,
273 2011) was used to rank individual animals according to their methane emission (g/d and g/kg
274 DMI) for each measurement technique using animal as a fixed effect. In addition, the GLM
275 procedure was used to regress GF measurements against RC or SF₆ measurements (g/d).

276

277 **3. Results**

278 *3.1 General observations*

279 *3.1.1 Experiment 1*

280 As stated previously, experiment 1 included data from four animals with $n = 8$
281 observations (only two periods out of a possible four were used). Dry matter intake during
282 GF and RC measurements was similar (Table 1). Average methane production (g/d) and yield
283 (g/kg DMI), determined using either GF or RC techniques, was similar for individual animals

284 (198 vs. 215 g/d, and 26.6 vs. 28.3 g/kg DMI, for GF vs. RC techniques, respectively) (Table
285 1). Individual animals had a similar methane output regardless of measurement technique
286 used, however methane data (g/d and g/kg DMI) generated by GF and RC techniques ranked
287 heifers differently in numerical order from high to low methane output (data not shown).

288 *3.1.2 Experiment 2*

289 There were four heifers used in experiment 2 with 16 observations (all four periods
290 included). Heifers had a similar DMI during GF and RC measurements (Table 1). Average
291 daily methane production (g/d) and yield (g/kg DMI) did not differ with measurement
292 technique for individual animals (208 vs. 209 g/d, and 27.8 vs. 27.7 g/kg DMI, for GF vs. RC
293 techniques, respectively; Table 1). For both GF and RC methods, animals were significantly
294 ($P = 0.05$) different to each other in their methane production but not methane yield. Both GF
295 and RC techniques ranked animals in numerical order, from low to high, the same for
296 methane production, but not for methane yield (data not shown).

297 *3.1.3 Experiment 3*

298 Experiment 3 used 12 heifers and had 136 observations. Approximately 88% of SF₆
299 canisters were accepted (478 measurements out of a possible 544), with 12% of
300 measurements unsuccessful due to sampling tube blockages, broken collection tubes, or
301 displacement of canisters from the heifer. Both GF and SF₆ techniques were used
302 simultaneously so DMI was the same with measurement technique. Daily methane
303 production determined by GF for individual heifers was lower ($P < 0.001$) than SF₆ (164 vs.
304 186 g/d, respectively; Table 3). For both GF and SF₆ methods, heifers were significantly ($P =$
305 0.05) different to each other in their methane production (g/d), and the ranking of animals,
306 from low to high methane production, was different for the two techniques.

307

308 *Insert Table 1 here*

309

310 *3.2 Technique comparisons*311 *3.2.1 GreenFeed vs. respiration chamber*

312 Combining data from experiments 1 and 2, Lin's Concordance Correlation
313 Coefficient between GF and RC, when used to measure methane production and yield of
314 individual heifers, was 0.1043 and 0.058, respectively, with a non-significant ($P > 0.50$)
315 association between the two techniques, based on the 95% confidence interval (CI) (Fig 1).
316 There were diurnal patterns of methane erucation over a 24 h period for animals in both
317 experiments 1 and 2, measured using GF and in RC (Fig 2). Emissions ranged from about 4
318 g/h immediately before their morning feeding to a maximum of about 15 g/h after feeding, on
319 both silage and haylage diets. The increase in methane production after 10:00 h in experiment
320 1 relates to once daily feeding, whereas the increases just after 10:00 and 16:00 h represent
321 the twice daily feeding regime. Based on all methane measurements, compared to the GF
322 data, there was less variability with the RC emission measurements (g/d) from both
323 experiments, in part because measurements for GF were much less frequent and fewer in
324 number than for RC (Fig 2).

325

326 *Insert Fig 1 here*327 *Insert Fig 2 here*

328

329 *3.2.2. GreenFeed vs. sulphur hexafluoride*

330 Lin's Concordance Correlation Coefficient between GF and SF₆ techniques, used to
331 measure methane production from individual heifers of experiment 3, was 0.602, with a
332 significant ($P < 0.01$) association between the two techniques, based on the 95% CI (Fig 3).

333

334 *Insert Fig 3 here*

335

336 *3.3 GreenFeed for detecting dietary treatment effects*

337 In experiment 2, DMI and methane production during RC measurements was affected
338 by haylage type ($P = 0.045$ and $P = 0.025$, respectively), but this was not evident for GF
339 measurements (Table 2). When methane was expressed in terms of DMI (yield, g/kg DMI),
340 RC detected significant differences ($P = 0.020$) between haylages, but GF did not. There was
341 no consistency in the relative difference between measurement techniques with dietary
342 treatment. Relative to RC, GF underestimated 15% of methane yield when heifers were fed a
343 ryegrass diet, compared to an overestimation of 12% for heifers on a flower diet (Table 2).

344 For heifers of experiment 3, methane production (g/d) differed significantly with both
345 GF and SF₆ techniques ($P = 0.019$ and $P < 0.001$, respectively) for all three forage treatments.
346 However, the ranking of mean estimates for the different forages differed with technique
347 (Table 2). When methane was expressed in terms of DMI (methane yield), noting that the
348 techniques estimated methane simultaneously, the ranking of treatment means was not the
349 same for GF ($P = 0.080$; flowers > clover = ryegrass) and SF₆ techniques ($P = 0.002$; clover =
350 ryegrass > flowers). For two out of three dietary treatments fed, GF underestimated methane
351 yield relative to SF₆ by up to 18% (Table 2).

352 *Insert Table 2 here*

353

354 *3.4 GreenFeed visitation*

355 During the 14 d of GF measurements in experiment 1, there were a total of 118 visits
356 to the GF unit, averaging 2.11 visits/d. For the 28 d measurement period in experiment 2,
357 total GF visitation was 262, averaging 2.34 visits/d. During the 48 d of measurements for
358 experiment 3, heifers visited the GF unit 880 times, averaging 1.60 visits/d (Table 3). The

359 average duration (min:sec) of GF measurements for experiments 1, 2, and 3 were 04:44,
360 04:43, and 04:58, respectively.

361 Figure 4 shows the pattern of visits to the GF, according to hour of the day. For all
362 experiments, animals frequented the GF most often between 07:00 and 08:00 h, and between
363 13:00 and 14:00 h, with fewer visits in early morning hours (between 01:00 and 06:00 h).
364 GreenFeed measurements were prevented if another animal was already using the unit, when
365 animals were yarded for other experimental activities such as SF₆ canister changes, and
366 during the allocation of new grazing. The type of diet offered affected GF visitation for
367 experiment 3, but not experiment 2 (Table 2). Heifers in experiment 3 made fewer (P <
368 0.001) visits overall to the GF when on the ryegrass (219) and clover (229) paddocks,
369 compared to the flower (432) paddock.

370

371 *Insert Table 3 here*

372 *Insert Fig 4 here*

373

374 **4. Discussion**

375 *4.1 Comparison of measurement techniques*

376 *4.1.1 GreenFeed vs. respiration chamber*

377 Based on the concordance analysis for methane emission from heifers of experiments
378 1 and 2, GF and RC techniques had a poor agreement, yet average methane emission overall
379 was similar for the two techniques. It is difficult to interpret these conflicting results;
380 however the large amount of variation about the line of equality (Fig 1) is a likely explanation
381 for overall methane means being similar between techniques (Table 1) but having low
382 concordance correlation. The lack of concordance between methods is in part attributable to
383 the relatively small number of short-term measurements obtained by GF on each day of

384 measurement. The concept behind the GF system is that although it is unknown what an
385 animal is eructating when not visiting the GF, the accumulation of data over 24 h can provide
386 a representative pattern (Fig 2). Thus, the GF technique relies on the animal visiting the unit
387 at different times during the day to characterise the daily pattern of methane emission over a
388 number of days. In contrast, RC measurements in this study were based on integrated
389 measurements every 4 min over four consecutive days.

390 The inability of GF to detect changes in methane production due to treatment or
391 animal effects compared to RC (and SF₆) is not unexpected given the methodology the
392 technique employs. Enteric methane production from ruminants typically exhibits a diurnal
393 pattern related to feeding and meal consumption, with methane emission rate varying by as
394 much as five-fold over the course of a day (Crompton et al., 2010). Peak enteric methane
395 production occurs approximately 120 and 60 min after the morning and afternoon feeds,
396 respectively, for a lactating dairy cow fed *ad libitum* twice daily (Crompton et al., 2010).
397 Frequent or continuous measurements over a 24 h period using RC or SF₆ account for any
398 diurnal variation in methane production, but intermittent short-term measurements may vary
399 significantly depending on when those measurements are taken during the day.

400 There was a greater range in absolute emissions for both measurement techniques
401 with experiment 2 data compared to experiment 1 that was associated with greater differences
402 in DMI and methane production. However, when emissions were expressed per unit of feed
403 intake (g/kg DMI), relationships between GF and RC measurements were still weak and
404 variation was large, especially for the GF measurements. The variable relationship suggests
405 that the differences in methane emission due to treatments and animal variation measured by
406 the RC are not correlated with differences measured by GF. In other words, ranking of the
407 animals according to methane production and yield differed between the two techniques,
408 despite substantial differences being observed. The absence of a significant correlation

409 between GF and RC measurements for individual animal observations (Fig 1) casts doubt on
410 the capability of GF to distinguish (and rank) individual animals under the conditions with
411 which GF was used in our experiments. With the exception of daily methane production in
412 experiment 2, GF and RC ranked heifers differently in their methane emission. Daily mean
413 methane production varied from about 160 to 270 g/d measured in RC, and although GF also
414 recorded a similar range, the range was for different heifers on different diets (data not
415 shown).

416 It is possible that the algorithms used by the GF system for the calculation of methane
417 output, or the timing of visits relative to daily patterns of methane emission, may account for
418 the discrepancies observed between GF and RC data. GreenFeed calculations are based on
419 differences in the concentration of the air exhaled and eructed by the animal, less background
420 air concentrations measured pre- and post-feeding. The GF is able to differentiate emissions
421 of methane in exhaled air above background, so exhaled air is included in the emission
422 calculation. The calculations are reliant on erucation events taking place within the
423 measurement period, and the algorithms may need to be modified to increase accuracy and
424 reduce variation. For the animals and diets used in our study, more GF measurements were
425 needed over a longer period, and at more frequent intervals, to better represent the diurnal
426 pattern of methane emissions over 24 h. Increased animal visitation to the GF may require
427 longer periods of measurement (more days), more visits per day (and thus greater feed
428 consumption), or the use of an alternative ‘bait’ (Hegarty, 2013). In addition to this, it has
429 been estimated that 1-2% of methane is voided as flatus (Murray et al., 1976), and it will
430 contribute to methane emissions (Ellis et al., 2008) measured in RC. These considerations for
431 GF measurements are also pertinent to other on-farm breath analysis techniques (*e.g.*
432 Garnsworthy et al., 2012).

433

434 *4.1.2 GreenFeed vs. sulphur hexafluoride*

435 GreenFeed and SF₆ techniques had moderate concordance (agreement), in part due to
436 the greater number of observations compared. However, overall methane emissions
437 determined using GF were significantly lower than those measured using the SF₆ technique.
438 Differences between GF and SF₆ techniques are likely due to the duration of methane
439 measurements obtained for each animal. The SF₆ technique is based on integrative sampling
440 with a sampling duration of nearly 1440 min/d (100% of 24 h). In comparison, the GF unit is
441 designed to take intermittent samples, and based on the average of the three experiments
442 presented here, sampling duration (5 min/visit at 2 visits/animal/d) was only 10 min/d (0.7%
443 of 24 h).

444 All tracers have weaknesses (Shipley and Clark, 1972) and the variation associated
445 with SF₆ estimates may be in part a consequence of factors affecting the technique itself
446 (Deighton et al., 2013, 2014a and 2014b), or alternatively the variation might be real. Recent
447 work has found that successful use of the SF₆ technique to detect differences in enteric
448 methane emissions due to diets or between animal species may be confounded by diet or
449 genetic effects on body temperature (Deighton et al., 2014b). In order to accurately determine
450 methane emissions, it is necessary that gases are collected continuously at a constant rate for
451 24 h, however; it has been recently shown by Deighton et al. (2014a) that capillary tubes are
452 unsuitable for use as flow restrictors to achieve this, causing a bias of up to 15.6% in
453 calculated methane emissions. Deighton et al. (2014a) has since proposed a ‘modified SF₆
454 technique’ which incorporates orifice plate flow restrictors for 24 h sample collection and has
455 found technique error associated with SF₆ release, sample collection and analysis to be
456 reduced.

457

458 *4.2 GreenFeed for detecting dietary treatment effects*

459 Although all three techniques measured significant treatment effects on methane
460 emissions, the ranking of these effects differed with measurement technique. Critically, both
461 the RC and SF₆ techniques found methane yield (g/kg DMI) to be the lowest in both
462 experiments 2 and 3 for animals fed flowers compared to the other dietary treatments
463 (Hammond et al., 2014). GreenFeed on the other hand, was unable to detect treatment effects
464 on methane yield in experiment 2, and ranked the treatments differently to SF₆ for experiment
465 3. This in part reflects the variability of GF measurements attributable to the timing and
466 limited number of short-term measurements obtained in the present experiments.

467

468 *4.3 GreenFeed visitation*

469 Although animals had few problems adapting to the GF and used it willingly, visits
470 were less frequent than permitted, particularly for grazing animals in experiment 3. The lack
471 of GF visits from animals both while out grazing and during early morning, may have
472 negatively biased methane production measured by GF (Fig 3). The low frequency of visits
473 between 09:00 and 13:00 h (Fig 4) is likely to be when peak methane emissions occur in a
474 once daily feeding system, as can be seen from the rise in methane production (g/h) in Fig 4.
475 Thus, the infrequent daily measurements made by the GF system in experiment 1 is a likely
476 explanation for numerically lower methane emissions from the GF compared to RC. In
477 experiment 2, the GF pattern of visitation was better distributed over the course of the day,
478 although a weaker relationship occurred between the two techniques (Fig 2).

479 In all experiments, fewer visits occurred in the early morning hours, and a lack of
480 methane data over this period may have affected the average estimate of daily emissions.
481 Every morning, heifers of experiment 3 were given a new allocation of feed at about 10:30 h
482 after SF₆ canisters were replaced. The allocation of new pasture is likely to have been
483 responsible for the drop in GF visits between 09:00 and 13:00 h. During this period of time,

484 methane emissions would have been at their highest, which must partly account for the 13%
485 greater daily methane emissions determined by the SF₆ technique, compared to the GF
486 system. The lower visitation to the GF by grazing heifers is cause for concern, especially as
487 the GF system relies on having enough daily measurements over the course of an ‘average’
488 day to estimate daily emissions. Further evaluations of the GF system should determine the
489 number of days and measurements per day required for GF to provide accurate and precise
490 measures of methane emissions.

491 It would appear that the number of visits to the GF is influenced by dietary treatment
492 (and possibly level of feed intake), with more visits made when heifers were grazing flowers,
493 compared to ryegrass and clover (experiment 3). For experiment 3, this may have been
494 attributable to the location of the GF relative to the paddocks, as well as the DM and nutrient
495 content of the swards grazed. It would appear that less favourable diets may contribute to
496 increased GF visitation when a favourable ‘treatment’ *i.e.* pelleted concentrates, is rewarded,
497 and this may have consequences for methane estimates on different treatments (different
498 numbers of samples and sampling times for each treatment). This is a concern for nutrition
499 experiments that investigate effects of diet composition on methane emission if diet
500 comparisons are affected by varying amounts of feed reward provided by GF.

501 One concern with the use of the GF under our conditions is the temporal distribution
502 of GF visitation and the potential for bias in methane emission measurements by the GF
503 system. This is because unlike both RC and SF₆ techniques the methane measurement
504 obtained from each individual animal by the GF system is voluntary and thus not completely
505 independent. The use of the GF unit by each individual animal within the group, and
506 therefore the temporal distribution of their methane measurement, is affected by their cohorts
507 and environmental circumstances. The inclusion of a given animal in the GF unit causes the
508 exclusion of all other animals within the group. Strictly speaking an individual animal is not

509 totally independent as an experimental unit when the GF system is used and therefore to
510 achieve completely independent experimental replication for individual animals housed as
511 groups in pens or paddocks multiple GF units may be required.

512

513 **5. Conclusions**

514 Overall, the GF system provided an estimate of methane emission by growing dairy
515 cattle that was not different from RC measurements, but significantly lower than for SF₆.
516 However, concordance analyses found no agreement between GF and RC, and only moderate
517 agreement with SF₆. We conclude that as used in our experiments, the GF system was unable
518 to detect significant treatment and individual animal differences in methane emissions that
519 were identified using both RC and SF₆ techniques. The successful use of the GF system is
520 reliant on the number and timing of measurements obtained relative to diurnal patterns of
521 methane emission. Therefore, animal and diet type, intake level and appetite (*e.g. ad libitum*
522 *vs. restrictive feeding*), total feed availability, accessibility of the GF unit relative to other
523 feeds and activities, as well as type, amount, and timing of feed used to elicit GF use all
524 affect GF visitation and thus measurements of methane emission using the GF system.
525 Multiple animals using a GF unit can alter the temporal distribution of measurements for
526 individual animals and this potential bias should also be considered in designing future
527 experiments.

528 Further evaluation of GF is needed to determine how best to deploy the system to
529 meet specific objectives, the number and timing of measurements required for specific
530 measurement conditions, as well as the capacity of the GF to detect significant changes in
531 methane emissions with individuals and treatments. We suggest that in addition to increased
532 frequency of daily GF visits future studies should include longer periods of measurement and
533 a greater number of animals per treatment than is required for RC studies.

534

535 **Acknowledgements**

536 This study was funded by Defra, the Scottish Government, DARD, and the Welsh
537 Government as part of the UK's Agricultural GHG Research Platform project
538 (www.ghgplatform.org.uk). Contributions from the technical staff of CEDAR in the daily
539 routine of experiments and care of cows, as well as assistance from C-Lock Inc. with regards
540 to GreenFeed operation and use, is gratefully acknowledged.

541

542 **References**

543 Berndt, A., Boland, T.M., Deighton, M.H., Gere, J.I., Grainger, C., Hegarty, R.S., Iwaasa,
544 A.D., Koolaard, J.P., Lassey, K.R., Luo, D., Martin, R.J., Martin, C., Moate, P.J., Molano,
545 G., Pinares-Patiño, C.S., Ribaux, B.E., Swainson, N.M., Waghorn, G.W., Williams, S.R.O.,
546 2014. Guidelines for use of sulphur hexafluoride (SF₆) tracer technique to measure enteric
547 methane emissions from ruminants. Pp 166. [M.G. Lambert, editor], New Zealand
548 Agricultural Greenhouse Gas Research Centre, New Zealand.

549

550 Cammell, S.B., Thomson, D.J., Beever, D.E., Haines, M.J., Dhanoa, M.S., Spooner, M.C.,
551 1986. The efficiency of energy utilization in growing cattle consuming fresh perennial
552 ryegrass (*Lolium perenne* cv. Melle) or white clover (*Trifolium repens* cv. Blanca). Br. J.
553 Nutr. 55, 669-680.

554

555 Crompton, L.A., Mills, J.A.N., Reynolds, C.K. and France, J., 2010. Fluctuations in methane
556 emission in response to feeding pattern in lactating dairy cows. In. Modelling Nutrient
557 Digestion and Utilization in Farm Animals, pp. 176-180 [D. Sauvant, J. Van Milgen, P.

558 Faverdin and N. Friggens, editors]. Wageningen, the Netherlands: Wageningen Academic
559 Publishers.

560

561 Deighton, M.H., O'Loughlin, B.M., Williams, S.R.O., Moate, P.J., Kennedy, E., Boland,
562 T.M., Eckard, R.J., 2013. Declining sulphur hexafluoride permeability of
563 polytetrafluoroethylene membranes causes overestimation of calculated ruminant methane
564 emissions using the tracer technique. *Anim. Feed Sci. Technol.* 183, 86-95.

565

566 Deighton, M.H., Williams, S.R.O., Hannah, M.C., Eckard, R.J., Boland, T.M., Wales, W.J.,
567 Moate, P.J., 2014a. A modified sulphur hexafluoride tracer technique enables accurate
568 determination of enteric methane emissions from ruminants. *Anim. Feed Sci. Technol.* 197,
569 47-63.

570

571 Deighton, M.H., Williams, S.R.O., Lassey, K.R., Hannah, M.C., Boland, T.M., Eckard, R.J.,
572 Moate, P.J., 2014b. Temperature, but not submersion or orientation, influences the rate of
573 sulphur hexafluoride release from permeation tubes used for estimation of ruminant methane
574 emissions. *Anim. Feed Sci. Technol.* 194, 71-80.

575

576 Ellis, J.L., Dijkstra, J., Kebreab, E., Bannink, A., Odongo, N.E., McBride, B.W., France, J.,
577 2008. Aspects of rumen microbiology central to mechanistic modelling of methane
578 production in cattle. *J. Agric. Sci.* 146, 213-233.

579

580 Garnsworthy, P.C., Craigon, J., Hernandez-Medrano, J.H., Saunders, N., 2012. On-farm
581 methane measurements during milking correlate with total methane production by individual
582 dairy cows. *J. Dairy Sci.* 95, 3166-3180.

583

584 Hammond, K.J., Hoskin, S.O., Burke, J.L., Waghorn, G.C., Koolaard, J.P., Muetzel, S., 2011.
585 Effects of feeding fresh white clover (*Trifolium repens*) or perennial ryegrass (*Lolium*
586 *perenne*) on enteric methane emissions from sheep. *Anim. Feed Sci. Technol.* 166-167, 398-
587 404.

588

589 Hammond, K.J., Humphries, D.J., Westbury, D.B., Thompson, A., Crompton, L.A., Kirton,
590 P., Green, C., Reynolds, C.K., 2014. The inclusion of forage mixtures in the diet of growing
591 dairy heifers: Impacts on digestion, energy utilisation, and methane emissions. *Agric. Eco.*
592 *Environ.* 197, 88-95.

593

594 Hammond, K.J., Muetzel, S., Waghorn, G.C., Pinares-Patiño, C.S., Burke, J.L., Hoskin, S.O.,
595 2009. The variation in methane emissions from sheep and cattle is not explained by the
596 chemical composition of ryegrass. *Proc. N. Z. Soc. Anim. Prod.* 69, 174-178.

597

598 Hegarty, R.S., 2013. Applicability of short-term emission measurements for on-farm
599 quantification of enteric methane. *Anim.* 7:s2, 401-408.

600

601 Johnson, K.A., Huyler, M., Westberg, H., Lamb, B., Zimmerman, P., 1994. Measurement of
602 methane emissions from ruminant livestock using a SF₆ tracer technique. *Environ. Sci.*
603 *Technol.* 28, 359-362.

604

605 Lin, L., Torbeck, L.D., 1998. Coefficient of accuracy and concordance correlation
606 coefficient: new statistics for methods comparison. *J. Pharm. Sci. Technol.* 52, 55-59.

607

- 608 Muñoz, C., Yan, T., Wills, D.A., Murray, S., Gordon, A.W., 2012. Comparison of the sulfur
609 hexafluoride tracer and respiration chamber techniques for estimating methane emissions and
610 correction for rectum methane output from dairy cows. *J. Dairy Sci.* 95, 3139-3148.
611
- 612 Murray, R.M., Bryant, A.M., Leng, R.A., 1976. Rates of production of methane in the rumen
613 and large intestine of sheep. *Brit. J. Nutr.* 36, 1-14.
614
- 615 Pinares-Patiño, C.S., Lassey, K.R., Martin, R.J., Molano, G., Fernandez, M., MacLean, S.,
616 Sandoval, E., Luo, D., Clark, H., 2011. Assessment of the sulphur hexafluoride (SF₆) tracer
617 technique using respiration chambers for estimation of methane emissions from sheep. *Anim.*
618 *Feed Sci. Technol.* 166-167, 201-209.
619
- 620 Pinares-Patiño, C.S., Clark, H., 2008. Reliability of the sulfur hexafluoride tracer technique
621 for methane emission measurement from individual animals: an overview. *Aust. J. Exp.*
622 *Agric.* 48, 223-229.
623
- 624 Pinares-Patiño, C.S., Machmuller, A., Molano, G., Smith, A., Vlaming, J.B., Clark, H., 2008.
625 The SF₆ tracer technique for measurements of methane emission from cattle - effect of tracer
626 permeation rate. *Can. J. Anim. Sci.* 88, 309-320.
627
- 628 Reynolds, C.K., Cammell, S.B., Humphries, D.J., Beever, D.E., Sutton, J.D., Newbold, J.R.,
629 2001. Effects of postrumen starch infusion on milk production and energy metabolism in
630 dairy cows. *J. Dairy Sci.* 84, 2250-2259.
631

- 632 Shipley, R.A., Clark, R.E., 1972. Tracer Methods for In Vivo Kinetics: Theory and
633 Applications. Academic Press, New York.
634
- 635 Sun, X.Z., Hoskin, S.O., Muetzel, S., Molano, G., Clark, H., 2011. Effect of chicory
636 (*Cichorium intybus*) and perennial ryegrass (*Lolium perenne*) on methane emissions *in vitro*
637 and from sheep. Anim. Feed Sci. Technol. 166-167, 391-397.
638
- 639 Sun, X.Z., Hoskin, S.O., Zhang, G.G., Molano, G., Muetzel, S., Pinares-Patiño, C.S., Clark,
640 H., Pacheco, D., 2012. Sheep fed forage chicory (*Cichorium intybus*) or perennial ryegrass
641 (*Lolium perenne*) have similar methane emissions. Anim. Feed Sci. Technol. 172, 217-225.
642
- 643 Vlaming, J.B., Brookes, I.M., Hoskin, S.O., Pinares-Patiño, C.S., Clark, H., 2007. The
644 possible influence of intra-ruminal sulphur hexafluoride release rates on calculated methane
645 emissions from cattle. Can. J. Anim. Sci. 87, 269-275.
646
- 647 Waghorn, G.C., Tavendale, M.H., Woodfield, D.R., 2002. Methanogenesis from forages fed
648 to sheep. Proc. N. Z. Grassland Assoc. 64, 167-171.
649
- 650 Zimmerman, P.R., 1993. System for measuring metabolic gas emissions from animals. U.S
651 Patent 5265618, U.S. Patent and Trademark Office, November 30, 1993.

652 **Table 1** Dry matter intake (DMI; kg/d), methane production (g/d), and methane yield (g/kg
 653 DMI) from growing dairy cattle within three different experiments using GreenFeed (GF),
 654 respiration chamber (RC) and sulphur hexafluoride tracer (SF₆) techniques.

| | Experiment 1 ^a | ± SD | Experiment 2 ^a | ± SD | Experiment 3 ^b | ± SD |
|---|------------------------------|------|------------------------------|------|------------------------------|------|
| DMI, kg/d | | | | | | |
| GF | 7.62 | 0.81 | 7.60 | 0.81 | 9.15 | 2.67 |
| RC ^a or SF ₆ ^b | 7.66 | 0.59 | 7.54 | 0.94 | 9.15 | 2.67 |
| <i>n</i> | 8 | | 16 | | 136 | |
| SEM | 0.132 | | 0.182 | | N/A | |
| P | 0.799 | | 0.747 | | N/A | |
| Methane, g/d | | | | | | |
| GF | 198 | 20.4 | 208 | 31.5 | 164 | 51.0 |
| RC ^a or SF ₆ ^b | 215 | 22.3 | 209 | 30.9 | 186 | 57.3 |
| <i>n</i> | 8 | | 16 | | 136 | |
| SEM | 9.230 | | 10.59 | | 2.900 | |
| P | 0.170 | | 0.940 | | < 0.001 | |
| Methane, g/kg DMI | | | | | | |
| GF | 26.6 | 2.80 | 27.8 | 5.62 | 18.8 | 6.94 |
| RC ^a or SF ₆ ^b | 28.3 | 3.01 | 27.7 | 1.81 | 21.5 | 7.60 |
| <i>n</i> | 8 | | 16 | | 136 | |
| SEM | 1.365 | | 1.459 | | 0.349 | |
| P | 0.255 | | 0.933 | | < 0.001 | |

^a Experiments 1 and 2 used RC for measurement of methane from dairy heifers.

^b Experiment 3 used SF₆ for estimate of methane from grazing dairy heifers.

^c DMI was measured using Calan gates for individual animals in experiments 1 and 2, however for experiment 3, DMI was estimated by pre- and post-herbage mass (hence same DMI for animals where both GF and SF₆ were used simultaneously).

655 **Table 2** The difference in methane emission between GreenFeed (GF), respiration chamber (RC), and sulphur hexafluoride tracer (SF₆)
 656 techniques with dairy heifers fed different dietary treatments.

658

| Experiments | <i>n</i> | Dry matter intake (DMI), kg/d ^c | | Methane production, g/d | | Methane yield, g/kg DMI | | | Relative difference between methods (%) |
|---------------------------------|----------|---|---|----------------------------|--|----------------------------|--|-------------------------|---|
| | | GF | RC ^a or SF ₆ ^b | GF | RC ^a or SF ₆ ^b | GF | RC ^a or SF ₆ ^b | Difference ^d | |
| Experiment 2^a | | | | | | | | | |
| Ryegrass | 4 | 8.28 | 8.13 | 196 | 230 | 24.1 | 28.4 | -4.32 | -15 |
| Clover | 4 | 6.86 ^d | 7.10 ^b | 202 | 200 ^c | 29.5 | 28.1 | 1.40 | 5 |
| Trefoil | 4 | 7.93 | 7.51 | 226 | 218 | 28.9 | 29.2 | -0.32 | -1 |
| Flowers | 4 | 7.34 | 7.42 ^c | 209 | 190 ^b | 28.8 | 25.7 ^c | 3.14 | 12 |
| SEM | | 0.377 | 0.255 | 17.33 | 8.890 | 3.013 | 0.662 | 2.844 | |
| P (haylage) | | 0.180 | 0.045 | 0.515 | 0.025 | 0.521 | 0.020 | 0.298 | |
| Experiment 3^b | | | | | | | | | |
| Ryegrass | 44 | 10.0 | 10.0 | 175 | 204 | 17.3 | 21.8 | -3.38 | -16 |
| Clover | 44 | 8.69 ^a | 8.69 ^a | 166 | 202 | 18.5 | 23.0 | -4.24 | -18 |
| Flowers | 48 | 8.78 ^b | 8.78 ^b | 159 ^b | 159 ^a | 19.7 ^c | 19.5 ^c | 0.48 | 2 |
| SEM | | 0.230 | 0.230 | 5.420 | 4.989 | 0.768 | 0.734 | 0.754 | |
| P (forage) | | 0.001 | 0.001 | 0.019 | < 0.001 | 0.080 | 0.002 | < 0.001 | |

For each parameter, different letters indicate significant differences from the ryegrass control according to Dunnetts test (^a P < 0.001, ^b P < 0.01, ^c P < 0.05, ^d P < 0.10).

^a Experiment 2 used RC for measurement of methane from dairy heifers

^b Experiment 3 used SF₆ for estimation of methane from grazing dairy heifers

^c DMI was measured using calan gates for individual animals in experiment 2, however for experiment 3, DMI was estimated by pre- and post-herbage mass (hence same DMI for animals in experiment 3 with measurement technique)


^d Difference is generated using GF value less corresponding RC or SF₆ value

659 **Table 3** Animal visitation to the GreenFeed (GF) unit across three different experiments

| 660 | Number of measurement days | Total number of GF visits (methane measurements) | Total number of GF visit (methane measurements) per animal per day |
|----------------------|-------------------------------|---|--|
| Experiment 1 | | | |
| | 14 | 120 | 2.11 |
| | | | SD 0.49 |
| | | | SEM 0.17 |
| Experiment 2 | | | |
| | | 76 | 2.71 |
| | | 60 | 2.14 |
| | | 68 | 2.43 |
| | | 58 | 2.07 |
| | 28 | 262 | 2.34 |
| | | | SD 1.05 |
| | | | SEM 0.26 |
| | | | P (haylage) 0.425 |
| Experiment 3* | | | |
| | | 219 | 1.24 |
| | | 229 | 1.30 ^d |
| | | 432 | 2.26 ^a |
| | 48 | 880 | 1.60 |
| | | | SD 1.09 |
| | | | SEM 0.07 |
| | | | P (forage) < 0.001 |


* For experiment 3, different letters indicate significant differences from ryegrass control according to Dunnetts test (^aP < 0.001, ^dP < 0.10)

661




662

663 **Fig 1.** Relationships between methane production (g/d), determined using GreenFeed (GF)
 664 and respiration chamber (RC) techniques, of individual dairy heifers in experiments 1 (open
 665 circle symbol; $n = 8$) and 2 (closed triangle symbol; $n = 16$). Solid line indicates $y = x$. Lin's
 666 Concordance value for both experiments combined = 0.1043.




667




668

669 **Fig 2.** Comparison of methane emission rate (g/h; minute average) measured using
 670 GreenFeed (GF; open square symbol) and respiration chambers (RC; closed diamond
 671 symbol) for all dairy heifers in experiments 1 ($n = 8$) and 2 ($n = 16$). There were 56 d GF and
 672 32 d RC measurements for experiment 1, and 112 d GF and 64 d RC for experiment 2.
 673 Arrows indicate time of feeding.



674

675 **Fig 3.** Relationship between methane production (g/d), determined using GreenFeed (GF)
 676 and sulphur hexafluoride tracer (SF₆) techniques, of individual dairy heifers in experiment 3
 677 ($n = 136$). Solid line indicates $y = x$. Lin's Concordance value = 0.6017.



678

679 **Fig 4.** Diurnal pattern of GreenFeed (GF) visitation (methane measurements) over 24 h, as a
 680 percentage of total visits, by growing dairy cattle of experiments 1 (120 GF visits/14 d), 2
 681 (262 GF visits/28 d) and 3 (880 GF visits/48 d).