



Metabolomics of fecal samples: a practical consideration

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Matysik, S., Le Roy, C. I., Liebisch, G. and Claus, S. P. (2016) Metabolomics of fecal samples: a practical consideration. Trends in Food Science and Technology, 57. pp. 244-255. ISSN 0924-2244 doi: <https://doi.org/10.1016/j.tifs.2016.05.011> Available at <http://centaur.reading.ac.uk/65839/>

It is advisable to refer to the publisher's version if you intend to cite from the work.

To link to this article DOI: <http://dx.doi.org/10.1016/j.tifs.2016.05.011>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online



Accepted Manuscript

Metabolomics of fecal samples: a practical consideration

Silke Matysik, Caroline Ivanne Le Roy, Gerhard Liebisch, Dr Sandrine Paule Claus



PII: S0924-2244(16)30198-4

DOI: [10.1016/j.tifs.2016.05.011](https://doi.org/10.1016/j.tifs.2016.05.011)

Reference: TIFS 1813

To appear in: *Trends in Food Science & Technology*

Received Date: 29 August 2015

Revised Date: 6 April 2016

Accepted Date: 16 May 2016

Please cite this article as: Matysik, S., Le Roy, C.I., Liebisch, G., Claus, S.P., Metabolomics of fecal samples: a practical consideration, *Trends in Food Science & Technology* (2016), doi: 10.1016/j.tifs.2016.05.011.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Metabolomics of fecal samples: a practical consideration**

2

3 Silke Matsyik¹, Caroline Ivanne Le Roy², Gerhard Liebisch¹, Sandrine Paule

4 Claus^{2*}

5

6 1 Institute of Clinical Chemistry and Laboratory Medicine, University of

7 Regensburg, Germany.

8 2 Department of Food and Nutritional Sciences, The University of Reading,

9 Whiteknights campus, PO Box 226, Reading RG6 6AP, UK.

10

11 * Corresponding author:

12 Dr Sandrine Claus, s.p.claus@reading.ac.uk, +44 (0) 118 378 8717

13

14 **Key words:** Metabolomics, Lipidomics, Feces, Methods

15

16

17 ABSTRACT

18

19 Background

20 Metabolic profiling is becoming increasingly popular to identify subtle metabolic
21 variations induced by diet alterations and to characterize the metabolic impact
22 of variations of the gut microbiota. In this context, fecal samples, that contain
23 unabsorbed metabolites, offer a direct access to the outcome of diet - gut
24 microbiota metabolic interactions. Hence, they are a useful addition to measure
25 the ensemble of endogenous and microbial metabolites, also referred to as the
26 hyperbolome.

27 Scope and Approach

28 Many reviews have focused on the metabolomics analysis of urine, plasma and
29 tissue biopsies; yet the analysis of fecal samples presents some challenges that
30 have received little attention. We propose here a short review of current
31 practices and some practical considerations when analyzing fecal material using
32 metabolic profiling of small polar molecules and lipidomics.

33 Key Findings and Conclusions

34 To allow for a complete coverage of the fecal metabolome, it is recommended to
35 use a combination of analytical techniques that will measure both hydrophilic
36 and hydrophobic metabolites. A clear set of guidelines to collect, prepare and
37 analyse fecal material is urgently needed.

38

39 **Highlights**

- 40 1. Untargeted metabolic profiling of fecal material is robustly achieved using
41 NMR-based metabolomics
- 42 2. Mass spectrometry is mostly used for targeted metabolic profiling of a
43 class of molecules for deep coverage and high sensitivity
- 44 3. Lipidomics profiles are extremely complex as they contain a mixture of
45 endogenous, diet-related and microbial lipids that may be of interest for
46 bacterial identification
47

48 **INTRODUCTION**

49 The gut microbiota is a highly metabolically active community of micro-
50 organisms inhabiting all niches along the intestine, that is now recognized as a
51 critical regulator of its host homeostasis.(O'Hara & Shanahan, 2006) It has been
52 estimated that the gut microbiota as a whole contains 100 times more genes
53 than human cells, hence its potential to be a key metabolic player. It is therefore
54 expected that modifying the gut microbial balance would induce a shift in the gut
55 metabolic environment that can in turn affect our own metabolism.

56 The gut microbiota composition varies considerably over a lifetime.(Yatsunenکو
57 et al., 2012) It takes approximately two years to a newborn to acquire a stable
58 GM population(Palmer, Bik, DiGiulio, & Relman, 2007) that will evolve through
59 life under the pressure of various factors such as, for instance, diet, lifestyle and
60 exposure to antibiotics, all commonly referred to as the 'exposome'. (Claesson et
61 al., 2011; Claus & Swann, 2013; Lozupone, Stombaugh, Gordon, & Jansson, 2012)
62 Later in life, a loss of microbial diversity is generally observed with senescence.
63 (Biagi, Candela, Fairweather-Tait, Franceschi, & Brigidi, 2011; Claesson et al.,
64 2011) As recently demonstrated, even perturbations of the circadian cycle have
65 been observed to affect the balance of the gut microbial community. (Mukherji,
66 Kobiita, Ye, & Chambon, 2013; Voigt et al., 2014)

67 Diet is the main factor influencing gut microbiota composition since it provides
68 microorganisms with their main organic carbon source.(Flint, Duncan, Scott, &
69 Louis, 2015) This connection was recently further evidenced by a study
70 demonstrating that a drastic change of diet such as switching from vegetarian to
71 carnivorous and inversely can profoundly reorient the GM ecosystem in a very
72 short period of time.(David et al., 2015) Thus, along genetic and other

73 environmental factors, diet strongly contributes to the unique character of every
74 individual's gut microbiota.

75 Bacteria have a high metabolic activity that generates a wide range of products
76 such as organic acids, alcohols and gas that may become available for the host or
77 other commensal bacteria for cross-feeding. This symbiotic activity shapes the
78 gut metabolic environment. Complex carbohydrates, that cannot be digested in
79 the upper gastrointestinal track and are a major source of carbon for colonic
80 bacteria.(Scott, Duncan, & Flint, 2008) Their fermentation results in the
81 production of short chain fatty acids (SCFAs: acetate, propionate, butyrate and
82 valerate) that play an important role in human health.(Besten, van Eunen, Groen,
83 & Venema, 2013) Other food components such as lipids and proteins can largely
84 impact the composition of the gut microbiota and its metabolic
85 activity.(Sonnenburg & Sonnenburg, 2014) Endogenous secretions such as the
86 bile acids contained in bile are important regulators of the gut microbiota.
87 Reciprocally, gut bacteria are known to extensively alter the structure of sterols
88 that leads to the formation of secondary and tertiary bile acids.(Sayin et al.,
89 2013) This is an example of a major gut microbiota-host interplay that
90 contributes to regulating the absorption of dietary lipids during digestion. Hence,
91 the diet-gut microbiota interaction plays a key role in the metabolic homeostasis
92 of its host. It is therefore of utmost importance to understand the biological
93 mechanisms that underlie this complex relationship. Systems biology
94 approaches that study a system as a whole (e.g. a micro-organism within a host,
95 the interactions occurring within a community of bacteria etc.) are increasingly
96 popular to decipher these interactions. In particular, metabolic profiling
97 techniques are tremendously useful to understand the metabolic pathways

98 regulated through the host-gut microbiota interaction. A variety of sample types
99 ranging from biofluids to tissue biopsies can be analyzed to capture the systemic
100 metabolic response to the exposome. Of particular interest, feces are easily
101 accessible and provide a non-invasive window to study the outcome of the diet-
102 gut microbiota-host interaction through the analysis of remaining unabsorbed
103 metabolites. Yet, the analysis of fecal samples for metabolic profiling has
104 received little attention. In this review, we will explore the dominant
105 technologies that are commonly applied to assess the fecal metabolome and
106 discuss about practical aspects that must be considered when dealing with this
107 material.

108

109 **NMR-based metabolomics of fecal samples**

110 Metabolic profiling, also referred to as metabolomics, is mostly achieved using
111 two analytical platforms: nuclear magnetic resonance (NMR) spectroscopy and
112 mass spectrometry (MS) to evaluate the metabolic composition of a chosen
113 biological matrix. These techniques allow the simultaneous measurement of a
114 wide range of metabolites in a sample, and when combined, offer a large
115 coverage of the metabolome (i.e. the set of metabolites in a sample).

116 Untargeted metabolic profiling by ^1H NMR spectroscopy measures all
117 metabolites with nonexchangeable protons that are present in a sample in a
118 relatively high concentration (in the micromolar range). Because it is highly
119 reproducible, is cost effective and usually requires only a few simple preparation
120 steps, NMR-based metabolic profiling has been widely applied to the analysis of
121 virtually all biological matrices, including feces. (Li et al., 2011; Martin et al.,
122 2010; Saric et al., 2008)

123 In humans, ^1H NMR-based metabolic profiling of fecal material has been
124 successfully applied to assess the impact of the composition of the gut
125 microbiota on the gut metabolic environment in the context of ulcerative colitis
126 (UC) and irritable bowel syndrome (IBS). (Le Gall et al., 2011) A similar approach
127 was applied to monitor the gut microbial metabolic activity in elderly (Claesson
128 et al., 2012). In this study, it was possible to cluster patients according to their
129 community setting (length of hospital care) based on fecal water profiling. A
130 recent study also demonstrated the possibility of evaluating independent
131 bacterial contributions at a species level to the gut metabolic environment using
132 this technique. (Le Roy et al., 2015). Applied to the monitoring of probiotic
133 consumption, it was possible to detect faecal metabolic modifications in
134 response to increased *Bifidobacterium* in the colon (Ndagijimana et al., 2009)

135 In animal models, profound reorientation of the gut microbial community
136 induced by antibiotics in mice was associated to modifications of fecal metabolic
137 profiles measured by the same technique. This was mainly associated to a
138 modification in the fecal content in amino acids and SCFAs. (Yap et al., 2008).
139 Similarly, Romick-Rosendale et al., (Romick-Rosendale et al., 2009) also showed
140 a modification of murine fecal metabolic profiles in response to antibiotic
141 treatments. NMR-based metabonomics analysis of fecal water also proved to be
142 able to differentiate age groups in mice. (Calvani et al., 2014) Finally NMR-based
143 metabonomics can be applied to nutrition (also referred to as
144 nutrimentabonomics) (Claus & Swann, 2013) to assess modification of the gut
145 metabolic environment in response to diet modulation. As an example, a study
146 by De Filippis et al., (De Filippis et al., 2015) used NMR-based metabonomics to
147 evaluate the impact of a Mediterranean diet on gut microbiota metabolic activity.

148 The study demonstrated that following a Mediterranean diet improved the
149 detection of SCFAs in fecal waters compared with a western diet. Similarly, fecal
150 metabolic modifications have been observed in response to food
151 supplementation investigated in *in vitro* gut models. (Frost et al., 2014).

152 Since fecal samples contain a complex mixture of metabolites, most NMR-based
153 metabolic profiling studies use a selective NOESY experiment with water
154 presaturation applied during recycle delay and mixing time to detect signals
155 caused by small molecular weight molecules as well as some lipids. This is often
156 referred to as the NOESYPR1D experiment [RD-90°-t₁-90°-t_M-90°-ACQ], where
157 RD is the recycle delay, t₁ a short interval of about 3 μs, t_M the mixing time of
158 approximately 100 ms and ACQ the FID acquisition period. Interestingly, a
159 number of studies investigating fecal samples have also used a CPMG (Carr-
160 Purcell-Meiboom-Gill) experiment, (Bjerrum et al., 2014; Li et al., 2011) which
161 uses t₂ filtering to reduce signal resonance from large macromolecules.
162 (Meiboom & Gill, 1958) However, this should be considered with care since the
163 number of loops and length of echo time that must be optimized for each CPMG
164 experiment would determine the signal/noise ratio, therefore preventing
165 absolute quantification. This is not an issue when only relative quantifications
166 are needed.

167 **Preparation of fecal material for metabolomics studies**

168

169 Recently, Deda et al. reviewed sample preparation methods for fecal samples for
170 metabolomic analysis.(Deda, Gika, Wilson, & Theodoridis, 2015) They provided a
171 comprehensive overview of fecal sample preparation for NMR, GC-MS and LC-MS
172 analysis including some critical aspects and specific requirement of the different
173 technologies. Therefore, we will not cover sample preparation in details here but

174 a summary of the protocols and methods used in previously published research
175 papers are presented in Table 1. However, it is noteworthy that this review
176 highlighted the lack of consensus about sample preparation for both metabolic
177 profiling technologies. For sample extraction, it seems that a dilution of 1 volume
178 of feces material for 2 volumes of PBS buffer is most commonly used. The buffer
179 is generally composed of a mixture of H₂O and D₂O (minimum 10 %) in various
180 amounts, with an adjusted pH of 7.4 and an internal standard to serve as NMR
181 reference. The most common internal reference is 3-(trimethylsilyl)-2,2,3,3-
182 propionate-d₄ (TSP). Deda et al. discuss that TSP signal intensity can be affected
183 by pH but so far, the only alternative is 2,2-dimethyl-2-silapentane-5-sulfo-
184 nate-d₆ (DSS). However, unlike TSP, DSS has multiple small resonances in addition to
185 the main resonance at 0 ppm that may interfere with other signals and therefore
186 it should be used at a very low concentration (0.01% would be recommended).
187 Homogenization of fecal material can be done directly in the NMR buffer that will
188 be used for NMR analysis, minimizing the number of sample processing steps
189 that may alter metabolic profiling. However, it also appears that a
190 water/methanol extraction tends to improve the overall recovery of fecal
191 metabolites. Nevertheless it was also argued in a publication by Jacobs et al.,
192 (Jacobs et al., 2007) that methanol extracts were less representative of the real
193 metabolic composition of the fecal water encounter in the colon and therefore of
194 the metabolite pool that interacts with the intestinal membrane.

195

196 **Mass spectrometry-based metabolomics of fecal samples**

197 Untargeted metabolic profiling using MS-based techniques is more sensitive than
198 NMR (in the nanomolar range) but often generates a large amount of unknown
199 signals and as a consequence, these techniques have been mostly used for

200 targeted metabolomics, where the method is optimized to the detection of a
201 specific class of samples. MS-based metabolic profiling is usually achieved using
202 either gas chromatography (GC-MS) or liquid chromatography (LC-MS).

203

204 *GC-MS based metabolomics*

205 The group of Sébédio presented two GC-MS methods to analyze the metabolome
206 of fecal water. In their first study they used an ethyl chloroformate derivatization
207 (Gao et al., 2009). In fecal water samples of healthy subjects 73 compounds were
208 identified and thereof 34 validated by reference standards. The second study
209 applied trimethylsilylation and identified 133 compounds (including amino
210 acids, carbohydrates short and long chain fatty acids and phenolics) in human
211 fecal water and the majority validated by authentic standards (Gao, Pujos-
212 Guillot, & Sébédio, 2010). In both studies several extraction conditions were
213 tested and the highest recovery of metabolites detected for neutral and basic pH
214 which was confirmed also by others (Deda et al., 2015). However, it has been
215 suggested that increasing pH from 6 to 7 may decrease the loss of volatile SCFA
216 during lyophilization (Gao, Pujos-Guillot, & Sébédio, 2010).

217 Phua et al. presented a GC-TOF-MS analysis of feces after freeze drying followed
218 by oximation and silylation (Phua, Koh, Cheah, Ho, & Chan, 2013). The authors
219 argue that removal of a variable content of water increased the reproducibility of
220 sample preparation. They identified 107 metabolites by matching with different
221 mass spectra libraries. However, only a few analytes were confirmed by
222 reference substances. This method was applied in detection of colorectal cancer
223 (Phua et al., 2014). Main markers for CRC differentiation include decreased level
224 of fructose, nicotinic acid and linoleic acid in CRC patients.

225 Using a similar methodology, Weir et al. analyzed the fecal metabolome in CRC
226 patients (Weir et al., 2013). Metabolite identification was based on database
227 matching. In agreement with Phua et al. they detected decreased linoleic acid in
228 CRC patients compared to healthy controls. Moreover, reduced level of oleic and
229 elaidic acids were found in CRC patients. In contrast to these fatty acids (FAs)
230 myristic acid and several amino acids were increased in CRC patients. It is
231 noteworthy that fatty acid identification does not differentiate double bond
232 positions.

233 A very efficient approach to analyze volatile organic compounds (VOC) of feces is
234 headspace solid-phase microextraction (SPME). Volatile metabolites are
235 adsorbed to polymer coated fibers which are analyzed by GC-MS. Typically the
236 analyte spectrum includes hydrocarbons, alcohols, aldehydes and organic acids
237 (primarily short chain) and their esters. Dixon et al. tested different fibers to get
238 a comprehensive coverage of VOCs (Dixon et al., 2011). Ahmed et al investigated
239 fecal VOCs in patients with irritable bowel syndrome, active Crohn's disease,
240 ulcerative colitis and healthy controls (Ahmed, Greenwood, de Lacy Costello,
241 Ratcliffe, & Probert, 2013). They identified 240 metabolites which allowed a
242 differentiation of patients with irritable bowel syndrome from patients with
243 inflammatory bowel diseases and healthy controls.

244

245 *LC-MS based metabolomics*

246 In contrast to GC-based methods, LC-MS usually does not require metabolite
247 derivatization but is restricted to analytes containing polar groups. Cao et al.
248 used UPLC-MS/TOF-MS to analyze the fecal metabolome in patients with liver
249 cirrhosis and hepatocellular carcinoma (HCC) (Cao et al., 2011). Fecal samples
250 were homogenized, centrifuged and injected after filtration. Metabolic features

251 were analyzed by multivariate data analysis. Chenodeoxycholic acid, 7-
252 ketolithocholic acid, urobilinogen, urobilin, lysophosphatidylcholine (LPC) 16:0
253 and 18:0 were found to discriminate between healthy controls and patients with
254 liver cirrhosis and HCC. Whereas LPC species were found in increased levels, the
255 other discriminatory markers were decreased in the patient samples. The
256 identities of these markers were confirmed by comparison of chromatographic
257 retention and product ion spectra with authentic standards.

258 A study by Jimenez-Girón investigated changes in the fecal metabolome related
259 to the consumption of red wine (Jiménez-Girón et al., 2015). Feces samples were
260 analyzed after mixing with saline solution, centrifugation and filtration by
261 UHPLC-TOF-MS. Mass features were subjected to statistical analysis and 37
262 metabolites were found to be related to wine intake. Metabolite identification
263 was performed by database searching and confirmation by authentic standards.
264 This way 14 metabolites could be identified tentatively, 6 mass features match to
265 standards (m/z and retention time).

266 The fecal metabolome of rats with chronic renal failure were analyzed by Zhao et
267 al. (Zhao, Cheng, Wei, Bai, & Lin, 2012). Homogenized fecal samples were
268 extracted with acetonitrile and analyzed by UPLC-Q-TOF-MS. Both polarities
269 including fragment ions were recorded and used for identification and validation
270 of mass features. Except an increase of adenine (used to induce kidney failure), 8
271 lipid metabolites were found decreased in rats with chronic renal failure.

272 An interesting approach to profile amine- and phenol-containing metabolites
273 was presented recently by Su and colleagues (Su et al., 2015). Dried fecal
274 samples are extracted sequentially with water and acetonitrile followed by
275 derivatization with dansyl chloride. As an internal standard an aliquot of a

276 pooled fecal extract was added which was derivatized with ^{13}C -labelled dansyl
277 chloride. Dansylation improves both LC separation efficiency and MS response of
278 the compounds. 6200 peaks were detected in 237 different samples and 67
279 metabolites (mainly amino acids) were identified based on mass and retention
280 time matching to a dansyl standard library.

281 In summary, both GC- and LC-based metabolomics studies are able to discover a
282 number of differentially regulated metabolic features in fecal samples in various
283 studies. However frequently, only a few of these features could be identified.
284 Moreover, several studies did not prove the identity by authentic standards but
285 only by matching of m/z values to database entries. In general, GC-MS based
286 studies identify an increased number of metabolites which may partly reflect the
287 superior chromatographic resolution and peak shape of GC compared to LC
288 methods. This allows a more reliable extraction and comparison of metabolic
289 features between different samples. Additionally, GC usually provides mass
290 spectra generated by ionization-induced fragmentation that are useful for
291 metabolite identification. A disadvantage of GC analyses is a more laborious
292 sample preparation including the need of derivatization as a potential source of
293 artifacts. Conversely, GC-MS clearly shows advantages compared to LC-MS in
294 terms of deleterious matrix effects. So it is generally accepted that quantification
295 of analytes by LC-MS requires internal standards, ideally stable isotope labeled
296 for each analyte. Therefore a major source of errors of LC-MS metabolic profiling
297 may be related to undiscovered matrix effects especially in heterogeneous
298 sample material like feces. Consequently, in order to provide solid data
299 metabolomics studies should validate their biomarkers by quantitative analysis

300 using authentic standards and internal standards especially when LC-MS is
301 applied at least in a representative cohort.

302

303 **Mass spectrometry-based targeted metabolic analysis**

304 In contrast to untargeted analysis, targeted analysis is confined to a limited set of
305 analytes. These methods are optimized for high analyte recoveries during
306 sample preparation and a reproducible, accurate quantification of the target
307 molecules. Most methods rely on internal standards (mandatory for LC-MS),
308 calibration lines and method validation shows their performance.

309

310 *SCFA*

311 An excellent example for targeted analysis of potential biomarkers is the
312 quantitation of short chain fatty acids (SCFA). A recent study by Han et al.
313 presented a LC-MS/MS method for SCFA quantification in human feces (Han, Lin,
314 Sequeira, & Borchers, 2014). SCFAs are converted to 3-nitrophenylhydrozones
315 (3NHPH) which are separated by reversed phase chromatography and detected
316 in negative ion mode. The method covers 10 straight- and branched chain SCFAs.
317 In an elegant way, this study introduced an internal standard for every analyte
318 by conversion of a standard mixture with a $^{13}\text{C}_6$ -labeled derivatization reagent.
319 This method showed a high reproducibility and analysis of human fecal samples
320 revealed an increased fraction of branched-chain SCFA in T2D patient compared
321 the other analyzed samples.

322 In contrast to LC-MS/MS analysis, GC-MS may rely on a fewer number of internal
323 standards since matrix effects are less pronounced. A study by Zheng et al.
324 quantified SCFAs and branched-chain amino acids (BCAAs) in feces and other
325 biological materials using D_3 -caproic acid as internal standard (Zheng et al.,
326 2013) after propyl chloroformate derivatization.

327

328 *Analysis of sterols and bile acids in faeces*

329 Sterols and bile acids belong to another class of analytes studied for a long time

330 in feces by targeted analysis. Cholesterol is an essential component of all

331 mammalian cell membranes and it is the precursor of steroid hormones and bile

332 acids. There has been a long interest in the intestinal metabolism of cholesterol

333 because the gut microbiome is highly involved in the balance between

334 absorption, excretion and metabolism. Between 34-57% of dietary cholesterol is

335 absorbed from the human intestine (Grundy et al., 1977). Fecal excretion of total

336 neutral sterols has been reported to range between 350-900 mg/day, of which

337 20% is cholesterol. There are several primary sources of fecal cholesterol:

338 unabsorbed from the diet, bile and intestinal epithelium. It is well-known that

339 the luminal cholesterol can be metabolized by the gut microbiota. This

340 cholesterol escaping intestinal absorption is degraded to coprostanol through

341 reduction of the double bond at C-5. (Figure 1) Coprostanone is also produced in

342 a lesser extent. (Eyssen et al., 1974; Lichtenstein, 1990; Gerard, 2013). In human

343 feces, cholesterol derivatives have been reported in the following proportions:

344 coprostanol 65%, cholesterol 20%, coprostanone 10 %. (Figure 1) Other minor

345 derivatives include cholestanone, cholestanol and epicoprostanol.

346 Many attempts have been made to isolate bacteria capable of reducing

347 cholesterol to coprostanol from human and animal faeces (Snog-Kjaer et al.,

348 1956; Crowther et al., 1973). Certain anaerobic bacteria from human faeces are

349 known to hydrogenate cholesterol in vitro. Cholesterol reduction by common

350 intestinal bacteria such as Bifidobacterium, Clostridium, and Bacteriodes has

351 also been reported and reviewed extensively (Gerard, 2013). In addition,

352 reference values have been generated for fecal excretion of cholesterol and

353 coprostanol (Benno et al., 2005) to differentiate between high-, low- and non-
354 converters.

355 Bile acids are derived from cholesterol and are produced by every class of
356 vertebrate animals and show substantial diversity across species (Hofmann et
357 al., 2010). In bile, bile acids rapidly form mixed micelles with secreted
358 cholesterol and phospholipids. Bile acids enter the intestine as di- and
359 trihydroxylated acyl conjugates, in mammals with the amino acids taurine and
360 glycine (Hofmann, 2009). In the intestinal lumen, conjugated bile acids directly
361 affect the microbiota because they exert antimicrobial properties besides
362 stimulating enterocytes to secrete undefined antimicrobial compounds
363 (Hofmann et al., 2006). Conversely, gut bacteria structurally alter bile acids
364 through deconjugation of taurine, glycine and sulfate moieties and hydroxylation
365 of the sterol backbone.

366 During digestion, bile acids facilitate lipid absorption by stabilizing lipid micelles.
367 Since bile acids have various degrees of hydrophobicity, and therefore various
368 stabilizing properties, the bile acid composition of the bile is an important factor
369 that regulates fat absorption. The enterohepatic cycle of bile acids is a key
370 regulator of hepatic bile acid *de novo* synthesis and cholesterol excretion. Indeed,
371 bile acids undergo extensive reabsorption by active and passive routes so that
372 95 % of secreted bile acids are reabsorbed daily. Since deconjugated bile acids
373 are less polar, passive diffusion is reduced and it has been shown that active
374 deconjugation by gut bacteria increases the overall excretion of bile acids in
375 feces, hence the role of bacteria in regulating fecal cholesterol loss (Claus et al.,
376 2011; 2008; Kellogg, Knight, & Wostmann, 1970; Sayin et al., 2013). In human, a
377 limited number of commensal bacteria are capable of removing the 7-hydroxyl

378 group from di- and trihydroxy bile acids and 7-deoxy species are formed. The
379 most common 7-deoxy bile acids are lithocholic and deoxycholic acid. Many
380 excellent reviews exist in the recent literature covering bile acid function,
381 signaling and therapeutic potential (Ridlon et al., 2006; Hofmann et al., 2008;
382 2009; Trauner et al., 2010; Hagey et al., 2013).

383

384 *Sample Preparation and Instrumental Strategies for Targeted Metabolomics*
385 *Analysis*

386 SCFAs are volatile and therefore freeze drying of stool samples may result in
387 lower recovery. Han et al. used homogenized samples and extracted SCFAs by
388 addition of 50% aqueous acetonitrile (Han, Lin, Sequeira, & Borchers, 2014).
389 These extracts were subjected directly to derivatization with 3-
390 nitrophenylhydrazine HCl. Zheng et al. used 0.005 M aqueous NaOH to
391 homogenize fecal samples at 4°C to protect the volatile SCFAs (Zheng et al.,
392 2013). The homogenates were derivatized with propyl chloroformate and
393 extracted with hexane extraction for GC-MS analysis.

394 Extensive overviews of state-of-the-art methods to analyze fecal steroids can be
395 found in the literature, e.g. (Story & Furumoto, 1991; Perwaiz et al., 2002;
396 Griffiths & Sjövall, 2010). The first and often referred methods for fecal steroid
397 and bile acid analysis were a combination of thin-layer chromatography and gas-
398 liquid chromatography published 1965 by Grundy, Miettinen and co-workers
399 (Grundy et al., 1965; Miettinen et al., 1965). Sample pretreatment included
400 homogenization, saponification and liquid/liquid extraction followed by thin-
401 layer chromatography and trimethylsilylation. Individual components are then
402 quantitatively measured by gas-liquid chromatography equipped with a flame
403 ionization detector. Bile acids underwent a methylation step before thin-layer

404 chromatography. This approach was applied in many studies for the analysis of
405 endogeneous and exogeneous (labelled) compounds.(Spritz et al., 1965; Grundy
406 et al., 1969). Evrard and Janssen suggested a method for bile acid analysis which
407 took advantage of a different extraction scheme and an alternate derivatization
408 (Evrard & Janssen, 1968). The method is based on heating in presence of acetic
409 acid and extraction with toluene. Quantification was done as methylketone
410 derivatives. In the 1980s methods with subfraction steps have been proposed to
411 isolate taurine and glycine conjugated and sulphated bile acids (Setchell et al.,
412 1983; Owen et al., 1984) using GC analysis. Nowadays, the most common
413 strategies to analyze sterol profiles including cholesterol, coprostanol and
414 coprostanone in faeces are based on GC-MS. The sample preparation protocol
415 includes repeated sampling for better representativeness, a dilution step,
416 hydrolyzation of esterified sterols, extraction with a mixture of hexane and
417 ethanol and derivatization (Lutjohann et al., 1993; Midtvedt et al., 1990; Andrasi
418 et al., 2011). For example, Korpela et al. suggested a very detailed protocol which
419 included a 72 h sampling of faeces, methanol-chloroform extraction, separation
420 of free and esterified sterols with a Lipidex-5000 column, saponification,
421 separation of hydroxylated and oxo-forms by a second column, and trimethylsilyl
422 derivatization followed by GCMS (Korpela, 1982). LC-MS and MS/MS have also
423 been widely exploited for bile acid analysis of human urine, plasma/serum, bile
424 and also feces (Perwaiz et al., 2002; Hago et al., 2009; Griffiths & Sjövall, 2010).
425 Quantification is best performed by addition of isotope labeled internal
426 standards. These should be added as early as possible in the analytical process so
427 as to account for analyte loss during sample preparation.

428 Bile acids can usually be extracted from fluids and tissues with ethanol, methanol
429 or acetonitrile. An ethanol fraction may be followed by extraction with a less
430 polar solvent, such as chloroform, to recover less polar bile acid derivatives (e.g.,
431 the fatty acid esters) and bile acids remaining in the lipophilic residue (Griffiths
432 et al., 2010). Batta et al. compared different methods to extract fecal bile acids
433 and sterols (Batta et al., 1999) to suggest a simplified method of extraction.
434 However, this is based on the assumption that fecal bile acids are unconjugated
435 (Setchell et al., 1983), which is not a valid assumption for many clinical
436 conditions, especially when subjects have been exposed to oral antibiotic
437 treatments that have affected the gut microbial ecosystem.

438 **Lipidomics**

439
440 Parallel to metabolomics, lipidomics emerged during the past decade as a
441 specialized discipline. Today virtually a full quantitative coverage of the lipidome
442 is possible by mass spectrometric methods (Wenk, 2010).

443 In contrast to global metabolomics, lipidomic analysis mostly relies on lipid
444 extracts prepared by extraction with apolar solvents like chloroform (Bligh &
445 Dyer, 1959) or MTBE (Matyash, et al., 2008). The complexity of these extracts is
446 greatly reduced compared to protein precipitates frequently applied for
447 untargeted metabolomics as polar analytes are removed.

448 Up to now there are only a few studies of the fecal lipidome. Gregory et al.
449 compared different extraction methods for lipidomics profiling by LC-HR-MS
450 (high resolution MS) (Gregory et al., 2013). Stools of preterm infants were
451 homogenized in water and lipophilic metabolites were extracted using either
452 dichloromethane or a MTBE/hexafluoroisopropanol mixture. Additionally, the
453 effect of pressure cycling on the extraction of lipid species was investigated.

454 Polar species like lyso-lipids showed a higher response after MTBE extraction
455 whereas increased responses were observed after dichloromethane extraction
456 for more hydrophobic species. The effect of pressure cycling does not show a
457 consistent increase of the lipid species response for both extraction methods.
458 Analysis by reversed phase chromatography couple to HR-MS was performed in
459 both positive and negative ion modes. 304 lipid species were identified by
460 unique monoisotopic m/z and retention time including 29
461 phosphatidylethanolamine (PE), 22 phosphatidylcholine (PC), 14
462 phosphatidylglycerol (PG), 88 triacylglycerol, 19 diacylglycerol and interestingly
463 50 ceramide species.

464 Recently, Davis and colleagues analyzed stools from mice fed a high fat or control
465 diet with or without induction of colitis-associated tumors (Davies et al., 2014).
466 Feces was homogenized by repeated freeze-thawing, minced to powder and
467 extracted with a modified Bligh and Dyer method. Samples were quantified by
468 shotgun lipidomics. However, mass spectra displayed very low intensities. The
469 selection of dioleoyl species of PE, PS, PI as internal standards has to be
470 considered as a potential source of error for quantification since these species
471 may be present in feces (PE and PI 36:2 were detected by Gregory et al. (Gregory
472 et al., 2013)). A general problem of this study and also a number of metabolomics
473 studies is the annotation of lipid species. Davis et al. showed detailed
474 annotations including even double bond positions of the fatty acyls. Such
475 structural differences usually may not be resolved by standard lipidomic or
476 metabolomics methods. Most methods determine the number of carbons and
477 number of double bonds with the acyl chains. Therefore, it is recommended to
478 annotate only structural details which are resolved by the analysis (Liebisch et

479 al., 2013). Moreover, common shorthand nomenclature provides a standard for
480 reporting and searching of lipid species including deposition in and retrieval
481 from databases.

482 Figure 2 presents the result of a lipidomic analysis of a fecal sample using a
483 method that has been applied to a variety of sample types such as plasma
484 (Siguener et al., 2014), lipoproteins (Scherer, Böttcher, & Liebisch, 2011), cells
485 (Leidl, Liebisch, Richter, & Schmitz, 2008), cell culture (Binder, Liebisch,
486 Langmann, & Schmitz, 2006) and tissues (Hebel et al., 2015). Fecal samples were
487 homogenized including bead-based grinding and subjected to liquid extraction
488 according to Bligh and Dyer (Bligh & Dyer, 1959). Analysis by flow injection ESI-
489 MS/MS using lipid class specific head group scans revealed huge differences
490 between the individual samples. For example, we could find substantial
491 concentrations of PG in some samples whereas other samples showed only
492 minor PG content (Figure 2). Similar observations were made for other lipid
493 classes. Additionally, the method of homogenization and extraction may greatly
494 influence lipid species recovery. So sample preparation may determine whether
495 analysis of lipids is confined to “easily accessible” lipids or includes also “hardly
496 extractable” bacterial lipid. Bacterial lipids could be of particular interest since
497 they are used as chemotaxonomic parameter to classify and identify bacterial
498 (Busse, Denner, & Lubitz, 1996). However, bacterial lipids also increase the
499 complexity of the fecal lipidome. So a number of additional fatty acids usually not
500 or only present at low concentrations in mammalian cells are found in bacteria
501 such as branched chain, cyclopropane and hydroxyl fatty acids. In summary, an
502 accurate analysis of the fecal lipidome poses a great challenge especially due to
503 its high complexity and high variability.

504 NMR has also been used for structural analyses and quantification of lipid
505 species including lipoproteins (Bou Khalil *et al.*, 2010; Sander *et al.*, 2013;
506 AlaKorpela *et al.*, 1996; Fernando *et al.*, 2010). ^{31}P -NMR is an attractive method
507 to investigate phospholipid molecules due to the fact that all phospholipids have
508 at least one phosphorous nucleus and this NMR active isotope occurs at a natural
509 abundance of 100% and has got a high gyromagnetic ratio. Therefore, high
510 resolution ^{31}P -NMR spectroscopy has successfully been employed to determine
511 the phospholipid composition of tissues and body fluids. Comprehensive reviews
512 with many applications can be found in the literature (Schiller & Arnold, 2002;
513 Schiller *et al.*, 2007). However, we could not identify any study that applied NMR
514 spectroscopy for lipid investigations in fecal samples.

515

516 **Final considerations**

517 It is noteworthy that although extremely useful and widely used, as illustrated by
518 this review, fecal materials reliably reflect the microbial activity of the distal
519 colon, which is moderately representative of the rest of the gastro-intestinal
520 tract. For example, although it is commonly accepted that SCFAs measured in
521 feces are an indicator of colonic production by gut bacteria, it is important to
522 remember that these metabolites are quickly absorbed by the intestinal
523 membrane and an increased detection in feces may also reflect a poorer
524 absorption. To overcome this issue, metabolic profiling techniques can be
525 applied on luminal content collected in various sections of the gastro-intestinal
526 track, although this implies a more invasive sample collection. Fecal water
527 profiling by NMR spectroscopy has been widely referenced for numerous animal
528 models such as rodents (Romick-Rosendale *et al.*, 2009) and horses (Escalona *et al.*,
529 *et al.*, 2014) as well as humans, (Jacobs *et al.*, 2008) providing a database for future

530 investigations. This is also extremely useful for similar evaluations performed on
531 fecal waters derived from *in vitro* batch cultures that simulate digestion by the
532 gut microbiota. Even if *in vitro* models are not a perfect representation of the
533 host-gut microbiota interplay, they provide a valuable overview of the microbial
534 activity in the gut in controlled conditions. For instance, metabolic profiling of
535 samples derived from such *in vitro* gut models has been recently applied to
536 compare the impact of diet on human and baboon gut microbial activity.(Frost,
537 Walton, Swann, & Psichas, 2014)

538 Another important consideration when analyzing feces metabolome, and
539 particularly fecal bile acids, is the irregularity of bile secretion and the
540 inhomogeneity of fecal samples as carefully studied by Setchell et al., who
541 demonstrated that this was strongly correlated with diet patterns (Setchell et al.,
542 1987). This is particularly relevant to human studies since humans tend to have
543 a defined regular feeding pattern with set time and number of meals per day
544 (unlike rodents that tend to feed all night, and from time to time during the day).
545 As a consequence, it is recommended to analyze aliquots of thoroughly
546 homogenized 4–5 day collections of feces.

547 Finally, since one of the main factors influencing the gut metabolic environment
548 is the microbiome, it is important to assess the microbial composition of the fecal
549 material when possible, using 16S rRNA sequencing or metagenomics when
550 possible, although these methods are not fully quantitative. Such metabolic
551 associations with gut bacteria should also be interpreted carefully as it is not
552 always possible to differentiate the host from the bacterial metabolic activity.
553 This is particularly true for amino acids that can be released by dead host cells or
554 be derived from protein digestion by host and bacterial enzymes. Typical

555 bacterial metabolites include SCFAs, some organic acids such as formate and by-
556 products of protein degradation such as indole. However, many fecal metabolites
557 derive from host-bacterial co-metabolism, which is typically the case for
558 secondary and tertiary bile acids.

559

560 To summarize, the measurement of the fecal metabolome is becoming
561 increasingly popular as it provides an easy estimate of the diet-gut microbiota-
562 host metabolic interaction. However, there is a need for establishing clear
563 guidelines for fecal sample collection, preparation and analysis for metabolic
564 profiling. Both NMR and MS-based metabolic profiling are complementary
565 techniques and none of them to date is able to holistically assess the fecal
566 metabolome. Instead, it is recommended that a combination of methods is used
567 to extend the metabolic coverage.

568

569 **Acknowledgements**

570

571 The authors would like to thank Dr Lesli Hingstrup Larsen who provided the
572 sample for lipidomics analysis. This work was supported by the EU FP7
573 programme of research MyNewGut (613979).

574

575

576 **References**

577

578 Ahmed, I., Greenwood, R., de Lacy Costello, B., Ratcliffe, N. M., & Probert, C. S.
579 (2013). An investigation of fecal volatile organic metabolites in irritable bowel
580 syndrome. *PLoS One*, *8*(3), e58204. doi:10.1371/journal.pone.0058204

581

582 Ala-Korpela, M., Posio, P., Mattila, S., Korhonen, A., & Williams, S. R. (1996).
583 Absolute quantification of phospholipid metabolites in brain-tissue extracts by 1
584 H NMR spectroscopy. *Journal of Magnetic Resonance, Series B*, *113*(2), 184-189.

585

586 András, N., Helenkár, A., Vasánits-Zsigrai, A., Zár, G., & Molnár-Perl, I. (2011).
587 The role of the acquisition methods in the analysis of natural and synthetic
588 steroids and cholic acids by gas chromatography-mass spectrometry. *Journal of*
589 *Chromatography*. *A*, *1218*(45), 8264–8272.
590 <http://doi.org/10.1016/j.chroma.2011.09.006>

591

592 Batta, A. K., Salen, G., Rapole, K. R., Batta, M., Batta, P., Alberts, D., & Earnest, D.
593 (1999). Highly simplified method for gas-liquid chromatographic quantitation of
594 bile acids and sterols in human stool. *Journal of Lipid Research*, *40*(6), 1148–
595 1154.

596

597 Benno, P., Midtvedt, K., Alam, M., Collinder, E., Norin, E., & Midtvedt, T. (2005).
598 Examination of intestinal conversion of cholesterol to coprostanol in 633 healthy
599 subjects reveals an age- and sex-dependent pattern. *Microbial Ecology in Health*
600 *and Disease*, *17*(4).

601

602 Besten, den, G., van Eunen, K., Groen, A. K., & Venema, K. (2013). The role of
603 short-chain fatty acids in the interplay between diet, gut microbiota, and host
604 energy metabolism. *Journal of Lipid Research*, 54(9), 2325-2340.

605

606 Bezabeh, T., Somorjai, R., Dolenko, B., Bryskina, N., Levin, B., Bernstein, C. N., &
607 Smith, I. C. P. (2009). Detecting colorectal cancer by ¹H magnetic resonance
608 spectroscopy of fecal extracts. *NMR in Biomedicine*, 22(6), 593-600.

609

610 Biagi, E., Candela, M., Fairweather-Tait, S., Franceschi, C., & Brigidi, P. (2012).
611 Ageing of the human metaorganism: the microbial counterpart. *Age*, 34(1), 247-
612 267.

613

614 Binder, M., Liebisch, G., Langmann, T., & Schmitz, G. (2006). Metabolic profiling of
615 glycerophospholipid synthesis in fibroblasts loaded with free cholesterol and
616 modified low density lipoproteins. *Journal of Biological Chemistry*, 281(31),
617 21869-21877.

618

619 Bjerrum, J. T., Wang, Y., Hao, F., Coskun, M., Ludwig, C., Günther, U., & Nielsen, O.
620 H. (2014). Metabonomics of human fecal extracts characterize ulcerative colitis,
621 Crohn's disease and healthy individuals. *Metabolomics*, 11(1), 122-133.
622 <http://doi.org/10.1007/s11306-014-0677-3>

623

624 Bligh, E. G., & Dyer, W. J. (1959). A rapid method of total lipid extraction and
625 purification. *Canadian journal of biochemistry and physiology*, 37(8), 911-917.

626

627 Bou Khalil, M., Hou, W., Zhou, H., Elisma, F., Swayne, L. A., Blanchard, A. P., ... &
628 Figeys, D. (2010). Lipidomics era: accomplishments and challenges. *Mass*
629 *spectrometry reviews*, 29(6), 877-929.

630

631 Busse, H. J., Denner, E. B., & Lubitz, W. (1996). Classification and identification of
632 bacteria: current approaches to an old problem. Overview of methods used in
633 bacterial systematics. *Journal of biotechnology*, 47(1), 3-38.

634

635 Calvani, R., Brasili, E., Praticò, G., Capuani, G., Tomassini, A., Marini, F., ... &
636 Miccheli, A. (2014). Fecal and urinary NMR-based metabolomics unveil an aging
637 signature in mice. *Experimental gerontology*, 49, 5-11.

638

639 Cao, G., Zhang, Y., Feng, J., Cai, H., Zhang, C., Ding, M., ... & Cai, B. (2011). A Rapid
640 and sensitive assay for determining the main components in processed *Fructus*
641 *corni* by UPLC-Q-TOF-MS. *Chromatographia*, 73(1-2), 135-141.

642

643 Claesson, M. J., Cusack, S., O'Sullivan, O., Greene-Diniz, R., de Weerd, H., Flannery,
644 E., et al. (2011). Composition, variability, and temporal stability of the intestinal
645 microbiota of the elderly. *Proceedings of the National Academy of Sciences*, 108
646 *Suppl 1*, 4586-4591. <http://doi.org/10.1073/pnas.1000097107>

647

648 Claesson, M. J., Jeffery, I. B., Conde, S., Power, S. E., O'Connor, E. M., Cusack, S., ... &
649 Fitzgerald, G. F. (2012). Gut microbiota composition correlates with diet and
650 health in the elderly. *Nature*, 488(7410), 178-184.

- 651 Claus, S. P., & Swann, J. R. (2013). Nutrimetabonomics: Applications for
652 Nutritional Sciences, with Specific Reference to Gut Microbial Interactions.
653 *Annual Review of Food Science and Technology*, 4, 381–399.
654 <http://doi.org/10.1146/annurev-food-030212-182612>
655
- 656 Claus, S. P., Ellero, S. L., Berger, B., Krause, L., Bruttin, A., Molina, J., et al. (2011).
657 Colonization-induced host-gut microbial metabolic interaction. *mBio*, 2(2).
658 <http://doi.org/10.1128/mBio.00271-10>
659
- 660 Claus, S. P., Tsang, T. M., Wang, Y., Cloarec, O., Skordi, E., Martin, F.-P., et al.
661 (2008). Systemic multicompartmental effects of the gut microbiome on mouse
662 metabolic phenotypes. *Molecular Systems Biology*, 4(1), 219.
663 <http://doi.org/10.1038/msb.2008.56>
664
- 665 Crowther, J. S., Drasar, B. S., Goddard, P., Hill, M. J., & Johnson, K. (1973). The
666 effect of a chemically defined diet on the faecal flora and faecal steroid
667 concentration. *Gut*, 14(10), 790–793.
668
- 669 David, L. A., Maurice, C. F., Carmody, R. N., Gootenberg, D. B., Button, J. E., Wolfe,
670 B. E., et al. (2015). Diet rapidly and reproducibly alters the human gut
671 microbiome. *Nature*, 505(7484), 559–563. <http://doi.org/10.1038/nature12820>
672
- 673 Davies, J. M., Hua, H. U., Dheer, R., Martinez, M., Bhattacharya, S. K., & Abreu, M. T.
674 (2014). Stool Phospholipid Signature is Altered by Diet and Tumors. *PloS one*,
675 9(12), e114352, doi:10.1371/journal.pone.0114352

676

677 Deda, O., Gika, H. G., Wilson, I. D., & Theodoridis, G. A. (2015). An overview of
678 fecal sample preparation for global metabolic profiling. *Journal of pharmaceutical
679 and biomedical analysis, 113*, 137-150.

680

681 De Filippis, F., Pellegrini, N., Vannini, L., Jeffery, I. B., La Storia, A., Laghi, L., ... &
682 Turrone, S. (2015). High-level adherence to a Mediterranean diet beneficially
683 impacts the gut microbiota and associated metabolome. *Gut*, gutjnl-2015.

684

685 Dixon, E., Clubb, C., Pittman, S., Ammann, L., Rasheed, Z., Kazmi, N., et al. (2011).
686 Solid-phase microextraction and the human fecal VOC metabolome. *PLoS One*,
687 6(4), e18471. doi:10.1371/journal.pone.0018471

688

689 Escalona, E. E., Leng, J., Dona, A. C., Merrifield, C. A., Holmes, E., Proudman, C. J., &
690 Swann, J. R. (2015). Dominant components of the Thoroughbred metabolome
691 characterised by ¹H-nuclear magnetic resonance spectroscopy: A metabolite
692 atlas of common biofluids. *Equine veterinary journal, 47*(6), 721-730.

693

694 Evrard, E., & Janssen, G. (1968). Gas-liquid chromatographic determination of
695 human fecal bile acids. *Journal of Lipid Research, 9*(2), 226-236.

696

697 Eyssen, H., & Parmentier, G. (1974). Biohydrogenation of sterols and fatty acids
698 by the intestinal microflora. *The American Journal of Clinical Nutrition, 27*(11),
699 1329-1340.

700

701 Fernando, H., Kondraganti, S., Bhopale, K. K., Volk, D. E., Neerathilingam, M.,
702 Kaphalia, B. S., ... & Shakeel Ansari, G. A. (2010). ^1H and ^{31}P NMR lipidome of
703 ethanol-induced fatty liver. *Alcoholism: Clinical and Experimental Research*,
704 34(11), 1937-1947.

705

706 Flint, H. J., Duncan, S. H., Scott, K. P., & Louis, P. (2015). Links between diet, gut
707 microbiota composition and gut metabolism. *Proceedings of the Nutrition Society*,
708 74(01), 13–22. <http://doi.org/10.1017/S0029665114001463>

709

710 Frost, G. S., Walton, G. E., Swann, J. R., Psichas, A., Costabile, A., Johnson, L. P., &
711 Barraclough, T. G. (2014). Impacts of plant-based foods in Ancestral Hominin
712 diets on the metabolism and function of gut microbiota in vitro. *Mbio*, 5(3),
713 e00853-14.

714

715 Gao, X., Pujos-Guillot, E., & Sébédio, J (2010). Development of a Quantitative
716 Metabolomic Approach to Study Clinical Human Fecal Water Metabolome Based
717 on Trimethylsilylation Derivatization and GC/MS Analysis. *Analytical Chemistry*,
718 82(15), 6447. doi:10.1021/ac1006552

719

720 Gao, X., Pujos-Guillot, E., Martin, J., Galan, P., Juste, C., Jia, W., & Sébédio, J (2009).
721 Metabolite analysis of human fecal water by gas chromatography/mass
722 spectrometry with ethyl chloroformate derivatization. *Analytical Biochemistry*,
723 393(2), 163. doi:10.1016/j.ab.2009.06.036

724

- 725 Gérard, P. (2013). Metabolism of cholesterol and bile acids by the gut microbiota.
726 *Pathogens*, 3(1), 14–24. <http://doi.org/10.3390/pathogens3010014>
727
- 728 Gregory, K. E., Bird, S. S., Gross, V. S., Marur, V. R., Lazarev, A. V., Walker, W. A., &
729 Kristal, B. S (2013). Method Development for Fecal Lipidomics Profiling.
730 *Analytical Chemistry*, 85(2), 1114. doi:10.1021/ac303011k
731
- 732 Griffiths, W. J., & Sjövall, J. (2010). Bile acids: analysis in biological fluids and
733 tissues. *Journal of Lipid Research*, 51(1), 23–41.
734 <http://doi.org/10.1194/jlr.R001941-JLR200>
735
- 736 Grundy, S. M., Ahrens, E. H., & Davignon, J. (1969). The interaction of cholesterol
737 absorption and cholesterol synthesis in man. *Journal of Lipid Research*, 10(3),
738 304–315.
739
- 740 Grundy, S. M., & Ahrens, E. H. (1969). Measurements of cholesterol turnover,
741 synthesis, and absorption in man, carried out by isotope kinetic and sterol
742 balance methods. *Journal of Lipid Research*, 10(1), 91–107.
743
- 744 Grundy, S. M., & Mok, H. Y. (1977). Determination of cholesterol absorption in
745 man by intestinal perfusion. *Journal of Lipid Research*, 18(2), 263–271.
746
- 747 Grundy, S. M., Ahrens, E. H., & Miettinen, T. A. (1965). Quantitative isolation and
748 gas-liquid chromatographic analysis of total fecal bile acids. *Journal of Lipid*
749 *Research*, 6, 397–410.

750

751 Hagey, L. R., & Krasowski, M. D. (2013). Microbial biotransformations of bile
752 acids as detected by electrospray mass spectrometry. *Advances in Nutrition*
753 *(Bethesda, Md.)*, 4(1), 29–35. <http://doi.org/10.3945/an.112.003061>

754

755 Hagio, M., Matsumoto, M., Fukushima, M., Hara, H., & Ishizuka, S. (2009).
756 Improved analysis of bile acids in tissues and intestinal contents of rats using
757 LC/ESI-MS. *Journal of Lipid Research*, 50(1), 173–180.
758 <http://doi.org/10.1194/jlr.D800041-JLR200>

759

760 Han, J., Lin, K., Sequeira, C., & Borchers, C. H. (2015). An isotope-labeled chemical
761 derivatization method for the quantitation of short-chain fatty acids in human
762 feces by liquid chromatography–tandem mass spectrometry. *Analytica chimica*
763 *acta*, 854, 86-94.

764

765 Hebel, T., Eisinger, K., Neumeier, M., Rein-Fischboeck, L., Pohl, R., Meier, E. M., ... &
766 Krautbauer, S. (2015). Lipid abnormalities in alpha/beta2-syntrophin null mice
767 are independent from ABCA1. *Biochimica et Biophysica Acta (BBA)-Molecular and*
768 *Cell Biology of Lipids*, 1851(5), 527-536.

769

770 Hofmann, A. F., & Eckmann, L. (2006). How bile acids confer gut mucosal
771 protection against bacteria. *Proceedings of the National Academy of Sciences of*
772 *the United States of America*, 103(12), 4333–4334.
773 <http://doi.org/10.1073/pnas.0600780103>

774

- 775 Hofmann, A. F., & Hagey, L. R. (2008). Bile acids: chemistry, pathochemistry,
776 biology, pathobiology, and therapeutics. *Cellular and Molecular Life Sciences*
777 (*CMLS*), 65(16), 2461–2483. <http://doi.org/10.1007/s00018-008-7568-6>
778
- 779 Hofmann, A. F. (2009). Bile acids: trying to understand their chemistry and
780 biology with the hope of helping patients. *Hepatology*, 49(5), 1403-1418.
781
- 782 Hofmann, A. F., Hagey, L. R., & Krasowski, M. D. (2010). Bile salts of vertebrates:
783 structural variation and possible evolutionary significance. *Journal of Lipid*
784 *Research*, 51(2), 226–246. <http://doi.org/10.1194/jlr.R000042>
785
- 786 Hong, Y. S., Ahn, Y. T., Park, J. C., Lee, J. H., Lee, H., Huh, C. S., ... & Hwang, G. S.
787 (2010). ¹H NMR-based metabonomic assessment of probiotic effects in a colitis
788 mouse model. *Archives of pharmacal research*, 33(7), 1091-1101.
789
- 790 Jacobs, D. M., Deltimple, N., van Velzen, E., Van Dorsten, F. A., Bingham, M.,
791 Vaughan, E. E., & van Duynhoven, J. (2008). ¹H NMR metabolite profiling of feces
792 as a tool to assess the impact of nutrition on the human microbiome. *NMR in*
793 *Biomedicine*, 21(6), 615–626. <http://doi.org/10.1002/nbm.1233>
794
- 795 Jiménez-Girón, A., Ibáñez, C., Cifuentes, A., Simó, C., Muñoz-González, I., Martín-
796 Álvarez, P. J., ... & Moreno-Arribas, M. (2015). Estudio del metaboloma fecal tras
797 el consumo moderado y continuado de vino tinto por individuos sanos.
798

- 799 Kellogg, T. F., Knight, P. L., & Wostmann, B. S. (1970). Effect of bile acid
800 deconjugation on the fecal excretion of steroids. *Journal of lipid research*, *11*(4),
801 362-366.
- 802
- 803 Korpela, J. T. (1982). Capillary gas-liquid chromatography of faecal free and
804 esterified neutral sterols. *Scandinavian Journal of Clinical and Laboratory*
805 *Investigation*, *42*(6), 529-534.
- 806
- 807 Le Gall, G., Noor, S. O., Ridgway, K., Scovell, L., Jamieson, C., Johnson, I. T., et al.
808 (2011). Metabolomics of Fecal Extracts Detects Altered Metabolic Activity of Gut
809 Microbiota in Ulcerative Colitis and Irritable Bowel Syndrome. *Journal of*
810 *Proteome Research*, *10*(9), 4208-4218. <http://doi.org/10.1021/pr2003598>
- 811
- 812 Le Roy, C. I., Štšepetova, J., Sepp, E., Songisepp, E., Claus, S. P., & Mikelsaar, M.
813 (2015). New insights into the impact of Lactobacillus population on host-bacteria
814 metabolic interplay. *Oncotarget*, *6*(31), 30545-30556.
- 815
- 816 Leidl, K., Liebisch, G., Richter, D., & Schmitz, G. (2008). Mass spectrometric
817 analysis of lipid species of human circulating blood cells. *Biochimica et*
818 *Biophysica Acta (BBA)-Molecular and Cell Biology of Lipids*, *1781*(10), 655-664.
- 819
- 820 Li, J. V., Saric, J., Wang, Y., Keiser, J., Utzinger, J., & Holmes, E. (2011).
821 Chemometric analysis of biofluids from mice experimentally infected with
822 *Schistosoma mansoni*. *Parasites & Vectors*, *4*, 179. [http://doi.org/10.1186/1756-](http://doi.org/10.1186/1756-3305-4-179)
823 3305-4-179

824

825 Lichtenstein, A. H. (1990). Intestinal cholesterol metabolism. *Annual medicine*
826 22(1)

827

828 Liebisch, G., Vizcaíno, J. A., Köfeler, H., Trötz Müller, M., Griffiths, W. J., Schmitz, G.,
829 & Wakelam, M. J. (2013). Shorthand notation for lipid structures derived from
830 mass spectrometry. *Journal of lipid research*, jlr-M033506.

831

832 Lozupone, C. A., Stombaugh, J. I., Gordon, J. I., Jansson, J. K., & Knight, R. (2012).
833 Diversity, stability and resilience of the human gut microbiota. *Nature*,
834 489(7415), 220-230.

835

836 Lütjohann, D., Meese, C. O., Crouse, J. R., & Bergmann, von, K. (1993). Evaluation
837 of deuterated cholesterol and deuterated sitostanol for measurement of
838 cholesterol absorption in humans. *Journal of Lipid Research*, 34(6), 1039–1046.

839

840 Marchesi, J. R., Holmes, E., Khan, F., Kochhar, S., Scanlan, P., Shanahan, F., ... &
841 Wang, Y. (2007). Rapid and noninvasive metabonomic characterization of
842 inflammatory bowel disease. *Journal of proteome research*, 6(2), 546-551.

843

844 Martin, F.-P. J., Sprenger, N., Montoliu, I., Rezzi, S., Kochhar, S., & Nicholson, J. K.
845 (2010). Dietary modulation of gut functional ecology studied by fecal
846 metabonomics. *Journal of Proteome Research*, 9(10), 5284–5295.
847 <http://doi.org/10.1021/pr100554m>

848

- 849 Meiboom, S., & Gill, D. (1958). Modified Spin-Echo Method for Measuring Nuclear
850 Relaxation Times. *Review of Scientific Instruments*, 29(8), 688.
851 <http://doi.org/10.1063/1.1716296>
852
- 853 Midtvedt, T., Lingaas, E., Carlstedt-Duke, B., Höverstad, T., Midtvedt, A. C.,
854 Saxerholt, H., et al. (1990). Intestinal microbial conversion of cholesterol to
855 coprostanol in man. Influence of antibiotics. *APMIS : Acta Pathologica,*
856 *Microbiologica, Et Immunologica Scandinavica*, 98(9), 839–844.
857
- 858 Miettinen, T. A., Ahrens, E. H., & Grundy, S. M. (1965). Quantitative isolation and
859 gas-liquid chromatographic analysis of total dietary and fecal neutral steroids.
860 *Journal of Lipid Research*, 6, 411–424.
861
- 862 Monleon, D., Morales, J. M., Barrasa, A., Lopez, J. A., Vazquez, C., & Celda, B.
863 (2009). Metabolite profiling of fecal water extracts from human colorectal
864 cancer. *NMR in Biomedicine*, 22(3), 342-348.
865
- 866 Mukherji, A., Kobiita, A., Ye, T., & Chambon, P. (2013). Homeostasis in intestinal
867 epithelium is orchestrated by the circadian clock and microbiota cues
868 transduced by TLRs. *Cell*, 153(4), 812–827.
869 <http://doi.org/10.1016/j.cell.2013.04.020>
870
- 871 Ndagijimana, M., Laghi, L., Vitali, B., Placucci, G., Brigidi, P., & Guerzoni, M. E.
872 (2009). Effect of a synbiotic food consumption on human gut metabolic profiles

873 evaluated by ^1H Nuclear Magnetic Resonance spectroscopy. *International*
874 *journal of food microbiology*, 134(1), 147-153.

875

876 O'Hara, A. M., & Shanahan, F. (2006). The gut flora as a forgotten organ. *EMBO*
877 *Reports*, 7(7), 688–693. <http://doi.org/10.1038/sj.embor.7400731>

878

879 Owen, R. W., Thompson, M. H., & Hill, M. J. (1984). Analysis of metabolic profiles
880 of steroids in faeces of healthy subjects undergoing chenodeoxycholic acid
881 treatment by liquid-gel chromatography and gas-liquid chromatography-mass
882 spectrometry. *Journal of Steroid Biochemistry*, 21(5), 593–600.

883

884 Palmer, C., Bik, E. M., DiGiulio, D. B., Relman, D. A., & Brown, P. O. (2007).
885 Development of the human infant intestinal microbiota. *PLoS Biol*, 5(7), e177.

886

887 Perwaiz, S., Mignault, D., Tuchweber, B., & Yousef, I. M. (2002). Rapid and
888 improved method for the determination of bile acids in human feces using MS.
889 *Lipids*, 37(11), 1093–1100.

890

891 Phua, L. C., Chue, X. P., Koh, P. K., Cheah, P. Y., Ho, H. K., & Chan, E. C. Y. (2014).
892 Non-invasive fecal metabonomic detection of colorectal cancer. *Cancer biology &*
893 *therapy*, 15(4), 389-397.

894

895 Phua, L. C., Koh, P. K., Cheah, P. Y., Ho, H. K., & Chan, E. C. Y. (2013). Global gas
896 chromatography/time-of-flight mass spectrometry (GC/TOFMS)-based

897 metabonomic profiling of lyophilized human feces. *Journal of Chromatography B*,
898 937, 103-113.

899

900 Ridlon, J. M., Kang, D.-J., & Hylemon, P. B. (2006). Bile salt biotransformations by
901 human intestinal bacteria. *Journal of Lipid Research*, 47(2), 241–259.
902 <http://doi.org/10.1194/jlr.R500013-JLR200>

903

904 Romick-Rosendale, L. E., Goodpaster, A. M., Hanwright, P. J., Patel, N. B., Wheeler,
905 E. T., Chona, D. L., & Kennedy, M. A. (2009). NMR-based metabonomics analysis
906 of mouse urine and fecal extracts following oral treatment with the broad-
907 spectrum antibiotic enrofloxacin (Baytril). *Magnetic Resonance in Chemistry :
908 MRC*, 47 Suppl 1, S36–46. <http://doi.org/10.1002/mrc.2511>

909

910 Saric, J., Wang, Y., Li, J., Coen, M., Utzinger, J., Marchesi, J. R., et al. (2008). Species
911 variation in the fecal metabolome gives insight into differential gastrointestinal
912 function. *Journal of Proteome Research*, 7(1), 352–360.
913 <http://doi.org/10.1021/pr070340k>

914

915 Sayin, S. I., Wahlström, A., Felin, J., Jäntti, S., Marschall, H.-U., Bamberg, K., et al.
916 (2013). Gut Microbiota Regulates Bile Acid Metabolism by Reducing the Levels of
917 Tauro-beta-muricholic Acid, a Naturally Occurring FXR Antagonist. *Cell
918 Metabolism*, 17(2), 225–235. <http://doi.org/10.1016/j.cmet.2013.01.003>

919

920 Scherer, M., Böttcher, A., & Liebisch, G. (2011). Lipid profiling of lipoproteins by
921 electrospray ionization tandem mass spectrometry. *Biochimica et Biophysica*
922 *Acta (BBA)-Molecular and Cell Biology of Lipids*, 1811(11), 918-924.

923

924 Schiller, J., & Arnold, K. (2002). Application of high resolution ³¹P NMR
925 spectroscopy to the characterization of the phospholipid composition of tissues
926 and body fluids-a methodological review. *Medical Science Monitor*, 8(11),
927 MT205-MT222.

928

929 Schiller, J., Muller, M., Fuchs, B., Arnold, K., & Huster, D. (2007). ³¹P NMR
930 spectroscopy of phospholipids: From micelles to membranes. *Current Analytical*
931 *Chemistry*, 3(4), 283-301.

932

933 SScott, K. P., Duncan, S. H., & Flint, H. J. (2008). Dietary fibre and the gut
934 microbiota. *Nutrition Bulletin*, 33(3), 201-211.

935

936 Setchell, K. D., Lawson, A. M., Tanida, N., & Sjövall, J. (1983). General methods for
937 the analysis of metabolic profiles of bile acids and related compounds in feces.
938 *Journal of Lipid Research*, 24(8), 1085-1100

939

940 Sigruener, A., Kleber, M. E., Heimerl, S., Liebisch, G., Schmitz, G., & Maerz, W.
941 (2014). Glycerophospholipid and sphingolipid species and mortality: the
942 Ludwigshafen Risk and Cardiovascular Health (LURIC) study. *PloS one*, 9(1),
943 e85724.

944

- 945 Sonnenburg, E. D., & Sonnenburg, J. L. (2014). Starving our Microbial Self: The
946 Deleterious Consequences of a Diet Deficient in Microbiota-Accessible
947 Carbohydrates. *Cell Metabolism*, 20(5), 779–786.
948 <http://doi.org/10.1016/j.cmet.2014.07.003>
949
- 950 Snog-Kjaer, A., Prange, I., & Dam, H. (1956). Conversion of cholesterol into
951 coprosterol by bacteria in vitro. *Journal of General Microbiology*, 14(2), 256–260.
952
- 953 Spritz, N., Ahrens, E. H., & Grundy, S. (1965). Sterol balance in man as plasma
954 cholesterol concentrations are altered by exchanges of dietary fats. *Journal of*
955 *clinical investigation*, 44(9), 1482–1493.
956
- 957 Story, J. A., & Furumoto, E. J. (1991). Bile acid analysis: methods and problems.
958 *European Journal of Cancer Prevention : the Official Journal of the European*
959 *Cancer Prevention Organisation (ECP)*, 1 Suppl 2, 29–33.
960
- 961 Su, X., Wang, N., Chen, D., Li, Y., Lu, Y., Huan, T., & Li, L. (2016). Dansylation
962 isotope labeling liquid chromatography mass spectrometry for parallel profiling
963 of human urinary and fecal submetabolomes. *Analytica chimica acta*, 903, 100-
964 109.
965
- 966 Tian, Y., Zhang, L., Wang, Y., & Tang, H. (2011). Age-related topographical
967 metabolic signatures for the rat gastrointestinal contents. *Journal of proteome*
968 *research*, 11(2), 1397-1411.
969

970 Trauner, M., Claudel, T., Fickert, P., Moustafa, T., & Wagner, M. (2010). Bile acids
971 as regulators of hepatic lipid and glucose metabolism. *Digestive Diseases*, 28(1),
972 220-224.

973

974 Voigt, R. M., Forsyth, C. B., Green, S. J., Mutlu, E., Engen, P., Vitaterna, M. H., et al.
975 (2014). Circadian disorganization alters intestinal microbiota. *PloS One*, 9(5),
976 e97500. <http://doi.org/10.1371/journal.pone.0097500>

977

978 Weir, T. L., Manter, D. K., Sheflin, A. M., Barnett, B. A., Heuberger, A. L., & Ryan, E.
979 P. (2013). Stool microbiome and metabolome differences between colorectal
980 cancer patients and healthy adults. *PloS one*, 8(8), e70803.

981

982 Wenk, M. R. (2010). Lipidomics: New Tools and Applications. *Cell*, 143(6), 888–
983 895. <http://doi.org/10.1016/j.cell.2010.11.033>

984

985 Wu, J., An, Y., Yao, J., Wang, Y., & Tang, H. (2010). An optimised sample
986 preparation method for NMR-based faecal metabonomic analysis. *Analyst*,
987 135(5), 1023-1030.

988

989 Yap, I. K. S., Li, J. V., Saric, J., Martin, F.-P., Davies, H., Wang, Y., et al. (2008).
990 Metabonomic and microbiological analysis of the dynamic effect of vancomycin-
991 induced gut microbiota modification in the mouse. *Journal of Proteome Research*,
992 7(9), 3718–3728. <http://doi.org/10.1021/pr700864x>

993

- 994 Yatsunenکو, T., Rey, F. E., Manary, M. J., Trehان, I., Dominguez-Bello, M. G.,
995 Contreras, M., et al. (2012). Human gut microbiome viewed across age and
996 geography. *Nature*, 486(7402), 222–227. <http://doi.org/10.1038/nature11053>
997
- 998 Zhao, Y. Y., Zhang, L., Long, F. Y., Cheng, X. L., Bai, X., Wei, F., & Lin, R. C. (2013).
999 UPLC-Q-TOF/HSMS/MS E-based metabonomics for adenine-induced changes in
1000 metabolic profiles of rat faeces and intervention effects of ergosta-4, 6, 8 (14),
1001 22-tetraen-3-one. *Chemico-biological interactions*, 201(1), 31-38.
1002
- 1003 Zhao, Y., Wu, J., Li, J. V., Zhou, N. Y., Tang, H., & Wang, Y. (2013). Gut microbiota
1004 composition modifies fecal metabolic profiles in mice. *Journal of proteome*
1005 *research*, 12(6), 2987-2999.
1006
- 1007 Zheng, X., Qiu, Y., Zhong, W., Baxter, S., Su, M., Li, Q., ... & Zeisel, S. H. (2013). A
1008 targeted metabolomic protocol for short-chain fatty acids and branched-chain
1009 amino acids. *Metabolomics*, 9(4), 818-827.
1010
1011

1012 **FIGURE CAPTIONS**

1013 **Figure 1:** Estimated proportion of fecal cholesterol derivatives in humans.

1014

1015 **Figure 2:** Neutral loss (NL) of 189 of lipid extracts prepared from fecal samples.
1016 Suspensions of human fecal samples in water/methanol (1/1) were subjected to
1017 bead-based homogenization. Homogenates were extracted according to Bligh
1018 and Dyer (Bligh & DYER, 1959). Crude lipid extracts were analyzed by direct
1019 flow injection analysis as described previously (Matyash et al., 2008). Displayed
1020 are NL 189 spectra, specific for phosphatidylglycerol (PG), of three different
1021 samples normalized to the highest intensity.

1022

1023 **Table 1:** Summary of published studies applying metabolomics to study fecal
1024 samples using NMR and MS platforms. Summary of protocols and main outcomes
1025 are included.


1026 Key: GC-FID: Gas chromatography- Flame ionization detector; GC-MS: Gas
1027 chromatography-Mass spectrometry; LC-MS: Liquid chromatography-Mass
1028 spectrometry; UPLC-MS: Ultra-Performance Liquid Chromatography-Mass
1029 spectrometry

Analytical technique	Study aim	Sample preparation overview	Material / buffer	Measurement	Main results	References
NMR 400 MHz	Aging in mice	1:4 (wN/Av) feces in deuterated PBS + two filtration step	60 μ L deuterated PBS containing 2 mM TSP in 600 μ L of extract	TOCSY & HSQC	Aging induces \uparrow 4-hydroxyphenylacetate, histidine, formate, succinate and \downarrow α -ketoisocaproate, α -ketoisovalerate, -hydroxybutyrate, bile salts, isoleucine, methionine	Calvani et al., 2014
	Colorectal cancer human	1:3 (wN/Av) feces in deuterated PBS + vortexing and centrifugation	50 μ L TSP (4 mM) in D2O added in 500 μ L supernatant	N/A	N/A	Bezabeh et al. 2009
NMR 500 MHz	Experimental optimisation, mice	1:10 mg. μ L ⁻¹ feces-to-buffer ratio, tissuelyser	N/A	COSY, TOCSY, HSQC, HMBC	Identification of 40 metabolites	Wu et al., 2010
NMR 600 MHz	Ulcerative colitis activity, humans	1:2 (WN/Av) feces to PBS, vortexing, centrifugation and filtration	4 μ L of D2O/A500 μ M TSP with 40 μ L of fecal extract	CPMG	Active UC induces \uparrow BCAAs, lysine, alanine, taurine	Bjerrum et al. 2014
	Antibiotic treatment (gentamicine, ceftriaxone), mice	1:10 feces to PBS, freeze-thaw treatment, tissuelysr, centrifugation	30% D2O, 0.002% TSP, 0.03% of Na3N (wN/Av)	COSY, TOCSY, JRES, HSQC, HMBC	Antibiotic induces \uparrow oligosacharides. phenolic acids and \downarrow SCFAs, uracil, hypoxanthine	Zhao et al., 2013
	Age, rat	1:10 feces to PBS, vortexing, freeze-thaw, tissuelyser, centrifugation	0.1 M K2HPO4N/ANaH2PO4, pH = 7.4, containing 10% D2O, 0.58 mM TSP	COSY, TOCSY, HSQC, HMBC, DOSY	Aging induces \downarrow arabinose, xylose, galactose, arabinoxylan, propionate and inosine and \uparrow taurine, xylose, arabinose, galactose, arabinoxylans	Tian et al., 2012
	Infection <i>Schistosoma</i>	2 fecal pellets homogenized in	PBS containing 0.01% TSP	CPMG, COSY, TOCSY	Infection induces \uparrow 5-Aminovalerate, SCFAs	Li et al., 2011


<i>mansoni</i> , mice	700 µL PBS, sonication, centrifugation				(propionate)	
IBS and UC, human	1:50 (wN/Av) feces to PBS, centrifugation, filtration	deuterated PBS containing 1 mM TSP		COSY, HSQC, HMBC	UC induces ↑ taurine and cadaverine, IBS induces ↑ bile acids and ↓ BCFAs	Le Gall et al., 2011
ProN/Aprebiotics, mice	1:12 (wN/Av) mashed feces to PBS, centrifugation	60 µL DSS (5 mM) in D2O added in 600 µL supernatant		TOCSY, HMBC, HSQC	Prebiotic induces ↓ threonine, alanine, glutamate, glutamine, aspartate, lysine, lycine, butyrate, uracil, hypoxanthine and ↑ monosaccharides, glucose, trimethylamine. Pre and probiotic ↓ trimethylamine and ↑ acetate, butyrate, glutamine	Hong et al., 2010
Colorectal cancer, human	1:2 feces to distilled water, homogenization, freezing, thawing, centrifugation	100 µL D2O to 500 µL fecal water		TOCSY	Cancer induces ↓ acetate, butyrate	Monleon et al., 2009
Grape juice and wine extract consumption, human	1:20 (wN/Av) feces to cold D2O or CD3OD, vortexing, centrifugation	D2O or CD3OD containing 1 mM TSP		CPMG	Grape juice consumption + wine induces ↓ isobutyrate	Jacobs et al., 2008
Effect of Species, storage, lyophilization, sonication, filtration and homogenisation	N/A	N/A		COSY, TOCSY, HSQC, HMBC	Storage: ↑ alanine, glutamate, threonine, aspartic acid, BCAAs, glucose. Lyophilization: ↑ BCAAs and ↓ succinate, SCFAs. Sonication: ↑ uracil, glucose and ↓ SCFAs	Saric et al., 2008
UC, human	1:2 (wN/Av) feces to PBS, vortexing, filtration, centrifugation	200 µL buffer (10% D2O & 0.01% TSP) in 400 µL fecal water		COSY, TOCSY	UC induces ↓ acetate, butyrate, methylamine, TMA and ↑ isoleucine, leucine, lysine	Marchesi et al., 2007


NMR 700 MHz	<i>Lactobacillus</i>	1:2 (wN/Av) feces to detarated PBS, tissue lyser, centrifugation	Fecal water extracted in 9:1 D2ON/AH2O and 0.05 % TSP	Noesy	<i>L. helveticus</i> induces ↑ butyrate, lactate and incresed <i>Lactobacillus</i> level induces ↑ phenylalanine, tyrosine, lysine, lactate, propionate, valine, leucine, isoleucine, butyrate, acetate.	Le Roy et al., 2015
NMR 850 MHz	Baytrill treatment, mice	1:2 (wN/Av) feces to PBS, vortexing, centrifugation	200 μL buffer 10% D2O and 0.01% TSP in 400 μL fecal water	CPMG	Treatment induces ↓ alanine, butyrate, isoleucine, leucine, propionate, threonine, valine and ↑ urea	Romick-Rosendale et al., 2009
GC-MS	Colorectal cancer human	oximation and silylation	lyophilized human feces	N/A	Cancer patients ↓ butyrate, poly and monounsaturated fatty acids, ursodeoxycholic acid and ↑ acetate, amino acids	Weir et al., 2013
	VOCs irritable bowel syndrome, active Crohn's disease, ulcerative colitis	SPME	human feces	N/A	240 metabolites; esters of short chain fatty acids, cyclohexanecarboxylic acid associated with irritable bowel syndrome	Ahmed, Greenwood, de Lacy Costello, Ratcliffe, & Probert, 2013
	technical paper	trimethylsilylation	human fecal water	N/A	133 compounds structurally confirmed; 33 quantified	Gao, Pujos-Guillot, & Sébédio, 2010
	technical paper	ethyl chloroformate derivatization	human fecal water	N/A	73 compounds identified; 34 validated by reference standards	Gao et al., 2009
GC-TOF-MS	technical paper	oximation and silylation	lyophilized human feces	N/A	107 metabolites matched with mass spectra libraries, influence of blood on fecal metabolome	Phua, Koh, Cheah, Ho, & Chan, 2013
	Colorectal cancer human	oximation and silylation	lyophilized human feces	N/A	fecal metabolomic profiles of patients clearly differ from healthy subjects	Phua et al., 2014
GC-MS, GC-FID	VOCs, technical paper	SPME	human feces	N/A	evaluation of eight different commercially available SPME	Dixon et al., 2011

					fibers		
UPLC-MSN/ATOF-MS	liver cirrhosis, hepatocellular carcinoma	homogenization, centrifugation, filtration	human feces	N/A	Cancer patients ↓ chenodeoxycholic acid, 7-ketolithocholic acid, urobilinogen, urobilin and ↑ lysophosphatidylcholine (LPC) 16:0 and 18:0	Cao et al., 2011	
UHPLC-TOF-MS	Effect of consumption of red wine	mixing with saline solution, centrifugation, filtration	human feces	N/A	37 metabolites related to wine intake	Jiménez-Girón et al., 2015	
UPLC-Q-TOF-MS	chronic renal failure	homogenization, extraction with acetonitrile	rat feces	N/A	renal failure ↑ chenodeoxycholic acid, palmitic acid, adenine, phytosphingosine, monoglycerol 24:1, 12-hydroxy-3-oxocholadienic acid, lysophosphatidylethanolamine 18:2 and 16:0 and ↓ 7-ketolithocholic acid	Zhao, Cheng, Wei, Bai, & Lin, 2012	
LC-MS, LC-UV	technical paper	Dried fecal samples extraction with water and acetonitrile, derivatization with dansyl chloride	human feces	N/A	67 metabolites (mainly amino acids) identified	Su et al., 2015	



Minor fecal cholesterol derivatives





ACCEPTED MANUSCRIPT