



# *Increased anticipatory but decreased consummatory brain responses to food in sisters of Anorexia Nervosa patients*

Article

Published Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Open Access

Horndasch, S., Okeefe, S. ., Lamond, A., Brown, K. and McCabe, C. (2016) Increased anticipatory but decreased consummatory brain responses to food in sisters of Anorexia Nervosa patients. *British Journal of Psychiatry Open*, 2 (4). pp. 255-261. ISSN 2056-4724 doi: <https://doi.org/10.1192/bjpo.bp.115.002550> Available at <http://centaur.reading.ac.uk/65961/>

It is advisable to refer to the publisher's version if you intend to cite from the work.

To link to this article DOI: <http://dx.doi.org/10.1192/bjpo.bp.115.002550>

Publisher: Royal College of Psychiatrists

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

## **CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

# Increased anticipatory but decreased consummatory brain responses to food in sisters of anorexia nervosa patients

Stefanie Horndasch, Sophie O’Keefe, Anneka Lamond, Katie Brown and Ciara McCabe

## Background

We have previously shown increased anticipatory and consummatory neural responses to rewarding and aversive food stimuli in women recovered from anorexia nervosa (AN).

## Aims

To determine whether these differences are trait markers for AN, we examined the neural response in those with a familial history but no personal history of AN.

## Method

Thirty-six volunteers were recruited: 15 who had a sister with anorexia nervosa (family history) and 21 control participants. Using fMRI we examined the neural response during an anticipatory phase (food cues, rewarding and aversive), an effort phase and a consummatory phase (rewarding and aversive tastes).

## Results

Family history (FH) volunteers showed increased activity in the caudate during the anticipation of both reward and aversive

food and in the thalamus and amygdala during anticipation of aversive only. FH had decreased activity in the dorsal anterior cingulate cortex, the pallidum and the superior frontal gyrus during taste consumption.

## Conclusions

Increased neural anticipatory but decreased consummatory responses to food might be a biomarker for AN. Interventions that could normalise these differences may help to prevent disorder onset.

## Declaration of interest

C.M. has acted as a consultant to P1VITAL, Givaudan, GWPharma, the British Broadcasting Corporation (BBC) and Channel 4.

## Copyright and usage

© The Royal College of Psychiatrists 2016. This is an open access article distributed under the terms of the Creative Commons Non-Commercial, No Derivatives (CC BY-NC-ND) license.

Anorexia nervosa (AN) tends to affect adolescents and young women and is characterised by restricted eating and emaciation. The aetiology of AN is still unclear.<sup>1</sup> However, recent research suggests abnormalities in reward and executive control as potential underlying biological mechanisms in the pathogenesis of AN.<sup>1</sup>

Neural correlates of reward processing in individuals suffering from AN seem to be particularly interesting, especially responses to food rewards. A recent review by Zhu *et al*<sup>2</sup> on fMRI studies concluded that emotional arousal (especially disgust) reflected by hyperactivation in, for example, the amygdala, the anterior cingulate cortex (ACC) and the caudate nucleus plays an important role in the processing of food information in AN. This is in line with the concept of increased food cue reactivity in AN patients.<sup>3,4</sup> Hypoactivity in somatosensory regions suggests reduced physical stimulation by food in patients.<sup>2</sup> The majority of studies, however, involve visual presentation of pictorial food stimuli; neural correlates of taste processing are less commonly investigated in AN.

During the application of sucrose or water, recovered AN patients have been found to have less neural activation in the insula and ventral striatum to both conditions compared with controls, which was interpreted as less pronounced reward reactions in those recovered from AN.<sup>5</sup> Further, when exposed to a pleasant chocolate drink during both hunger and satiety, it was found that AN patients had increased responses in the hunger condition in the right amygdala and left medial temporal gyrus, and in the inferior temporal gyrus (extrastriate body area) in the satiety condition compared with controls.<sup>6</sup> The authors report that this could reflect fear of weight gain and a mental focus on the body in AN patients.

In a more recent study using a reward-conditioning task, which has been associated with brain dopamine reward circuits, currently ill AN patients showed increased responses in the anteroventral striatum, insula and prefrontal cortex compared with the opposite response in an obese group. Together, this research points to aberrant brain reward activity in AN and blunted reward responses in obesity.<sup>7</sup>

We have developed a paradigm involving the sight and taste of food stimuli which enables us to look at two aspects of food responses: anticipatory and consummatory. The former is reflecting the concept of ‘wanting’ and the latter the concept of ‘liking’.<sup>8,9</sup> Affective neuroscience studies of ‘wanting’ and ‘liking’ have suggested that these psychological processes map onto distinct brain reward systems. For example, studies of pleasure identify hedonic impact in the ventral pallidum, nucleus accumbens and orbitofrontal cortex,<sup>10–13</sup> whereas ‘wanting’ or incentive salience is mediated by neural systems that include mesolimbic dopamine projections from the ventral tegmental area to the ventral striatum.<sup>14</sup> Furthermore, dopamine has been shown to be involved in the learning about rewards in prefrontal cortical regions such as the ACC and the orbitofrontal cortex.<sup>15</sup>

Using our paradigm, we have previously shown that females recovered from AN have increased neural responses to the anticipation of food in the occipital cortex, ACC and caudate and increased ventral striatum, insula and putamen activity to its consummation, compared with controls. As these are areas known to be involved in the processing of anticipatory and consummatory responses to rewarding stimuli in general and taste stimuli specifically, we suggested that our results might reflect a neural biomarker of risk

for AN.<sup>16</sup> However, as periods of malnutrition could lead to changes in the brain including microstructural alterations,<sup>17</sup> our results may have been a consequence of recurrent AN rather than a cause. One way to resolve this issue is to study the neural response in people at increased risk of AN before the onset of illness. One reliable risk factor for AN is family heritance with genetic heritability accounting for 50–80% of the risk of developing an eating disorder.<sup>18</sup> In a large controlled family study, 3.4% of female family members of women with AN had developed AN themselves.<sup>19</sup> AN was four times more common in siblings of AN patients compared with siblings of control participants of a nation-wide register-based Danish study.<sup>20</sup>

The present study therefore aimed to examine the neural response to food in those with a familial history of anorexia nervosa (family history) but with no personal history, compared to those with no family history (FH), to determine whether anticipatory and/or consummatory neural responses are possible trait markers for AN and not simply scars of starvation. Knowing whether anticipatory and/or consummatory responses to food stimuli are dysfunctional in those at risk of AN can help inform the development of psychological and pharmacological treatments that can directly target either both or one of the processes separately. Furthermore, we include both rewarding and aversive food stimuli to assess whether any aberrant neural responses to food in the FH group is valence specific.

We hypothesised that the FH group would have increased neural responses during both anticipatory and consummatory phases in areas such as the anterior cingulate, orbitofrontal cortex, and striatum similar to those found in our study with individuals recovered from AN and confirming that the neural response to food might be a biomarker for risk of AN.

## Method

### Participants

Fifteen women whose sisters met the DSM criteria<sup>21</sup> for AN and twenty-one controls were recruited for this study. Participants were recruited through advertisements both in the university and on social media sites asking, ‘Do you have a sister with anorexia nervosa?’ Ethics approval for the study was obtained from the Research Ethics Committee at the University of Reading. Written consent was obtained from all participants.

All participants underwent a screening process that involved a brief email screening checking that they had a sister with AN. Sisters then completed the Eating Disorder Diagnostic Scale standardised questionnaire.<sup>22</sup> We included participants where a diagnosis of clinical AN had been made by a general practitioner and/or psychiatrist and the symptoms described met criteria for AN. We did not attempt to assess the possible presence of other Axis I or II disorders in AN sisters. Study participants had a face-to-face assessment using the Structured Clinical Interview for DSM-IV.<sup>23</sup> All participants rated the following questionnaires: the Beck Depression Inventory-2nd Edition (BDI-II),<sup>24</sup> the Fawcett–Clark Pleasure Scale (FCPS),<sup>25</sup> the Snaith–Hamilton Pleasure Scale (SHAPS),<sup>26</sup> the Ruminative Thought Style Questionnaire,<sup>27</sup> the Temporal Experience of Pleasure Scale (TEPS),<sup>28</sup> the Behavioural Inhibition/Behavioural Activation Scale (BIS/BAS),<sup>29</sup> the Ruminative Response Scale for Eating Disorders<sup>30</sup> and the Eating Attitudes Test (EAT)<sup>31</sup> approximately 1 week before scanning. The participants completed a chocolate questionnaire to measure liking, craving and frequency of eating chocolate.<sup>16</sup> Current body mass index (BMI) was also recorded for each participant. Furthermore, IQ was tested using the National Adult Reading Test (NART).<sup>32</sup> Inclusion criteria for all participants included: (1) BMI between 18.5 and 25 kg/m<sup>2</sup>, (2) maintenance of a weight in the healthy range (defined by the World Health Organization) since menarche, (3) no

use of psychoactive medications in the previous 12 months, (4) no lifetime history of any Axis I psychiatric disorder on the Structured Clinical Interview for DSM-IV. Participants in the control group had no first-degree relative with a current or past eating disorder diagnosis. General exclusion criteria for all participants included a history of head injury, neurological or other severe medical illness, pregnancy, and any contradictions to MRI. All participants were fluent in English, right-handed, had normal or corrected to normal vision, and were not taking medication except for the contraceptive pill. On the scan day, participants completed a visual analogue scale (VAS) measuring mood-related items including happiness, anxiety and disgust; they also completed the mood questionnaire Befindlichkeits Scale (Bf-SR).<sup>33</sup> Participants were also asked to refrain from eating chocolate for 24 h prior to being scanned, which was checked verbally before testing. Finally participants were also asked to rate how hungry they were immediately before scanning on a VAS between 1 and 10 (Table DS2).

The task was adapted from our previous study<sup>34</sup> and similar to our recent study.<sup>35</sup> It consisted of 40 trials and had 4 conditions based on the trial type (reward or aversive) and its level of difficulty (easy/hard). Trial type was cued by a visual stimulus (2 s – anticipatory phase) which indicated either to work to win the chocolate taste or to avoid the aversive taste. Difficulty was determined by the amount of effort required to complete the effort stage (easy = 24, hard = 45 button presses). This required volunteers to press a button as fast as possible (< 6 s) to move a bar towards the pleasant picture (reward) or away from the unpleasant picture (aversive), allowing enough time to complete easy trials but not hard ones. If the volunteers were successful on reward trials, they received the chocolate taste (5 s delivery – consummatory phase, then 2 s swallow cue) and if not, they received the tasteless solution. If the volunteers were successful on aversive trials, they received the tasteless solution and if not, they received the unpleasant taste (consummatory phase). A grey image (2 s) was presented at the end of each trial as a control for the cues. Each condition was repeated 10 times, chosen by random permutation. To sustain motivation, four trials (two reward/two aversive) were longer at 9 s each. Volunteers also rated ‘wanting’ and ‘pleasantness’ (+2 to –2) on a VAS (Fig. 1).

### Stimuli

For the anticipation phase, we used either a picture of liquid chocolate (reward) or of a mouldy drink (aversive). The rewarding taste was a Belgian chocolate drink and the aversive taste was a combination of the chocolate drink mixed with beetroot juice, providing a similar texture and caloric content, but that was rated as unpleasant in valence. A tasteless solution ( $25 \times 10^{-3}$  mol/L KCl



**Fig. 1** Stimuli with jittered inter-trial intervals (layout for a reward – hard trial).

and  $2.5 \times 10^{-3}$  mol/L NaHCO<sub>3</sub> in distilled H<sub>2</sub>O) was used as a rinse between trials. This was subtracted from the effects of the other taste stimuli to allow somatosensory and mouth movement effects to be removed.<sup>34</sup> A picture of a grey box was presented at the end of each trial to use it as a control visual stimulus. Solutions were delivered through three Teflon tubes held together by a plastic mouthpiece and connected by a one-way syringe-activated check valve (Model 14044-5, World Precision Instruments, Inc.), allowing 0.5 mL of solution to be manually delivered.

### fMRI scan

The experimental protocol consisted of an event-related interleaved design. A Siemens Magnetom Trio 3T whole body MRI scanner and a 12-channel head coil were used at the Centre for Integrative Neuroscience and Neurodynamics at the University of Reading. 3T-weighted echo planar imaging slices were acquired every 2 s (repetition time 2 s). Thirty-six axial slices with in-plane resolution of  $3 \times 3$  mm and between-plane spacing of 3 mm were obtained. The matrix size was  $64 \times 64$ , and the field of view was  $192 \times 192$  mm. Acquisition was carried out during the task performance yielding, on average, 1130 volumes in total. A whole-brain 3T-weighted echo planar imaging volume of the above dimensions and an anatomic T1 volume with axial plane slice thickness of 1 mm and in-plane resolution of  $1 \times 1$  mm were also acquired.

### fMRI analysis

Statistical parametric mapping (SPM8; <http://www.fil.ion.ucl.ac.uk/spm/software/spm8/>) was used to analyse the imaging data as documented in our previous study.<sup>16</sup> Data were pre-processed using realignment, normalisation to the Montreal Neurological Institute (MNI) coordinate system and spatial smoothing with a 6-mm full-width-at-half-maximum Gaussian kernel and global scaling. The time series at each voxel was low-pass filtered with a haemodynamic response kernel. Time series non-sphericity at each voxel was estimated and corrected for, and a high-pass filter with a cut-off period of 128 s was applied.

In the single-event design, a general linear model was then applied to the time course of activation in which stimulus onsets were modelled as single-impulse response functions and then convolved with the canonical haemodynamic response function. Linear contrasts were defined to test specific effects. Time derivatives were included in the basis functions set. Following smoothness estimation, linear contrasts of parameter estimates were defined to test the specific effects of the four conditions (pleasant cue – grey image, unpleasant cue – grey image, pleasant taste – rinse, unpleasant taste – rinse) with each individual data-set. Voxel values for each contrast resulted in a statistical parametric map of the corresponding *t* statistic, which was then transformed into the unit normal distribution (SPM *z*). Movement parameters for each person were added as additional regressors in 1st-level analyses.

Second-level fMRI analyses first examined simple main effects of task with one-sample *t*-tests for all scans (Table DS2 in the data supplement). These results were thresholded at  $P=0.05$  uncorrected and whole-brain cluster corrected ( $P<0.05$  family-wise error (FWE) for multiple comparisons). To examine the effect of group, independent *t*-tests were implemented in SPM8 for each of the four conditions separately and all data were reported thresholded at  $P=0.05$  uncorrected and whole-brain cluster corrected ( $P<0.05$  FWE for multiple comparisons). Plots of contrast estimates were extracted using the plots tool in SPM8, and WFU Pick Atlas (<http://www.fmri.wfubmc.edu/cms/software>) was used to display neural activation, with error bars representing the standard error of the mean. Activation coordinates are listed in the stereotactic space of the MNI ICBM 152 brain (Table 2).

## Results

### Participant characteristics

The two groups were matched for age, BMI and chocolate eating/liking (Table 1). There were significant differences between the control group and the FH group in the measures of anhedonia (Snaith–Hamilton Pleasure Scale,  $F(31)=1.182$ ,  $P<0.05$ , FCPS,  $F(31)=1.914$ ,  $P<0.05$ ). There were no significant differences between groups on measures of depression (Beck Depression Inventory-II), attitudes to eating (Eating Attitudes Tests), pleasure (Temporal Experience of Pleasure Scale), motivation (Behavioural Inhibition/Behavioural Activation Scale) or rumination (Ruminative Response Scale for Eating Disorders, Ruminative Thought Style Questionnaire) (Table 1). On the day of the scan, there were no significant differences between groups on the VAS, the BFS and on subjective ratings of hunger (see Table DS1).

### Ratings of stimuli

#### Wanting

A repeated-measures ANOVA with one factor, condition, with two levels (chocolate/aversive), was used to analyse group (FH and controls) wanting of stimuli. As expected, there was a significant main effect of condition on wanting ratings ( $F(1,70)=1839.260$ ,  $P<0.001$ ), with all participants wanting the chocolate significantly more than the aversive stimuli ( $t(71)=40.725$ ,  $P<0.001$ ). The ANOVA revealed a significant interaction between condition and group on participants' wanting ratings ( $F(1,70)=13.298$ ,  $P=0.001$ ). *t*-Tests revealed that controls wanted the chocolate significantly more than the FH group ( $t(70)=2.636$ ,  $P=0.01$ ); the control group also wanted the aversive stimuli significantly less than the FH group ( $t(70)=-2.382$ ,  $P<0.05$ ) (Fig. DS1).

#### Liking

To analyse the groups' (FH and controls) liking of the stimuli, a repeated-measures ANOVA with one factor, condition with three levels (chocolate, tasteless rinse and aversive), was used. As expected, there was a significant main effect of condition on all participants' liking ratings ( $F(2,68)=172.295$ ,  $P<0.001$ ) with all

**Table 1** Group demographics and psychosocial measures

Measure	Group			
	FH ( <i>n</i> =15)		Controls ( <i>n</i> =21)	
	Mean	s.d.	Mean	s.d.
Age	23.71	9.43	24.06	7.37
BMI	22.37	1.94	22.10	2.44
IQ	116.6	4.16	112.5	6.74
BDI-II	4.03	4.15	4.95	4.27
FCPS	135.47	9.45	143.74	12.51 <sup>a</sup>
EAT	26.29	10.54	28.11	10.12
SHAPS	21.21	4.15	18.00	4.48 <sup>a</sup>
TEPS	85.50	8.62	87.74	9.30
RRS-ED	11.21	3.21	11.11	2.45
RSTQ	82.00	20.54	78.16	21.85
BIS-BAS	51.00	4.95	46.82	5.99
Chocolate				
Craving	5.69	2.32	6.42	2.36
Liking	7.46	1.76	7.68	2.31

a.  $P<0.05$  independent samples *t*-test. BMI, body mass index; BDI-II, Beck Depression Inventory – 2nd Edition; FCPS, Fawcett–Clarke Pleasure Scale; EAT, Eating Attitudes Test; SHAPS, Snaith–Hamilton Pleasure Scale; TEPS, Temporal Experience of Pleasure Scale; RRS-ED, Ruminative Response Scale for Eating Disorders; RSTQ, Ruminative Thought Style Questionnaire; BIS-BAS, Behavioural Inhibition–Behavioural Activation Scale; VAS, visual analogue scale.



participants liking the chocolate significantly more than the rinse ( $t(106)=11.10, P<0.001$ ) and the aversive stimuli ( $t(70)=19.528, P<0.001$ ). Despite no significant interaction between condition and group ( $F(2,68)=2.682, P=0.076$ ), *post hoc t*-tests revealed that the control group liked the chocolate stimuli significantly more than the FH group ( $t(34)=2.028, P=0.05$ ) (Fig. DS1). There were no group differences in participants' liking of tasteless rinse ( $t(70)=-1.761, P=0.083$ ) or of the aversive stimuli ( $t(34)=-1.405, P=0.169$ ).

**Button presses to complete reward effort phase**

Repeated-measures ANOVAs with two factors, difficulty (easy and hard) and group (FH and controls), were used to analyse button presses on the chocolate trials. There was a significant main effect of difficulty on participants' number of button presses ( $F(1,34) = 166, P<0.001$ ), with all participants pressing more on hard chocolate trials (Fig. DS2a). There was also a significant interaction between difficulty and group ( $F(1,34)=14.98, P<0.001$ ), with the controls pressing more than FH for the hard chocolate trials ( $t(34)=3.449, P=0.002$ ).

**Button presses to complete aversive effort phase**

Repeated-measures ANOVAs with two factors, difficulty (easy and hard) and group (FH and controls), were used to analyse button presses on the aversive trials. There was a significant main effect of difficulty on participants' number of button presses ( $F(1,34)=167, P<0.001$ ), with all participants pressing more on hard chocolate aversive trials (Fig. DS2b). There was also a main effect of group ( $F(1,34)=10.7, P=0.002$ ) and significant interaction between difficulty and group ( $F(1,34) = 10.6, P=0.003$ ), with the controls pressing more than FH for the hard aversive trials ( $t(34)=3.275, P=0.002$ ).

**Time taken to complete reward effort phase**

Repeated-measures ANOVAs with two factors, difficulty (easy and hard) and group (FH and controls) were used to analyse time taken on the chocolate trials. There was a significant main effect of difficulty on participants' number of button presses ( $F(1,34)=368, P<0.001$ ), with all participants being faster on chocolate easy trials (Fig. DS3a). There was also a main effect of group ( $F(1,34)=16.4, P<0.001$ ) and significant interaction between difficulty and group ( $F(1,34)=14.8, P<0.001$ ), with the FH taking longer than the controls on the chocolate easy trials ( $t(34)=-3.98, P<0.001$ ) and hard trials ( $t(34)=-3.67, P<0.001$ ) (Fig. DS3a).

**Time taken to complete aversive effort phase**

Repeated-measures ANOVAs with two factors, difficulty (easy and hard) and group (FH and controls), were used to analyse time taken on the aversive trials. There was a significant main effect of difficulty on participants' number of button presses ( $F(1,34)=334, P<0.001$ ), with all participants being faster on aversive easy trials (Fig. DS3b). There was also a main effect of group ( $F(1,34)=10.66, P=0.003$ ) and significant interaction between difficulty and group ( $F(1,34)=10.38, P=0.003$ ), with the FH taking longer than the controls on the aversive easy trials ( $t(34)=-3.26, P=0.003$ ) and hard trials ( $t(34)=-2.75, P=0.009$ ) (Fig. DS3b).

**Accuracy**

It was expected that easy trials would be successfully completed, and participants would achieve the chocolate taste and on aversive easy trials avoid the aversive taste and instead receive the tasteless rinse. It was also expected that the hard trials would not be completed, and participants would fail to achieve the chocolate and instead receive the tasteless rinse on chocolate trials and fail to avoid the aversive taste on hard aversive trials. Therefore, an error here means that participants did not perform as expected.

Using independent samples *t*-tests, we found that the FH group made more errors for both the chocolate easy trials ( $t(34)=1.221, P<0.001$ ) and the aversive easy trials ( $t(34)=10.437, P<0.05$ ) compared with the controls (Fig. DS4).

**fMRI response**

Main effects of task

Table DS2 provides a summary of the results for each contrast across all participants to indicate the main effect of task. Sights of the food stimuli (anticipation phase) activated regions including the occipital cortex, the ACC and lateral orbitofrontal cortex. As expected, the tastes activated areas such as the insula and ventral striatum (see Table DS2).

Main effects of group

Table 2 provides a summary of the results of the interaction with group (FH v. control). As expected, there were no significant differences in response to the taste of chocolate between the two groups in the primary taste cortex (i.e. anterior insula), confirming that the sensory experience of chocolate was associated with a similar neural response across groups (Table 2). As participants showed significant differences in measures of anhedonia (FCPS and SHAPS), and these measures were shown to be highly correlated:  $r=-0.492$  and  $P<0.01$ ; SHAPS scores were therefore entered as a covariate into the fMRI analysis. With only two exceptions, the fMRI results remained significant after global SHAPS scores were added as a covariate.

Anticipation

In support of our previous findings with recovered AN, we found increased activation during the anticipation phase (chocolate pictures and aversive pictures) in the FH group compared with the control group in the planum polare/amygdala region (Fig. 2) and the caudate/thalamus region (Fig. DS5) (Table 2).


Consummation

However, we found reduced activation in the FH group compared with the control group for the unpleasant taste in the pallidum/putamen region (Fig. 3; Table 2) the superior frontal gyrus (SFG) and the dorsal anterior cingulate cortex (dACC) (Fig. DS6). We found no differences between the groups for the chocolate taste.

**Table 2** Group differences in neural responses to reward and aversion

Brain region	x	y	z	Z-score	P
<i>Anticipatory</i>					
Chocolate cue: FH > controls					
Caudate	4	-6	6	3.43	0.004
Unpleasant cue: FH > controls					
Planum polare/amygdala	40	-6	-20	3.44	<0.001
Caudate	4	-6	6	3.36	<0.001
Thalamus	-2	-10	10	3.37	0.005
<i>Consummatory</i>					
Unpleasant taste: controls > FH					
Superior frontal gyrus	16	32	62	3.11	0.02
Pallidum/putamen	20	-2	6	2.86	0.02
dACC	10	4	28	2.96	0.02 <sup>a</sup>

a.  $P>0.05$  with SHAPS as a covariate. Coordinates are defined in the Montreal Neurological Institute (MNI) stereotaxic space. *P* values whole brain fully corrected at the cluster level (FWE  $P<0.05$  family wise error for multiple comparisons).



**Fig. 2** Unpleasant cue: (a) axial, sagittal and coronal image of increased amygdala activation in FH v. control group; (b) contrast estimates centred at  $-30, -2, -20$  (amygdala) in FH and control groups.


### Discussion

In this study, we examined the neural response during the anticipation and consumption of rewarding and aversive food stimuli in those with a family history of AN, by virtue of having a sister with AN, compared with controls with no family history. We found that there were increased activations to the anticipation of both rewarding and aversive food but decreased responses to the consumption of aversive food.

Specifically, we found that the FH group had increased caudate, thalamus and amygdala activations to the anticipation of both pleasant and unpleasant food. These results are consistent with our previous study which found that females with a personal history of AN also had increased neural responses during the anticipation of food.<sup>16</sup> As discussed in a recent review by Zhu *et al*, examining mostly visual food stimuli in AN, increased activation such as this could be related to increased emotional arousal in response to food.<sup>2</sup> Brain regions such as the caudate are thought to be involved in valence and salience coding of rewards, and have been found more active in weight-concerned women when choosing high-energy food from a selection of pictures<sup>36</sup> and in a meta-analysis of studies on food cue processing in AN patients suggested to be coding the

emotion of disgust.<sup>2</sup> Therefore, our result of increased activity in these regions during anticipation of food might suggest a mechanism for increased negative emotional reactions in those at risk of eating disorders. In fact, consistent with this notion, the FH group reported wanting the food less after seeing the cue than did the control participants, supporting a negative emotional response during the anticipatory phase; however, further exploratory correlational analysis did not find any relationship in the FH or control group between brain responses in the amygdala and liking or wanting ratings for the aversive taste and cue. This may be due both to the volunteers being healthy and a small sample size.

The increased amygdala response in our study during the anticipation phase of the unpleasant cue in the FH group, compared with the controls, might be due to increased disgust as amygdala activation has been found to correlate with disgust ratings in AN patients when confronted with food stimuli<sup>37</sup> and is known to be activated by aversive stimuli from multiple sensory modalities.<sup>38</sup> Taken together, our results show that even in those ‘at risk’ of AN but with no current symptoms or history of starvation, there are hyperactivations to the anticipation of food, supporting the idea that participants with AN may be highly responsive to food cues, perhaps as a mechanism to predict and control the anxiety



**Fig. 3** Unpleasant taste: (a) axial, sagittal and coronal image of decreased pallidum activation in FH v. control group; (b) contrast estimates centred at  $20, -2, 6$  (pallidum) in FH and control groups.

produced by food that may be associated with subjective unpleasantness.<sup>39,40</sup>

Interestingly, during the consummation phase of the task we found reduced neural responses in the FH group compared with the control group specifically for the unpleasant taste and in the dACC, pallidum/putamen and SFG. The ACC has been implicated in the coding of sensory-hedonic responses to taste and innervates the striatum, where behavioural responses are then computed.<sup>39</sup> Thus, reduced activity in this region in the FH group might indicate a mechanism by which there is a vulnerability towards reduced liking of food in AN; this was also supported by the behavioural responses of reduced liking for the chocolate in the FH group. Furthermore, as the SFG is involved in self-awareness and rumination in eating disorders<sup>41,42</sup> and the pallidum has been identified as a 'hedonic hotspot' with a pivotal role for food liking in the brain,<sup>9</sup> it seems that diminished responses in these regions in the FH group could explain a mechanistic predisposition towards reduced pleasure from food tastes.

As no differences between the FH group and controls were found in the activations of the primary taste cortex (anterior insula), our results are unlikely to be related to differences in physical sensitivity to taste and oral texture between the groups. Interestingly, in our previous study in those 'at risk' of anorexia, namely recovered AN patients, we found similar regions activated to the aversive taste such as the putamen and dACC. However, unlike the previous study of increased responses in those recovered, we find decreased response in the FH group. Therefore, we suggest that decreased activity in these regions to the unpleasant tastes is more like a trait marker which might only switch to hypersensitivity (increased activation) after experiencing AN (state) and is perhaps modulated by the amount of starvation experienced. Further studies, longitudinal in nature, are of course needed to clarify this.

Taken together our results suggest that the neural response during anticipatory and consummatory responses to rewarding and aversive food stimuli might be a biomarker for risk of developing AN in the future. Therefore, interventions with those at risk of AN that can normalise these differences in brain activity may be helpful in preventing the disorder onset. To clarify this, longitudinal studies investigating the neurobiological profiles of adolescents and young women at risk of AN are needed.

**Stefanie Horndasch**, MD, School of Psychology and Clinical Language Sciences, University of Reading, Reading, UK, and Department of Child and Adolescent Mental Health, University of Erlangen-Nuremberg, Erlangen, Germany; **Sophie O'Keefe**, MSc, School of Psychology and Clinical Language Sciences, University of Reading, Reading, UK; **Anneka Lamond**, MSc, School of Psychology and Clinical Language Sciences, University of Reading, Reading, UK; **Katie Brown**, MSc, School of Psychology and Clinical Language Sciences, University of Reading, Reading, UK; **Ciara McCabe**, PhD, School of Psychology and Clinical Language Sciences, University of Reading, Reading, UK

**Correspondence:** Ciara McCabe, School of Psychology and Clinical Language Sciences, University of Reading, Reading RG6 6AL, UK. Email: c.mccabe@reading.ac.uk

First received 8 Dec 2015, final revision 28 May 2016, accepted 24 Jun 2016

## Acknowledgements

We thank Ewelina Rzepa for helping with the scanning.

## Funding

This study was supported by the University of Reading start-up fund for C.M.

## References

- Kaye WH, Fudge JL, Paulus M. New insights into symptoms and neurocircuit function of anorexia nervosa. *Nat Rev Neurosci* 2009; **10**: 573–84.
- Zhu Y, Hu X, Wang J, Chen J, Guo Q, Li C, et al. Processing of food, body and emotional stimuli in anorexia nervosa: a systematic review and meta-analysis of functional magnetic resonance imaging studies. *Eur Eat Disord Rev* 2012; **20**: 439–50.
- Giel KE, Wittorf A, Wolkenstein L, Klingberg S, Drimmer E, Schonenberg M, et al. Is impaired set-shifting a feature of 'pure' anorexia nervosa? Investigating the role of depression in set-shifting ability in anorexia nervosa and unipolar depression. *Psychiatry Res* 2012; **200**: 538–43.
- Sanders N, Smeets PA, van Elburg AA, Danner UN, van Meer F, Hoek HW, et al. Altered food-cue processing in chronically ill and recovered women with anorexia nervosa. *Front Behav Neurosci* 2015; **9**: 46.
- Wagner A, Aizenstein H, Mazurkewicz L, Fudge J, Frank GK, Putnam K, et al. Altered insula response to taste stimuli in individuals recovered from restricting-type anorexia nervosa. *Neuropsychopharmacology* 2008; **33**: 513–23.
- Vocks S, Herpertz S, Rosenberger C, Senf W, Gizewski ER. Effects of gustatory stimulation on brain activity during hunger and satiety in females with restricting-type anorexia nervosa: an fMRI study. *J Psychiatr Res* 2011; **45**: 395–403.
- Frank GK, Reynolds JR, Shott ME, Jappe L, Yang TT, Tregellas JR, et al. Anorexia nervosa and obesity are associated with opposite brain reward response. *Neuropsychopharmacology* 2012; **37**: 2031–46.
- Berridge KC. 'Liking' and 'wanting' food rewards: brain substrates and roles in eating disorders. *Physiol Behav* 2009; **97**: 537–50.
- Berridge KC, Robinson TE, Aldridge JW. Dissecting components of reward: 'liking', 'wanting', and learning. *Curr Opin Pharmacol* 2009; **9**: 65–73.
- Wheeler RA, Carelli RM. The neuroscience of pleasure. Focus on 'ventral pallidum firing codes hedonic reward: when a bad taste turns good'. *J Neurophysiol* 2006; **96**: 2175–6.
- Pecina S, Berridge KC. Hedonic hot spot in nucleus accumbens shell: where do mu-opioids cause increased hedonic impact of sweetness? *J Neurosci* 2005; **25**: 11777–86.
- Smith KS, Berridge KC. The ventral pallidum and hedonic reward: neurochemical maps of sucrose 'liking' and food intake. *J Neurosci* 2005; **25**: 8637–49.
- Berridge KC, Kringelbach ML. Affective neuroscience of pleasure: reward in humans and animals. *Psychopharmacology (Berl)* 2008; **199**: 457–80.
- Berridge KC. The debate over dopamine's role in reward: the case for incentive salience. *Psychopharmacology (Berl)* 2007; **191**: 391–431.
- Dayan P, Balleine BW. Reward, motivation, and reinforcement learning. *Neuron* 2002; **36**: 285–98.
- Cowdrey FA, Park RJ, Harmer CJ, McCabe C. Increased neural processing of rewarding and aversive food stimuli in recovered anorexia nervosa. *Biol Psychiatry* 2011; **70**: 736–43.
- Yau WY, Bischoff-Grethe A, Theilmann RJ, Torres L, Wagner A, Kaye WH, et al. Alterations in white matter microstructure in women recovered from anorexia nervosa. *Int J Eat Disord* 2013; **46**: 701–8.
- Bulik CM, Sullivan PF, Tozzi F, Furberg H, Lichtenstein P, Pedersen NL. Prevalence, heritability, and prospective risk factors for anorexia nervosa. *Arch Gen Psychiatry* 2006; **63**: 305–12.
- Strober M, Freeman R, Lampert C, Diamond J, Kaye W. Controlled family study of anorexia nervosa and bulimia nervosa: evidence of shared liability and transmission of partial syndromes. *Am J Psychiatry* 2000; **157**: 393–401.
- Steinhausen HC, Jakobsen H, Helenius D, Munk-Jorgensen P, Strober M. A nationwide study of the family aggregation and risk factors in anorexia nervosa over three generations. *Int J Eat Disord* 2015; **48**: 1–8.
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders* (5th edn) (DSM-5). APA, 2013.
- Stice E, Telch CF, Rizvi SL. Development and validation of the Eating Disorder Diagnostic Scale: a brief self-report measure of anorexia, bulimia, and binge-eating disorder. *Psychol Assess* 2000; **12**: 123–31.
- First MB, Spitzer RL, Gibbon M, Williams JBW. *Structured Clinical Interview for DSM-IV-TR Axis I Disorders, Research Version, Non-patient Edition (SCID-I/NP)*. Biometrics Research, New York State Psychiatric Institute, 2002.
- Beck AT, Steer RA, Brown GK. *Manual for the Beck Depression Inventory-II*. Psychological Corporation, 1996.
- Fawcett J, Clark DC, Scheftner WA, Gibbons RD. Assessing anhedonia in psychiatric patients: The Pleasure Scale. *Arch Gen Psychiatry* 1983; **40**: 79–84.
- Snaith RP, Hamilton M, Morley S, Humayan A, Hargreaves D, Trigwell P. A scale for the assessment of hedonic tone: The Snaith–Hamilton Pleasure Scale. *Br J Psychiatry* 1995; **167**: 99–103.
- Brinker JK, Dozois DJ. Ruminative thought style and depressed mood. *J Clin Psychol* 2009; **65**: 1–19.
- Gard DE, Gard MG, Kring AM, John OP. Anticipatory and consummatory components of the experience of pleasure: a scale development study. *J Res Pers* 2006; **40**: 1086–102.



- 29 Jorm AF, Christensen H, Henderson AS, Jacomb PA, Korten AE, Rodgers B. Using the BIS/BAS scales to measure behavioural inhibition and behavioural activation: factor structure, validity and norms in a large community sample. *Pers Ind Diff* 1998; **26**: 49–58.
- 30 Cowdrey FA, Park RJ. Assessing rumination in eating disorders: principal component analysis of a minimally modified ruminative response scale. *Eat Behav* 2011; **12**: 321–4.
- 31 Garner DM, Olmsted MP, Bohr Y, Garfinkel PE. The Eating Attitudes Test: psychometric features and clinical correlates. *Psychol Med* 1982; **12**: 871–8.
- 32 Nelson HE. *The National Adult Reading Test (NART): Test Manual*. NFER-Nelson, 1982.
- 33 Zerssen D, Petermann F. *Bf-SR: Die Befindlichkeits-Skala - Revidierte Fassung* [Adjective Mood Scale – Revised version (AMS-R)]. Hogrefe, 2011.
- 34 McCabe C, Mishor Z, Cowen PJ, Harmer CJ. Diminished neural processing of aversive and rewarding stimuli during selective serotonin reuptake inhibitor treatment. *Biol Psychiatry* 2010; **67**: 439–45.
- 35 Dean Z, Horndasch S, Giannopoulos P, McCabe C. Enhanced neural response to anticipation, effort and consummation of reward and aversion during bupropion treatment. *Psychol Med* 18 May 2016 (doi: 10.1017/S003329171600088X).
- 36 van der Laan LN, de Ridder DT, Vieregger MA, Smeets PA. Activation in inhibitory brain regions during food choice correlates with temptation strength and self-regulatory success in weight-concerned women. *Front Neurosci* 2014; **8**: 308.
- 37 Joos AA, Saum B, van Elst LT, Perlov E, Glauche V, Hartmann A, et al. Amygdala hyperreactivity in restrictive anorexia nervosa. *Psychiatry Res* 2011; **191**: 189–95.
- 38 Zald DH. The human amygdala and the emotional evaluation of sensory stimuli. *Brain Res Brain Res Rev* 2003; **41**: 88–123.
- 39 Kaye WH, Wierenga CE, Bailer UF, Simmons AN, Bischoff-Grethe A. Nothing tastes as good as skinny feels: the neurobiology of anorexia nervosa. *Trends Neurosci* 2013; **36**: 110–20.
- 40 Kaye WH, Wierenga CE, Bailer UF, Simmons AN, Wagner A, Bischoff-Grethe A. Does a shared neurobiology for foods and drugs of abuse contribute to extremes of food ingestion in anorexia and bulimia nervosa? *Biol Psychiatry* 2013; **73**: 836–42.
- 41 Goldberg II, Harel M, Malach R. When the brain loses its self: prefrontal inactivation during sensorimotor processing. *Neuron* 2006; **50**: 329–39.
- 42 Brooks SJ, O'Daly O, Uher R, Friederich H-C, Giampietro V, Brammer M, et al. Thinking about eating food activates visual cortex with reduced bilateral cerebellar activation in females with anorexia nervosa: an fMRI study. *PLoS One* 2012; **7**: e34000.

