


Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

## Travel Medicine and Infectious Disease

journal homepage: [www.elsevier.com/locate/tmaid](https://www.elsevier.com/locate/tmaid)

Original article

## Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and Middle East Respiratory Syndrome Coronavirus (MERS-CoV) coinfection: A unique case series

Alyaa Elhazmi<sup>a</sup>, Jaffar A. Al-Tawfiq<sup>b,c,d,\*</sup>, Hend Sallam<sup>e</sup>, Awad Al-Omari<sup>f,g</sup>, Saad Alhumaid<sup>h</sup>, Ahmad Mady<sup>i,j</sup>, Abbas Al Mutair<sup>k,l,m</sup><sup>a</sup> Adult Critical Care Department, Dr. Sulaiman Alhabib Medical Group, Riyadh, Saudi Arabia<sup>b</sup> Infectious Disease Unit, Specialty Internal Medicine, Johns Hopkins Aramco Healthcare, Dhahran, Saudi Arabia<sup>c</sup> Infectious Disease Division, Department of Medicine, Indiana University School of Medicine, Indianapolis, IN, USA<sup>d</sup> Infectious Disease Division, Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, MD, USA<sup>e</sup> Adult Critical Care Medicine, King Faisal Specialist Hospital and Research Centre, Riyadh, Saudi Arabia<sup>f</sup> Research Center, Dr Sulaiman Al Habib Medical Group, Riyadh, Saudi Arabia<sup>g</sup> College of Medicine, Alfaisal University, Riyadh, Saudi Arabia<sup>h</sup> Administration of Pharmaceutical Care, Ministry of Health, Al-Ahsa, Saudi Arabia<sup>i</sup> Critical Care Department, King Saud Medical City, Riyadh, Saudi Arabia<sup>j</sup> Department of Anesthesiology and ICU, Tanta University Hospitals, Tanta, Egypt<sup>k</sup> Research Center, Almoosa Specialist Hospital, Al-Hasa, Saudi Arabia<sup>l</sup> College of Nursing, Prince Nora University, Riyadh, Saudi Arabia<sup>m</sup> School of Nursing, Wollongong University, Australia

## ARTICLE INFO

## Keywords:

COVID-19

SARS-CoV-2

MERS-CoV

Coinfection

Middle East Respiratory Syndrome Coronavirus

## ABSTRACT

**Introduction:** The emergence of the Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) had raised possibilities of coinfection with the Middle East Respiratory Syndrome Coronavirus (MERS-CoV) in countries where these two viruses were reported. In this study, we describe the clinical presentation and demographics of eight patients who were coinfecting with SARS-CoV-2 and MERS-CoV.

**Materials and methods:** This is a case series of hospitalized patients admitted to intensive care units (ICUs). We collected demographics, underlying conditions, presenting symptoms and clinical outcome from the patients' medical records.

**Results:** During the study period from March 14, 2020 to October 19, 2020, there was a total of 67 SARS-CoV-2 ICU admitted patients who underwent simultaneous SARS-CoV-2 and MERS-CoV testing by PCR. Of those patients, 8 (12%) tested positive for both SARS-CoV-2 and MERS-CoV. There were 6 (75%) males, the mean age  $\pm$  SD was  $44.4 \pm 11.8$  years, and 7 (87.5%) were obese. Of the patients, 7 (87.5%) were non-smokers, 1 (12.5%) had diabetes mellitus, 1 (12.5%) had heart failure, and 1 (12.5%) had been on anti-platelet therapy. The mean hospital length of stay (LOS) was  $21.1 \pm 11.6$  days and the average ICU LOS was  $10.9 \pm 6.03$  days. All patients received supportive therapy and all were treated with corticosteroid. Of all the patients, 4 (50%) were discharged home and 3 (37.5%) died.

**Conclusion:** This case series is an important addition to the medical knowledge as it showed the interaction of the coinfection of SARS-CoV-2 and MERS-CoV.

## 1. Introduction

The Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-

2) emerged at the end of 2019 and had caused the current Coronavirus Disease 19 (COVID-19) pandemic. COVID-19 had been associated with tremendous medical challenges as well as societal and industrial

\* Corresponding author. P.O. Box 76; Room A-428-2, Building 61, Dhahran Health Center, Saudi Aramco, Dhahran, 31311, Saudi Arabia.

E-mail addresses: [a.m.haz@live.com](mailto:a.m.haz@live.com) (A. Elhazmi), [jaffar.tawfiq@jhah.com](mailto:jaffar.tawfiq@jhah.com), [jaltawfi@yahoo.com](mailto:jaltawfi@yahoo.com) (J.A. Al-Tawfiq), [Hsallam@kfschrc.edu.sa](mailto:Hsallam@kfschrc.edu.sa) (H. Sallam), [saalhumaid@moh.gov.sa](mailto:saalhumaid@moh.gov.sa) (S. Alhumaid).

<https://doi.org/10.1016/j.tmaid.2021.102026>

Received 17 January 2021; Received in revised form 26 February 2021; Accepted 9 March 2021

Available online 13 March 2021

1477-8939/© 2021 Elsevier Ltd. All rights reserved.

interruption [1]. COVID-19 clinical presentation ranges from asymptomatic to severe disease [2–6]. And the case fatality rate of COVID-19 is variable [7,8]. The clinical characteristics of the disease in Saudi Arabia is similar to other reported presentations [4–6,8,9].

Recently, there had been few reports of coinfection of different micro-organisms with SARS-CoV-2. Coinfections with SARS-CoV-2 were reported with influenza, other respiratory pathogens, herpes simplex virus (HSV) and human immune deficiency virus (HIV) [10–15]. In the Kingdom of Saudi Arabia (KSA), the Middle East Respiratory Syndrome Coronavirus (MERS-CoV) is of particular concern as KSA is the country where MERS-CoV had been initially identified in 2012 [16–18]. One particular characteristic of MERS-CoV is the ability to cause healthcare associated outbreaks [19–21].

In addition, MERS-CoV is of particular importance as it has about 35% case fatality rate [22].

The first case of COVID-19 was reported in Saudi Arabia on March 2nd, 2020 [6,9]. However, we are not aware of any reports of coinfection of SARS-CoV-2 and MERS-CoV. In an initial cohort of 99 patients of SARS-CoV-2, none of them had coinfection with MERS-CoV [23]. Here, we describe the clinical and demographics of patients who were coinfecting with SARS-CoV-2 and MERS-CoV.

## 2. Materials and methods

This is a case series of hospitalized patients admitted to intensive care units (ICUs) from March 14, 2020 to October 19, 2020. We collected demographics, underlying conditions, presenting symptoms and clinical outcome from the patients' medical records. Nasopharyngeal swabs were tested for the presence of MERS-CoV *upE* and *orf1a* [24,

25] and for SARS-CoV-2 *E* and *S* genes using the RealStar® SARS-CoV-2 real-time Reverse Transcriptase PCR (RT-PCR) as described previously [6,23,26]. The study was approved by the IRB of the Central IRB of Ministry of Health (Log No: 20-80E).

## 3. Results

### 3.1. Case summaries

During the study period, there was a total of 67 SARS-CoV-2 patients who were admitted to the ICUs and underwent simultaneous SARS-CoV-2 and MERS-CoV testing by PCR at the discretion of the treating physicians. Of those patients, 8 (12%) tested positive for both SARS-CoV-2 and MERS-CoV.

### 3.2. Demographics and clinical characteristics

Of the eight patients, there were 6 (75%) males. The mean age  $\pm$ SD was  $44.4 \pm 11.8$  years and the mean Body Mass Index (BMI) was  $31 \pm 4.7$ . Based on BMI, 1 (12.5%) had normal BMI and 7 (87.5%) were obese and of these 2 (25%) were overweight (BMI 25–29.9), 3 (36.5%) were obese (BMI: 30–34.9), and 2 (25%) had extreme obesity (BMI: > 35) (Table 1).

One patient (12.5%) was Saudi and 1 (12.5%) was a healthcare worker. Of the patients, 7 (87.5%) were non-smokers, 1 (12.5%) had diabetes mellitus, 1 (12.5%) had heart failure, and 1 (12.5%) had anti-platelet therapy. None of the patients had hypertension, ischemic heart disease, chronic obstructive pulmonary disease (COPD), asthma, liver disease; hemoglobinopathy; chronic kidney disease (CKD); renal

**Table 1**  
Characteristics, clinical presentation and outcome of ICU MERS-CoV and SARS-CoV-2 coinfecting patients.

Patient Number	1	2	3	4	5	6	7	8
ICU LOS (days)	18	7	12	21	8	3	11	7
Hospital LOS (days)	18	19	39	31	29	4	20	9
Age (years)	48	36	36	39	33	52	69	42
Gender	Male	male	male	female	male	male	female	male
Was patient Saudi or non-Saudi?	non-Saudi	non-Saudi	non-Saudi	non-Saudi	non-Saudi	non-Saudi	non-Saudi	Saudi
BMI	32.8	38.1	31.9	23.9	29.0	31.1	35.2	26.0
Was the case in close contact with a person with fever and/or cough who has been to an affected area?	not indicated	not indicated	not indicated	not indicated	not indicated	No	No	Yes
Current smoker	No	no	no	no	unknown	no	no	no
Diabetes mellitus	No	no	no	no	no	no	yes	no
Hypertension	no	no	no	no	no	no	no	no
Ischemic heart disease	no	no	no	no	no	no	no	no
Heart failure (ejection fraction < 50%)	no	no	no	no	no	no	yes	no
Chronic Lung disease	no	no	no	no	unknown	no	no	no
Duration of symptoms prior to hospital admission	1	4	4	2	2	asymptomatic	4	4
Shortness of breath (Dyspnoea)	yes	yes	yes	yes	yes	yes	yes	yes
Runny nose (rhinorrhoea)	yes	unknown	no	no	unknown	no	yes	no
Gastrointestinal symptoms (Diarrhea/vomiting)	yes	unknown	no	no	unknown	yes	yes	yes
History of fever	yes	yes	yes	yes	yes	yes	no	yes
Altered consciousness/confusion	no	unknown	no	no	unknown	no	no	yes
Cough	yes	yes	yes	yes	yes	yes	no	yes
Hospital Admission Mean BP (mmHg)	101	85		73	129	82	109	91
Heart rate (beats/minute)	86	85		103	110	104	96	90
Respiratory rate (breaths/minute)	32	22		24	20	20	25	17
Oxygen saturation (on room air) (%)	75	81		88	96	88	94	98
Temperature (highest within the first 24 h) (°C)	38	38.3		38.1	39	39.6	37.2	38.6
Glasgow Coma Score (GCS)	15	15		15	10	15	15	15
Chest X-ray - Consolidation	yes	no	yes	yes	yes	no	no	yes
CXR infiltrate on admission	bilateral	no	bilateral	bilateral	bilateral	no	no	bilateral
ICU admission PH value	7.28	7.415	7.37	7.44	7.411		7.49	
PCO2 value (mmHg)	55.9	41.1	36.1	39.9	36.6		51	
PaO2 value (mmHg)	95.9	61.8	207	60	89		71	
O2 Saturation (%)	96.6	91.7	99.8	93	97.7		99	91
Intubated and mechanically ventilated	yes	yes	yes	yes	yes	yes	yes	no
Hospital discharge outcome	death	home	death	home	home	transferred out	death	home

ICU: intensive care unit; LOS: length of stay; BMI: body mass index; CXR: chest x-ray.

replacement; immunocompromised; transplant; malignancy; or recent surgery.

### 3.3. Symptoms

All patients had shortness of breath, 2 (25%) had runny nose, 4 (50%) had gastrointestinal (GI) symptoms, and 7 (87.5%) had history of fever. Altered consciousness/confusion was present in 1 (12.5%), cough in 7 (87.5%), abdominal pain in 6 (75%), headache in 1 (12.5%), chest pain in 1 (12.5%), arthralgia, myalgia, and fatigue each was present in 12.5%, and sore throat in 3 (37.5%).

### 3.4. Hospital course

The initial mean and SD of laboratory tests are shown in Table 2. The mean hospital length of stay (LOS) was  $21.1 \pm 11.6$  days, the average ICU LOS was  $10.9 \pm 6.03$  days. The initial mode of oxygen delivery was face mask (1; 12.5%), non-rebreather mask (NRM) (4; 50%), and mechanical ventilation (3; 37.5%). However, eventually 7 (87.5%) required mechanical ventilation and the average duration of mechanical ventilation was  $6.67 (\pm 5.9)$  days. All patients received supportive therapy and all were treated with corticosteroid. The patients did not receive specific anti-viral therapy. Of all the patients, 1 (12.5%) was transferred to another facility, 4 (50%) were discharged home and 3 (37.5%) died.

## 4. Discussion

This case series is a unique contribution to the medical literature by examining coinfection of two emerging respiratory pathogens, MERS-CoV and SARS-CoV-2. These two viruses had the chance to cause significant disease in the Kingdom of Saudi Arabia. The country had gained great experience from previous MERS-CoV infection and had adopted a significant infection control and preventative measures [27]. Coinfection of SARS-CoV-2 and other respiratory viruses had been reported [10–15]. In addition, there few cases of coinfection of MERS-CoV and influenza or tuberculosis [28,29]. Here, we report coinfection of MERS-CoV and SARS-CoV-2 in admitted patients. In an initial cohort of 99 cases who were tested simultaneously for MERS-CoV and SARS-CoV-2, none of them had SARS-CoV-2 and MERS-CoV coinfection [23]. In the current study, 12% of the tested patients were positive for both SARS-CoV-2 and MERS-CoV. It was reported that in the past few years, MERS-CoV positivity rates among suspected cases was 2–6% [30]. The difference in these two estimates is likely related to the population of patients being tested.

The included patients were relatively young with a mean age  $\pm$ SD of  $44.4 \pm 11.8$  years. Previously, the mean age of COVID-19 patients in Saudi Arabia was 36–50 years [6,31] and a median age 44 years [23]. The presence of comorbidities in patients with SARS-CoV-2 ranges from

**Table 2**  
Initial mean and SD of laboratory tests of patients coinfecting with SARS-CoV-2 and MERS-CoV.

Laboratory (normal range)	Mean and Standard Deviation of Data on Hospital Admission
Hgb (13–17 mg/dl)	13.12 (1.96)
WBC ( $4-11 \times 10^9/L$ )	8.29 (4.03)
Absolute Lymphocyte ( $0.90-2.90 \times 10^9/L$ )	1.55 (1.23)
Absolute Neutrophil ( $1.70-7 \times 10^9/L$ )	9.13 (3.26)
Neutrophils/Lymphocyte ratio (1–3)	10.14 (8.77)
Platelets ( $150-450 \times 10^9/L$ )	247.4 (76.9)
aPTT (30–40 s)	32.23 (5.56)

Hgb: hemoglobin; WBC: white blood cells; aPTT: activated partial thromboplastin time.

32% to 93% in different studies [32–34]. One half of the included patients had gastrointestinal symptoms. Previously, gastrointestinal symptoms were reported in 33% of MERS-CoV patients [24,25,35–38], and 29% of COVID-19 patients had gastrointestinal symptoms [39] and another study showed higher rate of diarrhea in elderly patients [40]. In addition, the majority of hospitalized MERS-CoV patients had fever, cough and shortness of breath in addition to pneumonia on chest x-ray [24,25,35,41].

This case series showed that 37.5% of ICU coinfecting patients had died. Previous studies of MERS-CoV showed a case fatality rate of 30% with increasing rate among those with comorbidities [24], critical ill patients, severe disease, those > 65 years of age, hospital-acquired infections and corticosteroid use [35,42–48]. However, the use of corticosteroid was associated with decreased case fatality rate in randomized controlled trials of COVID-19 patients from 25.7% in standard therapy to 22.9% in the dexamethasone group [49]. In addition, there is a variation in the case fatality rates among different countries and different time intervals and ranged from 60% in March 2020 to 42% in May 2020 [50]. The epidemiology of MERS-CoV infection in Saudi Arabia is well characterized by sporadic cases from camel exposure followed by either hospital [19–21] or community transmission [51–53]. However, we were not able to pinpoint the exact exposure of infection of the included patients.

The initial laboratory findings of the included patients showed normal WBC and lymphocyte counts. However, the mean  $\pm$  SD of the neutrophils to lymphocyte count ratio (NLCR) was 10.14 (8.77). The NLCR of  $\geq 3.13$  is a predictor of ICU admission and a marker of disease severity [32]. A previous study from Saudi Arabia showed that NLCR was significantly higher among ICU admissions than other patients [6]. We did not examine the duration of viral shedding in this study. A previous meta-analysis showed that the mean shedding duration of MERS-CoV was 15.3 days in the upper respiratory tract and 16.3 days in the lower respiratory tract. On the other hand, in COVID-19 patients the shedding of SARS-CoV-2 had a mean duration of 17 days in upper respiratory tract and 14.6 days in lower respiratory tract [54]. It was stated that SARS-CoV-2 viral loads peak about 10 days after symptoms onset [55,56].

In conclusion, this case series is important as it showed the interaction of the coinfection of SARS-CoV-2 and MERS-CoV. However, in this small case-series the combined infection was not associated with increased risk of death in comparison with mono-infection with MERS-CoV. However, as there was one patient who is transferred to another facility and lost to follow-up and if that patient died then the mortality would be 50%. Thus, additional studies are required to confirm or refute this finding and it is not possible to draw a firm conclusion about the mortality in co-infected MERS-CoV and SARS-CoV-2 patients. The presenting symptoms and laboratory data are not different from each of the infection alone. Although, 7 (87.5%) of the coinfection of MERS-CoV and SARS-CoV-2 were in the ICU, this is related to the fact that all tested and included patients were ICU patients. Thus, a firm conclusion could not be drawn in regard to the epidemiology, clinical presentation and outcome of this type of coinfection. With the continued COVID-19 pandemic, there is a continued need to continue precautionary measures including social distancing, hand hygiene and universal masking. These measures continued to be enforced in the Kingdom of Saudi Arabia [9,57]. Moreover, KSA implemented universal masking in public and common areas as these measures are important for the control of COVID-19 [58,59].

### Funding source

None.

### CRediT authorship contribution statement

**Alyaa Elhazmi:** conceptual design, Data curation, Formal analysis,

Helped with the first draft of the manuscript, All authors finalized and approved the final draft of the manuscript. **Jaffar A. Al-Tawfiq**: conceptual design, Formal analysis, drafted the first manuscript. All authors finalized and approved the final draft of the manuscript. **Hend Sallam**: Data curation, Formal analysis, All authors finalized and approved the final draft of the manuscript. **Awad Al-Omari**: conceptual design, Data curation, Formal analysis, All authors finalized and approved the final draft of the manuscript. **Saad Alhumaid**: Data curation, Formal analysis, All authors finalized and approved the final draft of the manuscript. **Ahmad Mady**: Data curation, Formal analysis, All authors finalized and approved the final draft of the manuscript. **Abbas Al Mutair**: conceptual design, Data curation, Formal analysis, Helped with the first draft of the manuscript, All authors finalized and approved the final draft of the manuscript.

## Declaration of competing interest

None.

## References

- Al-Tawfiq JA, Al-Yami SS, Rigamonti D. Changes in healthcare managing COVID and non-COVID-19 patients during the pandemic: striking the balance. *Diagn Microbiol Infect Dis* 2020;98. <https://doi.org/10.1016/j.diagmicrobio.2020.115147>.
- Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;6736:1–9. [https://doi.org/10.1016/S0140-6736\(20\)30566-3](https://doi.org/10.1016/S0140-6736(20)30566-3).
- Nicastri E, D'Abramo A, Faggioni G, De Santis R, Mariano A, Lepore L, et al. Coronavirus disease (COVID-19) in a paucisymptomatic patient: epidemiological and clinical challenge in settings with limited community transmission, Italy. *Euro Surveill* 2020;25. <https://doi.org/10.2807/1560-7917.ES.2020.25.11.2000230>.
- Al-Omari A, Alhuqbani WN, Zaidi ARZ, Al-Subaie MF, AlHindi AM, Abogosh AK, et al. Clinical characteristics of non-intensive care unit COVID-19 patients in Saudi Arabia: a descriptive cross-sectional study. *J Infect Public Health* 2020. <https://doi.org/10.1016/j.jiph.2020.09.003>.
- Al Mutair A, Alhumaid S, Alhuqbani WN, Zaidi ARZ, Alkoraisi S, Al-Subaie MF, et al. Clinical, epidemiological, and laboratory characteristics of mild-to-moderate COVID-19 patients in Saudi Arabia: an observational cohort study. *Eur J Med Res* 2020;25:61. <https://doi.org/10.1186/s40001-020-00462-x>.
- AlJishi JM, Alhajjaj AH, Alkhabbaz FL, AlAbduljabar TH, Alsaif A, Alsaif H, et al. Clinical characteristics of asymptomatic and symptomatic COVID-19 patients in the Eastern Province of Saudi Arabia. *J Infect Public Health* 2021;14:6–11. <https://doi.org/10.1016/j.jiph.2020.11.002>.
- Tirupathi R, Muradova V, Shekhar R, Salim SA, Al-Tawfiq JA, Palabindala V. COVID-19 disparity among racial and ethnic minorities in the US: a cross sectional analysis. *Trav Med Infect Dis* 2020;38:101904. <https://doi.org/10.1016/j.tmaid.2020.101904>.
- Al-Tawfiq JA, Leonardi R, Fasoli G, Rigamonti D. Prevalence and fatality rates of COVID-19: what are the reasons for the wide variations worldwide? *Trav Med Infect Dis* 2020;35:101711. <https://doi.org/10.1016/j.tmaid.2020.101711>.
- Al-Tawfiq JA, Sattar A, Al-Khadra H, Al-Qahntani S, Al-Mulhim M, Al-Omouh O, et al. Incidence of COVID-19 among returning travelers in quarantine facilities: a longitudinal study and lessons learned. *Trav Med Infect Dis* 2020;38. <https://doi.org/10.1016/j.tmaid.2020.101901>.
- Mi T, Yq L, X C, H L, Zc J, Dl G, et al. Co-infection with common respiratory pathogens and SARS-CoV-2 in patients with COVID-19 pneumonia and laboratory biochemistry findings: a retrospective cross-sectional study of 78 patients from a single center in China. *Med Sci Mon Int Med J Exp Clin Res* 2021;27. <https://doi.org/10.12659/MSM.929783>.
- Thuy-Boun PS, Mehta S, Gruening B, McGowan T, Nguyen A, Rajczewski AT, et al. Metaproteomics analysis of SARS-CoV-2-infected patient samples reveals presence of potential coinfecting microorganisms. *J Proteome Res* 2021. <https://doi.org/10.1021/acs.jproteome.0c00822>.
- Hashemi SA, Safamanesh S, Zadeh-moghaddam HG, Ghafouri M, Azimian A. High prevalence of SARS-CoV-2 and influenza A virus (H1N1) coinfection in dead patients in Northeastern Iran. *J Med Virol* 2020. <https://doi.org/10.1002/jmv.26364>.
- Lew S, Manes P, Smith B. Coinfection with sars-cov-2 and influenza a virus in a 32-year-old man. *Am J Case Rep* 2020;21:1–5. <https://doi.org/10.12659/AJCR.926092>.
- Alharthy A, Faqih F, Noor A, Memish Z, Karakitsos D. Co-infection of human immunodeficiency virus, herpes simplex virus-2 and SARS-CoV-2 with false-negative real-time polymerase chain reaction. *Singap Med J* 2020. <https://doi.org/10.11622/smedj.2020158>.
- Byrd KM, Beckwith CG, Garland JM, Johnson JE, Aung S. Cu-Uvin S, et al. SARS-CoV-2 and HIV coinfection: clinical experience from Rhode Island, United States. *J Int AIDS Soc* 2020;23. <https://doi.org/10.1002/jia2.25573>.
- de Groot RJ, Baker SC, Baric RS, Brown CS, Drosten C, Enjuanes L, et al. Middle East respiratory syndrome coronavirus (MERS-CoV): announcement of the coronavirus study group. *J Virol* 2013;87:7790–2. <https://doi.org/10.1128/JVI.01244-13>.
- Zaki AM, van Boheemen S, Bestebroer TM, Osterhaus ADME, Fouchier RAM. Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med* 2012;367:1814–20. <https://doi.org/10.1056/NEJMoa1211721>.
- Corman VM, Eckerle I, Bleicker T, Zaki A, Landt O, Eschbach-Bludau M, et al. Detection of a novel human coronavirus by real-time reverse-transcription polymerase chain reaction. *Euro Surveill* 2012;17.
- Al-Tawfiq JA, Auwaerter PG. Healthcare-associated infections: the hallmark of Middle East respiratory syndrome coronavirus with review of the literature. *J Hosp Infect* 2019;101:20–9. <https://doi.org/10.1016/j.jhin.2018.05.021>.
- Hastings DL, Tokars JI, Abdel Aziz IZAM, Alkhalidi KZ, Bensadek AT, Alraddadi BM, et al. Outbreak of Middle East respiratory syndrome at tertiary care hospital, Jeddah, Saudi Arabia. *Emerg Infect Dis* 2014;22:794–801. <https://doi.org/10.3201/eid2205.151797>.
- Garout MA, Jokhdar HAA, Aljahdali IA, Zein AR, Goweda RA, Hassan-Hussein A. Mortality rate of ICU patients with the middle east respiratory syndrome – coronavirus infection at king Fahad hospital, Jeddah, Saudi Arabia. *Cent Eur J Publ Health* 2018;26:87–91. <https://doi.org/10.21101/cejph.a4764>.
- Al-Tawfiq JA, Memish ZA. Middle East respiratory syndrome coronavirus and severe Acute respiratory syndrome coronavirus. *Semin Respir Crit Care Med* 2020;41:568–78. <https://doi.org/10.1055/s-0040-1709160>.
- Barry M, AlMohaya AE, AlHijji A, Akkiel L, AlRajhi A, Almajid F, et al. Clinical characteristics and outcome of hospitalized COVID-19 patients in a MERS-CoV endemic area. *J Epidemiol Glob Health* 2020;10:214–21. <https://doi.org/10.2991/jegh.k.200806.002>.
- Assiri A, Al-Tawfiq JA, Al-Rabeeah AA, Al-Rabiah FA, Al-Hajjar S, Al-Barrak A, et al. Epidemiological, demographic, and clinical characteristics of 47 cases of Middle East respiratory syndrome coronavirus disease from Saudi Arabia: a descriptive study. *Lancet Infect Dis* 2013;13:752–61. [https://doi.org/10.1016/S1473-3099\(13\)70204-4](https://doi.org/10.1016/S1473-3099(13)70204-4).
- Al-Tawfiq JA, Hinedi K, Ghandour J, Khairalla H, Musleh S, Ujayli A, et al. Middle East Respiratory Syndrome-Coronavirus (MERS-CoV): a case-control study of hospitalized patients. *Clin Infect Dis* 2014;59:160–5. <https://doi.org/10.1093/cid/ciu226>.
- AlJishi JM, Al-Tawfiq JA. Intermittent viral shedding in respiratory samples of patients with SARS-CoV-2: observational analysis with infection control implications. *J Hosp Infect* 2020. <https://doi.org/10.1016/j.jhin.2020.09.011>.
- Al-Tawfiq JA, Garout MA, Gautret P. Preparing for emerging respiratory pathogens such as SARS-CoV, MERS-CoV, and SARS-CoV-2. *Le Inf Med* 2020;28:64–70.
- Alfaraj SH, Al-Tawfiq JA, Altuwaijri TA, Memish ZA. Middle East respiratory syndrome coronavirus and pulmonary tuberculosis coinfection: implications for infection control. *Intervirology* 2017;60:53–5. <https://doi.org/10.1159/000477908>.
- Alfaraj SH, Al-Tawfiq JA, Alzahrani NA, Altuwaijri TA, Memish ZA. The impact of co-infection of influenza A virus on the severity of Middle East Respiratory Syndrome Coronavirus. *J Infect* 2017;74:521–3. <https://doi.org/10.1016/j.jinf.2017.02.001>.
- Ebrahim SH. Lack of MERS-CoV co-infection among hospitalized COVID-19 patients in Saudi Arabia. *J Epidemiol Glob Health* 2020;10:191. <https://doi.org/10.2991/jegh.k.200819.001>.
- Alsafayan YM, Althunayyan SM, Khan AA, Hakawi AM, Assiri AM. Clinical characteristics of COVID-19 in Saudi Arabia: a national retrospective study. *J Infect Public Health* 2020;13:920–5. <https://doi.org/10.1016/j.jiph.2020.05.026>.
- Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5).
- Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in wuhan, China. *JAMA, J Am Med Assoc* 2020;323:1061–9. <https://doi.org/10.1001/jama.2020.1585>.
- Al Bahrani SY, Al-Tawfiq JA, Alshaer AR, Shilash A, Alswefy K, Al-Zayer RS, et al. A case series of severe hospitalized COVID-19 patients treated with tocilizumab and glucocorticoids: a report from Saudi arabian hospital. *J Epidemiol Glob Health* 2021. <https://doi.org/10.2991/jegh.k.210112.001>.
- Saad M, Omrani AS, Baig K, Bahloul A, Elzein F, Matin MA, et al. Clinical aspects and outcomes of 70 patients with Middle East respiratory syndrome coronavirus infection: a single-center experience in Saudi Arabia. *Int J Infect Dis* 2014;29:301–6. <https://doi.org/10.1016/j.ijid.2014.09.003>.
- Assiri A, McGeer A, Perl TM, Price CS, Al Rabeeah AA, Cummings DAT, et al. Hospital outbreak of Middle East respiratory syndrome coronavirus. *N Engl J Med* 2013;369:407–16. <https://doi.org/10.1056/NEJMoa1306742>.
- Arabi YM, Arifi AA, Balkhy HH, Najm H, Aldawood AS, Ghabashi A, et al. Clinical course and outcomes of critically ill patients with Middle East respiratory syndrome coronavirus infection. *Ann Intern Med* 2014;160:389–97. <https://doi.org/10.7326/M13-2486>.
- Shalhoub S, Farahat F, Al-Jiffri A, Simhairi R, Shamma O, Siddiqi N, et al. IFN- $\alpha$ 2a or IFN- $\beta$ 1a in combination with ribavirin to treat Middle East respiratory syndrome coronavirus pneumonia: a retrospective study. *J Antimicrob Chemother* 2015;70:2129–32. <https://doi.org/10.1093/jac/dkv085>.
- Docherty AB, Harrison EM, Green CA, Hardwick HE, Pius R, Norman L, et al. Features of 20 133 UK patients in hospital with covid-19 using the ISARIC WHO Clinical Characterisation Protocol: prospective observational cohort study. *BMJ* 2020;369. <https://doi.org/10.1136/bmj.m1985>.



- [40] Lechien JR, Chiesa-Estomba CM, Place S, Van Laethem Y, Cabaraux P, Mat Q, et al. Clinical and epidemiological characteristics of 1420 European patients with mild-to-moderate coronavirus disease 2019. *J Intern Med* 2020;288:335–44. <https://doi.org/10.1111/joim.13089>.
- [41] Fagbo SF, Skakni L, Chu DKW, Garbati MA, Joseph M, Peiris M, et al. Molecular epidemiology of hospital outbreak of Middle East respiratory syndrome, Riyadh, Saudi Arabia. *Emerg Infect Dis* 2014;21:1981. <https://doi.org/10.3201/eid2111.150944>, 2015.
- [42] Al-Tawfiq JA, Alfaraj SH, Altuwaijri TA, Memish ZA. A cohort-study of patients suspected for MERS-CoV in a referral hospital in Saudi Arabia. *J Infect* 2017;75:378–9. <https://doi.org/10.1016/j.jinf.2017.06.002>.
- [43] Ahmed AE. The predictors of 3- and 30-day mortality in 660 MERS-CoV patients. *BMC Infect Dis* 2017;17:615. <https://doi.org/10.1186/s12879-017-2712-2>.
- [44] Arabi YM, Mandourah Y, Al-Hameed F, Sindi AA, Almekhlafi GA, Hussein MA, et al. Corticosteroid therapy for critically ill patients with Middle East respiratory syndrome. *Am J Respir Crit Care Med* 2018;197:757–67. <https://doi.org/10.1164/rccm.201706-1172OC>.
- [45] Alfaraj SH, Al-Tawfiq JA, Assiri AY, Alzahrani NA, Alanazi AA, Memish ZA. Clinical predictors of mortality of Middle East Respiratory Syndrome Coronavirus (MERS-CoV) infection: a cohort study. *Trav Med Infect Dis* 2019. <https://doi.org/10.1016/j.tmaid.2019.03.004>.
- [46] Nam H-S, Park JW, Ki M, Yeon M-Y, Kim J, Kim SW. High fatality rates and associated factors in two hospital outbreaks of MERS in Daejeon, the Republic of Korea. *Int J Infect Dis* 2017. <https://doi.org/10.1016/j.ijid.2017.02.008>.
- [47] Kim KH, Tandil TE, Choi JW, Moon JM, Kim MS. Middle East respiratory syndrome coronavirus (MERS-CoV) outbreak in South Korea, 2015: epidemiology, characteristics and public health implications. *J Hosp Infect* 2017;95:207–13. <https://doi.org/10.1016/j.jhin.2016.10.008>.
- [48] Choi WS, Kang C-I, Kim Y, Choi J-P, Joh JS, Shin H-S, et al. Clinical presentation and outcomes of Middle East respiratory syndrome in the Republic of Korea. *Infect Chemother* 2016;48:118–26. <https://doi.org/10.3947/ic.2016.48.2.118>.
- [49] RECOVERY Collaborative Group. Dexamethasone in hospitalized patients with covid-19 — preliminary report. *N Engl J Med* 2020. <https://doi.org/10.1056/nejmoa2021436>.
- [50] Armstrong RA, Kane AD, Cook TM. Outcomes from intensive care in patients with COVID-19: a systematic review and meta-analysis of observational studies. *Anaesthesia* 2020;75:1340–9. <https://doi.org/10.1111/anae.15201>.
- [51] Omrani AS, Matin MA, Haddad Q, Al-Nakhli D, Memish ZA, Albarrak AM. A family cluster of middle east respiratory syndrome coronavirus infections related to a likely unrecognized asymptomatic or mild case. *Int J Infect Dis* 2013;17:e668–72. <https://doi.org/10.1016/j.ijid.2013.07.001>.
- [52] Memish Z a, Zumla AI, Al-Hakeem RF, Al-Rabeeh A a, Stephens GM. Family cluster of Middle East respiratory syndrome coronavirus infections. *N Engl J Med* 2013;368:2487–94. <https://doi.org/10.1056/NEJMoa1303729>.
- [53] Memish ZA, Cotten M, Watson SJ, Kellam P, Zumla A, Alhakeem RF, et al. Community case clusters of Middle East respiratory syndrome coronavirus in hafir Al-batin, kingdom of Saudi Arabia: a descriptive genomic study. *Int J Infect Dis* 2014;23:63–8. <https://doi.org/10.1016/j.ijid.2014.03.1372>.
- [54] Cevik M, Tate M, Lloyd O, Maraolo AE, Schafers J, Ho A. SARS-CoV-2, SARS-CoV, and MERS-CoV viral load dynamics, duration of viral shedding, and infectiousness: a systematic review and meta-analysis. *The Lancet Microbe* 2021;2:e13–22. [https://doi.org/10.1016/s2666-5247\(20\)30172-5](https://doi.org/10.1016/s2666-5247(20)30172-5).
- [55] Xiao AT, Tong YX, Zhang S. Profile of RT-PCR for SARS-CoV-2: a preliminary study from 56 COVID-19 patients. *Clin Infect Dis* 2020. <https://doi.org/10.1093/cid/ciaa460>.
- [56] Tirupathi R, Ramparas TR, Wadhwa G, Areti S, Kaur J, Salim S, et al. Viral dynamics in the upper respiratory tract (URT) of SARS-CoV-2. *Le Infez Med* 2020;28:486–99.
- [57] Al-Tawfiq JA, Memish ZA. COVID-19 in the eastern mediterranean region and Saudi Arabia: prevention and therapeutic strategies. *Int J Antimicrob Agents* 2020;55. <https://doi.org/10.1016/j.ijantimicag.2020.105968>.
- [58] Tirupathi R, Bharathidasan K, Palabindala V, Salim SA, Al-Tawfiq JA. Comprehensive review of mask utility and challenges during the COVID-19 pandemic. *Le Infez Med* 2020;28:57–63.
- [59] Liang M, Gao L, Cheng C, Zhou Q, Uy JP, Heiner K, et al. Efficacy of face mask in preventing respiratory virus transmission: a systematic review and meta-analysis. *Trav Med Infect Dis* 2020;36. <https://doi.org/10.1016/j.tmaid.2020.101751>.



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.