

brought to you by CORE

IUScholarWorks at Indiana University South Bend

Observation and Properties of the X(3872) Decaying to $J/\psi\pi+\pi-$ in p⁻p Collisions at $\sqrt{s}=1.96$ TeV

Abazov, V. M., Abbott, B., Abolins, M., Acharya, B. S., Adams, D. L., Adams, M., Adams, T., Agelou, M., Agram, J.-L., Ahmed, S. N., Ahn, S. H., Alexeev, G. D., Alkhazov, G., Alton, A., Alverson, G., Alves, G. A., Anderson, S., Andrieu, B., Arnoud, Y., Lynker, M. ... Zylberstejn, A.

To cite this post-print: Abazov, V. M., et al. "Observation and Properties of the X(3872) Decaying to $J/\psi\pi+\pi-$ in p⁻p Collisions at $\sqrt{s}=1.96$ TeV." *Phys. Rev. Lett.*, vol. 93, no. 16, American Physical Society, Oct. 2004, p. 162002, doi:10.1103/PhysRevLett.93.162002.

This document has been made available through IUScholarWorks repository, a service of the Indiana University Libraries. Copyrights on documents in IUScholarWorks are held by their respective rights holder(s). Contact <u>iusw@indiana.edu</u> for more information.

Observation and Properties of the X(3872) Decaying to $J/\psi \pi^+\pi^$ in $p\bar{p}$ Collisions at $\sqrt{s}= 1.96$ TeV

V.M. Abazov,³² B. Abbott,⁶⁹ M. Abolins,⁶⁰ B.S. Acharya,²⁶ D.L. Adams,⁶⁷ M. Adams,⁴⁷ T. Adams,⁴⁵ M. Agelou,¹⁶ J.-L. Agram,¹⁷ S.N. Ahmed,³¹ S.H. Ahn,²⁸ G.D. Alexeev,³² G. Alkhazov,³⁶ A. Alton,⁵⁹ G. Alverson,⁵⁸ G.A. Alves,² S. Anderson,⁴¹ B. Andrieu,¹⁵ Y. Arnoud,¹² A. Askew,⁷² B. Åsman,³⁷ C. Autermann,¹⁹ C. Avila,⁷ L. Babukhadia,⁶⁶ T.C. Bacon,³⁹ A. Baden,⁵⁶ S. Baffioni,¹³ B. Baldin,⁴⁶ P.W. Balm,³⁰ S. Banerjee,²⁶ E. Barberis,⁵⁸ P. Bargassa,⁷² P. Baringer,⁵³ C. Barnes,³⁹ J. Barreto,² J.F. Bartlett,⁴⁶ U. Bassler,¹⁵ D. Bauer,⁵⁰ A. Bean,⁵³ S. Beauceron,¹⁵ F. Beaudette,¹⁴ M. Begel,⁶⁵ S.B. Beri,²⁵ G. Bernardi,¹⁵ I. Bertram,³⁸ M. Besançon,¹⁶ A. Besson,¹⁷ R. Beuselinck,³⁹ V.A. Bezzubov,³⁵ P.C. Bhat,⁴⁶ V. Bhatnagar,²⁵ M. Bhattacharjee,⁶⁶ M. Binder,²³ A. Bischoff,⁴⁴ K.M. Black,⁵⁷ I. Blackler,³⁹ G. Blazey,⁴⁸ F. Blekman,³⁰ D. Bloch,¹⁷ U. Blumenschein,²¹ A. Boehnlein,⁴⁶ T.A. Bolton,⁵⁴ P. Bonamy,¹⁶ F. Borcherding,⁴⁶ G. Borissov,³⁸ K. Bos,³⁰ T. Bose,⁶⁴ C. Boswell,⁴⁴ A. Brandt,⁷¹ G. Briskin,⁷⁰ R. Brock,⁶⁰ G. Brooijmans,⁶⁴ A. Bross,⁴⁶ D. Buchholz,⁴⁹ M. Buehler,⁴⁷ V. Buescher,²¹ S. Burdin,⁴⁶ T.H. Burnett,⁷⁴ E. Busato,¹⁵ J.M. Butler,⁵⁷ J. Bystricky,¹⁶ F. Canelli,⁶⁵ W. Carvalho,³ B.C.K. Casey,⁷⁰ D. Casey,⁶⁰ N.M. Cason,⁵¹ H. Castilla-Valdez,²⁹ S. Chakrabarti,²⁶ D. Chakraborty,⁴⁸ K.M. Chan,⁶⁵ A. Chandra,²⁶ D. Chapin,⁷⁰ F. Charles,¹⁷ E. Cheu,⁴¹ L. Chevalier,¹⁶ D.K. Cho,⁶⁵ S. Choi,⁴⁴ S. Chopra,⁶⁷ T. Christiansen,²³ L. Christofek,⁵³ D. Claes,⁶²
A.R. Clark,⁴² C. Clément,³⁷ Y. Coadou,⁵ D.J. Colling,³⁹ L. Coney,⁵¹ B. Connolly,⁴⁵ W.E. Cooper,⁴⁶ D. Coppage,⁵³
M. Corcoran,⁷² J. Coss,¹⁸ A. Cothenet,¹³ M.-C. Cousinou,¹³ S. Crépé-Renaudin,¹² M. Cristetiu,⁴⁴
M.A.C. Cummings,⁴⁸ D. Cutts,⁷⁰ H. da Motta,² B. Davies,³⁸ G. Davies,³⁹ G.A. Davis,⁶⁵ K. De,⁷¹ P. de Jong,³⁰ S.J. de Jong,³¹ E. De La Cruz-Burelo,²⁹ C. De Oliveira Martins,³ S. Dean,⁴⁰ K. Del Signore,⁵⁹ F. Déliot,¹⁶ P.A. Delsart,¹⁸ M. Demarteau,⁴⁶ R. Demina,⁶⁵ P. Demine,¹⁶ D. Denisov,⁴⁶ S.P. Denisov,³⁵ S. Desai,⁶⁶ H.T. Diehl,⁴⁶ M. Diesburg,⁴⁶ M. Doidge,³⁸ H. Dong,⁶⁶ S. Doulas,⁵⁸ L. Duflot,¹⁴ S.R. Dugad,²⁶ A. Duperrin,¹³ J. Dyer,⁶⁰
A. Dyshkant,⁴⁸ M. Eads,⁴⁸ D. Edmunds,⁶⁰ T. Edwards,⁴⁰ J. Ellison,⁴⁴ J. Elmsheuser,²³ J.T. Eltzroth,⁷¹ V.D. Elvira,⁴⁶ S. Eno,⁵⁶ P. Ermolov,³⁴ O.V. Eroshin,³⁵ J. Estrada,⁴⁶ D. Evans,³⁹ H. Evans,⁶⁴ A. Evdokimov,³³ V.N. Evdokimov,³⁵ J. Fast,⁴⁶ S.N. Fatakia,⁵⁷ D. Fein,⁴¹ L. Feligioni,⁵⁷ T. Ferbel,⁶⁵ F. Fiedler,²³ F. Filthaut,³¹ H.E. Fisk,⁴⁶ F. Fleuret,¹⁵ M. Fortner,⁴⁸ H. Fox,²¹ W. Freeman,⁴⁶ S. Fu,⁶⁴ S. Fuess,⁴⁶ C.F. Galea,³¹ E. Gallas,⁴⁶ E. Galyaev,⁵¹ M. Gao,⁶⁴ C. Garcia,⁶⁵ A. Garcia-Bellido,⁷⁴ J. Gardner,⁵³ V. Gavrilov,³³ D. Gelé,¹⁷ R. Gelhaus,⁴⁴ K. Genser,⁴⁶ C.E. Gerber,⁴⁷ Y. Gershtein,⁷⁰ G. Geurkov,⁷⁰ G. Ginther,⁶⁵ K. Goldmann,²⁴ T. Golling,²⁰
B. Gómez,⁷ K. Gounder,⁴⁶ A. Goussiou,⁵¹ G. Graham,⁵⁶ P.D. Grannis,⁶⁶ S. Greder,¹⁷ J.A. Green,⁵² H. Greenlee,⁴⁶ Z.D. Greenwood,⁵⁵ E.M. Gregores,⁴ S. Grinstein,¹ J.-F. Grivaz,¹⁴ L. Groer,⁶⁴ S. Grünendahl,⁴⁶ M.W. Grünewald,²⁷ W. Gu,⁶ S.N. Gurzhiev,³⁵ G. Gutierrez,⁴⁶ P. Gutierrez,⁶⁹ A. Haas,⁷⁴ N.J. Hadley,⁵⁶ H. Haggerty,⁴⁶ S. Hagopian,⁴⁵ I. Hall,⁶⁹ R.E. Hall,⁴³ C. Han,⁵⁹ L. Han,⁴⁰ K. Hanagaki,⁴⁶ P. Hanlet,⁷¹ K. Harder,⁵⁴ J.M. Hauptman,⁵² R. Hauser,⁶⁰ C. Hays,⁶⁴ J. Hays,⁴⁹ C. Hebert,⁵³ D. Hedin,⁴⁸ J.M. Heinmiller,⁴⁷ A.P. Heinson,⁴⁴ U. Heintz,⁵⁷ C. Hensel,⁵³ G. Hesketh,⁵⁸ M.D. Hildreth,⁵¹ R. Hirosky,⁷³ J.D. Hobbs,⁶⁶ B. Hoeneisen,¹¹ M. Hohlfeld,²² S.J. Hong,²⁸ R. Hooper,⁵¹ S. Hou,⁵⁹ Y. Hu,⁶⁶ J. Huang,⁵⁰ Y. Huang,⁵⁹ I. Iashvili,⁴⁴ R. Illingworth,⁴⁶ A.S. Ito,⁴⁶ S. Jabeen,⁵³ M. Jaffré,¹⁴ S. Jain,⁶⁹ V. Jain,⁶⁷ K. Jakobs,²¹ A. Jenkins,³⁹ R. Jesik,³⁹ Y. Jiang,⁵⁹ K. Johns,⁴¹ M. Johnson,⁴⁶ P. Johnson,⁴¹ A. Jonckheere,⁴⁶ P. Jonsson,³⁹ H. Jöstlein,⁴⁶ A. Juste,⁴⁶ M.M. Kado,⁴² D. Käfer,¹⁹ W. Kahl,⁵⁴ S. Kahn,⁶⁷ E. Kajfasz,¹³ A.M. Kalinin,³² J. Kalk,⁶⁰ D. Karmanov,³⁴ J. Kasper,⁵⁷ D. Kau,⁴⁵ Z. Ke,⁶ R. Kehoe,⁶⁰ S. Kermiche,¹³ S. Kesisoglou,⁷⁰ A. Khanov,⁶⁵ A. Kharchilava,⁵¹ Y.M. Kharzheev,³² K.H. Kim,²⁸ B. Klima,⁴⁶ M. Klute,²⁰ J.M. Kohli,²⁵ M. Kopal,⁶⁹ V. Korablev,³⁵ J. Kotcher,⁶⁷ B. Kothari,⁶⁴ A.V. Kotwal,⁶⁴ A. Koubarovsky,³⁴ A. Kouchner,¹⁶ O. Kouznetsov,¹² A.V. Kozelov,³⁵ J. Kozminski,⁶⁰ J. Krane,⁵² M.R. Krishnaswamy,²⁶ S. Krzywdzinski,⁴⁶ M. Kubantsev,⁵⁴ S. Kuleshov,³³ Y. Kulik,⁴⁶ S. Kunori,⁵⁶ A. Kupco,¹⁶ T. Kurča,¹⁸ V.E. Kuznetsov,⁴⁴ S. Lager,³⁷ N. Lahrichi,¹⁶ G. Landsberg,⁷⁰ J. Lazoflores,⁴⁵ A.-C. Le Bihan,¹⁷ P. Lebrun,¹⁸ S.W. Lee,²⁸ W.M. Lee,⁴⁵ A. Leflat,³⁴ C. Leggett,⁴² F. Lehner,^{46,*} C. Leonidopoulos,⁶⁴ P. Lewis,³⁹ J. Li,⁷¹ Q.Z. Li,⁴⁶ X. Li,⁶ J.G.R. Lima,⁴⁸ D. Lincoln,⁴⁶ S.L. Linn,⁴⁵ J. Linnemann,⁶⁰ R. Lipton,⁴⁶ L. Lobo,³⁹ A. Lobodenko,³⁶ M. Lokajicek,¹⁰ A. Lounis,¹⁷ J. Lu,⁶ H.J. Lubatti,⁷⁴ A. Lucotte,¹² L. Lueking,⁴⁶ C. Luo,⁵⁰ M. Lynker,⁵¹ A.L. Lyon,⁴⁶ A.K.A. Maciel,⁴⁸ R.J. Madaras,⁴² P. Mättig,²⁴ A.-M. Magnan,¹² M. Maity,⁵⁷ P.K. Mal,²⁶ S. Malik,⁵⁵ V.L. Malyshev,³² V. Manankov,³⁴ H.S. Mao,⁶ Y. Maravin,⁴⁶ T. Marshall,⁵⁰ M. Martens,⁴⁶ M.I. Martin,⁴⁸ S.E.K. Mattingly,⁷⁰ A.A. Mayorov,³⁵ R. McCarthy,⁶⁶ R. McCroskey,⁴¹ T. McMahon,⁶⁸ D. Meder,²² H.L. Melanson,⁴⁶ A. Melnitchouk,⁷⁰ X. Meng,⁶ M. Merkin,³⁴ K.W. Merritt,⁴⁶ A. Meyer,¹⁹ C. Miao,⁷⁰ H. Miettinen,⁷² D. Mihalcea,⁴⁸ C.S. Mishra,⁴⁶ J. Mitrevski,⁶⁴ N. Mokhov,⁴⁶ J. Molina,³ N.K. Mondal,²⁶ H.E. Montgomery,⁴⁶ R.W. Moore,⁵ M. Mostafa,¹ G.S. Muanza,¹⁸ M. Mulders,⁴⁶ Y.D. Mutaf,⁶⁶ E. Nagy,¹³

F. Nang,⁴¹ M. Narain,⁵⁷ V.S. Narasimham,²⁶ N.A. Naumann,³¹ H.A. Neal,⁵⁹ J.P. Negret,⁷ S. Nelson,⁴⁵ P. Neustroev,³⁶ C. Noeding,²¹ A. Nomerotski,⁴⁶ S.F. Novaes,⁴ T. Nunnemann,²³ E. Nurse,⁴⁰ V. O'Dell,⁴⁶ D.C. O'Neil,⁵ V. Oguri,³ N. Oliveira,³ B. Olivier,¹⁵ N. Oshima,⁴⁶ G.J. Otero y Garzón,⁴⁷ P. Padley,⁷² K. Papageorgiou,⁴⁷ N. Parashar,⁵⁵ J. Park,²⁸ S.K. Park,²⁸ J. Parsons,⁶⁴ R. Partridge,⁷⁰ N. Parua,⁶⁶ A. Patwa,⁶⁷ P.M. Perea,⁴⁴ E. Perez,¹⁶ O. Peters,³⁰ P. Pétroff,¹⁴ M. Petteni,³⁹ L. Phaf,³⁰ R. Piegaia,¹ P.L.M. Podesta-Lerma,²⁹ V.M. Podstavkov,⁴⁶ B.G. Pope,⁶⁰ E. Popkov,⁵⁷ W.L. Prado da Silva,³ H.B. Prosper,⁴⁵ S. Protopopescu,⁶⁷ M.B. Przybycien,^{49,†} J. Qian,⁵⁹ A. Quadt,²⁰ B. Quinn,⁶¹ K.J. Rani,²⁶ P.A. Rapidis,⁴⁶ P.N. Ratoff,³⁸ N.W. Reay,⁵⁴ J.-F. Renardy,¹⁶ S. Reucroft,⁵⁸ J. Rha,⁴⁴ M. Ridel,¹⁴ M. Rijssenbeek,⁶⁶ I. Ripp-Baudot,¹⁷ F. Rizatdinova,⁵⁴ C. Royon,¹⁶ P. Rubinov,⁴⁶ R. Ruchti,⁵¹ B.M. Sabirov,³² G. Sajot,¹² A. Sánchez-Hernández,²⁹ M.P. Sanders,⁴⁰ A. Santoro,³ G. Savage,⁴⁶ L. Sawyer,⁵⁵ T. Scanlon,³⁹ R.D. Schamberger,⁶⁶ H. Schellman,⁴⁹ P. Schieferdecker,²³ C. Schmitt,²⁴ A. Schukin,³⁵ A. Schwartzman,⁶³ R. Schwienhorst,⁶⁰ S. Sengupta,⁴⁵ E. Shabalina,⁴⁷ V. Shary,¹⁴ W.D. Shephard,⁵¹ D. Shpakov,⁵⁸ R.A. Sidwell,⁵⁴ V. Simak,⁹ V. Sirotenko,⁴⁶ D. Skow,⁴⁶ P. Slattery,⁶⁵ R.P. Smith,⁴⁶ K. Smolek,⁹ G.R. Snow,⁶² J. Snow,⁶⁸ S. Snyder,⁶⁷ S. Söldner-Rembold,⁴⁰ X. Song,⁴⁸ Y. Song,⁷¹ L. Sonnenschein,⁵⁷ A. Sopczak,³⁸ V. Sorín,¹ M. Sosebee,⁷¹ K. Soustruznik,⁸ M. Souza,² N.R. Stanton,⁵⁴ J. Stark,¹² J. Steele,⁶⁴ G. Steinbrück,⁶⁴ K. Stevenson,⁵⁰ V. Stolin,³³ A. Stone,⁴⁷ D.A. Stoyanova,³⁵ J. Strandberg,³⁷ M.A. Strang,⁷¹ M. Strauss,⁶⁹ R. Ströhmer,²³ M. Strovink,⁴² L. Stutte,⁴⁶ A. Sznajder,³ M. Talby,¹³ P. Tamburello,⁴¹ W. Taylor,⁶⁶ P. Telford,⁴⁰ J. Temple,⁴¹ S. Tentindo-Repond,⁴⁵ E. Thomas,¹³ B. Thooris,¹⁶ M. Tomoto,⁴⁶ T. Toole,⁵⁶ J. Torborg,⁵¹ S. Towers,⁶⁶ T. Trefzger,²² S. Trincaz-Duvoid,¹⁵ T.G. Trippe,⁴² B. Tuchming,¹⁶ C. Tully,⁶³ A.S. Turcot,⁶⁷ P.M. Tuts,⁶⁴ L. Uvarov,³⁶ S. Uvarov,³⁶ S. Uzunyan,⁴⁸ B. Vachon,⁴⁶ R. Van Kooten,⁵⁰ W.M. van Leeuwen,³⁰ N. Varelas,⁴⁷ E.W. Varnes,⁴¹ I. Vasilyev,³⁵ M. Vaupel,²⁴ P. Verdier,¹⁴ L.S. Vertogradov,³² M. Verzocchi,⁵⁶ F. Villeneuve-Seguier,³⁹ J.-R. Vlimant,¹⁵ E. Von Toerne,⁵⁴ M. Vreeswijk,³⁰ T. Vu Anh,¹⁴ M. verzocch, ⁵⁵ F. Villeneuve-Seguer, ⁵⁵ J.-R. Vilmant, ¹⁵ E. Von Toerne, ⁵⁴ M. Vreeswijk, ⁵⁰ T. Vu Anh, ¹⁴
H.D. Wahl, ⁴⁵ R. Walker, ³⁹ N. Wallace, ⁴¹ Z.-M. Wang, ⁶⁶ J. Warchol, ⁵¹ M. Warsinsky, ²⁰ G. Watts, ⁷⁴ M. Wayne, ⁵¹
M. Weber, ⁴⁶ H. Weerts, ⁶⁰ M. Wegner, ¹⁹ A. White, ⁷¹ V. White, ⁴⁶ D. Whiteson, ⁴² D. Wicke, ²⁴ D.A. Wijngaarden, ³¹
G.W. Wilson, ⁵³ S.J. Wimpenny, ⁴⁴ J. Wittlin, ⁵⁷ T. Wlodek, ⁷¹ M. Wobisch, ⁴⁶ J. Womersley, ⁴⁶ D.R. Wood, ⁵⁸
Z. Wu, ⁶ T.R. Wyatt, ⁴⁰ Q. Xu, ⁵⁹ N. Xuan, ⁵¹ R. Yamada, ⁴⁶ T. Yasuda, ⁴⁶ Y.A. Yatsunenko, ³² Y. Yen, ²⁴
K. Yip, ⁶⁷ S.W. Youn, ²⁸ J. Yu, ⁷¹ A. Yurkewicz, ⁶⁰ A. Zabi, ¹⁴ A. Zatserklyaniy, ⁴⁸ M. Zdrazil, ⁶⁶ C. Zeitnitz, ²²
B. Zhang, ⁶ D. Zhang, ⁴⁶ X. Zhang, ⁶⁹ T. Zhao, ⁷⁴ Z. Zhao, ⁵⁹ H. Zheng, ⁵¹ B. Zhou, ⁵⁹ Z. Zhou, ⁵² J. Zhu, ⁵⁶ M. Zielinski,⁶⁵ D. Zieminska,⁵⁰ A. Zieminski,⁵⁰ R. Zitoun,⁶⁶ V. Zutshi,⁴⁸ E.G. Zverev,³⁴ and A. Zylberstejn¹⁶ (DØ Collaboration) ¹Universidad de Buenos Aires, Buenos Aires, Argentina ²LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil ³Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil ⁴Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil ⁵University of Alberta and Simon Fraser University, Canada ⁶Institute of High Energy Physics, Beijing, People's Republic of China ⁷Universidad de los Andes, Bogotá, Colombia ⁸Charles University, Center for Particle Physics, Prague, Czech Republic ⁹Czech Technical University, Prague, Czech Republic

¹⁰Institute of Physics, Academy of Sciences, Center for Particle Physics, Prague, Czech Republic

¹¹ Universidad San Francisco de Quito, Quito, Ecuador

¹²Laboratoire de Physique Subatomique et de Cosmologie, IN2P3-CNRS, Universite de Grenoble 1, Grenoble, France

³CPPM, IN2P3-CNRS, Université de la Méditerranée, Marseille, France

¹⁴Laboratoire de l'Accélérateur Linéaire, IN2P3-CNRS, Orsay, France

¹⁵LPNHE, Universités Paris VI and VII, IN2P3-CNRS, Paris, France

¹⁶DAPNIA/Service de Physique des Particules, CEA, Saclay, France

¹⁷ IReS, IN2P3-CNRS, Univ. Louis Pasteur Strasbourg, and Univ. de Haute Alsace, France

¹⁸Institut de Physique Nucléaire de Lyon, IN2P3-CNRS, Université Claude Bernard, Villeurbanne, France

¹⁹ RWTH Aachen, III. Physikalisches Institut A, Aachen, Germany

²⁰ Universität Bonn, Physikalisches Institut, Bonn, Germany

²¹ Universität Freiburg, Physikalisches Institut, Freiburg, Germany

²² Universität Mainz, Institut für Physik, Mainz, Germany

²³Ludwig-Maximilians-Universität München, München, Germany

²⁴ Fachbereich Physik, University of Wuppertal, Wuppertal, Germany

²⁵Panjab University, Chandigarh, India

²⁶ Tata Institute of Fundamental Research, Mumbai, India

²⁷ University College Dublin, Dublin, Ireland

²⁸Korea Detector Laboratory, Korea University, Seoul, Korea

²⁹CINVESTAV, Mexico City, Mexico

³⁰ FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands

³¹ University of Nijmegen/NIKHEF, Nijmegen, The Netherlands

³² Joint Institute for Nuclear Research, Dubna, Russia

³³Institute for Theoretical and Experimental Physics, Moscow, Russia

³⁴Moscow State University, Moscow, Russia

³⁵Institute for High Energy Physics, Protvino, Russia

³⁶Petersburg Nuclear Physics Institute, St. Petersburg, Russia

³⁷Lund University, Royal Institute of Technology, Stockholm University, and Uppsala University, Sweden

³⁸Lancaster University, Lancaster, United Kingdom

³⁹Imperial College, London, United Kingdom

⁴⁰ University of Manchester, Manchester, United Kingdom

⁴¹ University of Arizona, Tucson, Arizona 85721

⁴²Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720

⁴³California State University, Fresno, California 93740

⁴⁴ University of California, Riverside, California 92521

⁴⁵ Florida State University, Tallahassee, Florida 32306

⁴⁶ Fermi National Accelerator Laboratory, Batavia, Illinois 60510

⁴⁷ University of Illinois at Chicago, Chicago, Illinois 60607

⁴⁸Northern Illinois University, DeKalb, Illinois 60115

⁴⁹Northwestern University, Evanston, Illinois 60208

⁵⁰Indiana University, Bloomington, Indiana 47405

⁵¹ University of Notre Dame, Notre Dame, Indiana 46556 ⁵² Iowa State University, Ames, Iowa 50011

⁵³University of Kansas, Lawrence, Kansas 66045

⁵⁴Kansas State University, Manhattan, Kansas 66506

⁵⁵Louisiana Tech University, Ruston, Louisiana 71272

⁵⁶ University of Maryland, College Park, Maryland 20742

⁵⁷Boston University, Boston, Massachusetts 02215

⁵⁸Northeastern University, Boston, Massachusetts 02115

⁵⁹ University of Michigan, Ann Arbor, Michigan 48109

⁶⁰ Michigan State University, East Lansing, Michigan 48824

⁶¹ University of Mississippi, University, Mississippi 38677

⁶²University of Nebraska, Lincoln, Nebraska 68588
 ⁶³Princeton University, Princeton, New Jersey 08544

⁶⁴ Columbia University, New York, New York 10027

⁶⁵ University of Rochester, Rochester, New York 14627

⁶⁶State University of New York, Stony Brook, New York 11794

⁶⁷Brookhaven National Laboratory, Upton, New York 11973

⁶⁸Langston University, Langston, Oklahoma 73050

⁶⁹ University of Oklahoma, Norman, Oklahoma 73019

⁷⁰Brown University, Providence, Rhode Island 02912

⁷¹University of Texas, Arlington, Texas 76019

⁷²Rice University, Houston, Texas 77005

⁷³ University of Virginia, Charlottesville, Virginia 22901

⁷⁴ University of Washington, Seattle, Washington 98195

We report the observation of the X(3872) in the $J/\psi \pi^+\pi^-$ channel, with J/ψ decaying to $\mu^+\mu^-$, in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV. Using approximately 230 pb⁻¹ of data collected with the Run II DØ detector, we observe $522 \pm 100 X(3872)$ candidates. The mass difference between the X(3872)state and the J/ψ is measured to be 774.9 ± 3.1 (stat) ± 3.0 (syst) MeV/ c^2 . We have investigated the production and decay characteristics of the X(3872), and find them to be similar to those of the $\psi(2S)$ state.

PACS numbers: 14.40.Gx, 14.40.Lb, 13.25.Gv, 13.25.Jx, 13.85.Ni

A new particle, the X(3872), was recently discovered by the Belle Collaboration [1] in the decay mode $B^{\pm} \to XK^{\pm} \ (X \to J/\psi \pi^{+}\pi^{-})$ where the mass of the X(3872) was measured to be 3872.0 ± 0.6 (stat) ± 0.5 (syst) MeV/ c^{2} . The existence of the X(3872) state has been confirmed (also decaying in the $J/\psi \pi^{+}\pi^{-}$ mode) in $p\bar{p}$ collisions by the CDF Collaboration [2]. At this time, it is still unclear whether this particle is a $c\bar{c}$ state, or a more complex object. See for example [3, 4, 5, 6].

The charmonium state $\psi(2S)$, with mass $m_{\psi(2S)} = 3685.96 \pm 0.09 \text{ MeV}/c^2$ [7], has the same decay mode, providing a good benchmark for comparison with the

X(3872). The $\psi(2S)$ mesons produced in $p\bar{p}$ collisions can originate either from decays of *B* hadrons or from direct production. The $\psi(2S)$ mesons from *B* decays have longer effective decay lengths and tend to be less isolated than directly produced $\psi(2S)$ mesons [8].

We examine the production rate of the X(3872) relative to $\psi(2S)$ as a function of the transverse momentum with respect to the beam axis (p_T) , isolation and decay length, as well as a function of rapidity $(y = \frac{1}{2} \log \frac{E+P_L}{E-P_L})$, where E is the energy and P_L is the longitudinal momentum with respect to the beam axis), to determine whether the production characteristics of the X(3872)are similar to those of the $\psi(2S)$. We also compare the angular decay distributions of the $\pi^+\pi^-$ and $\mu^+\mu^-$ systems in X(3872) decays with those from $\psi(2S)$, to check for any differences in helicities of these two states.

The data set used in this Letter was collected in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ TeV between April 2002 and January 2004, and corresponds to an integrated luminosity of approximately 230 pb⁻¹. The DØ detector is described elsewhere [9]. The components most important to this analysis include the vertex, central tracking and muon systems. The DØ tracking system consists of a silicon microstrip tracker (SMT) and a central fiber tracker (CFT), both within a 2 T solenoidal magnetic field.

The SMT has approximately 800, 000 individual strips, with typical pitch of 50–80 μ m, and a design optimized for tracking and vertexing over the range $|\eta| < 3$, where $\eta = -\ln[\tan(\theta/2)]$ is the pseudorapidity and θ is the polar angle measured relative to the proton beam direction. The system has a six-barrel longitudinal structure, each with a set of four layers arranged axially around the beam pipe, and interspersed with 16 radial disks. The system provides a resolution, in the plane transverse to the beam axis, for the distance of closest approach of a charged particle relative to the primary vertex of $\approx 50 \ \mu$ m for tracks with $p_T \approx 1 \ {\rm GeV}/c$, improving asymptotically to 15 μ m for tracks with $p_T \geq 10 \ {\rm GeV}/c$.

The CFT comprises eight thin coaxial barrels, each supporting two doublets of overlapping scintillating fibers of 0.835 mm diameter, one doublet being parallel to the collision axis, and the other alternating by $\pm 3^{\circ}$ to provide information along the beam axis.

The muon system is located outside the calorimeters, and consists of a layer of tracking detectors and scintillation trigger counters in front of 1.8 T toroidal magnets, followed by two similar layers behind the toroids. Tracking in the muon system in the range $|\eta| < 1$ relies on 10 cm wide drift tubes [10], while 1 cm mini-drift tubes are used for $1 < |\eta| < 2$.

 $J/\psi \rightarrow \mu^+\mu^-$ decays are selected by triggering on dimuons using a three-tier trigger system. The first trigger level uses hardware to form roads defined by hits in two layers of the muon scintillator system. The second trigger level uses digital signal processors to form track stubs defined by hits in the muon drift-chamber and muon scintillator systems. The third level comprises a farm of computer processors with access to the entire event. Events passing the third-level trigger are recorded for analysis.



FIG. 1: $\Delta M = M(\mu^+\mu^-\pi^+\pi^-) - M(\mu^+\mu^-)$ for all candidates satisfying the selection requirements. The solid curve is a fit to the data and the dashed curve represents the background under each peak. The insert shows the mass distribution of the J/ψ candidates used in the analysis.

Muons are identified by extrapolating charged particle tracks from the central tracking system that match with muon track segments formed from hits in the muon system. Oppositely charged muons are combined to form J/ψ candidates, which are then combined with two oppositely charged particles assumed to be pions. At least two of these four tracks are required to have at least one hit in the SMT. To reduce background from combinatorics, events are required to satisfy the following selection criteria. An event is required to have less than 100 tracks. J/ψ candidates are selected by requiring the invariant mass of the $\mu^+\mu^-$ system to be between 2.80 and 3.35 GeV/c^2 , and the transverse momentum with respect to the beam axis (p_T) of the J/ψ is required to be greater than 5 GeV/c. In addition, the p_T of each of the two pions must be greater than 0.7 GeV/c, and the spatial separation, ΔR , between the momentum vector of the $J/\psi \pi^+\pi^-$ system and each pion momentum vector is required to have $\Delta R < 0.4$, where ΔR is defined as $\sqrt{(\Delta \phi)^2 + (\Delta \eta)^2}$, with ϕ being the azimuthal angle. The invariant mass of the two pions, $M(\pi^+\pi^-)$, is required to be greater than 0.52 GeV/ c^2 , and the χ^2 of a fit to the $\mu^+\mu^-\pi^+\pi^-$ vertex is required to be less than 16 (for five degrees of freedom).

Figure 1 shows the distribution in the mass difference $\Delta M = M(\mu^+\mu^-\pi^+\pi^-) - M(\mu^+\mu^-)$, after all selections.

Superimposed is a fit to the data, where Gaussians are used to describe the $\psi(2S)$ and the X(3872), and a thirdorder polynomial is used to account for the background. The fitted width of the X(3872) peak is $17 \pm 3 \text{ MeV}/c^2$, which is consistent with detector resolution. The results of the Gaussian fit yield $522 \pm 100 X(3872)$ candidates with $\Delta M = 771.9 \pm 3.1$ (stat) MeV/ c^2 . The yield of the X(3872) is dependent on the value of the $M(\pi^+\pi^-)$ requirement. When we tighten the requirement on $M(\pi^+\pi^-)$ to be greater than 0.60 GeV/ c^2 , 0.65 GeV/ c^2 and 0.70 GeV/ c^2 and fix the mean of the X(3872) to its nominal value, we obtain yields of 433 ± 98 , 336 ± 78 and 170 ± 54 events, respectively.

The position of the J/ψ mass in Fig. 1 is shifted relative to the currently accepted value [7]. This is within the uncertainties of our momentum scale, but we correct the measured ΔM by the ratio of the currently accepted mass difference between the $\psi(2S)$ and the J/ψ [7] and our corresponding measured mass difference. We assign a 100% systematic uncertainty to this mass scale correction, giving a final measurement of $\Delta M = 774.9 \pm 3.1$ (stat) \pm 3.0 (syst) MeV/c². We investigated systematic effects by using different background parameterizations and bin widths in our fits, and found the largest change in the central value of the mass difference to be 0.6 MeV/c².

To investigate the characteristics of the X(3872) state, we study its production and decay properties, and compare the signal yields of the X(3872) to the $\psi(2S)$. For example, Fig. 2 shows the ΔM distribution in two separate regions of rapidity of the X(3872). The data is also separated into regions of effective proper decay length, transverse momentum of the X(3872) candidate, isolation, and decay angle.

The effective proper decay length, dl, is defined as the distance in the transverse plane from the primary vertex to the decay vertex of the J/ψ scaled by the mass of the $\mu^+\mu^-\pi^+\pi^-$ system divided by the $\mu^+\mu^-\pi^+\pi^-$ system p_T .

The isolation is defined as the ratio of the X(3872) momentum to the sum of the momentum of the X(3872) and the momenta of all other reconstructed charged particles within a cone of radius $\Delta R = 0.5$, about the direction of the X(3872) momentum.

The helicity of the $\pi^+\pi^-(\mu^+\mu^-)$ system can be inferred by boosting one of the pions (or muons) and the X(3872) into the dipion (dimuon) rest frame, and measuring the angle $\theta_{\pi}(\theta_{\mu})$ between them. The cosine of this angle is used for the comparison between the X(3872)and $\psi(2S)$.

The widths of the $\psi(2S)$ and X(3872) are fixed to the values obtained in the fit to the full sample, and the fitted number of $\psi(2S)$ and X(3872) candidates in each region of chosen variables is given in Table I. The data in all regions are well represented by our choice of fitting function, with fit probabilities all greater than 15%. In Fig. 3, we present the results of Table I in a graphical



FIG. 2: ΔM distribution for two ranges of rapidity of the $\mu^+\mu^-\pi^+\pi^-$ system.

form. For example, we compare the event yield fraction:

$$f_{X(3872)} = \frac{N(X(3872)) |y| < 1}{N(X(3872)) |y| < 2}$$

to that for $\psi(2S)$:

$$f_{\psi(2S)} = \frac{N(\psi(2S)) |y| < 1}{N(\psi(2S)) |y| < 2}.$$

From the ratios of isolation and decay lengths, it is apparent that the production of the X(3872) has a similar mixture of prompt and long-lived contributions as the $\psi(2S)$.

In summary, we observe the X(3872) particle in approximately 230 pb⁻¹ of data with a significance of 5.2 standard deviations. The mass difference between the X(3872) particle and the J/ψ is $\Delta M = 774.9 \pm 3.1$ (stat) ± 3.0 (syst) MeV/ c^2 . When the data are separated according to production and decay variables, we find no significant differences between the X(3872) and the $c\bar{c}$ state $\psi(2S)$.

We thank Estia Eichten, Ken Lane, and Chris Quigg, for fruitful discussions. We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the Department of Energy and National Science Foundation (USA), Commissariat à l'Energie Atomique and CNRS/Institut National de Physique Nucléaire et de Physique des Particules (France), Ministry of Education and Science, Agency for Atomic Energy and RF President Grants Program (Russia), CAPES, CNPq, FAPERJ, FAPESP and FUNDUNESP (Brazil), Departments of Atomic Energy and Science and Technology

TABLE I: Number of $\psi(2S)$ and X(3872) candidates for different ranges of variables. The fitted widths of the X(3872)and $\psi(2S)$ are not constrained in the initial selection, but in the different ranges they are fixed to the values obtained from the full sample. Apart from the full sample, uncertainties are only from the normalization of the fitted Gaussian function and do not include small contributions from uncertainties in the background.

Regions	Number	Number
	of $\psi(2S)$	of $X(3872)$
Initial selection	1192 ± 55	522 ± 100
(a) $p_T(J/\psi \pi^+\pi^-) > 15 \text{ GeV}/c$	396 ± 26	179 ± 39
$p_T(J/\psi \pi^+\pi^-) \le 15 \text{ GeV}/c$	796 ± 41	358 ± 64
(b) $ y < 1$	741 ± 37	316 ± 57
$1 \le y \le 2$	449 ± 31	204 ± 49
(c) $\cos(\theta_{\pi}) < 0.4$	589 ± 34	288 ± 53
$\cos(\theta_{\pi}) \ge 0.4$	606 ± 34	$244~\pm~53$
(d) $dl < 0.01 \text{ cm}$	838 ± 41	351 ± 66
$dl \ge 0.01 \text{ cm}$	359 ± 26	164 ± 41
(e) isolation $= 1$	257 ± 20	85 ± 22
isolation < 1	942 ± 44	438 ± 72
(f) $\cos(\theta_{\mu}) < 0.4$	593 ± 33	232 ± 46
$\cos(\theta_{\mu}) \ge 0.4$	602 ± 35	288 ± 60



FIG. 3: Comparison of event-yield fractions for X(3872) and $\psi(2S)$ in the regions: (a) $p_T(J/\psi \pi^+\pi^-) > 15 \text{ GeV}/c$; (b) |y| of $J/\psi \pi^+\pi^- < 1$; (c) $\cos(\theta_\pi) < 0.4$; (d) effective proper decay length, dl < 0.01 cm; (e) isolation = 1; (f) $\cos(\theta_\mu) < 0.4$.

- [*] Visitor from University of Zurich, Zurich, Switzerland.
- [†] Visitor from Institute of Nuclear Physics, Krakow, Poland.

91, 262001 (2003).

- [2] D. Acosta *et al.* (CDF II Collaboration), submitted to Phys. Rev. Lett., hep-ex/0312021.
- [1] S. K. Choi et al. (Belle Collaboration), Phys. Rev. Lett.
- [3] F. E. Close, P. R. Page, "The $D^{*0}\bar{D^0}$ threshold reso-

(India), Colciencias (Colombia), CONACyT (Mexico), KRF (Korea), CONICET and UBACyT (Argentina), The Foundation for Fundamental Research on Matter (The Netherlands), PPARC (United Kingdom), Ministry of Education (Czech Republic), Natural Sciences and Engineering Research Council and WestGrid Project (Canada), BMBF (Germany), A.P. Sloan Foundation, Civilian Research and Development Foundation, Research Corporation, Texas Advanced Research Program, and the Alexander von Humboldt Foundation. nance", Phys. Lett. B578, 119 (2004), hep-ph/0309253.

- [4] E. J. Eichten, K. Lane, and C. Quigg, "Charmonium levels near threshold and the narrow state $X(3872) \rightarrow \pi^+\pi^- J/\psi$," FERMILAB–Pub–04/001–T, hep-ph/0401210, and references therein.
- [5] C. Quigg, "Quarkonium: New developments," hep-ph/0403187.
- [6] N. Törnqvist, "Isospin breaking of the narrow charmonium state of Belle at 3872 MeV as a deuson", hep-ph/0402237.
- [7] K. Hagiwara *et al.*, Phys. Rev. D 66, 010001 (2002) and 2003 off-year partial update for the 2004 edition available

on the PDG WWW pages (URL: http://pdg.lbl.gov/).

- [8] S. Abachi *et al.* (DØ Collaboration), Phys. Lett. B370, 239 (1996); F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. 79, 572 (1997).
- [9] V. Abazov, et al., in preparation for submission to Nucl. Instrum. Methods Phys. Res. A, and T. LeCompte and H.T. Diehl, "The CDF and D0 Upgrades for Run II", Ann. Rev. Nucl. Part. Sci. 50, 71 (2000).
- [10] S. Abachi, *et al.*, Nucl. Instrum. Methods Phys. Res. A 338, 185 (1994).