

Research

In vitro and in vivo effects of the PPAR-alpha agonists fenofibrate and retinoic acid in endometrial cancer

Samir A Saidi*, Cathrine M Holland, D Stephen Charnock-Jones and Stephen K Smith

Address: University Department of Obstetrics & Gynaecology, The Rosie Hospital, Robinson Way, Cambridge, CB2 2SW, UK

Email: Samir A Saidi* - samsaidi@obgyn.cam.ac.uk; Cathrine M Holland - cath.holland@obgyn.cam.ac.uk; D Stephen Charnock-Jones - dscj1@cam.ac.uk; Stephen K Smith - s.k.smith@imperial.ac.uk

* Corresponding author

Published: 28 March 2006

Received: 23 November 2005

Molecular Cancer 2006, **5**:13 doi:10.1186/1476-4598-5-13

Accepted: 28 March 2006

This article is available from: <http://www.molecular-cancer.com/content/5/1/13>

© 2006 Saidi et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

: Fenofibrate, an agonist of PPAR-alpha, in doses above 25 μ M, inhibits proliferation and induces apoptosis in Ishikawa endometrial cancer cells. We show that these effects are potentiated by retinoic acid, an agonist of the retinoid-X-receptor. DNA content analysis shows that G₁/S phase progression through the cell cycle is inhibited. Independent Component Analysis of gene microarray experiments demonstrated downregulation of Cyclin D1 (CCND1) and associated changes in cell cycle gene expression. Expression of PPAR-alpha mRNA was reduced by >75% using RNA-interference but this resulted in only minor changes in biological effects. A nude mouse model of endometrial carcinoma was used to investigate the effect of fenofibrate *in vivo* but failed to show consistent inhibition of tumour growth.

Conclusion: The combination of fenofibrate and retinoic acid is a potent inhibitor of Ishikawa endometrial cancer cell growth *in vitro*.

Introduction

Endometrial cancer is the most prevalent gynaecological cancer in the UK, and represents the third commonest cancer affecting women in the Western World [1]. By contrast, the incidence in the non-Western world is approximately ten-fold lower [2]. The excellent prognosis of early stage endometrial cancers belies the impact of the disease on mortality, being of similar magnitude to that of cervical cancer [2]. Indeed, the long-term survival of advanced stage endometrial cancer, at approximately 10%, is similar to that of ovarian cancer.

Established risk factors for sporadic endometrial cancer mainly involve hormonal factors, with the unopposed estrogen hypothesis believed to be the central pathoge-

netic mechanism [3,4]. Although this theory is strongly supported, it does not satisfactorily account for all the risk factors associated with endometrial cancer risk. Obesity is a significant independent risk factor, with relative risks in the 2–10 range [5,6]. The mechanism for this has not yet been elucidated but postulates include the collateral involvement of estrogen and insulin-like growth factor (IGF) receptor pathways [5,7]. Improving understanding of the carcinogenesis of endometrial cancer is essential in the development of targeted therapy.

The potential of gene array methods and systems biology has been exploited in recent years for the investigation of a number of tumour types [8,9]. The aim of the new biology is to provide a global overview of carcinoma at the

molecular level, whilst focusing on biologically relevant data. Although oncology has received a great deal of attention from computational biology, a limited number of gene array studies have been applied solely to endometrial cancer [10-12].

Using gene array methods within a computational biology environment, we have previously demonstrated that lipid metabolism is likely to play an important role in endometrial carcinogenesis [12,13]. Consequentially, we identified fenofibrate, a ligand of the peroxisome proliferator-activated receptor alpha (PPAR α), as a potential therapeutic agent in endometrial cancer [12]. PPARs comprise a group of transcription factors belonging to the nuclear hormone receptor subfamily and consist of subtypes α , γ and β/δ [14]. Their main actions regulate the metabolism of fatty acids and are therefore closely involved with prostanoid pathways [14]. Furthermore, receptor-mediated transcription is dependent upon heterodimerisation with the retinoid-X receptors (RXRs). Following activation by their ligands (eg fenofibrate and fatty acids in the case of PPAR α) and heterodimerization with RXR, PPARs bind to the peroxisome-proliferator response element (PPRE) in the promoter of their target genes and activate their transcription [14]. PPREs are most commonly found in genes that are involved in lipid metabolism and energy homeostasis, including lipid storage or catabolism (β -oxidation and ω -oxidation), fatty-acid transport, uptake and intracellular binding. In recent years there has been interest and some success in the use of retinoids, synthetic ligands of the RXR, in the treatment of hormonally derived cancers such as those of the breast and endometrium [15,16].

Our previous work demonstrated upregulation of PPAR α transcript in association with downregulation of its heterodimerisation partner RXR β [12,13] in endometrial cancer. We also showed that the PPAR α agonist fenofibrate, in doses above 25 μ M, inhibits Ishikawa and ECC-1 endometrial cancer cell growth *in vitro*, in association with increased apoptosis and PPAR α receptor activation [12]. In this study, attention was focussed on the Ishikawa cell line in view of its endometrioid-like characteristics, estrogen receptor positivity [17] and suitability for xenografting [18].

Having identified PPAR α as a potential therapeutic target in endometrial cancer, the aim of this study was to further investigate the biological effects of fenofibrate, from a molecular to a cellular level and finally to an animal model. We further aimed to investigate whether targeting the PPAR α receptor using retinoid-X-receptor ligands would increase the growth-inhibitory effects of this agent. Finally, a systems biology approach was used to help understand the mode of action of fenofibrate by identify-

ing the global transcription changes induced in the treatment of endometrial cancer *in vitro*.

Materials and methods

In vitro studies

Cell culture & proliferation assays

Ishikawa cells were obtained from the European Collection of Cell Cultures (Cat. No. 99040201) [19] and were grown in DMEM/F-12 Ham medium (Cat. No. D6421, Sigma-Aldrich, UK) supplemented with L-glutamine and 10% fetal calf serum in 96-well plates (proliferation assays), 6-well plates (FACS analysis, luciferase reporter assays) or cell culture flasks (RNA extraction, tumour explant preparation). Cells were cultured at 37°C and 5%CO₂ with varying doses of fenofibrate (Cat. No. F6020), 9-cis retinoic acid (9-cRA) (Cat. No. R4643) or all-trans retinoic acid (ATRA) (Cat. No. R6265, all Sigma-Aldrich, UK) dissolved in DMSO (Sigma-Aldrich). Control cells were treated with DMSO and the concentration of DMSO was kept the same throughout each experiment, to a maximum of 1% v/v. A minimum of five replicates per dose was performed for automated assays (BrdU and MTS) and all other cell culture experiments were performed independently in duplicate or triplicate. Cell proliferation was measured by the uptake of 5-bromo-2'-deoxyuridine (BrdU) using the BrdU Labelling and Detection Kit (Cat. No. 1444611, Roche, UK) according to the manufacturer's instructions. Relative cell abundance was measured either by cell counting, or using an MTS assay (CellTiter 96 Aqueous Cell Proliferation Assay, Cat. No. TB169, Promega, UK). Absorbance values for each well were measured using a microtiter plate reader (Anthos Labtech Instruments, Salzburg, Austria)

FACS analysis

All FACS analysis was performed on a FACSCalibur analyser (Becton Dickinson, USA). Computational analysis was performed using FCSPress software for Macintosh [20]. Gating parameters were based on untreated cells in each experiment, to exclude subcellular fragments and conglomerate cells.

Apoptosis analysis

Cells in 6-well plates at approximately 80% confluence were washed in PBS, and the medium replaced with medium containing DMSO (control) or drug in DMSO solution to a maximum of 1% v/v DMSO. After 3 hours' exposure, the cells were harvested by trypsinisation, spun, and resuspended in 500 μ l buffer (containing per ml of serum-free medium: 4 μ l Propidium Iodide (Cat. No. P3566, Invitrogen, USA), 3.5 μ l Annexin-V FITC (Cat. No. A13199, Invitrogen, USA), 10 μ l 0.1 M Calcium chloride). Cells were transported on ice and FACS analysis was performed within one hour.

DNA content analysis

Cells were similarly treated with drug solution for 12 hours, and harvested as above. Approximately 10^6 cells were fixed in 0.3 ml of PBS and 0.7 ml of ice cold 70% EtOH at 4°C for 1 hr. After fixation, the cells were resuspended in 0.25 ml of PBS containing 12.5 µl of PI and 6.25 µl RNase A (20 mg/ml, SIGMA, UK). Cells were incubated at 37°C for 15 minutes prior to FACS analysis.

Cell counting

Cell counting was independently verified using FACS analysis. Samples harvested by trypsinisation were fixed in ethanol as above. 20 µl of a 1:1000 aqueous solution of Fluoresbrite plain YG 2 µm beads (Cat. No. 18338, Polysciences Inc, Germany) was added to 1 ml of cells resuspended in PBS containing 10 µl Propidium Iodide solution, giving a final bead concentration of 5.68×10^5 /ml. Cell concentration was then calculated as: (Cell count) \times (Bead Count)⁻¹ \times (5.68×10^5)

RNA interference

RNAi was performed through a lipid-based transfection method, with either PPAR α -RNAi (Cat. No. M-003434-00) or Control-RNAi (Cat. No. D-001206-13), using SmartPool reagents (Dharmacon, Lafayette CO, USA) according to the manufacturers' instructions. Cells were harvested at confluence from a 75 cm² culture flask, and plated at approximately 2×10^6 cells per plate into 6 well plates with 2 ml/well of medium as described. For transfection, 5 µl/well of 20 µM RNAi (control or PPAR), 4 µl/well of lipofectamine 2000 (Cat. No. 18324-111, Invitrogen, UK) and 100 µl/well of serum-free medium were incubated at room temperature. 100 µl/well of transfection mixture was then added to each well for overnight transfection, giving a transfection rate $\geq 30\%$, as measured by beta-galactosidase staining. The medium was replaced with fresh medium and cells grown to confluence. Cells were then harvested and pooled in each group (control RNAi/PPAR RNAi) before replating to 6-well plates. Cells later harvested for RT-PCR analysis were plated from the same batches. For cell counting, cells were allowed to grow in drug-containing media for 48 hours, then harvested. For FACS analysis (apoptosis assays), drug-based media was added to cells for 2 hours before harvesting.

Microarray studies

RNA preparation

Cells in the logarithmic growth phase were plated into 75 cm² culture flasks containing culture medium as described, containing DMSO or fenofibrate solution (in DMSO at 10 µM and 100 µM) as required. Five replicate cultures were grown at each drug dose. Cells were cultured for 48 hours, and then harvested by cell scraper using 1 ml Trizol reagent (Cat. No. 15596-026, Invitrogen, UK) per flask, and stored overnight at -70°C. RNA was then pre-

cipitated and resuspended using an isopropanol precipitation method [21]. cDNA was generated using random hexamer primers as described previously [22].

Sequencing

cDNA generated as above was used as template DNA for 35 cycles of PCR using the following conditions: 95°C (30s), 57°C (30s), 72°C (60s). To facilitate sequencing of large fragments, 2 overlapping fragments were amplified, covering the entire coding sequence, using the following primers:

Forward: 5'-GGCACAACCAGCACCATCT-3'

Reverse: 5'-CTCCACAGCAAATGATAGCAGC-3'

(Amplicon length 1185 bp, position 191-1375)

Forward: 5'-GCCAGTAACAATCCACCTTTT-3'

Reverse: 5'-AAGGTGTGGCTGATCTGAAGG-3'

(Amplicon length 735 bp, position 913-1647)

The PCR products were then subjected to electrophoresis on a 2% agarose gel. Bands corresponding to the above amplicons were then extracted from the gel using the Qiaquick kit according to the manufacturer's instructions (Cat. No. 28704, Qiagen, UK). This product was then used as template for a further 25 cycles of PCR, from which the relevant bands were again extracted and sequenced using the above primers on an ABI Prism 310 genetic analyzer (Applied Biosystems, UK). Sequence analysis and comparison was performed using DNASTar software (DNASTar Inc, USA).

Preparation of cDNA for microarray hybridisation

Following precipitation from Trizol, RNA was re-extracted using RNAeasy mini-columns (Cat. No. 74106, Qiagen, UK) and assayed for purity and concentration using spectrophotometry and an Agilent 2100 Bioanalyzer (Agilent, USA). Glass microarrays sourced from within the University of Cambridge Department of Pathology were printed on 2 slides and comprised over 10000 clones from various sources as previously described [22,23]. Confirmation of the array's performance had been conducted by in-house analysis of reproducibility [24], in addition to its use in prior validated experiments [23]. Further quality control of the arrays in this experiment was ensured by reproducibility analysis of the array data generated from the pooled DNA samples (see additional file 1). cDNA synthesis and labelling for hybridization was carried out as previously described with minor modifications [25]. 1 µg total RNA was used to synthesise double-strand cDNA (ds-cDNA) using the SMART PCR cDNA synthesis kit (Clontech, UK),

according to the manufacturer's instructions, over 15 cycles of PCR.

Spotted oligonucleotide microarray hybridisation and scanning

The ds-cDNA was labelled by Cy3-deoxyuridine triphosphate or Cy5- deoxyuridine triphosphate (Amersham-Pharmacia, UK) using the Bioprime DNA labelling kit (Cat. No. 18094-011, Invitrogen, UK) with random hexamers. To counter the effect of dye bias, a reference cDNA consisting of amplified cDNA pooled from the five control-treated samples was labelled with Cy3. Each of the 15 sample cDNAs were labelled with Cy5. Paired samples (Pooled-Cy3 + Sample-Cy5) were purified using Autoseq G50 columns (Amersham, UK), mixed with 5 µg/ml Human Cot-1 DNA (Cat. No. 15279-011, Invitrogen, UK), 1 µg/ml Poly-dA (Amersham-Pharmacia, UK). Labelled targets were resuspended in 50 µl of hybridisation buffer (40% formamide, 5× SSC, 5× Denhardt's solution, 1 mM sodium pyrophosphate, 50 mM Tris pH 7.4, 0.1% SDS), denatured at 95°C for 5 min, incubated at 50°C for 5 min and then centrifuged at 8000 rpm for 5 min before being applied to the cover-slipped array. Hybridisations were performed at 50°C in a humidified environment for 16 h. Following hybridisation, slides were washed twice in 2× SSC for 10 min, twice in 0.1× SSC/0.1% SDS for 5 min and finally twice in 0.1× SSC for 5 min; all washes were performed at room temperature. After washing, slides were dried by centrifugation at 1000 rpm for 2 min, and then scanned using a Genepix 4100 microarray scanner (Axon Instruments, Foster City, CA). The scanned images were processed by using GenePix Pro 4.1 software (Axon).

Analysis of gene array data

Raw Cy3- and Cy5- channel data extracted using Imagene were normalized against a reference array using a LOESS-based algorithm with a span of 0.4 as previously described [13]. Empty spots and known housekeepers were removed from the data. Independent Component Analysis [26] was performed on the remaining data set. Components extracted were then compared against the array grouping using ANOVA in the R-statistical environment, to identify components related to drug dosage, where $p < 0.01$. Significant genes relating to high-dose fenofibrate were those with the top 1% of component loadings in the relevant component. Analysis of gene ontology (GO) classification was performed as previously described [13]

Quantitative Real-Time PCR (Taqman)

RNA samples analysed by RT-PCR were the same as those used for microarray analysis. Total RNA (2 µg) was incubated with random hexamer primers at 70°C and reverse-transcribed at 42°C using Super Reverse Transcriptase (HT Biotechnology Ltd, Cambridge, UK). 0.6 µl of template DNA was amplified using PCR MasterMix (Cat. No. AB-

1142, ABgene, UK). All RT-PCR experiments were performed in an ABI Prism 7700 Sequence Detector (Applied Biosystems) in triplicate. Results for gene abundance in each sample were normalised to abundance of 18S RNA when appropriate. Normalised log-transformed transcript levels were compared across samples using one-way ANOVA and Student's t-tests.

(i) PPAR α

Due to the low abundance of the PPAR α transcript, 20 cycles of PCR (see below for conditions) were performed prior to the quantitative PCR cycles. The following primers were used to amplify a 553 bp region of PPAR α which was subsequently amplified with Taqman reagents.

Forward: 5'-AGGGCCCTGTCTGCTCTGTG-3'

Reverse: 5'-CGGGTGGACTCCGTAATGATAG-3'

PCR product from this round of PCR was used as the template for subsequent quantitative PCR. Primer/probe sequences used for RT-PCR were:

Forward primer: 5'-GACGTGCTTCCTGCTTCATAGA-3'

Reverse primer: 5'-CACCATCGCGACCAGATG-3'

Probe: 5'-6FAM-TGGAGCTCGGCGCACAACCA-TAMRA-3'

(ii) Verification of gene arrays

RT-PCR for three genes selected from microarray analysis was performed using the assays-on-demand primers below according to the manufacturers instructions (Applied Biosystems, USA).

(a) Cyclin D1 (CCND1) (Cat. No. Hs00277039_m1)

(b) Methionine adenosyltransferase II alpha (MAT2A) (Cat. No. Hs00428515_g1)

(c) Phosphoenolpyruvate carboxykinase 2 (PCK2) (Cat. No. Hs00356436_m1)

Animal studies

All animal care and experimental protocols were approved by the animal ethics committee of the University of Cambridge and the Home Office of the United Kingdom government. 6-week old female cd-1 nude mice (Charles River Inc) were ovariectomised and implanted in the flank with one-half of a 1.5 mg 60-day release estradiol pellet (Cat. No. SE-121, Innovative Research of America, USA). After one week, mice that were to receive retinoic acid were implanted with a further pellet in the neck containing 5 mg ATRA (Cat. No. V-111, Innovative

Research). At the same time, the diet was changed to a diet containing 0.1% or 0.25% fenofibrate (constituted by Lillico Biotechnology, Surrey, UK) or the same diet without fenofibrate. Approximately 1 million Ishikawa cells in 100 µl Matrigel (Cat. No. 35-6234, BD Biosciences, UK) were injected subdermally into the left flank, raising a bleb. Mice were checked daily and weighed twice-weekly to ensure good health. At the end of the experiment the mice were culled, the tumours were excised from surrounding tissue and weighed. Immunohistochemistry for PPARα was performed on selected harvested tumour samples as previously described [12]

Statistics

All statistical analyses were performed under the R statistical environment [27]. Where appropriate, Student's t-test, Wilcoxon Rank-Sum test or one-way ANOVA was used for comparisons as indicated. Methods particular to gene array analysis are described above.

Results

Growth inhibitory effect of fenofibrate is precipitated by retinoic acid

Treatment of Ishikawa cells with 50 µM fenofibrate over 48 hours significantly reduced cell growth (27%, $p < 0.05$). Although treatment with 10 µM all-trans retinoic acid (ATRA) had no significant effect on cell growth, combined treatment with 50 µM fenofibrate caused a greater reduction in cell number than produced by fenofibrate alone (49%, $p < 0.05$). Treatment with high dose ATRA combined with fenofibrate caused >90% reduction in cell number ($p < 0.05$) (figure 1a). In addition, a dose-response was demonstrated when 10, 30 or 50 µM fenofibrate was combined with 30 µM ATRA (figure 1b). Similar effects were seen with retinoic acid, where no inhibition of cell growth was detected below 50 µM in the absence of fenofibrate (data not shown). BrdU assays




Figure 1
Effect of fenofibrate and retinoic acid (48 hours exposure) on endometrial cancer cell growth, using cell counting (Top) and MTS assay (Bottom).




Figure 2
Top – DNA content analysis. The bars represent the ratio of cells in G1:G2M phase of the cell cycle after 12 hours of treatment with drug (RA+ = ATRA 10 µM. RA++ = ATRA 30 µM, FN+ = Fenofibrate 30 µM, FN++ = Fenofibrate 100 µM). Bottom – Apoptosis by FACS analysis using Annexin-FITC staining. Cells were treated for 3 hours. FN = fenofibrate, RG = rosiglitazone, St = staurosporine, a potent apoptogen.

again demonstrated inhibition of cell proliferation with fenofibrate but failed to show a similar effect with high dose rosiglitazone, a PPAR α agonist. Other PPAR agonists and antagonists were tried but failed to show a significant effect in the doses used (data not shown)

Fenofibrate induces cell cycle arrest and apoptosis in Ishikawa cells

The proportion of cells in G₁ and G₂M phases of the cell cycle were measured by FACS analysis after 12-hour exposure to fenofibrate. Application of 50 μ M or 100 μ M fenofibrate increased the ratio of cells in G₁ phase to G₂M phase of the cell cycle (data not shown, $p < 0.05$, Student's t-test). Treatment of ISK cells with 100 μ M fenofibrate combined with 30 μ M ATRA, however, had a greater effect on cell cycle progression than either drug alone (figure 2a). The effect of fenofibrate and retinoic acid on apoptosis was also measured by FACS analysis, and similarly demonstrated an increase in apoptosis induction when the drugs were combined (figure 2b).

Although our previous results had suggested a direct effect of fenofibrate on the PPAR α receptor, experiments conducted using RNAi for PPAR α failed to show a dramatic reduction in fenofibrate effect. This was despite achieving consistent downregulation of >75% in PPAR α expression as measured by RT-PCR (Figure 3a). The effect of ATRA combined with fenofibrate on cell growth was similar in the presence or absence of PPAR α RNAi (figure 3b). However, the total viable cell number in the PPAR α transfected control-treated cells was approximately 40% lower than the control RNAi transfected cells. After correcting for this difference, the effect of 75 μ M fenofibrate (48 hours treatment) on cell growth inhibition was less in the RNAi transfected cells ($p = 0.05$, t-test) (figure 3b).

Genes affected by fenofibrate in Ishikawa cells

Using Independent Component analysis (ICA), a component relating to treatment with high-dose fenofibrate (100 μ M) was identified in both gene array experiments (slide 1 & slide 2) ($p < 0.01$ after correction for multiple testing). Other significant components identified from ICA related to batch effects (hybridisation date, RNA preparation date, hybridisation batch) ($p < 0.01$, ANOVA). Batch effects were so strong that the "dose" component was only the 9th largest of 30 components. No component could be identified relating to "low dose" (10 μ M) fenofibrate treatment.

Significant genes from the identified component were selected as those with absolute component loading values greater than 3 standard deviations, treating positive and negative values separately. This was an arbitrary value providing a reasonable number of genes all of which had component loadings outside the inflexion of the plotted




Figure 3

Top – mRNA expression as quantified using Taqman RT-PCR following RNAi for PPAR α using RNAi concentration shown. Middle – Cell growth following RNAi and treatment with fenofibrate (FN) and ATRA (RA) for 48 hours. Cell numbers were corrected to adjust for the reduced cell growth following PPAR α RNAi compared to control RNAi. Bottom – Apoptosis measured by FACS analysis using Annexin-FITC staining following RNAi and treatment with fenofibrate and retinoic acid for 3 hours.




Figure 4
 Genes identified as differentially expressed after treatment with high-dose (100 μM) fenofibrate. The plots show upregulation (red) and downregulation (green) of expression compared to DMSO control-treated cells. Cells were treated with 10 μM ("low dose") or 100 μM ("high dose") fenofibrate for 48 hours. *Left* – Normalised uncorrected data. *Right* – Data corrected using ICA-based filtering and removal of artefactual components. The data is improved but reveals an aberrant sample in the high-dose group (the first).

curve (see additional file 1). At this threshold, 425 gene spots representing 213 annotated unique genes from Slide 1 were identified as differentially expressed following treatment with high dose fenofibrate. A similar number of differentially expressed genes were additionally identified from the second microarray experiment (using Slide 2 which contained a greater number of non-annotated clones). A subset of genes from the Slide 1 geneset is displayed as an expression plot in figure 4a.

Three genes were chosen for RT-PCR validation of the gene array (figure 5): Cyclin D1 (CCND1), Phosphoenolpyruvate carboxykinase 2 (PCK2) and Methionine Adenosyltransferase 2A. These genes were chosen due to a combination of their loading value in the component of interest (suggesting a significant effect) as well as their biological interest. Both CCND1 and MAT2A were down-regulated following treatment with fenofibrate 100 μ M. RT-PCR analysis confirmed downregulation in gene expression (CCND1 3.2-fold, MAT2A 5.6-fold, $p < 0.05$). RT-PCR analysis confirmed a small increase in PCK2 expression (1.3-fold, $p = \text{NS}$) in the treated cells but this was less than that seen with the array data (figure 5c).

Gene Ontology analysis revealed a number of functional groups which were significantly over-represented in the transcripts identified as altered by fenofibrate. A functional group was considered to be over-represented in the gene list when the number of gene spots belonging to that group was higher than expected from the distribution of genes on the array (chi-squared test). The following functional groups of interest were identified as being over-represented by the genes altered with fenofibrate treatment (see table 1): nucleotide binding (GO:0000166), MAP kinase-related (GO:0017017, GO:0000188, GO:0000185), cell growth (GO:0016049, GO:0008283), cyclin-dependent protein kinase inhibitor (GO:0004861), insulin-like growth factor binding (GO:0005520), platelet-derived growth factor receptor ligand (GO:0005161), metalloprotease inhibitor (GO:0008191) and cell cycle arrest (GO:0007050). The full table of GO classifications, by significance level, is provided in additional file 2.

Effect of fenofibrate on tumour growth *in vivo*

Two separate *in vivo* experiments were performed (figure 6). In the first, mice were allocated to each of three groups – (1) normal diet (N = 6), (2) low-dose fenofibrate (0.1%) (N = 4), (3) high-dose fenofibrate (0.25%) (N = 6). The median tumour weights at 21 days in the three groups were (1) 684 g (2) 1280 g (3) 439 g. There was a trend towards lower tumour weights in the high-dose-treated group compared to those receiving normal diet ($p = 0.046$, Mann-Whitney test, one-sided). Surprisingly, however, the group fed low-dose fenofibrate demon-




Figure 5
Comparison between array data and gene expression (as measured using Taqman RT-PCR). The "532 control" column represents the green-channel array data from the pooled cDNA (from DMSO-treated control samples) and is used as an internal control. Three genes were chosen for verification of the array: *top* – Cyclin D1, *middle* – Methionine Adenosyltransferase 2-alpha, *bottom* – Phosphoenolpyruvate carboxykinase 2.

Table 1: Selected genes derived from ICA & GO analysis of gene array experiments. Genes are combined into ontologies. Average Fold Changes are calculated from normalised values over the number of spots shown for each gene. As can be seen, not all genes are differentially expressed according to t-test p-values. Genes are instead selected based on their ICA loading (see discussion)

GO Description	Gene symbol	Description	No of spots	Ave FC	Ave ttest p	
activation of MAPKKK (GO:0000185) cell cycle arrest (GO:0007050)	GADD45B	growth arrest and DNA-damage-inducible, beta	2	2.6	0.090	
	EIF4G2	eukaryotic translation initiation factor 4 gamma, 2	2	1.2	0.249	
	GADD45A	growth arrest and DNA-damage-inducible, alpha	2	5.1	0.003	
	PPP1R15A	protein phosphatase 1, regulatory (inhibitor) subunit 15A	2	1.4	0.085	
	PR48	protein phosphatase 2A 48 kDa regulatory subunit	1	-1.1	0.427	
cell death (GO:0008219)	SESN3	sestrin 3	2	-1.3	0.161	
	TP53	tumor protein p53 (Li-Fraumeni syndrome)	4	1.7	0.036	
	EIF4G2	eukaryotic translation initiation factor 4 gamma, 2	2	1.2	0.249	
	EMP3	epithelial membrane protein 3	2	1.7	0.111	
	cell growth (GO:0016049)	P8	p8 protein (candidate of metastasis 1)	2	3.1	0.005
SLC3A2		solute carrier family 3 (activators of dibasic and neutral amino acid transport), member 2	4	2.0	0.012	
cell proliferation (GO:0008283)	EGFR	epidermal growth factor receptor (erythroblastic leukemia viral (v-erb-b) oncogene homolog, avian)	2	2.2	0.009	
	FTH1	ferritin, heavy polypeptide 1	1	1.2	0.301	
	IGFBP4	insulin-like growth factor binding protein 4	2	1.8	0.002	
	OSMR	oncostatin M receptor	2	1.7	0.088	
	PDGFA	platelet-derived growth factor alpha polypeptide	2	1.5	0.032	
	PDGFB	platelet-derived growth factor beta polypeptide (simian sarcoma viral (v-sis) oncogene homolog)	4	2.2	0.083	
	SYK	spleen tyrosine kinase	2	-1.1	0.631	
	TGFA	transforming growth factor, alpha	2	-1.0	0.907	
	TGFBI	transforming growth factor, beta-induced, 68kDa	2	2.1	0.164	
	TP53	tumor protein p53 (Li-Fraumeni syndrome)	4	1.7	0.036	
	VEGF	vascular endothelial growth factor	2	6.6	0.175	
	G1/S transition of mitotic cell cycle (GO:0000082) insulin-like growth factor binding (GO:0005520)	CCND1	cyclin D1 (PRAD1: parathyroid adenomatosis 1)	2	-1.3	0.168
		CYR61	cysteine-rich, angiogenic inducer, 61	2	2.9	0.013
KAZALD1		Kazal-type serine protease inhibitor domain 1	2	-1.1	0.770	
IGFBP4		insulin-like growth factor binding protein 4	2	1.8	0.002	
CTGF		connective tissue growth factor	2	2.9	0.006	
IGFBP5		insulin-like growth factor binding protein 5	4	2.7	0.041	
metalloprotease inhibitor (GO:0008191)		TIMPI	tissue inhibitor of metalloproteinase 1 (erythroid potentiating activity, collagenase inhibitor)	3	1.4	0.250
		MAT2A	methionine adenosyltransferase II, alpha	4	-2.9	0.001
methionine adenosyltransferase (GO:0004478)		MARS	methionine-tRNA synthetase	2	6.7	0.006


strated the largest tumours at 21 days ($p < 0.05$, Mann-Whitney test).

In the second experiment, 8 mice were allocated to each of four groups, to receive (1) normal diet, (2) high-dose fenofibrate (0.25%), (3) ATRA, (4) ATRA + fenofibrate. All mice treated with ATRA were culled at day 14 due to weight loss (>10% starting weight) and the majority displayed a local reaction to the ATRA pellet. The remaining mice were culled at day 28. There were no significant differences between the tumour weights in the groups compared (fenofibrate vs normal diet; ATRA vs ATRA + fenofibrate) (Mann-Whitney test, $p > 0.05$)

Discussion

To our knowledge, we are the only group to have published on the effects of fenofibrate in endometrial cancer *in vitro* [12]. Our previous data demonstrated a moderate inhibitory effect of fenofibrate on endometrial cancer cell growth but with limited understanding of the mechanism of this effect. This study significantly expands on this knowledge.

We have confirmed, using additional methods, that Ishikawa endometrial cancer cell growth is inhibited by fenofibrate. We have also confirmed, by using FACS analysis, that fenofibrate increases apoptosis in ISK cells *in*

**Figure 6**

Tumour weights in nude mice following endometrial cancer cell xenografting and treatment with "high dose" (0.25%) or "low dose" (0.1%) fenofibrate. *Top* – n = 6 mice per group treated for 21 days. *Middle* – n = 8 mice per group treated for 14 days and *Bottom* – n = 8 mice per group treated for a further 10 days with fenofibrate alone. Tumour weights were lower in the high-dose fenofibrate-treated mice in the first experiment, but this was not repeated in the second experiment.

in vitro. These effects appear not to be associated with excess toxicity. More importantly, we were able to show, using drugs acting on the same pathway, that the inhibitory

effects on cell growth can be potentiated using agents acting on the same pathway. The use of either all-trans-retinoic acid (ATRA) or its metabolite 9-cis retinoic acid in this regard demonstrated similar results, as might be expected. Furthermore it might be anticipated that these agents potentiate the effect of fenofibrate via their action on the PPAR α receptor [28]. Similarly, a synergistic effect of RXR and PPAR agonists has previously been demonstrated in breast cancer [29] and bladder cancer [30] cell lines. However, although retinoic acid derivatives alone have been demonstrated to show activity in breast cancer [31,32] and other tumours, particularly leukaemias [33,34], there is limited evidence that retinoids are effective in endometrial cancer [15,35]. Indeed, our data suggest that, even at comparatively high doses of retinoic acid, no inhibition of Ishikawa cell growth occurs.

The mechanism of action of fenofibrate is, as yet, unknown. A parallel might be drawn between the effect of fenofibrate (a PPAR α agonist) in endometrial cancer and thiazolidenediones (TZDs, PPAR γ agonists), such as rosiglitazone and troglitazone, in breast cancer. TZD's have been demonstrated to cause inhibition of cancer cell growth *in vitro* and/or *in vivo* in a number of cancers, particularly breast [29,36], colon [37] and salivary gland [38]. Although yet to be fully determined, it would appear that at least one mechanism for this action involves inhibition of translation and transition through G₁-S phase of the cell cycle [39]. These effects are associated with decreased expression of D1/E cyclins in the absence of a change in p21, or cyclin-dependent kinases (CDK's) and are independent of PPAR γ activation [39].

Anti-tumour effects of PPAR α -specific agonists appear to be less commonly reported. One reason for this may be the perception that PPAR α agonists are carcinogenic due to their tumorigenic effects in rodent liver [14]. Furthermore, agonism of PPAR α has been shown to increase proliferation in MCF-7 breast cancer cells [40]. This effect, however, is reported inconsistently and PPAR α agonists have also demonstrated efficacy *in vitro* against melanoma [41] and breast cancer cell lines [42]. These studies outline various mechanisms which explain the effect of fenofibrate on cancer cells. Firstly, PPAR α agonist effects on the cell cycle and apoptosis are mediated by environmental agents and the MAP kinase pathway [43]. PPAR activity is also closely linked to the eicosanoid pathway of inflammation and PPAR α induces cyclo-oxygenase-2 (PTGS2) expression, which is linked to the development of colon cancer [44]. Furthermore, there is significant cross-talk between PPAR receptors and many other nuclear hormone receptor subfamilies [45-47]. The most obvious of these is the link to the retinoid receptor family, but even within this subfamily, sex-steroid hormone influence is evident. Examples of this are the estrogen-dependence of

retinoic acid metabolism [48] and the fact that PPAR α -mediated anti-inflammatory action stimulated by TNF α is interrupted by the anti-progesterone RU486 in HUVEC's [49]. Probably the strongest nuclear hormone interrelationship with the PPAR receptors is to insulin and the IGF-receptor [47,50,51]. Although the primary clinical function of TZD's is to increase insulin sensitivity, PPAR α agonists also reduce insulin resistance and insulin concentration *in vivo* [52]. Furthermore, the strong association between PPARs and lipid and glucose homeostasis appears to be a primary factor in the development of obesity [47,53]. Taken together, these features support the hypothesis that the PPAR pathway might harbour anti-neoplastic targets in endometrial cancer, which is known to be promoted by obesity and hyperinsulinaemia [5]. Indeed, the inhibition of G₁-S phase progression, demonstrated in this study with fenofibrate, is consistent with the anticipated effects of targeting of PPAR α in endometrial cancer.

We attempted to demonstrate that the inhibitory effect of fenofibrate on Ishikawa cell growth was specific for PPAR α . Support for this hypothesis was provided by the synergistic effects demonstrated from combined treatment with fenofibrate and retinoic acid, and from the lack of similar effect seen with therapeutic doses of rosiglitazone. We have previously demonstrated a functional PPAR α receptor using a luciferase reporter assay [12]. In this study we also demonstrated absence of mutation within the coding region of the hPPAR α gene in the cell lines tested, as it is known that up to 15% of endometrial cancers harbour variants of PPAR γ [54].

The gene array experiments were designed to better understand the mechanism of action of this drug, at the transcript level. The traditional approach to identification of differentially expressed genes involves the use of first-order statistics such as the t-test (eg Cyber-T [55]) or permutation-based methods (eg SAM [56]), often in combination with a "threshold of interest" applied to fold-change values in gene expression. Here we have applied Independent Component Analysis (ICA) [26,57] to identify a "gene signature" which correlates with our biological feature of interest, ie response to high-dose fenofibrate. Using this method we are not only able to filter out stochastic noise from the microarray data, but also to exclude components of the data associated with experimental artefact such as batch effects. Using ICA we could not discern a gene signature associated with the response to low-dose (10 μ M) fenofibrate, suggesting that the effects on the transcriptome at this dose were small, consistent with the lack of biological effect seen. Of more interest, however, was the response to high-dose (100 μ M) fenofibrate, which was seen in a significant proportion of the transcriptome (see figure 4). One advantage of

using a factor analysis method is the avoidance of supervision bias when identifying differentially expressed genes. Here we combined ICA with Gene Ontology [58] analysis to identify groups of genes with shared function which were differentially expressed with high-dose fenofibrate.

Amongst the identified gene ontology groups, ontologies related to cell proliferation were over-represented in the list of differentially expressed genes. These included cell proliferation, cell growth, cell cycle arrest, and ontologies related to MAPkinase signalling. Within these functional groups, genes that were up- or down-regulated were generally consistent with the experimental findings of inhibition of G₁-S phase progression. Downregulation of Cyclin D1 (CCND1) was accompanied by upregulation of p8, p21, GADD45A, GADD45B and TP53. Cyclin D1 is known to be upregulated in endometrial cancer and is a common feature of endometrial carcinogenesis [59,60]. Cyclin D1 plays a major role in cell cycle progression and is therefore central to the proliferative ability of many cancers [61]. In view of its importance in cell cycle progression, we chose Cyclin D1 as one of the genes with which to verify the array findings and confirmed its downregulation (see figure 4b). Conversely, and as might be expected, p21 (cyclin-dependent kinase inhibitor 1a, CDKN1A) was upregulated following administration of high-dose fenofibrate. P8 (COM-1) was also upregulated. P8 encodes for a protein which inhibits cancer cell growth in estrogen-dependent breast cancer cell lines [62]. TP53 is another protein central to the regulation of cancer cell growth [63] and its gene expression was also upregulated.

Although cell cycle-related genes are important anti-neoplastic targets, other pathways are particularly relevant in endometrial carcinogenesis. There is overwhelming evidence that obesity plays a pathogenetic role in endometrial cancer and is most likely mediated by IGF [5,7]. In particular, the IGF binding proteins appear to play a pivotal role in estrogen-dependent endometrial cancer cell growth [64]. Transcript levels of both IGFBP4 and IGFBP5 as well as Connective Tissue Growth Factor (CTGF, IGFBP8) were increased in response to fenofibrate, suggesting that this lipid-lowering agent may exert some direct action on pathways known to be relevant to endometrial carcinogenesis.

The gene array experiments also revealed a further pathway which may be involved in the inhibition of Ishikawa endometrial cancer cell growth. Two of the genes that were most differentially regulated by fenofibrate were methionine adenosyltransferase 2-alpha (MAT2A) (FC = -2.7, $p < 0.01$) and methionine-tRNA synthetase (FC = 6.7, $p < 0.01$). These two genes form part of the methionine metabolism pathway, central to normal cellular function

and DNA synthesis. In particular, MAT2A catalyses the conversion of L-Methionine to S-Adenosyl methionine, the principal methyl donor. Induction of MAT2A is mediated by TNF α and reduced expression is associated with inhibition of cell growth [65,66]. Induction of MAT2A has also been correlated with disease progression in colon cancer [67]. MAT2A may also prove to be an important therapeutic target in view of the increasingly realised epigenetic role of methylation in endometrial carcinogenesis [68,69]. By contrast, whether MARS plays a role in carcinogenesis is not known, and the induction of this gene may simply be a reflection of the degree of translational change taking place under the influence of fenofibrate as suggested earlier [39].

In addition to gene array analysis, we applied RNA-interference [70] in order to establish whether the inhibitory effects of fenofibrate seen were due to activation of the PPAR α receptor. Although our results demonstrated some diminution of effect at lower doses of fenofibrate, the main effects were unchanged despite confirmation of at least 75% knock-down of PPAR α expression. A further unexpected effect encountered during the RNAi experiments was the significant and consistent reduction in cell viability seen following knock-down of PPAR expression. This might be explained by the central role PPAR α plays in cellular metabolism and may also account for a blunting of any direct effect on cell growth induced by PPAR α knock-down.

Furthermore, it was expected that some of the differentially expressed genes identified in the array experiments would comprise those known to be regulated by fenofibrate. Comparison of our gene list to previously published data [71,72] failed to demonstrate a similar pattern of differentially expressed genes, although there were occasional similarities (eg \uparrow Stearoyl-CoA desaturase). One explanation for this may be that the small changes in metabolic gene expression remained undetectable in an environment of apoptosis and cell death. A second explanation, however, could be that the majority of published gene expression data relating to fenofibrate in rodents is less applicable to human cells.

This latter possibility may also explain the absence of a consistent demonstrable effect on tumour growth *in vivo*. Results from our initial experiment in mice were encouraging, but failed to show reduction in tumour growth, or a difference when retinoic acid was added, in the second experiment. Furthermore, immunohistochemistry failed to identify significant levels of PPAR α protein expression in the harvested tumour samples, raising the possibility that the receptor level is not maintained within the animal model environment (data not shown). There are known species differences in the effects of fenofibrate: humans

have a lower level of liver PPAR α expression than rodents [73], the promoter for human Acyl CoA Oxidase is different from that in rodents [74] and effects on the cell cycle in mice liver are different from that in humans [75]. Taken together, these findings may explain the unexpected results from our *in vivo* experiments. A further unexpected finding from the animal experiments was the increased tumour size seen in the low dose fenofibrate arm compared to normal diet ($p = 0.04$). It was postulated that the serum level of fenofibrate in the low dose arm may have been well below the anticipated dose, and that a low fenofibrate dose may have had a stimulatory effect on ISK cell growth. A further experiment was therefore conducted to investigate the effect of prolonged (>96 hours) low-dose (0.1 μ M – 3 μ M) exposure to fenofibrate, but failed to show any effect on cell growth (data not shown). The increased tumour size in this group was therefore considered to be a statistical, rather than a true biological effect. A further possibility explaining the absence of expected effect may have been inadequate dosing, although the doses chosen were based on previously published data to achieve desired serum fenofibrate levels [76]. One further point to note was that the mechanism for delivery of ATRA ultimately proved to be a problem resulting in toxicity, either due to the dose (although this was well within established dosage according to the manufacturer) or to local reaction to the pellet. Our preference would have been to use Liposomal ATRA (ATRA-iv, Antigenics Inc, New York) but it proved impossible to obtain this from the manufacturer. It is possible that the toxicity encountered from the pellets prevented a demonstrable response to fenofibrate in the final experiment.

PPAR γ agonists have been demonstrated to show tumour inhibition *in vitro* and *in vivo*, in a number of tumour types such as breast and colon. PPAR α agonists, however, have so far failed to show similar results, despite inhibitory effects on tumour growth in breast cancer [42] and melanoma [41] *in vitro*. It is highly plausible that the species differences in PPAR α activity [72,77] is at least in part responsible for the difficulties in demonstrating efficacy of PPAR α agonists in a rodent model of human tumours. The promising effects of these agents *in vitro*, however, suggest that the lack of a consistent effect on endometrial cancer in mice warrants further investigation, most likely requiring the use of an alternative animal model.

The full mechanism of action of fenofibrate remains to be clarified but it appears to exert effects on cell cycle regulators in addition to pathways pertinent to endometrial carcinogenesis. The data presented suggest that PPAR α remains a potential therapeutic target in endometrial cancer, whilst the addition of retinoic acid to fenofibrate creates a potent therapeutic combination *in vitro*. The development of an appropriate animal model, however,

remains essential to demonstrating the applicability of this promising combination in the treatment of endometrial cancer.

Competing interests

The author(s) declare that they have no competing interests.

Additional material

Additional File 1

Contains supplemental figures referred to in the main text. The main text is self-contained and does not require these figures which are provided for further reference for the interested reader

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1476-4598-5-13-S1.pdf>]

Additional File 2

Contains a supplemental table referred to in the text and is an expansion of table 1 (from the main paper).

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1476-4598-5-13-S2.xls>]

Acknowledgements

We thank Dr Andrew Sharkey, Dr Roberto Catalano and Amanda Evans for their assistance in design and construction of the arrays used in these studies. This work was carried out in the MRC Reproductive Angiogenesis Cooperative Group and was supported in part by program grant G9623012.

References

- Pisani P, Bray F, et al.: **Estimates of the world-wide prevalence of cancer for 25 sites in the adult population.** *Int J Cancer* 2002, **97(1)**:72-81.
- Ferlay J, Bray F, Pisani P, Parkin DM: **GLOBOCAN 2002: Cancer Incidence, Mortality and Prevalence Worldwide, IARC CancerBase No. 5, version 2.0.** 2004 [<http://www-dep.iarc.fr/>]. IARC-Press, Lyon
- Key TJ, Pike MC: **The dose-effect relationship between 'unopposed' oestrogens and endometrial mitotic rate: its central role in explaining and predicting endometrial cancer risk.** *Br J Cancer* 1988, **57(2)**:205-12.
- Hale GE, Hughes CL, et al.: **Endometrial cancer: hormonal factors, the perimenopausal "window of risk," and isoflavones.** *J Clin Endocrinol Metab* 2002, **87(1)**:3-15.
- Kaaks R, Lukanova A, et al.: **Obesity, endogenous hormones, and endometrial cancer risk: a synthetic review.** *Cancer Epidemiol Biomarkers Prev* 2002, **11(12)**:1531-43.
- Bray GA: **The underlying basis for obesity: relationship to cancer.** *J Nutr* 2002, **132(11 Suppl)**:3451S-3455S.
- Kaaks R, Lukanova A: **Energy balance and cancer: the role of insulin and insulin-like growth factor-I.** *Proc Nutr Soc* 2001, **60(1)**:91-106.
- Crnici I, Christofori G: **Novel technologies and recent advances in metastasis research.** *Int J Dev Biol* 2004, **48(5-6)**:573-81.
- Ramaswamy S, Ross KN, et al.: **A molecular signature of metastasis in primary solid tumors.** *Nat Genet* 2002, **33(1)**:49-54.
- Ferguson SE, Olshen AB, et al.: **Gene expression profiling of tamoxifen-associated uterine cancers: evidence for two molecular classes of endometrial carcinoma.** *Gynecol Oncol* 2003, **92(2)**:719-725.
- Risinger JI, Maxwell GL, et al.: **Microarray Analysis Reveals Distinct Gene Expression Profiles among Different Histologic Types of Endometrial Cancer.** *Cancer Res* 2003, **63(1)**:6-11.
- Holland CM, Saidi SA, et al.: **Transcriptome analysis of endometrial cancer identifies peroxisome proliferator-activated receptors as potential therapeutic targets.** *Mol Cancer Ther* 2004, **3(8)**:993-1001.
- Saidi SA, Holland CM, et al.: **Independent component analysis of microarray data in the study of endometrial cancer.** *Oncogene* 2004, **23(39)**:6677-83.
- Michalik L, Desvergne B, et al.: **Peroxisome-proliferator-activated receptors and cancers: complex stories.** *Nat Rev Cancer* 2004, **4(1)**:61-70.
- Siddiqui NA, Thomas EJ, et al.: **Retinoic acid receptors and retinoid binding proteins in endometrial adenocarcinoma: differential expression of cellular retinoid binding proteins in endometrioid tumours.** *Int J Cancer* 1995, **64(4)**:253-63.
- Dawson MI, Chao WR, et al.: **Effects of trans-retinoic acid, 9-cis-retinoic acid, 1alpha,25-(dihydroxy)vitamin D3 and a novel apoptosis-inducing retinoid on breast cancer and endothelial cell growth.** *Cancer Lett* 1999, **133(1)**:1-8.
- Nishida M, Kasahara K, et al.: **Establishment of eighteen clones of Ishikawa cells.** *Hum Cell* 1996, **9(2)**:109-16.
- Gong Y, Murphy LC, et al.: **Hormonal regulation of proliferation and transforming growth factors gene expression in human endometrial adenocarcinoma xenografts.** *J Steroid Biochem Mol Biol* 1994, **50(1-2)**:13-9.
- [<http://www.ecacc.org.uk>].
- [<http://www.fcspress.com>].
- Chomczynski P: **A reagent for the single-step simultaneous isolation of RNA, DNA and proteins from cell and tissue samples.** *Biotechniques* 1993, **15(3)**:532-4. 536-7
- Evans AL, Sharkey AS, et al.: **Generation and use of a tailored gene array to investigate vascular biology.** *Angiogenesis* 2004, **6(2)**:93-104.
- Rossi M, Sharkey AM, et al.: **Identification of genes regulated by interleukin-1beta in human endometrial stromal cells.** *Reproduction* 2005, **130(5)**:721-9.
- British Standards Institution: **ISO 5725: Accuracy (trueness and precision) of measurement methods and results.** 1994.
- Petalidis L, Bhattacharyya S, et al.: **Global amplification of mRNA by template-switching PCR: linearity and application to microarray analysis.** *Nucleic Acids Res* 2003, **31(22)**:e142.
- Martoglio AM, Miskin JW, et al.: **A decomposition model to track gene expression signatures: preview on observer-independent classification of ovarian cancer.** *Bioinformatics* 2002, **18(12)**:1617-1624.
- [<http://www.r-project.org>].
- Kliwer SA, Umesono K, et al.: **Convergence of 9-cis retinoic acid and peroxisome proliferator signalling pathways through heterodimer formation of their receptors.** *Nature* 1992, **358(6389)**:771-4.
- Mehta RG, Williamson E, et al.: **A ligand of peroxisome proliferator-activated receptor gamma, retinoids, and prevention of preneoplastic mammary lesions.** *J Natl Cancer Inst* 2000, **92(5)**:418-23.
- Fauconnet S, Lascombe I, et al.: **Differential regulation of vascular endothelial growth factor expression by peroxisome proliferator-activated receptors in bladder cancer cells.** *J Biol Chem* 2002, **277(26)**:23534-43.
- Elstner E, Müller C, et al.: **Ligands for peroxisome proliferator-activated receptor gamma and retinoic acid receptor inhibit growth and induce apoptosis of human breast cancer cells in vitro and in BXH mice.** *Proc Natl Acad Sci USA* 1998, **95(15)**:8806-11.
- Zhu WY, Jones CS, et al.: **Retinoic acid inhibition of cell cycle progression in MCF-7 human breast cancer cells.** *Exp Cell Res* 1997, **234(2)**:293-9.
- Lallemant-Breitenbach V, Guillemain MC, et al.: **Retinoic acid and arsenic synergize to eradicate leukemic cells in a mouse model of acute promyelocytic leukemia.** *J Exp Med* 1999, **189(7)**:1043-52.
- Simoni D, Tolomeo M: **Retinoids, apoptosis and cancer.** *Curr Pharm Des* 2001, **7(17)**:1823-37.
- Carter CA, Pogribny M, et al.: **Effects of retinoic acid on cell differentiation and reversion toward normal in human**

- endometrial adenocarcinoma (RL95-2) cells. *Anticancer Res* 1996, **16**(1):17-24.
36. Clay CE, Namen AM, et al.: **Influence of J series prostaglandins on apoptosis and tumorigenesis of breast cancer cells.** *Carcinogenesis* 1999, **20**(10):1905-11.
 37. Sarraf P, Mueller E, et al.: **Differentiation and reversal of malignant changes in colon cancer through PPARgamma.** *Nat Med* 1998, **4**(9):1046-52.
 38. Begum NM, Nakashiro K, et al.: **Expression of peroxisome proliferator-activated receptor gamma and the growth inhibitory effect of its synthetic ligands in human salivary gland cancer cell lines.** *Int J Oncol* 2002, **20**(3):599-605.
 39. Palakurthi SS, Aktas H, et al.: **Anticancer effects of thiazolidinediones are independent of peroxisome proliferator-activated receptor gamma and mediated by inhibition of translation initiation.** *Cancer Res* 2001, **61**(16):6213-8.
 40. Suchanek KM, May FJ, et al.: **Peroxisome proliferator-activated receptor alpha in the human breast cancer cell lines MCF-7 and MDA-MB-231.** *Mol Carcinog* 2002, **34**(4):165-71.
 41. Kang HY, Chung E, et al.: **Expression and function of peroxisome proliferator-activated receptors in human melanocytes.** *Br J Dermatol* 2004, **150**(3):462-8.
 42. Avis I, Hong SH, et al.: **Five-lipoxygenase inhibitors can mediate apoptosis in human breast cancer cell lines through complex eicosanoid interactions.** *FASEB J* 2001, **15**(11):2007-9.
 43. Roberts RA, Chevalier S, et al.: **PPAR alpha and the regulation of cell division and apoptosis.** *Toxicology* 2002, **181-182**:167-70.
 44. Roberts-Thomson SJ: **Peroxisome proliferator-activated receptors in tumorigenesis: targets of tumour promotion and treatment.** *Immunol Cell Biol* 2000, **78**(4):436-41.
 45. Hiji AK, Michalik L, et al.: **PPARs: transcriptional effectors of fatty acids and their derivatives.** *Cell Mol Life Sci* 2002, **59**(5):790-8.
 46. Chu R, Madison LD, et al.: **Thyroid hormone (T3) inhibits ciprofibrate-induced transcription of genes encoding beta-oxidation enzymes: cross talk between peroxisome proliferator and T3 signaling pathways.** *Proc Natl Acad Sci USA* 1995, **92**(25):11593-7.
 47. Moller DE, Berger JP: **Role of PPARs in the regulation of obesity-related insulin sensitivity and inflammation.** *Int J Obes Relat Metab Disord* 2004, **27**(Suppl 3):S17-21.
 48. Okamoto K, Andreola F, et al.: **Differences in uptake and metabolism of retinoic acid between estrogen receptor-positive and -negative human breast cancer cells.** *Cancer Chemother Pharmacol* 2000, **46**(2):128-34.
 49. Xu X, Otsuki M, et al.: **RU486 antagonizes the inhibitory effect of peroxisome proliferator-activated receptor alpha on interleukin-6 production in vascular endothelial cells.** *J Steroid Biochem Mol Biol* 2002, **81**(2):141-6.
 50. Lewitt MS, Brismar K, et al.: **Responses of insulin-like growth factor (IGF)-I and IGF-binding proteins to nutritional status in peroxisome proliferator-activated receptor-alpha knockout mice.** *Growth Horm IGF Res* 2001, **11**(5):303-13.
 51. Shalev A, Siegrist-Kaiser CA, et al.: **The peroxisome proliferator-activated receptor alpha is a phosphoprotein: regulation by insulin.** *Endocrinology* 1996, **137**(10):4499-502.
 52. Guerre-Millo M, Gervois P, et al.: **Peroxisome proliferator-activated receptor alpha activators improve insulin sensitivity and reduce adiposity.** *J Biol Chem* 2000, **275**(22):16638-42.
 53. Evans RM, Barish GD, et al.: **PPARs and the complex journey to obesity.** *Nat Med* 2004, **10**(4):355-61.
 54. Smith WM, Zhou XP, et al.: **Opposite association of two PPARG variants with cancer: overrepresentation of H449H in endometrial carcinoma cases and underrepresentation of P12A in renal cell carcinoma cases.** *Hum Genet* 2001, **109**(2):146-51.
 55. Baldi P, Long AD: **A Bayesian framework for the analysis of microarray expression data: regularized t-test and statistical inferences of gene changes.** *Bioinformatics* 2001, **17**(6):509-19.
 56. Tusher VG, Tibshirani R, et al.: **Significance analysis of microarrays applied to the ionizing radiation response.** *Proc Natl Acad Sci USA* 2001, **98**(9):5116-21.
 57. Liebermeister W: **Linear modes of gene expression determined by independent component analysis.** *Bioinformatics* 2002, **18**(1):51-60.
 58. Ashburner M, Ball CA, et al.: **Gene ontology: tool for the unification of biology. The Gene Ontology Consortium.** *Nat Genet* 2000, **25**(1):25-9.
 59. Skomedal H, Kristensen GB, et al.: **TP53 alterations in relation to the cell cycle-associated proteins p21, cyclin D1, CDK4, RB, MDM2, and EGFR in cancers of the uterine corpus.** *J Pathol* 1999, **187**(5):556-62.
 60. Smid-Koopman E, Blok LJ, et al.: **Gene expression profiles of human endometrial cancer samples using a cDNA-expression array technique: assessment of an analysis method.** *Br J Cancer* 2000, **83**(2):246-51.
 61. Lundberg AS, Weinberg RA: **Control of the cell cycle and apoptosis.** *Eur J Cancer* 1999, **35**(4):531-9.
 62. Jiang WG, Davies G, et al.: **Com-1/P8 in oestrogen regulated growth of breast cancer cells, the ER-beta connection.** *Biochem Biophys Res Commun* 2005, **330**(1):253-62.
 63. Sherr CJ: **Cancer cell cycles.** *Science* 1996, **274**(5293):1672-7.
 64. Hana V, Murphy LJ: **Expression of insulin-like growth factors and their binding proteins in the estrogen responsive Ishikawa human endometrial cancer cell line.** *Endocrinology* 1994, **135**(6):2511-6.
 65. Yang H, Satta MR, et al.: **Induction of human methionine adenosyltransferase 2A expression by tumor necrosis factor alpha. Role of NF-kappa B and AP-1.** *J Biol Chem* 2003, **278**(51):50887-96.
 66. Cai J, Mao Z, et al.: **Differential expression of methionine adenosyltransferase genes influences the rate of growth of human hepatocellular carcinoma cells.** *Cancer Res* 1998, **58**(7):1444-50.
 67. Ito K, Ikeda S, et al.: **Correlation between the expression of methionine adenosyltransferase and the stages of human colorectal carcinoma.** *Surg Today* 2000, **30**(8):706-10.
 68. Risinger JI, Maxwell GL, et al.: **Promoter hypermethylation as an epigenetic component in Type I and Type II endometrial cancers.** *Ann N Y Acad Sci* 2003, **983**:208-12.
 69. Sasaki M, Kaneuchi M, et al.: **Hypermethylation can selectively silence multiple promoters of steroid receptors in cancers.** *Mol Cell Endocrinol* 2003, **202**(1-2):201-7.
 70. Hannon GJ: **RNA interference.** *Nature* 2002, **418**(6894):244-51.
 71. Gould Rothberg BE, Sundseth SS, et al.: **The characterization of PPAR alpha ligand drug action in an in vivo model by comprehensive differential gene expression profiling.** *Funct Integr Genomics* 2002, **1**(5):294-304.
 72. Vanden Heuvel JP, Kreder D, et al.: **Comprehensive analysis of gene expression in rat and human hepatoma cells exposed to the peroxisome proliferator WY14,643.** *Toxicol Appl Pharmacol* 2003, **188**(3):185-98.
 73. Palmer CN, Hsu MH, et al.: **Peroxisome proliferator activated receptor-alpha expression in human liver.** *Mol Pharmacol* 1998, **53**(1):14-22.
 74. Roberts RA, James NH, et al.: **Apoptosis and proliferation in non-genotoxic carcinogenesis: species differences and role of PPARalpha.** *Toxicol Lett* 2000, **112-113**:49-57.
 75. Peters JM, Aoyama T, et al.: **Role of peroxisome proliferator-activated receptor alpha in altered cell cycle regulation in mouse liver.** *Carcinogenesis* 1998, **19**(11):1989-94.
 76. Murai T, Yamada T, et al.: **Fenofibrate inhibits reactive amyloidosis in mice.** *Arthritis Rheum* 2002, **46**(6):1683-8.
 77. Holden PR, Tugwood JD: **Peroxisome proliferator-activated receptor alpha: role in rodent liver cancer and species differences.** *J Mol Endocrinol* 1999, **22**(1):1-8.