

Cerebrospinal Fluid Research

Open Access

Oral Presentation

Neuropsychological profile of patients with normal pressure hydrocephalus and Alzheimer's disease

JL Iddon, C Loveday, L Gelling, DJR Morgan and JD Pickard*

Address: Department of Academic Neurosurgery, Addenbrooke's Hospital, Cambridge, UK

Email: JD Pickard* - prof.jdp@medschl.cam.ac.uk

* Corresponding author

from 48th Annual Meeting of the Society for Research into Hydrocephalus and Spina Bifida
Dublin, Ireland, 23–26 June 2004
Published: 23 December 2004

Cerebrospinal Fluid Research 2004, 1(Suppl 1):S17 doi:10.1186/1743-8454-1-S1-S17

This article is available from: <http://www.cerebrospinalfluidresearch.com/content/1/S1/S17>

Normal pressure hydrocephalus (NPH) is a potentially treatable neurological disorder if diagnosed accurately and early. If misdiagnosed, or left untreated, the pattern is one of continuing physical and cognitive decline, which becomes increasingly irreversible as more diffuse brain damage is caused. Cognitive profiling is a technique, which can aid early and accurate detection of NPH.

This paper presents the results of two studies showing specific contrasting patterns of neuropsychological dysfunction in subgroups of patients with either early or late stage NPH with patients either in the prodromal stages (mild cognitive impairment) or later stages of Alzheimer's disease. The results demonstrate that there are differential patterns that can be observed in patients with NPH, which can be used effectively within a clinic-context to enhance accurate and early diagnosis and improve long-term prognosis.

References

1. Iddon JL, Pickard JD, et al.: **Specific patterns of cognitive impairment in patients with idiopathic normal pressure hydrocephalus and Alzheimer's disease.** *JNNP* 1999, **67**:723-732.
2. Iddon JL: **Cognitive dysfunction in hydrocephalus.** In *Cognitive Deficits in Brain Disorders* Edited by: Harrison J, Owen AM. Martin Dunitz Publishers; 2001.