



POSTER PRESENTATION

Open Access

A novel 9 kDa phosphoprotein is a component of the primary cilium and interacts with polycystin-1

DPS Osborn^{1*}, C Boucher², P Wilson³, V Gattone⁴, PL Beales¹, I Drummond⁵, R Sandford²*From* First International Cilia in Development and Disease Scientific Conference (2012)
London, UK. 16-18 May 2012

Polycystin-1 (PKD1) forms a mechanosensitive cation channel complex with polycystin-2 (PKD2) in the primary cilium. Functional defects in this complex caused by mutation of PKD1 and PKD2 result in autosomal dominant polycystic kidney disease (ADPKD). The mechanisms by which this complex regulates normal cell physiology remain elusive. In particular the proteins that interact directly with polycystin-1 remain poorly characterised. A Y-2H screen using the C-terminus of polycystin-1 as bait identified a novel binding partner that we have called PIP9 (polycystin interacting protein 9). PIP9 is evolutionary conserved, ubiquitously expressed, and predicted to be a 9KDa phosphoprotein containing a single coiled-coil domain that interacts with the coiled-coil domain of PC-1. Specific anti-PIP9 antisera confirmed the widespread and developmentally regulated expression of PIP9 and identified phosphorylated isoforms. PIP9 co-localised to the renal primary cilium with PC1 and immuno-EM studies revealed localisation to IFT particles. Knockdown of *zPip9*, using morpholino oligonucleotides, resulted in developmental defects in gastrulation movements. In addition, evidence of cell detachment was observed, common to the phenotype described for double *bbs4;bbs10* morphant zebrafish which suggests *pip9* may act in the same genetic pathway. *zpip9* morphants also displayed variable head and eye degeneration, including coloboma and hydrocephalus, pronephric cysts and shortening of the body axis. Loss of *Pip9* function in mice is lethal before embryonic day E11.5. This preliminary analysis suggests that PIP9 plays a central role in mediating polycystin-1 dependant signalling.

Author details¹UCL Institute of Child Health, UK. ²Department of Medical Genetics, University of Cambridge, UK. ³The Royal Free Hospital, London, UK. ⁴Indiana University School of Medicine, USA. ⁵Massachusetts General Hospital, USA.

Published: 16 November 2012

doi:10.1186/2046-2530-1-S1-P74

Cite this article as: Osborn *et al.*: A novel 9 kDa phosphoprotein is a component of the primary cilium and interacts with polycystin-1. *Cilia* 2012 **1**(Suppl 1):P74.**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit* Correspondence: dan.osborn@ucl.ac.uk¹UCL Institute of Child Health, UK

Full list of author information is available at the end of the article