



Stoumpos, S., Jardine, A. G., and Mark, P. B. (2014) Cardiovascular morbidity and mortality after kidney transplantation. *Transplant International* . ISSN 0934-0874

Copyright © 2014 Steunstichting ESOT

A copy can be downloaded for personal non-commercial research or study, without prior permission or charge

Content must not be changed in any way or reproduced in any format or medium without the formal permission of the copyright holder(s)

When referring to this work, full bibliographic details must be given

<http://eprints.gla.ac.uk/96020>

Deposited on: 25 August 2014

Enlighten – Research publications by members of the University of Glasgow_
<http://eprints.gla.ac.uk>

Received Date : 17-Jun-2014
Accepted Date : 28-Jul-2014
Article type : Review Article

Cardiovascular morbidity and mortality after kidney transplantation

Sokratis Stoumpos¹

Alan G Jardine^{1,2}

Patrick B Mark^{*1,2}

1. The Glasgow Renal & Transplant Unit, Western Infirmary, Dumbarton Road, Glasgow, United Kingdom

2. Institute of Cardiovascular and Medical Sciences, BHF Glasgow Cardiovascular Research Centre, University of Glasgow, Glasgow, United Kingdom

***Corresponding author:** Patrick B Mark, Institute of Cardiovascular and Medical Sciences, BHF Glasgow Cardiovascular Research Centre, University of Glasgow, Glasgow, G12 8TA, United Kingdom

Email: patrick.mark@glasgow.ac.uk

Contribution of authors. SS and PBM wrote the manuscript. AGJ critically revised the manuscript and wrote the manuscript.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/tri.12413

This article is protected by copyright. All rights reserved.

Conflict of interest. AGJ has received honoraria and consultancy fees from Novartis, Astellas, Pfizer, Bristol-Myers Squibb, AstraZeneca, and Merck Sharp & Dohme in the specialties of cardiovascular disease and transplantation. The other authors have no conflicts to declare.

Funding sources: No funding

Keywords. Kidney transplant, cardiovascular, lipids, statins, arrhythmia, hypertension, atherosclerosis

Abstract

Kidney transplantation is the optimal treatment for patients with end stage renal disease (ESRD) who would otherwise require dialysis. Patients with ESRD are at dramatically increased cardiovascular (CV) risk compared to the general population. As well as improving quality of life, successful transplantation accords major benefits by reducing cardiovascular risk in these patients. Worldwide, cardiovascular disease remains the leading cause of death with a functioning graft and therefore is a leading cause of graft failure. This review focuses on the mechanisms underpinning excess cardiovascular morbidity and mortality and current evidence for improving cardiovascular risk in kidney transplant recipients. Conventional cardiovascular risk factors such as hypertension, diabetes mellitus, dyslipidaemia, and pre-existing ischaemic heart disease are all highly prevalent in this group. In addition, kidney transplant recipients exhibit a number of risk factors associated with pre-existing renal disease. Furthermore, complications specific to transplantation may ensue including

Accepted Article

reduced graft function, side effects of immunosuppression and post transplantation diabetes mellitus. Strategies to improve cardiovascular outcomes post transplantation may include pharmacological intervention including lipid lowering or antihypertensive therapy, optimisation of graft function, lifestyle intervention and personalising immunosuppression to the individual patients risk profile.

Background

Transplantation confers the highest survival benefit among all the different renal replacement therapies. Multiple studies have shown that patient survival is better with renal transplantation than with maintenance dialysis after an increased risk of death in the early period after transplantation[1-5]. This is true for patient groups who are otherwise at increased cardiovascular (CV) risk including diabetics, African Americans, all age groups[3], obese patients[6], as well as recipients of marginal kidneys[7-9] and following repeat renal transplantation after failed primary transplantation[10, 11]. Although long-term allograft survival has improved, death with a functioning graft remains the leading cause of late renal allograft loss[12, 13]. Cardiovascular disease (CVD) persists as the leading cause of premature death in most kidney transplant registries[14].

Knowledge of the incidence, risk factors and the natural history of CVD in renal transplantation derives from registry data, observational population-based studies, clinical trials and extrapolation from studies on non-transplant cohorts. This review focuses on describing the nature of CVD in renal transplant recipients (RTR), including dissecting the various components,

which combine as the syndrome of CVD in RTR. We highlight evidence based treatments for reducing cardiovascular risk where this exists.

Epidemiology of cardiovascular disease in kidney transplantation

CVD mortality in haemodialysis (HD) patients is 10 to 20 times greater than in the general population[15]. Renal transplant recipients have lower risk for CVD than patients who remain on the transplant waiting list, but higher CVD risk when compared with the general population[16], particularly those aged 25–55 years who have substantially more CVD mortality than their age-, gender-, and race-matched non-dialysis counterparts. Registry data show that cardiac disease is the cause of death for 18-30% of prevalent transplant patients[17, 18]. Recent UK Registry data of 566 deaths (3.0%) within the first year post transplant (from 19,103 kidney transplants performed over an 11-year period) demonstrates that whilst infection was the single leading cause of death (21.6% of deaths) in the first year post transplant, cardiovascular events combined with cerebrovascular disease accounted for the greatest proportion of deaths at 22.9%[19]. With longer follow up CVD continues to accumulate and accounts for 31% of deaths with a functioning graft in the 2013 United States Renal Data System (USRDS) Report [17].

Clinical aspects of cardiovascular disease in kidney transplantation

In the general population CVD predominantly relates to underlying coronary artery atherosclerosis and is associated with conventional cardiovascular risk factors such as hypertension, dyslipidaemia, diabetes, cigarette smoking and family history. This paradigm does not hold true for patients with ESRD where

sudden, presumed arrhythmic, cardiac death rather than myocardial infarction (MI) is the predominant mode of cardiovascular mortality. The paradox of reverse epidemiology is acknowledged in ESRD patients on dialysis where J-shaped (rather than linear) relationships are seen between blood pressure[20], cholesterol[21] and body mass index[22] and mortality risk. Following successful transplantation patients with ESRD have more conventional relationships between cardiovascular risk factors and outcome, as illustrated recently by *post hoc* analysis of the Folic Acid for Vascular Outcome Reduction in Transplantation [23] [23] trial[24], where there was a linear relationship between increasing systolic blood pressure and mortality. Although, more conventional relationships between CVD and risk factors evolve post transplantation, legacy of time spent on dialysis remains. Along with diabetes and age, evidence of left ventricular hypertrophy with 'strain' on the ECG, often associated with longstanding ESRD was associated with increased risk of cardiac death in the Assessment of Lescol in Renal Transplantation (ALERT) trial[25].

Whilst the relationship between risk factors and CV events reverts towards the general population, the outcomes following a CV event in RTR are not comparable with the general population. In the ALERT study[26], there was a similar rate of fatal and non-fatal CV events to other randomised controlled trials of lipid lowering therapy(WOSCOPS[27] and 4S[28]). In non-transplant populations non-fatal CV events are more common than fatal events. Therefore in RTR, risk of CVD is increased and there is a high prevalence of CV risk factors. There are dichotomous patterns of CVD in RTR, including

both atheromatous coronary artery disease (CAD) as seen in the general population and sudden cardiac death as observed in dialysis patients. When they occur, CV events are more likely to be fatal than in the general population.

Coronary artery disease

Coronary artery disease influences listing for transplantation. At the time of transplantation, prior MI suggesting occlusive CAD is reported in 2.6% of transplant recipients in the UK [19]. Estimating prevalence of non-occlusive coronary artery atherosclerosis prior to transplantation is more difficult as most reports favour only performing coronary angiography in higher risk transplant candidates (e.g. over aged 50, diabetes mellitus, prior MI). Nonetheless, it appears that coronary artery atheroma is present in approximately 50% of higher risk transplant candidates[29]. Performing unselected coronary angiography appears to lead to low rates of coronary intervention in renal transplant candidates and is unlikely to be a useful strategy for risk reduction pre-transplantation[30, 31].

Once transplanted, prevalent CVD accumulates and was reported in 20% (14% previous MI or CAD) of participants at study entry in the FAVORIT trial[32] compared to 11.5% (4.7% previous MI, 6.8% revascularisation procedure) of participants in the observational PORT study[33]. US data[17] show that hospitalisations for coronary atherosclerosis increases from 5.5% in year one to 9% in year two. Lentine *et al*[34] concluded that post transplantation MI is common, affecting approximately 11.1% of patients by 3 years post transplantation, and that much of this risk is experienced early,

within the first 6 months of transplantation. MI risk was linked with modifiable factors, including delayed graft function, post transplantation diabetes, and graft failure, and in turn, occurrence of MI predicted graft failure and death. Additionally, *post hoc* analyses of the ALERT trial[25] demonstrated that, determinants of non-fatal myocardial infarction in RTR include total cholesterol level, prior CAD and previous acute rejection. Combined, these data suggest that whilst RTR share common risk factors with the general population for CAD and MI post transplantation; there are further graft-specific aspects to post transplant CAD.

Congestive heart failure

Congestive heart failure (CHF) and renal dysfunction form a “vicious circle” that augur poor prognosis. At commencement of dialysis, up to 70% of patients with ESRD may have abnormal cardiac structure or function[35]. Transplantation may decrease the risk for CHF specifically compared with dialysis therapy[36]. However, CHF remains a clinical concern after transplantation. Wali *et al*[37] showed an improvement of left ventricular (LV) systolic function in more than 86% of patients following kidney transplantation which was associated with an improvement in NYHA functional status in more than two-thirds of patients. Duration of dialysis therapy before transplantation was the only factor that predicted normalisation of LV systolic dysfunction. Lentine *et al*[38] examined incidence of *de novo* CHF in 27,011 transplant recipients. Cumulative incidences of CHF were 10.2% and 18.3% at 1 and 3 years post transplantation and beyond the early post transplantation period, incidence of new onset CHF decreased progressively to less than the

Accepted Article

incidence in transplant candidates (18.3% versus 32.3% at 3 years). US data[17] show that cardiovascular hospitalisations due to congestive heart failure rise from 21% in the first post transplant year to 25% in the second year.

Arrhythmia

Despite the high incidence of sudden cardiac death, surprisingly little is known about arrhythmias in RTR. This reflects the difficulty in capturing short-lived arrhythmic episodes in asymptomatic patients. Using 24-hour ECG monitoring, RTR have been shown to have higher rates of ventricular arrhythmia, usually ventricular extra systoles, compared to patients with mild chronic kidney disease (CKD)[39]. Ventricular repolarisation is also abnormal, usually due to underlying left ventricular hypertrophy (LVH). Whilst ventricular arrhythmia when sustained is life threatening, atrial arrhythmia, in particular atrial fibrillation and flutter are relatively common (6.4% of US transplant recipients[40]), and confers increased risk of ischaemic stroke. Patients with atrial fibrillation are likely to be older and have greater co-morbidity burden[40]. There are no specific guidelines or studies to inform therapeutic strategies in RTR with atrial fibrillation.

Risk factors for development of CVD in kidney transplantation

The burden of CVD in RTR is not entirely explained by traditional risk factors such as hypertension, dyslipidaemia, and diabetes[16]. Other factors may be involved, particularly those that influence systemic inflammation including graft rejection, infection, and use of immunosuppressive medications[13, 41].

Standard CVD risk calculators for the general population are poorly predictive in RTR. Soveri *et al* developed a formula for 7-year CVD and mortality risk calculation for prevalent RTR[42] using models with different variables including age, CAD, diabetes, low-density lipoprotein (LDL), creatinine, number of transplants, time on renal replacement therapy, and smoking (<http://www.anst.uu.se/insov254/calculator/>). Carpenter *et al*[32] demonstrated that traditional CVD risk factors are inadequately managed in RTR. Using baseline data from the FAVORIT study, they showed that almost a third of the patients did not meet blood pressure (BP) target of <130/80mmHg, one to five had borderline or elevated LDL cholesterol and a third of the participants with prevalent CVD were not using an antiplatelet agent for secondary prevention.

Hypertension

Hypertension is common after kidney transplantation and is present in 50% to 90% of RTR [43, 44]. Multiple factors induce susceptibility to high blood pressure (BP) after transplantation including recipient, donor and transplant factors, immunotherapy, transplant dysfunction, renal artery stenosis and obstruction[45]. Hypertension is a leading cause for both decline in graft function and development of CVD[13, 16, 41, 46]. The influence of blood pressure on long-term kidney graft outcomes was demonstrated in the Collaborative Transplant Study (CTS)[47]. Increased blood pressure at different time intervals post transplantation was associated with late graft failure. Subsequent studies[43, 48] showed that this strong, graded relationship between post transplant BP and renal allograft failure was

independent of acute rejection and baseline renal function, thus suggesting that progressive renal dysfunction was the result of elevated BP.

BP control in RTR is particularly challenging. Kidney Disease: Improving Global Outcomes (KDIGO) guidelines[49] recommend a BP target of $\leq 130/80$ mmHg irrespectively of level of proteinuria. This is based on data from patient subgroups in the general population rather than data in RTR. Retrospective data[50] showed that systolic BP of ≤ 140 mm Hg at 3 years after transplantation is associated with improved graft survival and reduced cardiovascular mortality at 10 years and this effect remained even after lowering systolic BP several years post transplantation.

Vasoconstriction is the dominant mechanism by which calcineurin inhibitors (CNI) induce acute nephrotoxicity and hypertension, thus, dihydropyridine calcium-channel blockers (CCB) are an attractive option at least for the early management of hypertension after transplant. The beneficial effect of CCB on kidney function, compared with either placebo or angiotensin converting enzyme inhibitors (ACEi), was shown in short-term randomised controlled trials (RCT)[51-54] although effects of CCB on long-term kidney function in CNI-treated RTR have been reported with variable efficacy[55-57]. A recent meta-analysis of randomised controlled trials[58] indicated that use of CCB, versus placebo or no treatment (plus additional agents in either arm, as needed) was associated with 25% lower rate of graft loss and higher glomerular filtration rate (GFR) and in direct comparison with ACEi, CCB significantly improved GFR by approximately 12ml/min.

ACEi may reverse post transplant erythrocytosis, decrease proteinuria and have a theoretical effect in mitigating antibody-mediated rejection mediated by antibody to AT1 receptor. In a RCT of renal transplant recipients with LVH[59], patients administered ACEi had significantly better general and cardiovascular outcome after 10-year follow-up, suggesting that the effect by renin-angiotensin-system (RAS) blockade on clinical outcome can only be observed with longer follow-up. Another RCT that was adequately powered to assess hard outcomes in comparison between RAS blockade and placebo in RTR[60] was prematurely discontinued after 2 years because the incidence of events was considerably lower than expected in both arms of the study.

Finally, two recent meta-analyses[61, 62] pointed out advantages of adopting calcium channel blockers for blood pressure control in RTR because RAS blockers are associated with progressive worsening of renal graft function without benefit in CV risk. Patients on ACEi or angiotensin receptor blockers (ARB) had a decrease in glomerular filtration rate (5.8ml/min), lower haematocrit (3.5% translating to haemoglobin lowering of approximately 1.2 g/dl) and reduction in proteinuria[61]. The rate of cardiovascular death was similar in patients who received ACEi/ARB therapy or other antihypertensive treatment overall and in subpopulations of patients known to be at high cardiovascular risk[62].

Dyslipidaemia

Almost half of the RTR have low density lipoprotein (LDL) cholesterol levels >2.6mmol/L and 41% are on statin treatment six months post transplantation[63]. Dyslipidaemia is common after transplantation, partly due

Accepted Article

to the hyperlipidaemic effect of corticosteroids, cyclosporine, tacrolimus, and mammalian target of rapamycin (mTOR) inhibitors. In the CONVERT trial[64], conversion from cyclosporine or tacrolimus to rapamycin was associated with higher prevalence of hypertriglyceridemia (54 versus 26%) and hypercholesterolemia (42 versus 12%) by month 24, even in the context of more common use of lipid-lowering therapy (78 versus 55%). On the contrary, converting from cyclosporine to tacrolimus may provide significant benefits in serum lipid levels[65].

The Assessment of Lescol in Renal Transplantation (ALERT) trial[26] was the first large study to address cardiac and renal outcomes in transplantation. Treatment with fluvastatin (40-80mg) failed to reach statistical significance in the primary composite end points (major adverse cardiac events defined as cardiac death, non-fatal MI, and coronary intervention) despite 32% lowering of LDL cholesterol during a mean follow-up of 5.1 years. On the “hard” cardiovascular endpoints, treatment with fluvastatin demonstrated a reduction of 38, 32 and 35% in the risk of cardiac death, non-fatal MI and in the cumulative incidence of cardiac death or first non-fatal MI respectively. Nonetheless, a 2-year extension to the original study[66] demonstrated significant long-term benefits in the primary composite outcome. *Post hoc* analyses of the ALERT study demonstrated that early initiation of lipid-lowering therapy had a more favourable effect on cardiac events than late intervention[67] and lowering of LDL cholesterol by 39mg/dL reduced cardiac death or myocardial infarction by approximately 30%[68].

Long-term treatment with fluvastatin was well tolerated and had no harmful effects on renal function. This was opposed with previous reports[69] highlighting increased risk of myopathy and rhabdomyolysis, especially when co-administered with cyclosporine, which often results in several-fold increases in statin blood level.

Additionally, a Cochrane meta-analysis of 22 studies[70] including 3,465 RTR confirmed the CV benefits of statins. The 2013 KDIGO guidelines[71] suggest initial evaluation of all transplant patients with a lipid profile, without follow-up lipid levels for the majority of patients since the indication for treatment is the higher cardiovascular risk, rather than LDL concentration.

Post transplantation diabetes mellitus

Post transplantation diabetes mellitus (PTDM) is increasing in incidence and is a major challenge following solid organ transplantation[72, 73]. Approximately one-third of non-diabetic kidney transplant recipients develop persistently impaired glucose metabolism by six months post transplantation[74]. At 3 years, the cumulative incidence of post transplant diabetes amongst RTR is between 24.0% and 42.0%[17, 75]. Risk factors for PTDM include age, obesity, African American race and Hispanic ethnicity, family history and impaired glucose tolerance[76]. In addition, risk factors that are unique to transplantation include immunosuppressive agents, HLA mismatch, donor gender, type of underlying renal disease and viral infections (HCV and CMV)[76, 77]. Both PTDM and impaired glucose tolerance [65] confer a higher risk of developing CVD. The increased relative risk for death from CVD ranges from 1.5 to 3 among those who develop PTDM versus

those without diabetes[75, 78]. In a cohort study of 37,448 RTR[79], pre-existing diabetes was associated with higher cardiovascular and overall mortality compared with post transplantation diabetes mellitus, at least shortly after transplantation.

Renal impairment

Reduced kidney function is a risk factor for CVD in the general population, in part reflecting the close association of CVD risk factors and GFR. Population data[80] suggest that even minor kidney dysfunction is associated with increased cardiovascular risk. The relationship of GFR with CVD risk in transplant recipients may differ following transplantation, because the level of GFR may no longer reflect lifelong exposure to CVD risk factors[13]. A *post hoc* analysis of 1,052 participants in the ALERT Study[25], showed that renal dysfunction was associated with fatal CVD. Mild renal insufficiency was independently associated with increased risk of acute coronary syndromes and CHF[81], and 15% higher risk of CVD and death for each 5ml/min/1.73m² lower eGFR (at levels below 45ml/min/1.73m²)[82].

Left ventricular hypertrophy

LVH is present in 50-70% of patients following renal transplantation[16] and is a significant risk factor for CHF and death in RTR[83]. Correction of the uraemic state by transplantation leads to a fall in LV mass with echocardiographic examination[84, 85] though when LV mass was measured by the more accurate cardiac magnetic resonance[86], renal transplantation

was not associated with significant regression of LV mass suggesting that improvement in fluid balance leads to apparent improvement in LV mass.

LVH is primarily an adaptive response to volume and pressure overload with the aim of minimising ventricular wall stress. Multiple risk factors contribute to LVH development including age, hypertension, hypercholesterolemia, tobacco smoking, obesity, or diabetes, as well as transplant-specific risk factors including anaemia, the arteriovenous fistula flow, and immunosuppressive therapy[87]. Resistance to left ventricular outflow produced by aortic valve calcification during dialysis[88], anaemia and high blood pressure[89] appear to be leading contributors to development, progression and persistence of LVH in RTR. ACEi are effective in reversing LVH persisting despite successful renal transplantation, probably by reducing BP[90] but also through mechanisms that are at least partially independent of hemodynamic effects on BP[91]. In one study[91], ACEi were effective in regressing post transplantation LVH only in patients on cyclosporine therapy, perhaps because of an interaction effect between the two treatments. This indicates that immunosuppressive agents might modulate the effect of antihypertensive therapy on the left ventricular mass of RTR.

The mTOR inhibitors play a role in regulating cell growth and may be a therapeutic tool to regress established cardiac hypertrophy. In two small studies, both sirolimus[92] and everolimus[93] regressed LVH in RTR regardless of BP changes, mainly by decreasing left ventricular wall thickness, suggesting non-hemodynamic effect mechanisms of mTOR inhibitors on left ventricular mass.

Effects of immunosuppression on cardiovascular risk

Corticosteroids

Corticosteroids have been a cornerstone of transplant immunosuppression for over 50 years, both as maintenance immunosuppression and for treatment of acute rejection. However, adverse effects of corticosteroids, mainly cardiovascular, have led to attempts to find maintenance immunosuppression regimens that do not include corticosteroids. Different protocols have been developed including 'steroid-free' protocols which do not use steroids as initial or maintenance immunosuppression, 'steroid avoidance' protocols in which steroids are initially used and are then withdrawn during the first week after transplantation, and 'steroid withdrawal' protocols in which steroids are discontinued weeks to months after transplantation.

Cardiovascular risk in RTR varies with comorbidities such as pre-existing metabolic syndrome, race and age and in addition, many of the adverse effects attributed to corticosteroids were observed with high doses. Whether the low doses commonly used for maintenance immunosuppression are associated with major adverse effects is less clear and is difficult to dissociate the CV profile of steroids from other factors, such as underlying renal function and CNI use.

Calcineurin inhibitors

CNI-sparing maintenance immunosuppression regimens have been applied to help maintain the balance between allograft survival and nephrotoxicity. CNI raise arterial blood pressure in transplant recipients by several mechanisms, including arteriolar vasoconstriction, activation of the renin-angiotensin

system, direct effects on juxtaglomerular cells, and increased tubular sodium reabsorption[94, 95]. The beneficial effects of late CNI withdrawal on ambulatory blood pressure were documented in a recently published RCT of 119 stable RTR on a triple-drug regimen[96].

Complete CNI withdrawal after the initial period of high immunological risk is attractive, however, this has been associated with an increased incidence of late acute rejection and a possible reduction in long-term allograft survival[97]. Trials that explored the switching from CNI to mTOR inhibitors 3 to 6 months post transplant[98-100] have shown the relative safety of this approach with improvement in renal function and blood pressure despite increased risk of acute rejection.

Lifestyle and other cardiovascular risk factors in kidney transplant recipients

Cigarette smoking increases the risk for graft failure[101, 102], ischaemic heart disease[102], and CHF[38] in renal transplant recipients. Prevalence of cigarette smoking at time of transplantation varies between 25% and 50%[101, 102]. Between smokers RTR, graft failure is largely due to death with a functioning graft[102] and having quit smoking more than 5 years before transplantation reduced the relative risk of graft failure by 34%[102]. In the general population, there is strong evidence that screening patients for tobacco use and implementing prevention and treatment measures are effective. Guidelines suggest that the same approach should be applied for the RTR[103, 104].

Obesity among transplant recipients is associated with the metabolic syndrome, which is present in two-thirds of RTR at six years post transplant[105]. Registry data show that obesity is associated with adverse cardiovascular outcomes including increased risk of cardiac death[106], CHF[81] and atrial fibrillation[107]. Lifestyle changes based in diet and exercise, with dietary counselling as needed, are first-line strategies to achieve normal body weight among obese RTR. There is a paucity of data on the safety and efficacy of post transplantation gastric banding or bypass surgery in ameliorating comorbid conditions such as hypertension, diabetes mellitus, and dyslipidaemia.

Although low physical activity is strongly associated with increased risk for cardiovascular and all-cause mortality in RTR[108], the effect of exercise training in the cardiovascular risk profiles of RTR are unclear. A recent meta-analysis of exercise training in solid organ transplant recipients[109] (including two RCT with 164 RTR) showed no significant improvements in exercise capacity or cardiovascular risk factors such as incidence of PTDM, indicating that exercise training is a promising but unproven intervention for improving the CV outcomes in RTR.

Homocysteine is implicated to be an atherogenic amino acid and fasting hyperhomocysteinaemia has been shown to be an independent predictor of cardiovascular events among RTR[110]. However, in the Folic Acid for Vascular Outcome Reduction in Transplantation [23] trial[111], lowering homocysteine levels has not been shown to decrease cardiovascular risk among 4,110 RTR treated with vitamin B6 and vitamin B12 and with either

high or low dose folic acid, despite the fact that homocysteine was effectively lowered with high dose folic acid.

Multiple other non-traditional risk factors have been associated with increased cardiovascular risk in various studies including anaemia, dialysis vintage prior to transplantation, elevated levels of lipoprotein a, elevated C-reactive protein and interleukin-6 levels[46, 112-114].

Conclusions

Renal transplantation is the single most effective intervention for reducing cardiovascular risk in appropriately selected patients with ESRD. Nonetheless, CVD is common and is the leading cause of death with a functioning graft and hence graft loss. Strategies targeting modifiable conventional cardiovascular risk factors (diabetes, hypertension, dyslipidaemia and lifestyle) are crucial to reducing post transplant CVD. However further strategies to address transplant specific cardiovascular risk factors should also be employed. These should include optimisation of renal function, limiting risk of rejection, avoidance of PTDM and anticipation of cardiovascular side effects of immunosuppression. Further studies are required to address how each of these strategies is tailored to the requirements of the individual patient and graft.

Table 1

Modifiable risk factors for post transplant CVD	Pre transplant hazard	Post transplant exacerbation	Potential strategies to address	Rationale	Studies
Hypertension	Recipient, donor and transplant factors	Immunotherapy, transplant dysfunction, renal artery stenosis	Target BP \leq 130/80, avoid ACEi/ARB within the first 3 months post transplantation, consider in the long-term especially if persistent albuminuria, LVH or other indications	No evidence for benefit of any particular antihypertensive agent	Small RCT[51-54, 57, 59] Observational data[43, 46-48, 50]
Post transplantation diabetes mellitus	African American or American Hispanic ethnicity, obesity	Immunotherapy, HLA mismatch, viral infections	Target HbA1c 7.0-7.5%, early basal insulin, incretin-based therapy, steroid reduction or withdrawal, switch from tacrolimus to CsA	Chances of reversing or ameliorating PTDM may be improved by early detection and intervention	Small RCT[115, 116] Observational data[75, 78, 79]
Left ventricular hypertrophy	Hypertension, aortic valve calcification	Immunotherapy	ACEi, CsA, mTOR inhibitors	Regression of LVH with BP decrease, other	Small RCT[90, 91, 93]

	on, anaemia, dialysis, AVF flow			mechanisms independent of haemodynamic effects on BP	
Dyslipidaemia	Pre transplant lipid abnormalities	Hyperlipemic effect of immunotherapy	Statins, replace CsA by tacrolimus, avoid sirolimus, low dose prednisolone	Statins improve CV outcomes, no benefit in overall mortality, remarkably safe	ALERT[26, 66] Post hoc analyses[67, 68]
Renal impairment	Previous graft loss, transplant factors	CNI nephrotoxicity, AR episodes, proteinuria	Optimisation of graft function, CNI minimisation to achieve adequate IS	Mild renal insufficiency is associated with adverse CV outcomes	Post hoc analyses[25, 82] Observational data[81]
Calcineurin inhibitors	Transplant factors	Nephrotoxic, PTDM, Hypertension, Dyslipidaemia	CNI dose reduction after the first 3 months, conversion from CNI to mTOR inhibitors 3 to 6 months post transplant is relatively safe	CNI withdrawal leads to increased acute rejection	RCT[96, 100]
Corticosteroids	Diabetes, race	PTDM, Hypertension, Dyslipidaemia	Low dose steroids in the long-term, discontinuation could be attempted in patients at	No clear benefit for improved patient or graft outcomes after early or late discontinuation of	Short- and medium-term RCT[117-121]

			increased CV risk with low immunological risk and stable allograft function	steroids	
Other CV risk factors	Smoking, obesity, low physical activity, anaemia	Immunotherapy, systemic inflammation	Quit smoking, diet, correction of anaemia	Unclear benefits of exercise and vitamins	RCT[111] Observational data[102, 106, 108] Extrapolation from studies in the general population[103]

Figure 1. Pre and post transplant factors conferring increased cardiovascular risk after kidney transplantation.

BMI, body mass index; CNI, calcineurin inhibitors; CHF, congestive heart failure; CAD, coronary artery disease; CyA, cyclosporine; CV, cardiovascular; eGFR, estimated glomerular filtration rate; LVH, left ventricular hypertrophy; PTDM, post transplantation diabetes mellitus;

Table 1. Modifiable risk factors for post transplant cardiovascular disease and strategies to address.

AR, acute rejection; ACEi, angiotensin converting enzyme inhibitors; ARB, angiotensin receptor blockers; AVF, arteriovenous fistula, CVD, cardiovascular disease; BP, blood pressure; CsA, cyclosporine; IS, immunosuppression; HLA, human leukocyte antigen; mTOR, mammalian target of rapamycin; LVH, left ventricular hypertrophy; RTR, renal transplant

recipients; RCT, randomised controlled trials; PTDM, post transplantation diabetes mellitus

1. Port FK, Wolfe RA, Mauger EA, Berling DP, Jiang K. Comparison of survival probabilities for dialysis patients vs cadaveric renal transplant recipients. *JAMA*. 1993; **270**: 1339-43.
2. Ojo AO, Port FK, Wolfe RA, Mauger EA, Williams L, Berling DP. Comparative mortality risks of chronic dialysis and cadaveric transplantation in black end-stage renal disease patients. *Am J Kidney Dis*. 1994; **24**: 59-64.
3. Wolfe RA, Ashby VB, Milford EL, et al. Comparison of mortality in all patients on dialysis, patients on dialysis awaiting transplantation, and recipients of a first cadaveric transplant. *N Engl J Med*. 1999; **341**: 1725-30.
4. Rabbat CG, Thorpe KE, Russell JD, Churchill DN. Comparison of mortality risk for dialysis patients and cadaveric first renal transplant recipients in Ontario, Canada. *J Am Soc Nephrol*. 2000; **11**: 917-22.
5. Oniscu GC, Brown H, Forsythe JL. Impact of cadaveric renal transplantation on survival in patients listed for transplantation. *J Am Soc Nephrol*. 2005; **16**: 1859-65.
6. Glanton CW, Kao TC, Cruess D, Agodoa LY, Abbott KC. Impact of renal transplantation on survival in end-stage renal disease patients with elevated body mass index. *Kidney Int*. 2003; **63**: 647-53.
7. Ojo AO, Hanson JA, Meier-Kriesche H, et al. Survival in recipients of marginal cadaveric donor kidneys compared with other recipients and wait-listed transplant candidates. *J Am Soc Nephrol*. 2001; **12**: 589-97.
8. Merion RM, Ashby VB, Wolfe RA, et al. Deceased-donor characteristics and the survival benefit of kidney transplantation. *JAMA*. 2005; **294**: 2726-33.
9. Snoeijs MG, Schaubel DE, Hené R, et al. Kidneys from donors after cardiac death provide survival benefit. *J Am Soc Nephrol*. 2010; **21**: 1015-21.
10. Ojo A, Wolfe RA, Agodoa LY, et al. Prognosis after primary renal transplant failure and the beneficial effects of repeat transplantation: multivariate analyses from the United States Renal Data System. *Transplantation*. 1998; **66**: 1651-9.
11. Rao PS, Schaubel DE, Wei G, Fenton SS. Evaluating the survival benefit of kidney retransplantation. *Transplantation*. 2006; **82**: 669-74.
12. Pascual M, Theruvath T, Kawai T, Tolckoff-Rubin N, Cosimi AB. Strategies to improve long-term outcomes after renal transplantation. *N Engl J Med*. 2002; **346**: 580-90.
13. Kasiske BL, Chakkera HA, Roel J. Explained and unexplained ischemic heart disease risk after renal transplantation. *J Am Soc Nephrol*. 2000; **11**: 1735-43.
14. Wheeler DC, Steiger J. Evolution and etiology of cardiovascular diseases in renal transplant recipients. *Transplantation*. 2000; **70**: SS41-5.
15. Levey AS, Beto JA, Coronado BE, et al. Controlling the epidemic of cardiovascular disease in chronic renal disease: what do we know? What

- do we need to learn? Where do we go from here? National Kidney Foundation Task Force on Cardiovascular Disease. *Am J Kidney Dis.* 1998; **32**: 853-906.
16. Sarnak MJ, Levey AS, Schoolwerth AC, et al. Kidney disease as a risk factor for development of cardiovascular disease: a statement from the American Heart Association Councils on Kidney in Cardiovascular Disease, High Blood Pressure Research, Clinical Cardiology, and Epidemiology and Prevention. *Circulation.* 2003; **108**: 2154-69.
 17. Collins AJ, Foley RN, Chavers B, et al. US Renal Data System 2013 Annual Data Report. *Am J Kidney Dis.* 2014; **63**: A7.
 18. Pruthi R, Steenkamp R, Feest T. UK Renal Registry 16th Annual Report: Chapter 8 Survival and Cause of Death of UK Adult Patients on Renal Replacement Therapy in 2012: National and Centre-specific Analyses. *Nephron Clinical Practice.* 2013; **125**: 139-69.
 19. Farrugia D, Cheshire J, Begaj I, Khosla S, Ray D, Sharif A. Death within the first year after kidney transplantation - an observational cohort study. *Transpl Int.* 2013.
 20. Zager PG, Nikolic J, Brown RH, et al. "U" curve association of blood pressure and mortality in hemodialysis patients. Medical Directors of Dialysis Clinic, Inc. *Kidney Int.* 1998; **54**: 561-9.
 21. Liu Y, Coresh J, Eustace JA, et al. Association between cholesterol level and mortality in dialysis patients: role of inflammation and malnutrition. *JAMA.* 2004; **291**: 451-9.
 22. Fleischmann E, Teal N, Dudley J, May W, Bower JD, Salahudeen AK. Influence of excess weight on mortality and hospital stay in 1346 hemodialysis patients. *Kidney Int.* 1999; **55**: 1560-7.
 23. Lobo ML, Favorito LA, Abidu-Figueiredo M, Sampaio FJ. Renal pelvic diameters in human fetuses: anatomical reference for diagnosis of fetal hydronephrosis. *Urology.* 2011; **77**: 452-7.
 24. Carpenter MA, John A, Weir MR, et al. BP, Cardiovascular Disease, and Death in the Folic Acid for Vascular Outcome Reduction in Transplantation Trial. *J Am Soc Nephrol.* 2014.
 25. Jardine AG, Fellström B, Logan JO, et al. Cardiovascular risk and renal transplantation: post hoc analyses of the Assessment of Lescol in Renal Transplantation (ALERT) Study. *Am J Kidney Dis.* 2005; **46**: 529-36.
 26. Holdaas H, Fellström B, Jardine AG, et al. Effect of fluvastatin on cardiac outcomes in renal transplant recipients: a multicentre, randomised, placebo-controlled trial. *Lancet.* 2003; **361**: 2024-31.
 27. Shepherd J, Cobbe SM, Ford I, et al. Prevention of coronary heart disease with pravastatin in men with hypercholesterolemia. West of Scotland Coronary Prevention Study Group. *N Engl J Med.* 1995; **333**: 1301-7.
 28. Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet.* 1994; **344**: 1383-9.
 29. Kumar N, Baker CS, Chan K, et al. Cardiac survival after pre-emptive coronary angiography in transplant patients and those awaiting transplantation. *Clin J Am Soc Nephrol.* 2011; **6**: 1912-9.

30. Patel RK, Mark PB, Johnston N, et al. Prognostic value of cardiovascular screening in potential renal transplant recipients: a single-center prospective observational study. *Am J Transplant.* 2008; **8**: 1673-83.
31. De Lima JJ, Gowdak LH, de Paula FJ, et al. Treatment of coronary artery disease in hemodialysis patients evaluated for transplant-a registry study. *Transplantation.* 2010; **89**: 845-50.
32. Carpenter MA, Weir MR, Adey DB, House AA, Bostom AG, Kusek JW. Inadequacy of cardiovascular risk factor management in chronic kidney transplantation - evidence from the FAVORIT study. *Clin Transplant.* 2012; **26**: E438-46.
33. Pilmore HL, Skeans MA, Snyder JJ, Israni AK, Kasiske BL. Cardiovascular disease medications after renal transplantation: results from the Patient Outcomes in Renal Transplantation study. *Transplantation.* 2011; **91**: 542-51.
34. Lentine KL, Brennan DC, Schnitzler MA. Incidence and predictors of myocardial infarction after kidney transplantation. *J Am Soc Nephrol.* 2005; **16**: 496-506.
35. Collins AJ. Cardiovascular mortality in end-stage renal disease. *Am J Med Sci.* 2003; **325**: 163-7.
36. Meier-Kriesche HU, Schold JD, Srinivas TR, Reed A, Kaplan B. Kidney transplantation halts cardiovascular disease progression in patients with end-stage renal disease. *Am J Transplant.* 2004; **4**: 1662-8.
37. Wali RK, Wang GS, Gottlieb SS, et al. Effect of kidney transplantation on left ventricular systolic dysfunction and congestive heart failure in patients with end-stage renal disease. *J Am Coll Cardiol.* 2005; **45**: 1051-60.
38. Lentine KL, Schnitzler MA, Abbott KC, et al. De novo congestive heart failure after kidney transplantation: a common condition with poor prognostic implications. *Am J Kidney Dis.* 2005; **46**: 720-33.
39. Stewart GA, Gansevoort RT, Mark PB, et al. Electrocardiographic abnormalities and uremic cardiomyopathy. *Kidney Int.* 2005; **67**: 217-26.
40. Lenihan CR, Montez-Rath ME, Scandling JD, Turakhia MP, Winkelmayr WC. Outcomes after kidney transplantation of patients previously diagnosed with atrial fibrillation. *Am J Transplant.* 2013; **13**: 1566-75.
41. Ojo AO. Cardiovascular complications after renal transplantation and their prevention. *Transplantation.* 2006; **82**: 603-11.
42. Soveri I, Holme I, Holdaas H, Budde K, Jardine AG, Fellström B. A cardiovascular risk calculator for renal transplant recipients. *Transplantation.* 2012; **94**: 57-62.
43. Kasiske BL, Anjum S, Shah R, et al. Hypertension after kidney transplantation. *Am J Kidney Dis.* 2004; **43**: 1071-81.
44. Premasathian NC, Muehrer R, Brazy PC, Pirsch JD, Becker BN. Blood pressure control in kidney transplantation: therapeutic implications. *J Hum Hypertens.* 2004; **18**: 871-7.
45. Mangray M, Vella JP. Hypertension after kidney transplant. *Am J Kidney Dis.* 2011; **57**: 331-41.
46. Ducloux D, Kazory A, Chalopin JM. Predicting coronary heart disease in renal transplant recipients: a prospective study. *Kidney Int.* 2004; **66**: 441-7.

47. Opelz G, Wujciak T, Ritz E. Association of chronic kidney graft failure with recipient blood pressure. Collaborative Transplant Study. *Kidney Int.* 1998; **53**: 217-22.
48. Mange KC, Cizman B, Joffe M, Feldman HI. Arterial hypertension and renal allograft survival. *JAMA.* 2000; **283**: 633-8.
49. Group KBPW. KDIGO clinical practice guideline for the management of blood pressure in chronic kidney disease. *Kidney Int.* 2012; **Supplement 2**: 337-414.
50. Opelz G, Döhler B, Study CT. Improved long-term outcomes after renal transplantation associated with blood pressure control. *Am J Transplant.* 2005; **5**: 2725-31.
51. Midtvedt K, Hartmann A, Foss A, et al. Sustained improvement of renal graft function for two years in hypertensive renal transplant recipients treated with nifedipine as compared to lisinopril. *Transplantation.* 2001; **72**: 1787-92.
52. van Riemsdijk IC, Mulder PG, de Fijter JW, et al. Addition of isradipine (Lomir) results in a better renal function after kidney transplantation: a double-blind, randomized, placebo-controlled, multi-center study. *Transplantation.* 2000; **70**: 122-6.
53. Kuypers DR, Neumayer HH, Fritsche L, et al. Calcium channel blockade and preservation of renal graft function in cyclosporine-treated recipients: a prospective randomized placebo-controlled 2-year study. *Transplantation.* 2004; **78**: 1204-11.
54. Rahn KH, Barenbrock M, Fritschka E, et al. Effect of nitrendipine on renal function in renal-transplant patients treated with cyclosporin: a randomised trial. *Lancet.* 1999; **354**: 1415-20.
55. McCulloch TA, Harper SJ, Donnelly PK, et al. Influence of nifedipine on interstitial fibrosis in renal transplant allografts treated with cyclosporin A. *J Clin Pathol.* 1994; **47**: 839-42.
56. Ladefoged SD, Andersen CB. Calcium channel blockers in kidney transplantation. *Clin Transplant.* 1994; **8**: 128-33.
57. Mourad G, Ribstein J, Mimran A. Converting-enzyme inhibitor versus calcium antagonist in cyclosporine-treated renal transplants. *Kidney Int.* 1993; **43**: 419-25.
58. Cross NB, Webster AC, Masson P, O'connell PJ, Craig JC. Antihypertensives for kidney transplant recipients: systematic review and meta-analysis of randomized controlled trials. *Transplantation.* 2009; **88**: 7-18.
59. Paoletti E, Bellino D, Marsano L, Cassottana P, Rolla D, Ratto E. Effects of ACE inhibitors on long-term outcome of renal transplant recipients: a randomized controlled trial. *Transplantation.* 2013; **95**: 889-95.
60. Philipp T, Martinez F, Geiger H, et al. Candesartan improves blood pressure control and reduces proteinuria in renal transplant recipients: results from SECRET. *Nephrol Dial Transplant.* 2010; **25**: 967-76.
61. Hiremath S, Fergusson D, Doucette S, Mulay AV, Knoll GA. Renin angiotensin system blockade in kidney transplantation: a systematic review of the evidence. *Am J Transplant.* 2007; **7**: 2350-60.
62. Opelz G, Döhler B. Cardiovascular Death in Kidney Recipients Treated With Renin-Angiotensin System Blockers. *Transplantation.* 2013.

63. Gaston RS, Kasiske BL, Fieberg AM, et al. Use of cardioprotective medications in kidney transplant recipients. *Am J Transplant.* 2009; **9**: 1811-5.
64. Schena FP, Pascoe MD, Alberu J, et al. Conversion from calcineurin inhibitors to sirolimus maintenance therapy in renal allograft recipients: 24-month efficacy and safety results from the CONVERT trial. *Transplantation.* 2009; **87**: 233-42.
65. Artz MA, Boots JM, Ligtenberg G, et al. Improved cardiovascular risk profile and renal function in renal transplant patients after randomized conversion from cyclosporine to tacrolimus. *J Am Soc Nephrol.* 2003; **14**: 1880-8.
66. Holdaas H, Fellström B, Cole E, et al. Long-term cardiac outcomes in renal transplant recipients receiving fluvastatin: the ALERT extension study. *Am J Transplant.* 2005; **5**: 2929-36.
67. Holdaas H, Fellström B, Jardine AG, et al. Beneficial effect of early initiation of lipid-lowering therapy following renal transplantation. *Nephrol Dial Transplant.* 2005; **20**: 974-80.
68. Jardine AG, Holdaas H, Fellström B, et al. fluvastatin prevents cardiac death and myocardial infarction in renal transplant recipients: post-hoc subgroup analyses of the ALERT Study. *Am J Transplant.* 2004; **4**: 988-95.
69. Ballantyne CM, Corsini A, Davidson MH, et al. Risk for myopathy with statin therapy in high-risk patients. *Arch Intern Med.* 2003; **163**: 553-64.
70. Palmer SC, Navaneethan SD, Craig JC, et al. HMG CoA reductase inhibitors (statins) for kidney transplant recipients. *Cochrane Database Syst Rev.* 2014; **1**: CD005019.
71. Group KW. KDIGO Clinical Practice Guideline for Lipid Management in Chronic Kidney Disease. *Kidney Int.* 2013; **Supplement 3**: 259-305.
72. Woodward RS, Schnitzler MA, Baty J, et al. Incidence and cost of new onset diabetes mellitus among U.S. wait-listed and transplanted renal allograft recipients. *Am J Transplant.* 2003; **3**: 590-8.
73. Cosio FG, Pesavento TE, Osei K, Henry ML, Ferguson RM. Post-transplant diabetes mellitus: increasing incidence in renal allograft recipients transplanted in recent years. *Kidney Int.* 2001; **59**: 732-7.
74. Valderhaug TG, Jenssen T, Hartmann A, et al. Fasting plasma glucose and glycosylated hemoglobin in the screening for diabetes mellitus after renal transplantation. *Transplantation.* 2009; **88**: 429-34.
75. Kasiske BL, Snyder JJ, Gilbertson D, Matas AJ. Diabetes mellitus after kidney transplantation in the United States. *Am J Transplant.* 2003; **3**: 178-85.
76. Sharif A, Baboolal K. Complications associated with new-onset diabetes after kidney transplantation. *Nat Rev Nephrol.* 2012; **8**: 34-42.
77. Gaston RS, Basadonna G, Cosio FG, et al. Transplantation in the diabetic patient with advanced chronic kidney disease: a task force report. *Am J Kidney Dis.* 2004; **44**: 529-42.
78. Ducloux D, Kazory A, Chalopin JM. Posttransplant diabetes mellitus and atherosclerotic events in renal transplant recipients: a prospective study. *Transplantation.* 2005; **79**: 438-43.
79. Kuo HT, Sampaio MS, Vincenti F, Bunnapradist S. Associations of pretransplant diabetes mellitus, new-onset diabetes after transplant, and

- acute rejection with transplant outcomes: an analysis of the Organ Procurement and Transplant Network/United Network for Organ Sharing (OPTN/UNOS) database. *Am J Kidney Dis.* 2010; **56**: 1127-39.
80. Fellström B, Jardine AG, Soveri I, et al. Renal dysfunction is a strong and independent risk factor for mortality and cardiovascular complications in renal transplantation. *Am J Transplant.* 2005; **5**: 1986-91.
81. Abbott KC, Yuan CM, Taylor AJ, Cruess DF, Agodoa LY. Early renal insufficiency and hospitalized heart disease after renal transplantation in the era of modern immunosuppression. *J Am Soc Nephrol.* 2003; **14**: 2358-65.
82. Weiner DE, Carpenter MA, Levey AS, et al. Kidney function and risk of cardiovascular disease and mortality in kidney transplant recipients: the FAVORIT trial. *Am J Transplant.* 2012; **12**: 2437-45.
83. Rigatto C, Foley R, Jeffery J, Negrijn C, Tribula C, Parfrey P. Electrocardiographic left ventricular hypertrophy in renal transplant recipients: prognostic value and impact of blood pressure and anemia. *J Am Soc Nephrol.* 2003; **14**: 462-8.
84. Parfrey PS, Harnett JD, Foley RN, et al. Impact of renal transplantation on uremic cardiomyopathy. *Transplantation.* 1995; **60**: 908-14.
85. Dudziak M, Debska-Slizień A, Rutkowski B. Cardiovascular effects of successful renal transplantation: a 30-month study on left ventricular morphology, systolic and diastolic functions. *Transplant Proc.* 2005; **37**: 1039-43.
86. Patel RK, Mark PB, Johnston N, McGregor E, Dargie HJ, Jardine AG. Renal transplantation is not associated with regression of left ventricular hypertrophy: a magnetic resonance study. *Clin J Am Soc Nephrol.* 2008; **3**: 1807-11.
87. Rigatto C, Parfrey P. Therapy insight: management of cardiovascular disease in the renal transplant recipient. *Nat Clin Pract Nephrol.* 2006; **2**: 514-26.
88. Ventura JE, Tavella N, Romero C, Petraglia A, Báez A, Muñoz L. Aortic valve calcification is an independent factor of left ventricular hypertrophy in patients on maintenance haemodialysis. *Nephrol Dial Transplant.* 2002; **17**: 1795-801.
89. Ibernón M, Moreso F, Ruiz-Majoral A, et al. Contribution of anemia and hypertension to left ventricular hypertrophy during the initial 2 years after renal transplantation. *Transplant Proc.* 2011; **43**: 2199-204.
90. Hernández D, Lacalzada J, Salido E, et al. Regression of left ventricular hypertrophy by lisinopril after renal transplantation: role of ACE gene polymorphism. *Kidney Int.* 2000; **58**: 889-97.
91. Paoletti E, Cassottana P, Amidone M, Gherzi M, Rolla D, Cannella G. ACE inhibitors and persistent left ventricular hypertrophy after renal transplantation: a randomized clinical trial. *Am J Kidney Dis.* 2007; **50**: 133-42.
92. Paoletti E, Amidone M, Cassottana P, Gherzi M, Marsano L, Cannella G. Effect of sirolimus on left ventricular hypertrophy in kidney transplant recipients: a 1-year nonrandomized controlled trial. *Am J Kidney Dis.* 2008; **52**: 324-30.

- Accepted Article
93. Paoletti E, Marsano L, Bellino D, Cassottana P, Cannella G. Effect of everolimus on left ventricular hypertrophy of de novo kidney transplant recipients: a 1 year, randomized, controlled trial. *Transplantation*. 2012; **93**: 503-8.
 94. Kurtz A, Della Bruna R, Kühn K. Cyclosporine A enhances renin secretion and production in isolated juxtaglomerular cells. *Kidney Int*. 1988; **33**: 947-53.
 95. Mason J, Müller-Schweinitzer E, Dupont M, et al. Cyclosporine and the renin-angiotensin system. *Kidney Int Suppl*. 1991; **32**: S28-32.
 96. Mourer JS, de Koning EJ, van Zwet EW, Mallat MJ, Rabelink TJ, de Fijter JW. Impact of late calcineurin inhibitor withdrawal on ambulatory blood pressure and carotid intima media thickness in renal transplant recipients. *Transplantation*. 2013; **96**: 49-57.
 97. Mulay AV, Hussain N, Fergusson D, Knoll GA. Calcineurin inhibitor withdrawal from sirolimus-based therapy in kidney transplantation: a systematic review of randomized trials. *Am J Transplant*. 2005; **5**: 1748-56.
 98. Ekberg H, Bernasconi C, Tedesco-Silva H, et al. Calcineurin inhibitor minimization in the Symphony study: observational results 3 years after transplantation. *Am J Transplant*. 2009; **9**: 1876-85.
 99. Egbuna OI, Davis RB, Chudinski R, et al. Outcomes with conversion from calcineurin inhibitors to sirolimus after renal transplantation in the context of steroid withdrawal or steroid continuation. *Transplantation*. 2009; **88**: 684-92.
 100. Lebranchu Y, Thierry A, Toupance O, et al. Efficacy on renal function of early conversion from cyclosporine to sirolimus 3 months after renal transplantation: concept study. *Am J Transplant*. 2009; **9**: 1115-23.
 101. Cosio FG, Falkenhain ME, Pesavento TE, et al. Patient survival after renal transplantation: II. The impact of smoking. *Clin Transplant*. 1999; **13**: 336-41.
 102. Kasiske BL, Klinger D. Cigarette smoking in renal transplant recipients. *J Am Soc Nephrol*. 2000; **11**: 753-9.
 103. Group KDIGOKTW. KDIGO clinical practice guideline for the care of kidney transplant recipients. *Am J Transplant*. 2009; **9 Suppl 3**: S1-155.
 104. Baker R, Jardine A, Andrews P. Renal Association Clinical Practice Guideline on post-operative care of the kidney transplant recipient. *Nephron Clin Pract*. 2011; **118 Suppl 1**: c311-47.
 105. de Vries AP, Bakker SJ, van Son WJ, et al. Metabolic syndrome is associated with impaired long-term renal allograft function; not all component criteria contribute equally. *Am J Transplant*. 2004; **4**: 1675-83.
 106. Meier-Kriesche HU, Arndorfer JA, Kaplan B. The impact of body mass index on renal transplant outcomes: a significant independent risk factor for graft failure and patient death. *Transplantation*. 2002; **73**: 70-4.
 107. Abbott KC, Reynolds JC, Taylor AJ, Agodoa LY. Hospitalized atrial fibrillation after renal transplantation in the United States. *Am J Transplant*. 2003; **3**: 471-6.
 108. Zelle DM, Corpeleijn E, Stolk RP, et al. Low physical activity and risk of cardiovascular and all-cause mortality in renal transplant recipients. *Clin J Am Soc Nephrol*. 2011; **6**: 898-905.

109. Didsbury M, McGee RG, Tong A, et al. Exercise training in solid organ transplant recipients: a systematic review and meta-analysis. *Transplantation*. 2013; **95**: 679-87.
110. Ducloux D, Motte G, Challier B, Gibey R, Chalopin JM. Serum total homocysteine and cardiovascular disease occurrence in chronic, stable renal transplant recipients: a prospective study. *J Am Soc Nephrol*. 2000; **11**: 134-7.
111. Bostom AG, Carpenter MA, Kusek JW, et al. Homocysteine-lowering and cardiovascular disease outcomes in kidney transplant recipients: primary results from the Folic Acid for Vascular Outcome Reduction in Transplantation trial. *Circulation*. 2011; **123**: 1763-70.
112. Rigatto C, Parfrey P, Foley R, Negrijn C, Tribula C, Jeffery J. Congestive heart failure in renal transplant recipients: risk factors, outcomes, and relationship with ischemic heart disease. *J Am Soc Nephrol*. 2002; **13**: 1084-90.
113. Abedini S, Holme I, März W, et al. Inflammation in renal transplantation. *Clin J Am Soc Nephrol*. 2009; **4**: 1246-54.
114. Meier-Kriesche HU, Baliga R, Kaplan B. Decreased renal function is a strong risk factor for cardiovascular death after renal transplantation. *Transplantation*. 2003; **75**: 1291-5.
115. Haidinger M, Werzowa J, Hecking M, et al. Efficacy and safety of vildagliptin in new-onset diabetes after kidney transplantation--a randomized, double-blind, placebo-controlled trial. *Am J Transplant*. 2014; **14**: 115-23.
116. Hecking M, Haidinger M, Döller D, et al. Early basal insulin therapy decreases new-onset diabetes after renal transplantation. *J Am Soc Nephrol*. 2012; **23**: 739-49.
117. Woodle ES, First MR, Pirsch J, et al. A prospective, randomized, double-blind, placebo-controlled multicenter trial comparing early (7 day) corticosteroid cessation versus long-term, low-dose corticosteroid therapy. *Ann Surg*. 2008; **248**: 564-77.
118. Vincenti F, Schena FP, Paraskevas S, et al. A randomized, multicenter study of steroid avoidance, early steroid withdrawal or standard steroid therapy in kidney transplant recipients. *Am J Transplant*. 2008; **8**: 307-16.
119. Rizzari MD, Suszynski TM, Gillingham KJ, et al. Ten-year outcome after rapid discontinuation of prednisone in adult primary kidney transplantation. *Clin J Am Soc Nephrol*. 2012; **7**: 494-503.
120. Vanrenterghem Y, Lebranchu Y, Hené R, Oppenheimer F, Ekberg H. Double-blind comparison of two corticosteroid regimens plus mycophenolate mofetil and cyclosporine for prevention of acute renal allograft rejection. *Transplantation*. 2000; **70**: 1352-9.
121. Ahsan N, Hricik D, Matas A, et al. Prednisone withdrawal in kidney transplant recipients on cyclosporine and mycophenolate mofetil--a prospective randomized study. Steroid Withdrawal Study Group. *Transplantation*. 1999; **68**: 1865-74.

