


## Accepted Manuscript

Title: Original Research Article: Carotenoid stability during storage of yellow gari made from biofortified cassava or with palm oil

Author: Aurelie Bechoff Ugo Chijioke Keith I. Tomlins Pesila Govinden Paul Ilona Andrew Westby Erick Boy



PII: S0889-1575(15)00143-X  
DOI: <http://dx.doi.org/doi:10.1016/j.jfca.2015.06.002>  
Reference: YJFCA 2598

To appear in:

Received date: 29-10-2014  
Revised date: 1-6-2015  
Accepted date: 8-6-2015

Please cite this article as: Bechoff, A., Chijioke, U., Tomlins, K. I., Govinden, P., Ilona, P., Westby, A., and Boy, E., Original Research Article: Carotenoid stability during storage of yellow gari made from biofortified cassava or with palm oil, *Journal of Food Composition and Analysis* (2015), <http://dx.doi.org/10.1016/j.jfca.2015.06.002>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Original Research Article: Carotenoid stability during storage of yellow**  
2 **gari made from biofortified cassava or with palm oil**

3  
4 **Aurelie Bechoff<sup>a\*</sup>, Ugo Chijioke<sup>b</sup>, Keith I. Tomlins<sup>a</sup>, Pesila Govinden<sup>a</sup>,**  
5 **Paul Ilona<sup>c</sup>, Andrew Westby<sup>a</sup>, Erick Boy<sup>d</sup>**

6 <sup>a</sup> *Natural Resources Institute, University of Greenwich, Central Avenue, Chatham Maritime, Kent,*  
7 *ME4 4TB, United Kingdom*

8 <sup>b</sup> *National Root Crop Research Institute, Umudike, PMB 7006, Umuahia, Abia State, Nigeria.*

9 <sup>c</sup> *HarvestPlus Nigeria, c/o International Institute of Tropical Agriculture (IITA) PMB 5320, Ibadan,*  
10 *Oyo State, Nigeria*

11 <sup>d</sup> *HarvestPlus Headquarters, c/o IFPRI, 2033 K Street, NW, Washington, DC 20006-1002, USA*

12 \*Corresponding author. Tel: +44 (0)1634883071; fax +44 (0)1634 88 3386. E-mail address:  
13 [a.bechoff@gre.ac.uk](mailto:a.bechoff@gre.ac.uk) (A. Bechoff).

Field Code Changed

14  
15 **ABSTRACT**

16 The carotenoid composition of gari made from biofortified cassava (BG) was compared to  
17 that of existing gari of similar appearance but made from white cassava with added red palm  
18 oil (RPG). Storage of both yellow gari products was modelled at ambient temperatures  
19 typical of tropical areas (19-40°C) over a 3 month-period at constant relative humidity.  
20 Carotenoid content and hence vitamin A activity of the gari products decreased markedly  
21 with time and temperature. Trans-β-carotene degradation fitted well the kinetics predicted by  
22 the Arrhenius model, in particular for BG. Activation energies for trans-β-carotene were  
23 ~~63.160.4~~ and ~~82.381.0~~ kJ.mol<sup>-1</sup> for BG and RPG respectively (R<sup>2</sup> = 0.998 and 0.988-997  
24 respectively): hence the minimum energy to cause degradation of trans-β-carotene in gari was  
25 lower with BG. Rates of degradation of 9-cis β-carotene in gari were of the same order as

26 with trans- $\beta$ -carotene. Although the initial content of trans- $\beta$ -carotene was twice as high in  
27 the BG compared to RPG, trans- $\beta$ -carotene in BG degraded much faster. Results showed that  
28 the average shelf life at ambient temperature for BG was significantly shorter than for RPG  
29 and therefore carotenoids in BG were less stable than in RPG.

30

31 **Key-words:** carotenoid degradation, biofortified cassava, gari, palm oil, storage, temperature,  
32 model prediction, vitamin A deficiency, nutritional impact, food composition

33

34 Abbreviations: BG: gari from biofortified yellow cassava; RPG: gari from white cassava with added palm oil.  
35 FW: fresh weight basis; DW: dry weight basis; R: retention; RAE: Retinol Activity Equivalent; EAR: Estimated  
36 average requirement; VAD: vitamin A deficiency; pVACs: provitamin A carotenoids

37

## 38 1. Introduction

39 Cassava (*Manihot esculenta* Crantz), a tropical root crop, is a starch staple and an important  
40 crop for food security for millions of people in sub-Saharan Africa. The short shelf life (2-3  
41 days) of the crop however is a major drawback because it limits its transportation and  
42 consumption in its fresh form (Westby, 2002). Hence processed forms of cassava have been  
43 developed. Gari is a dried granulated food product with a slight acidic taste is one of the most  
44 common processed forms of cassava in West Africa. Gari is produced by grating (to remove  
45 cyanide inherent to the root) followed by fermentation (to produce flavour and make the  
46 product sour – by lowering pH value, the shelf life is also increased) and drying (to extend  
47 the shelf life). Gari presents as a dried form, which makes it stable under ambient conditions,  
48 easy to transport and can be stored for many months.

49

50 Recently biofortified varieties of cassava that contain significant levels of provitamin A  
51 carotenoids (pVACs) have been developed by conventional plant breeding methods and

Formatted: Font: Italic

52 released for use by the local populations. These biofortified varieties could be used to help  
53 tackle vitamin A deficiency (VAD) (Saltzman et al., 2009) an important public health  
54 problem in sub-Saharan Africa and in the world. In some countries higher mortality rates,  
55 susceptibility to infections and blindness can clearly be attributed to VAD occurrence. The  
56 new varieties have a different visible colour to the traditional varieties because they are  
57 yellow compared to traditional cassava that is white and very low in provitamin A. These  
58 biofortified varieties produce a gari that is very similar in colour to gari made with added  
59 crude palm oil that also contains vitamin A (Abu et al. 2006). However, there are some  
60 disadvantages in adding palm oil. Firstly it is not widely consumed, and also, the addition of  
61 palm oil adds to the production costs, and finally darkening of gari occurs when added in  
62 excess and rancidity can happen during storage (Burri et al. 2012). The use of gari made from  
63 biofortified cassava would therefore solve the issue of rancidity and without the additional  
64 cost of palm oil help tackle vitamin A deficiency on a wider scale. Nigeria is the most  
65 densely populated country in Africa and an emerging country with a fast growing population  
66 that could reach 300 million by 2050 (Oshikoya 2008) and the impact of such a product could  
67 potentially impact millions. But the challenges are to measure the stability of carotenoids in  
68 gari made with biofortified cassava and also to compare it with gari made with crude palm oil  
69 that also contains provitamin A.  
70  
71 Understanding how pVACs degrade during storage of vitamin A- containing-gari is critical  
72 because it will affect its nutritional impact. Storage of gari at ambient temperatures is a  
73 current practice in Nigeria. Gari storage is not only generally practiced at household level but  
74 also at commercial level. Periods of storage are on average around six months but some  
75 processors can store up to a year. Stability of carotenoids in gari made from white cassava  
76 varieties with added palm oil has been studied during processing and storage at ambient

77 temperature (Abu et al., 2006; Gouado et al., 2008; Uzomah et al., 2006). Gouado et al.  
78 (2008) showed that during gari processing the product retained a significant amount of the  
79 carotenoids from palm oil. In contrast, authors reported a significant loss in carotenoid during  
80 storage that followed processing: Abu et al. (2006) measured a loss of 57% in total carotenoid  
81 after 4 months at 28°C in Nigeria. However Uzomah et al. (2006) working also at 28°C in  
82 Nigeria reported different results: an average loss of 25% after the 2<sup>nd</sup> week of storage and of  
83 50% by the 3<sup>rd</sup> week. Some gari samples that had lower levels of palm oil lost most of  
84 vitamin A activity after only 2 weeks of storage (Uzomah et al., 2006). These dissimilar  
85 findings appeal for more research to understand the carotenoid degradation in palm oil gari  
86 during storage under controlled conditions.

87  
88 The effect of storage on carotenoid in products made from biofortified crops has also been  
89 studied. Stability of carotenoids during storage of biofortified maize has been studied by  
90 Mugode et al. (2014). It was shown that most of the carotenoid degradation occurred in the  
91 first weeks of storage and the degradation rate then lowered. Bechoff et al. (2011a) working  
92 on biofortified orange-fleshed sweet potato similarly reported that storage of dried chips had  
93 a dramatic effect on carotenoid stability (~80% loss in 4 months). Furthermore, the authors  
94 demonstrated that the carotenoid degradation followed a first order degradation (logarithmic  
95 curve), which explains why the degradation was higher in the first weeks of storage and then  
96 stabilised with time. Temperature and oxygen were the main factors that caused the loss in  
97 carotenoids whilst water activity only had a minor effect (Bechoff et al., 2010). A  
98 mathematical model was developed to predict the degradation of trans- $\beta$ -carotene, the main  
99 carotenoid in sweet potato, under controlled conditions of temperature, oxygen and humidity  
100 and the model was validated by field data (Bechoff et al., 2010).

101 Little research has been done on biofortified cassava with relation to storage of gari. Ukenye,  
102 et al. (2013) observed that the gari made with biofortified cassava was similar in appearance  
103 to the gari made with palm oil (traditional gari). Onadipe Olapeju (2011) studied the  
104 degradation of total carotenoids in gari from the varieties of biofortified yellow cassava  
105 developed in Nigeria (01/1371; 01/1368 and 01/1412). According to the data presented by the  
106 author, 50% on average of total carotenoids were lost after 3 month-storage at  $30\pm 2^{\circ}\text{C}$ .  
107 However there was minimal information on the degradation rate and the influence of  
108 temperature.  
109 There appears to be little published studies on the prediction of carotenoid degradation during  
110 storage although it is a critical issue for gari containing carotenoids and hence gari's potential  
111 impact on tackling VAD. More research is needed to understand the stability of gari from  
112 biofortified cassava (BG). Traditional gari made with crude "red" palm oil (RPG) is a  
113 common product in Southern Nigeria and should also be tested for stability to compare with  
114 gari made from biofortified cassava. This information will be useful to understand the  
115 potential for the promotion and marketing of gari made from biofortified yellow cassava in  
116 Nigeria and its contribution to reducing VAD.

117

## 118 2. Materials and Methods

119

### 120 2.1. Description of samples

121 Biofortified yellow cassava roots (TMS 01/1371) were harvested from Ikenne (about 2h drive  
122 south from IITA, Ibadan, Nigeria). White cassava roots (variety IITA 3303, locally called  
123 Oko-Iyawo) were harvested from the Army Barracks field in Ibadan, Nigeria. Cassava roots  
124 had a growing period of approximately 12 months after planting. Roots (50kg per variety)  
125 were processed into gari by commercial processors based at the Army Barracks, Ibadan:

126 biofortified yellow cassava variety and white cassava variety with added red palm oil  
127 (approximately 0.45L/32.6kg or 0.328g/32.6kg of grated mash). The amount of red palm  
128 oil to add to the mash was selected by the commercial processors. All the processing  
129 parameters (time, temperature, pH, quantities etc.) were monitored and the gari produced was  
130 of commercial quality. A representative sample of gari was stored in the freezer (-20°C) and  
131 maintained frozen during transport and up to start of the storage experiment.

132

### 133 *2.2. Storage experiment and sample collection for carotenoid analysis*

134 Gari samples (about 1kg of from either BG or from RPG) were divided into equal portions  
135 using a riffle divider. Representative gari sub-samples (50g) were wrapped in a sewed cotton  
136 bag and stored in Kilner jars (having metal lever catch and rubber seal) with a saturated salt  
137 solution (Sodium Bromide (NaBr) that has a water activity ( $a_w$ ) of about 0.5). The saturated  
138 salt solution was used to maintain the ambient relative humidity constant around the gari  
139 product so that only the effect of temperature could be measured. Jars in triplicate were  
140 placed in incubators (LMS Cooled Incubator, Sevenoaks, UK) at the Natural Resources  
141 Institute (NRI), University of Greenwich, UK and set at four different ambient temperatures  
142 ( $19\pm 1$ ,  $26\pm 1$ ,  $33\pm 1$  and  $40\pm 1$  °C). The range of temperatures was comprised between the  
143 minimum and maximum ambient temperatures in Nigeria. Hence the degradation of  
144 carotenoids during storage of gari could be predicted under similar temperature conditions as  
145 those found in Nigeria. Samples were stored in jars for 80 days (20<sup>th</sup> November 2012- 7<sup>th</sup>  
146 February 2013). The storing system used in the incubators was similar to the one used with  
147 dried sweet potato (Bechoff, 2010). Stored gari samples (about 5g) were collected in a  
148 representative manner by using a riffle divider and moisture content was checked at the  
149 beginning and the end of storage. Sample collections at 19°C were on 24<sup>th</sup>; 49<sup>th</sup>; 60<sup>th</sup>; 80<sup>th</sup>  
150 day; collections at 26°C were on 18<sup>th</sup>; 24<sup>th</sup>; 49<sup>th</sup>; 60<sup>th</sup>; 80<sup>th</sup> day; collections at 33°C were on

151 10<sup>th</sup>; 18<sup>th</sup>; 24<sup>th</sup>; 49<sup>th</sup>; 60<sup>th</sup>; 80<sup>th</sup> day.; collections at 40°C were on 10<sup>th</sup>; 18<sup>th</sup>; 24<sup>th</sup>; 31<sup>st</sup>; 49<sup>th</sup>; 60<sup>th</sup>  
152 day. Collected gari samples were immediately stored at -80°C.

153

### 154 2.3. Carotenoid analysis

155 The extraction stage was based on Rodriguez-Amaya and Kimura (2004) and described in  
156 Bechoff et al. (2011a). Analyses were carried out at NRI, UK. In brief, gari samples (0.6-2.0g  
157 depending on the carotenoid content in sample) were rehydrated for 10 minutes in 10 mL  
158 tepid deionised water (water was heated at 30°C to facilitate extraction). The samples were  
159 homogenised with 50mL methanol:tetrahydrofuran (THF) (1:1) for 1 minute and filtered. The  
160 homogenised extract was rinsed with methanol:THF (1:1) until there was no yellow colour  
161 left in the residue. Partition between the aqueous phase and organic phase containing the  
162 carotenoids was achieved by the addition of petroleum ether (PE 40-60°C) and sodium  
163 chloride (NaCl) solution (10%). The PE phase was further washed with deionised water,  
164 dried by addition of anhydrous sodium sulphate, then filtered and made up to volume (50  
165 mL). For the determination of individual carotenoids by HPLC, the carotenoid extracts in PE  
166 | (~~20mL~~20mL) were dried by flushing with nitrogen in a dry block system at 35° C. Extracts  
167 were then dissolved in 500 µL THF: Methanol (1:1). After vortexing, dissolved samples were  
168 collected into a vial with septum for HPLC analysis. A reverse-phase high performance liquid  
169 chromatography using an Agilent 1200 system (UK) was used with a polymeric C30 reverse  
170 phase column (250 x 4.6 mm i.d. 5µm YMC (EUROP GmbH Germany) having a flow rate of  
171 1 ml.min<sup>-1</sup>, a temperature of 25°C, a running time of 40 minutes and an injection volume of  
172 10µL. The isocratic mix consisted of Methanol: MTBE (80:20). Detection of compounds was  
173 performed at 450nm. Concentrations on a fresh weight basis were determined by comparing  
174 with standard curve of pure trans-β-carotene (Sigma, UK). Percentages of cis-isomers and  
175 other minor compounds such as epoxides were also determined (Bechoff et al. 2011b). Minor



176 compounds (epoxides of  $\beta$ -carotene) were tentatively identified when these were in very  
 177 small amounts. Trans- $\alpha$ -carotene was identified by injection of a mixture of carotenoids from  
 178 carrot extract (Sigma, UK).

179

#### 180 2.4. Retention

181 Retention of trans- $\beta$ -carotene (TR) was calculated using a simplified equation of the true  
 182 retention assuming that the dry matter content was constant and therefore the weight of gari  
 183 did not vary in the samples stored (at constant humidity in incubators) (Equation 1).

$$R(\%) = 100 \times \frac{\text{trans-}\beta\text{-carotene content per kg of stored gari}}{\text{trans-}\beta\text{-carotene per kg of unstored gari}} \quad (\text{Equation 1})$$

#### 186 2.5. Kinetics modelling and statistical analysis

187

188 Carotenoid content was determined on a fresh weight basis (FW) at different storage times  
 189 and temperatures in triplicate. Carotenoid degradation followed a first order kinetics. Hence  
 190 logarithms of carotenoid content were linear as a function of storage time (Excel, Windows  
 191 2007) (Equation 2):

$$\ln C = \ln C_0 - kt \quad (\text{Equation 2})$$

193 Where: C: Carotenoid content of gari ( $\mu\text{g}\cdot\text{g}^{-1}$ ) at storage time t;  $C_0$ : Carotenoid content of  
 194 food ( $\mu\text{g}\cdot\text{g}^{-1}$ ) at initial time (before storage); t: storage time (day); k: degradation constant rate  
 195 ( $\text{day}^{-1}$ ). k was determined graphically and using linear regression (XLStat 2014 software,  
 196 <http://www.xlstat.com>) for the three replicate data pooled together.

197

198 Carotenoid degradation kinetics can be evaluated using different models as in Bechoff et al.  
 199 (2010). The most common model is Arrhenius model. The Arrhenius model (Equation 3) is

200 an empirical collision model that describes the relationship between reaction constant rates  
 201 and temperature using activation energy ( $E_a$ ) and a pre-exponential factor ( $k_\infty$ ).

$$k = k_\infty e^{-\frac{E_a}{RT}} \quad (\text{Equation 3})$$

204 Where: T: temperature (K)

205  $k_\infty$ : value of k at infinite time  $t = \infty$  ( $\text{day}^{-1}$ )

206  $E_a$ : Activation energy ( $\text{kJ}\cdot\text{mol}^{-1}$ )

207 R: gas constant =  $8.314 \text{ J} \cdot \text{K}^{-1} \cdot \text{mol}^{-1}$

208

209 The prediction model (Equation 4) is calculated by the equation based on the Arrhenius  
 210 model and using temperature (T) expressed in Kelvin:

$$C = C_0 e^{-k_\infty \int_0^t e^{\frac{RT}{212}} dt} \quad (\text{Equation 4})$$

214 Determination of  $E_a$  and  $k_\infty$  parameters helps predict carotenoid degradation for known

215 storage temperatures and times. [Ea and k<sub>∞</sub> were determined using linear regression \(XLStat](#)  
 216 [2014 software\).](#)

217 ~~Other models however exist. The Eyring model (Equation 5) is based on the transition state~~  
 218 ~~theory in which enthalpy of activation ( $\Delta H^*$ ) and entropy of activation ( $\Delta S^*$ ) are the model's~~  
 219 ~~parameters. The model's parameters were identified from experimental data measured in~~  
 220 ~~triplicate using linear regressions.~~

$$k = \frac{k_B}{h} T \cdot e^{-\frac{\Delta H^*}{RT}} \quad (\text{Equation 5})$$

223 ~~Where:  $k_B$ : Boltzmann constant =  $1.381 \cdot 10^{-23} \text{ J}\cdot\text{K}^{-1}$~~

224  ~~$h$ : Planck constant =  $6.626 \cdot 10^{-34} \text{ J}\cdot\text{s}$~~

225  ~~$\Delta H^*$ : activation enthalpy ( $\text{kJ}\cdot\text{mol}^{-1}$ )~~

226 |  ~~$\Delta S^\ddagger$ : activation entropy ( $\text{J}\cdot\text{mol}^{-1}\cdot\text{K}^{-1}$ )~~

227

228 | Data were processed on SPSS 20.0 software using ~~One-Way Analysis of Variance to a t-test~~  
229 | ~~to~~ determine if there were significant differences ( $p < 0.05$ ) between model parameters of the  
230 | two gari products (BG and RPG).

231

### 232 | 3. Results and discussion

233

#### 234 | 3.2.1. Nutritional value of stored gari

235

236 | The nutritional value (vitamin A activity) of the gari products at different storage times and  
237 | ambient temperatures was determined (Table 1). Vitamin A activities per daily “100 g”  
238 | portion of BG and RPG based on the classical estimate were ~~125.6126~~ and ~~84.885~~ RAE,  
239 | respectively. On the other hand unstored BG and RPG had substantial average vitamin A  
240 | activity based on the new estimate (301.4 and 203.5 RAE, respectively). The Estimated  
241 | Average Requirement (EAR) for a child is 200 RAE based on FAO/WHO (2002) and 275  
242 | RAE based on the National Academy of Sciences/Institute of Medicine (2001) s’  
243 | recommendations. According to the standard estimate for food,  $1 \mu\text{g}\cdot\text{g}^{-1}$  Retinol Activity  
244 | Equivalent (RAE) corresponds to  $12 \mu\text{g}\cdot\text{g}^{-1}$  all-*trans*-BC or  $24 \mu\text{g}\cdot\text{g}^{-1}$  minor carotenoids  
245 | (National Academy of Sciences / Institute of Medicine, 2001). Recent studies on the  
246 | bioconversion of provitamin A from cassava products indicated that the factor might be  
247 | lower: working with women, Liu et al (2010) showed that the bioefficacy of the BC from  
248 | porridge made with biofortified cassava was as good as that of a BC supplement (2:1). Later,  
249 | La Frano et al. (2013) calculated a bioconversion factor of 4.5:1 for biofortified cassava  
250 | meals and Phorbee et al. (2013), a factor of 6:1. Therefore the conversion factor 5:1 was

251 suggested here as 'new' estimated bioconversion factor. With either estimate (classical or  
252 new), the potential daily contribution of yellow gari to a child's vitamin A intake is close or  
253 superior to 50% of EAR.

254

255 However, during storage, vitamin A activity of gari products sharply decreased. Our results  
256 using the classical estimate showed that after 60 days vitamin A activity was 54.1 and 64.2  
257 RAE per 100g at 26°C; 24.3 and 51.4 RAE at 33°C, and 24.3 and 8.69 and 33.4 RAE at 40°C  
258 for BG and RPG, respectively. Calculations on estimation of vitamin A activity presented  
259 here highlighted that the choice of adequate bioconversion factor (classical or new) is critical  
260 because it will be determinant to provide advice on shelf life of the gari product.

261 Using the new estimate, values superior to 50 RAE would only be achieved with BG stored at  
262 33°C for up to 60 days or at 40°C for 35 days. The decrease was temperature dependent.

263 Uzomah et al. (2006) analysed gari products from six different locations in Eastern parts of  
264 Nigeria where red palm oil gari is the most common form of gari consumption. Their results  
265 similarly reported that vitamin A activity of palm oil-enriched gari significantly decreased  
266 when stored at ambient temperature. Initial vitamin A activity in freshly made gari was very  
267 variable ranging between 13.2-723 RAE per 100g (using classical estimate), which shows a  
268 wide variation in levels of palm oil added by different communities. Uzomah et al. (2006)  
269 reported a loss of 25% in activity after 2 weeks and 50% after 3 weeks of storage at ambient  
270 temperature of 28°C. This loss was higher than the results presented here (10% for RPG after  
271 15 days) but other factors under field storage such as light and humidity might have  
272 contributed to additional loss (Uzomah et al., 2006).

273

274 ~~Calculations on estimation of vitamin A activity presented here highlighted that the choice of~~  
275 ~~adequate bioconversion factor (classical or new) is critical because it will be determinant to~~  
276 ~~provide advice on shelf life of the gari product.~~

277

278 ~~3.12. Carotenoid composition of unstored gari from yellow cassava and white cassava with~~  
279 ~~palm oil~~

280

281 Trans- $\beta$ -carotene and other carotenoid (9-cis, 13-cis, 5,6 epoxy, 5,8- $\beta$ -carotene and trans- $\alpha$ -  
282 carotene) contents were determined on a fresh weight basis at different storage durations and  
283 temperatures (Fig. 2).

284

285 Trans- $\beta$ -carotene content on a fresh basis (FW) in unstored gari from TMS 01/1371 variety  
286 was  $10.9 \mu\text{g}\cdot\text{g}^{-1}$  on average and  $13.0 \mu\text{g}\cdot\text{g}^{-1}$  as a maximum. Carotenoid content of gari from  
287 yellow cassava (BG) obtained by conventional cross breeding techniques has been  
288 determined (Chavez et al., 2007; La Frano et al., 2014; Maziya-Dixon et al., 2009; Onadipe  
289 Olapeju et al., 2011; Thakkar et al., 2009). Chavez et al. (2007); Frano et al. (2014) reported  
290 lower trans- $\beta$ -carotene content in gari, between 3 and  $4 \mu\text{g}\cdot\text{g}^{-1}$  on a dry basis (DW) whilst  
291 Maziya-Dixon et al. (2009) and Onadipe Olapeju et al. (2011) working on the variety TMS  
292 01/1371 indicated total carotenoid content of 16 and  $20 \mu\text{g}\cdot\text{g}^{-1}$  FW, respectively that was in  
293 accordance with our data: in this study, maximal total carotenoid content determined by  
294 spectrophotometer was approximately  $18 \mu\text{g}\cdot\text{g}^{-1}$  FW (data not shown). In addition, Thakkar  
295 et al. (2009) found a trans- $\beta$ -carotene content in gari from TMS 01/1371 variety around 15  
296  $\mu\text{g}\cdot\text{g}^{-1}$  DW, which would be approximately  $13 \mu\text{g}\cdot\text{g}^{-1}$  FW for a product with approximately  
297 10% moisture content. Our data on unstored gari from yellow cassava TMS 01/1371 is  
298 therefore mostly in agreement with previously published work with the same variety.

299 Carotenoid content of gari from genetically modified cassava has also been measured (Failla  
300 et al., 2012). Nonetheless levels of trans- $\beta$ -carotene reported between 3-8  $\mu\text{g}\cdot\text{g}^{-1}$  DW were  
301 lower than in our study.

302

303 Differences in initial carotenoid content of gari described by different authors may be  
304 explained by cassava varietal differences but also by variations in processing steps that  
305 influences carotenoid retention from roots into gari.

306

307 Average trans- $\beta$ -carotene and trans- $\alpha$ -carotene content in unstored RPG were 5.60  $\mu\text{g}\cdot\text{g}^{-1}$  and  
308 3.10  $\mu\text{g}\cdot\text{g}^{-1}$  respectively, on a fresh basis (FW). Trans- $\alpha$ -carotene content in RPG was about  
309 six times more than in BG. The higher concentration of trans- $\alpha$ -carotene in RPG can be  
310 explained by the composition of red palm oil (Fig. 1A and B): red palm oil is known to  
311 contain both trans- $\beta$ -carotene and trans- $\alpha$ -carotene (Bonnie Tay & Choo, 2000). The  
312 carotenoid content of this gari would depend on the amount of palm added during the  
313 process, which can be variable according to practices of gari processors (Gouado et al.,  
314 2008). Alpha and  $\beta$ -carotene contents of gari reported by Gouado et al. (2008) were at least  
315 100 times greater than ours (trans- $\alpha$ -carotene: 352.6-1572.5  $\mu\text{g}\cdot\text{g}^{-1}$  and trans- $\beta$ -carotene:  
316 309.7-1624.3  $\mu\text{g}\cdot\text{g}^{-1}$ , for 2 and 8 ml of oil respectively for 210g of gari) and therefore indicate  
317 that the analysis was done on the palm oil and not on the gari product itself (2mL for 210g is  
318 actually quite close to the amount of oil added in this study). Mortensen (2005) reported that  
319 palm oil identified on a C30 HPLC column mainly contained trans- $\beta$ -carotene and trans- $\alpha$ -  
320 carotene and the other compound identified was 13-cis- $\beta$ -carotene. Although many minor  
321 carotenoids (about 15, including lycopene and  $\gamma$ -carotene) are present in palm oil (Bonnie  
322 Tay & Choo, 2000; Mortensen, 2005) they were not visible on the present chromatogram  
323 (Fig. 1B). Andreu-Sevilla et al. (2009) equally reported that the main carotenoids absorbed in

324 potato fried in palm oil were  $\alpha$  and  $\beta$ -carotene, which was in accordance with this present  
325 study. In addition, Andreu-Sevilla et al. (2009) reported amounts of lutein, 5,6 epoxy- $\alpha$ -  
326 carotene,  $\gamma$ -carotene;  $\delta$ -carotene;  $\epsilon$ -carotene; 15-cis and 9-cis- $\beta$ -carotene in palm oil that were  
327 partially absorbed in fried potato. The minor carotenoids in common for BG and RPG that  
328 were identified in our study were all from  $\beta$ -carotene, being 13-cis, 9-cis, 5,6 and 5,8 epoxy-  
329  $\beta$ -carotene. In some cases, minor carotenoids were in very small amount, which made their  
330 identification more difficult (Fig. 21).

331

### 332 3.3. Kinetics of carotenoid degradation in gari during storage

333

334 Globally major and minor carotenoids compounds degraded during storage (Fig. 2). The  
335 degradation of the minor compounds was more difficult to model than that of major  
336 carotenoids because of the very low concentrations recorded. Degradation kinetics of 13-cis-  
337  $\beta$ -carotene content was globally similar to that of trans- $\beta$ -carotene and 9-cis- $\beta$ -carotene but  
338 slightly more irregular. According to Achir et al. (2013) 5,6 and 5,8 epoxy- $\beta$ -carotene were  
339 formed from cis-isomers on dried and stored sweet potato; therefore irregular pattern for  
340 epoxides may be explained by their formation that precedes their oxidation. Besides, 5,6 and  
341 5,8-epoxy-  $\beta$ -carotene were in very small concentrations, which made mathematical  
342 modelling difficult. Overall it appears that all minor carotenoids decreased quite sharply  
343 during storage and degradation was increased with temperature and storage duration.

344

345 Trans- $\beta$ -carotene and 9-cis- $\beta$ -carotene were both the main carotenoids present in BG and  
346 RPG (Fig. 2). Degradation of trans- $\beta$ -carotene followed logarithmic first order kinetics during  
347 storage between 19 and 40°C. Similarly, the degradation of 9-cis- $\beta$ -carotene also followed a  
348 first order kinetics equation.

349

350 Degradation rates (k) for major carotenoids are found in Table 2. Rates of carotenoid  
351 degradation were determined from the first order kinetics linear curves for BG and RPG  
352 stored in incubators. The higher the temperature and the longer the storage time, the greater  
353 was the trans- $\beta$ -carotene degradation and this was greatest for the gari made from yellow  
354 cassava (BG) and least for the gari made with palm oil (RPG). Coefficients of determination  
355 ( $R^2$ ) showed that ~~overall globally~~ the first order kinetics equation fitted carotenoid  
356 degradation well ( $R^2 \geq 0.8$ ). RPG at lower temperatures (*i.e.* 19°C) however did not fit the  
357 first order degradation as well as BG ( $R^2$  around 0.5-0.7) but the reasons are not clear.

358

359 Degradation rates (k) of trans- $\beta$ -carotene and 9-cis- $\beta$ -carotene clearly differ in BG and RPG  
360 (Table 2). ~~In addition, trans- $\alpha$  carotene was identified in significant concentrations in the  
361 palm oil gari samples (RPG) (Table 2). Trans- $\alpha$  carotene followed a first order kinetics and  
362 had close rates of degradation as those of trans  $\beta$  carotene and 9-cis  $\beta$  carotene in RPG.~~

363 Degradation rates of trans- $\beta$ -carotene and 9-cis- $\beta$ -carotene were greater for BG compared to  
364 RPG stored under the same conditions. Although the initial trans- $\beta$ -carotene content and the  
365 initial vitamin A activity was about twice higher in BG compared to RPG, trans- $\beta$ -carotene in  
366 BG degraded much faster.

367

368 More research would be needed to understand the difference in kinetics of carotenoid  
369 degradation in these two types of gari. The more complex matrix of palm oil gari including  
370 several different carotenoids having different types of kinetics might explain why the  
371 degradation of some minor carotenoids did not fit a first order degradation and why the major  
372 carotenoids (trans- $\beta$ ; 9-cis- $\beta$ -carotene ~~and trans- $\alpha$  carotene~~) did not fit well a first order  
373 degradation at lower temperature. Possibly the lower concentrations of carotenoids in palm

Formatted: Superscript



374 oil gari might also make the linearisation more difficult. Overall, the fitness of the models  
 375 demonstrates that mathematical modelling for the major carotenoids present is possible when  
 376 working with yellow cassava in spite of the relatively lower concentrations in cassava  
 377 compared to orange-fleshed sweet potato (Bechoff et al., 2010).

378

### 379 3.4. Model prediction of $\beta$ -carotene degradation in gari products

#### 380 3.4.1. Activation energy of $\beta$ -carotene degradation based on model prediction

381 The degradation of trans- $\beta$ -carotene was further modelled using an Arrhenius and Eyring  
 382 models (Table 3). The models fitted well trans and 9-cis- $\beta$ -carotene degradation in incubators  
 383 between 19 and 40°C for both BG and RPG gari preparations. Degradation kinetics of trans-  
 384  $\alpha$ -carotene and also more minor carotenoid compounds present in very small amounts in gari  
 385 (Fig. 2) could not be clearly described by the above models. Concentrations might be too  
 386 weak to be accurately predicted.

387

388 Activation energy for trans- $\beta$ -carotene in BG (~~6360.14~~  $6360.14 \text{ kJ.mol}^{-1}$ ) was of the same order as in  
 389 dried orange-fleshed sweet potato ( $64.2 \text{ kJ.mol}^{-1}$ ) during ambient storage (Bechoff et al.,  
 390 2010) though the carotenoid content in yellow cassava was much less than in orange-fleshed  
 391 sweet potato. ~~Interestingly, Eyring parameters—activation enthalpy and entropy being 60.5~~  
 392  ~~$\text{kJ.mol}^{-1}$  and  $77.1 \text{ J.mol}^{-1}$ , respectively in this study and  $61.7 \text{ kJ.mol}^{-1}$  and  $74.3 \text{ J.mol}^{-1}$ ,~~  
 393 ~~respectively in Bechoff et al. (2010) were also similar.~~

394

395 Activation energy ( $E_a$ ) was greater for RPG compared to BG (~~82.381.0~~ and ~~6360.14~~  $\text{kJ.mol}^{-1}$   
 396 for trans- $\beta$ -carotene, respectively, and ~~75.673.7~~ and ~~64.161.2~~  $\text{kJ.mol}^{-1}$  for 9-cis- $\beta$ -carotene,  
 397 respectively). It is therefore confirmed that the energy needed to degrade trans- $\beta$ -carotene

398 was higher in RPG. Palm oil might have coated trans- $\beta$ -carotene present in gari from white  
399 cassava and this might have limited degradation. Oxidation of fatty acids and carotenoids was  
400 similarly reported to be a free radical mechanism (Lieber 1993). It is therefore possible that  
401 fatty acids present in palm oil might have been oxidised in the place of trans- $\beta$ -carotene and  
402 hence acted as a protection against carotenoid degradation. Achir *et al.* (2010) reported that  
403 Ea of trans- $\beta$ -carotene in pure palm oil was comparable, being 86 kJ.mol<sup>-1</sup>, and this was in  
404 accordance with this present study. More research shall be required to understand how  
405 different matrices (i.e. different oil types; vegetable cells) can affect activation energy of  
406 carotenoids.

#### 407 3.4.2. Model prediction of retention during storage of gari

408

409 Relationships between storage time, storage temperature and predicted retention of trans- $\beta$ -  
410 carotene are described in Fig. 3. Predicted nutritional values for gari (vitamin A activity)  
411 were also calculated. According to the Arrhenius model predictions, if BG with an initial  
412 nutritional value of 301.4 RAE (based on the new conversion factor) was stored for 60 days  
413 (2 months) at an ambient temperature of 25°C and constant humidity, about 32.44% of the  
414 initial trans- $\beta$ -carotene would be retained (equivalent to 135.136 RAE in the product).

415 Periods of storage are 5-6 months on average under tropical temperatures (about 25°C in the  
416 daytime) in sub-Saharan Africa (i.e. Nigeria). If BG was stored for 5 months at 25°C, only  
417 12.13% of trans- $\beta$ -carotene (equivalent to 41 RAE) would be retained and if it was stored for  
418 6 months, about 8% of trans- $\beta$ -carotene would be retained (equivalent to 27.28 RAE) and the  
419 nutritional value would be negligible. If the same gari was stored at a lower temperature of  
420 20°C about 25.26% would be preserved after 5 months leading to a nutritional value of about  
421 79 RAE. If the initial carotenoid content were the same as for BG (with further addition of

422 | palm oil), RPG could be stored three times longer, for up to 50 days according to the  
423 | Arrhenius model predictions. Hence the traditional practices of storage for gari in Nigeria  
424 | should not be recommended when working with BG. One option may be to lower storage  
425 | temperature but this would require facilities such as a fridge or freezer.

426 |  
427 | On the other hand, RPG with an initial nutritional value of 203.5 RAE would retain about  
428 | 80% of the initial trans- $\beta$ -carotene after 60 days (2 months) at an ambient temperature of  
429 | 25°C and constant humidity (corresponding to 142 RAE). About ~~50.57~~ % would be retained if  
430 | the same gari was stored for 5 months at 25°C (about ~~8.32~~ RAE). Trans- $\beta$ -carotene contained  
431 | in RPG was therefore more stable during storage. ~~If the initial carotenoid content were the~~  
432 | ~~same as for BG (with further addition of palm oil), RPG could be stored three times longer,~~  
433 | ~~for up to 50 days according to the Arrhenius model predictions.~~

434 |  
435 | While provitamin A was more stable in RPG, the presence of palm oil created quality issues  
436 | such as rancidity (Abu *et al.* 2006; Burri *et al.*, 2012): Abu *et al.* (2006) reported that whilst  
437 | half of the initial carotenoid content was lost during a 2 month-ambient storage there was a  
438 | concomitant increase in the peroxide index (six times more than the initial value) of the gari  
439 | product.

#### 440 | 441 | 442 | **4. Conclusions**

443 |  
444 | The effect of temperature on carotenoid stability was measured in the two types of gari; from  
445 | yellow cassava (BG) and from white cassava with palm oil (RPG). Carotenoid content was  
446 | temperature and storage time sensitive. Trans- and 9-cis- $\beta$ -carotene contents in BG or RPG

447 followed a first order (logarithmic) degradation equation during storage. Although the initial  
448 content of trans- $\beta$ -carotene and the initial vitamin A activity was about twice higher in BG  
449 compared to RPG, trans- $\beta$ -carotene BG degraded much faster. Trans- and 9-cis- $\beta$ -carotene  
450 degradation in BG and RPG was accurately described by the Arrhenius and Eyring models.  
451 The mathematical model can therefore be used to predict storage times at various storage  
452 temperatures. The addition of red palm oil significantly increased the shelf life of gari in  
453 terms of carotenoid retention: fatty acids present in palm may have protected carotenoids  
454 against degradation. More research would be needed to understand the role of oil in  
455 preserving carotenoids in gari over time. Although gari made with added palm oil could be  
456 stored longer, issues of oil rancidity of the latter within the typical market turn-around time  
457 for such product will need to be explored. In conclusion, this work has proven that gari made  
458 with biofortified cassava has a limited carotenoid stability under ambient temperature  
459 conditions and breeders, processors and marketers or supporters should be aware of this  
460 constraint.

461

#### 462 **Acknowledgments**

463

464 This research was supported by HarvestPlus. The views expressed are however those of the  
465 authors. We wish to thank HarvestPlus staff in Nigeria who helped with monitoring the  
466 processing of gari product: Mary Okocha, Joy Okoh, Christiana Okon Essien and trainees  
467 Osanyinro Oluwatosin, Onyiba Cosmos Ifeanyi, Tobi Olaniyi and Bose and the workers at  
468 Barrack Centre who processed the gari. We would like to thank Dudley Farman, (NRI) for  
469 carrying out the HPLC analyses at NRI. We express our gratitude to [Dr. Corinne Rumney](#),  
470 [\(NRI\)](#) for helping with the sample collection. [We also would like to thank Dr. Stephen Young](#)  
471 [\(NRI\) for his help and insights in statistical analysis.](#)

472

473 **References**

474

475 Abu, J.O., Badifu, G.I.O., Akpapunam, M.A., 2006. Effect of crude palm oil inclusion on  
476 some physic-chemical properties of gari – a fermented cassava food product. *Nigerian Food*  
477 *Journal* 24, 73-79.

478

479 Achir, N., Randrianatoandro, V. A., Bohuon, P., Laffargue, A., & Avallone, S., 2010. Kinetic  
480 study of  $\beta$ -carotene and lutein degradation in oils during heat treatment. *European Journal of*  
481 *Lipid Science and Technology* 112, 349-361.

482

483 Achir, N., Pénicaud, C., Bechoff, A., Boulanger, R., Dornier, M., Dhuique-Mayer, C., 2013.  
484 Use of Multi-response Modelling to investigate mechanisms of  $\beta$ -carotene degradation in  
485 dried orange-fleshed sweet potato during storage: from carotenoids to aroma compounds.  
486 *Food and Bioprocess Technology* 1-14.

487

488 Andreu-Sevilla, A.J. Hartmann, A. Burló, F. Poquet, N., Carbonell-Barrachina, A.A., 2009.  
489 Health benefits of using red palm oil in deep-frying potatoes: low acrolein emissions and high  
490 intake of carotenoids. *Food Science and Technology International* 15, 15-20.

491

492 Bechoff, A. (2010). Investigating carotenoid losses after drying and storage of orange-fleshed  
493 sweet potato. PhD thesis, University of Greenwich, UK.

494

495 Bechoff, A., Dhuique-Mayer, C., Dornier, M., Tomlins, K., Boulanger, R., Dufour, D. ,  
496 Westby, A., 2010. Relationship between the kinetics of  $\beta$ -carotene degradation and

497 norisoprenoid formation in the storage of dried sweet potato chips. Food Chemistry 121, 348–  
498 357.

499

500 Bechoff, A., Tomlins, K.I., Dhuique-Mayer, C., Dove, R., Westby, A. , 2011a. On-farm  
501 evaluation of the impact of drying and subsequent storage on the carotenoid content of  
502 orange-fleshed sweet potato. International Journal of Food Science & Technology, 46, 52-60.

503

504 Bechoff, A. Poulaert, M., Tomlins, K.I., Westby, A., Menya, G., Young S., Dhuique-Mayer,  
505 C., 2011b. Retention and bioaccessibility of  $\beta$ -carotene in blended food from sub-Saharan  
506 Africa containing orange-fleshed sweet potato. Journal of Agricultural & Food Chemistry 59,  
507 10373-10380.

508

509 Bonnie Tay, Y. P., & Choo, Y. M., 2000. Practical guide to establishing palm carotenoids  
510 profiles by HPLC with three-dimensional diode array detector. Palm Oil Developments 33,  
511 13-17.

512 Burri, B.J., 2012. Evaluating global barriers to the use of red palm oil as an intervention food  
513 to prevent vitamin A deficiency. Comprehensive Reviews in Food Science and Food Safety  
514 11(2), 221-232.

515

516 Chavez, A.L., Sanchez, T., Ceballos, H., Rodriguez-Amaya, D.B., Nestel, P., Tohme, J.,  
517 Ishitani, M., 2007. Retention of carotenoids in cassava roots submitted to different processing  
518 methods. Journal of the Science of Food and Agriculture 87, 388-393.

519

Formatted: English (U.K.)

520 Failla, M.L., Chitchumroonchokchai, C., Siritunga, D., De Moura, F., Fregene, M., Sayre,  
521 R.T., 2012. Retention during processing and bioaccessibility of  $\beta$ -carotene in high  $\beta$ -carotene  
522 transgenic cassava root. *Journal of Agricultural & Food Chemistry* 60, 3861–3866.

523

524 FAO/WHO, 2002. Vitamin A. In *Human Vitamin and Mineral Requirements. Report of joint*  
525 *FAO/WHO expert consultation: Bangkok, Thailand (pp. 87-107)*. Rome, Italy: Food and  
526 Nutrition Division FAO.

527

528 Gouado, I., Demasse Mawamba, A., Meyimbo Ouambo, R. S., Some, I. T., Félicité, T. M.,  
529 2008. Provitamin A carotenoid content of dried fermented cassava flour: The effect of palm  
530 oil addition during processing. *International Journal of Food Engineering* 4(4) 1556-3758.

531

532 National Academy of Sciences/Institute of Medicine, 2001. Vitamin A. In *Dietary Reference*  
533 *Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron,*  
534 *Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc*. Washington, DC. National  
535 Academy Press.

536

537 La Frano, M. R., Woodhouse, L. R., Burnett, D. J., Burri, B. J., 2013. Biofortified cassava  
538 increases  $\beta$ -carotene and vitamin A concentrations in the TAG-rich plasma layer of American  
539 women. *British Journal of Nutrition* 110, 310-320.

540

541 La Frano, M., Zhu, C., Burri, B., 2014. Effects of processing, cooking, and storage on  $\beta$ -  
542 carotene retention and bioaccessibility in biofortified cassava (*Manihot esculenta*) (646.4).  
543 *The FASEB Journal* 28 (1 Supplement), 646-4.

544

545 Liu, W., Zhou, Y., Sanchez, T., Ceballos, H., & White, W. S., 2010. The vitamin A  
546 equivalence of beta-carotene in beta-carotene-biofortified cassava ingested by women. The  
547 FASEB Journal, 24 (meeting abstract supplement), abstract No. 92.7  
548

549 Lieber, D.C., 1993. Antioxidant reactions of carotenoids. In L.M. Canfield, & J.A., Olson  
550 (Eds), Carotenoids in Human Health. (pp. 20-31). New York, NY, USA: The New York  
551 Academy of Sciences.  
552

553 Maziya-Dixon, B., Dixon, A.G.O., Ssemakula, G., 2009. Changes in total carotenoid content  
554 at different stages of traditional processing of yellow-fleshed cassava genotypes. International  
555 Journal of Food Science & Technology 44, 2350-2357.  
556

557 Mortensen, A., 2005. Analysis of a complex mixture of carotenes from oil palm (*Elaeis*  
558 *guineensis*) fruit extract. Food Research International, 38, 847–853.  
559

560 Mugode, L., Ha, B., Kaunda, A., Sikombe, T., Phiri, S., Mutale, R., De Moura, F. F., 2014.  
561 Carotenoid retention of biofortified provitamin A maize (*Zea mays* L.) after Zambian  
562 traditional methods of milling, cooking and storage. Journal of Agricultural and Food  
563 Chemistry (In press).  
564

565 National Academy of Sciences / Institute of Medicine, 2001. Vitamin A. In Dietary  
566 Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine,  
567 Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc. Washington, DC,  
568 USA: National Academy Press.  
569



570 Oshikoya, T. W. 2008. Nigeria in the global economy. *Business Economics*, 43(1), 31-43.

571

572 Onadipe Olapeju, Olubusola, 2011. Total carotenoid content, retention, bioavailability and  
573 consumer acceptability of gari from bio-fortified cassava roots. PhD thesis, University of  
574 Agriculture, Abeokuta, Nigeria.

575

576 Phorbee, O. O., Olayiwola, I. O., Sanni, S.A., 2013. Bioavailability of Beta Carotene in  
577 Traditional Fermented, Roasted Granules, Gari from Bio-Fortified Cassava Roots. *Food and*  
578 *Nutrition Sciences* 4(12), 1247.

579

580 Rodriguez Amaya, D.B. & Kimura, M., 2004. HarvestPlus handbook for carotenoid analysis.  
581 In HarvestPlus Technical monograph series 2. HarvestPlus handbook for carotenoid analysis  
582 (pp. 1-58) Washington DC, USA: HarvestPlus c/o International Food Policy Research  
583 Institute. [Online] <http://www.harvestplus.org/pdfs/tech02.pdf>

Field Code Changed

584

585 Saltzman, A., Birol, E., Bouis, H. E., Boy, E., De Moura, F. F., Islam, Y., Pfeiffer, W. H. ,  
586 2013. Biofortification: progress toward a more nourishing future. *Global Food Security* 2(1),  
587 9-17.

588

589 Thakkar, S. K., Huo, T., Maziya-Dixon, B., Failla, M. L., 2009. Impact of style of processing  
590 on retention and bioaccessibility of  $\beta$ -Carotene in Cassava (*Manihot esculanta*, Crantz).  
591 *Journal of Agricultural and Food Chemistry* 57(4), 1344-1348.

592

593 Ukenye, E., Ukpabi, U. J., Chijoke, U., Egesi, C., Njoku, S., 2013. Physicochemical,  
594 nutritional and processing properties of promising newly bred white and yellow fleshed  
595 cassava genotypes in Nigeria. *Pakistan Journal of Nutrition*, 12(3), 302-305.  
596  
597 Uzomah, A., Ubbaonu, C. N., Osuji, C. M., 2006. Vitamin A retention in palm oil-gari during  
598 processing and storage. *Nigerian Food Journal* 23(1), 69-73.  
599  
600 Westby, A. 2002. Cassava utilization, storage and small-scale processing. In R.J. Hillocks,  
601 J.M. Thresh & A.C. Bellotti (Eds.), *Cassava biology, production and utilization* (pp 281-300).  
602 Oxon: CABI Publishing.  
603  
604  
605

Accepted Manuscript

**Table 1.** Estimation of vitamin A activity ( $\mu\text{g}$  retinol equivalent) for a 100g portion of gari made from yellow biofortified cassava (BG) and from white cassava with palm oil (RPG) ( $\mu\text{g}$  RAE<sup>a</sup>) during storage and maintained at constant humidity ( $a_w = 0.5$  for NaBr)

RAE <sup>a</sup> estimate	Storage days	5:1 <sup>b</sup>		12:1 <sup>b</sup>	
		BG	RPG	BG	RPG
T (°C)	Initial	301.45±7.36	203.49±5.886	425.61260±3.07	84.7985±2.452
19	24	256.5±11.4	208.9209±10.811	406.9107±4.75	87.0±4.5
	49	205.2±5.45	161.5±24.0	85.5±2.2	67.3±10.0
	60	199.7200±7.3	174.8175±12.713	83.2±3.0	72.873±5.35
	80	155.5156±9.4	139.8140±18.2	64.865±3.84	58.2±7.68
26	18	244.8245±23.523	181.4±8.2	102.0±9.810	75.4±3.4
	24	250.5±25.726	190.3±10.911	104.4±10.711	79.3±4.5
	49	150.3±29.2	135.4±7.0	62.663±12.2	56.3±2.93
	60	129.8130±20.4	154.0±8.79	54.1±8.4	64.2±3.64
33	80	99.2±15.4	130.4±12.3	41.3±6.3	54.2±5.4
	10	200.8201±14.4	147.6148±15.716	83.784±5.96	61.5±6.5
	18	203.9204±11.612	199.6200±5.76	85.0±4.85	83.2±2.4
	24	172.4±14.815	188.7189±8.2	71.972±6.2	78.679±3.4
	49	87.4±6.77	137.3±9.910	36.4±2.83	57.2±4.4
	60	58.3±1.82	123.3±5.86	24.3±0.81	51.4±2.4
40	80	40.841±2.83	92.993±3.84	17.0±1.2	38.739±1.62
	10	147.8148±25.926	164.0±4.75	61.662±10.811	68.3±2.0
	18	139.6140±7.4	182.7183±5.2	58.2±3.0	76.4±2.2
	24	99.4±3.4	157.0±9.5	41.4±1.4	65.4±3.94
	31	67.668±1.72	143.7144±6.67	28.1±0.71	59.960±2.73
	49	28.629±1.52	99.3±3.5	11.912±0.61	41.4±1.5
	60	20.621±0.91	80.1±6.67	8.69±0.4	33.4±2.83
	80	12.5±0.2	42.3±0.61	5.2±0.1	17.618±0.3

Mean of triplicate determinations ± standard deviation. <sup>a</sup> Retinol Activity Equivalent. <sup>b</sup> RAE was calculated for a bioconversion factor of 5:1 (La Frano et al., 2013) estimate = {All-trans- $\beta$ -carotene content / 5 + minor  $\beta$ -carotene content / 10} x unit (g) or for a bioconversion factor of 12:1 (National Academy of Sciences / Institute of Medicine, 2001). Classical estimate = {All-trans- $\beta$ -carotene content / 12 + minor  $\beta$ -carotene content / 24} x unit (g). Minor compounds are epoxy and cis  $\beta$ -carotene that are estimated to possess half of trans  $\beta$ -carotene activity.

Formatted Table

**Table 2.** Carotenoid rate of degradation ( $k$ )<sup>a</sup> in gari made from biofortified cassava (BG) (A) and from white cassava with palm oil (RPG)

Gari	Temperature (°C)		19	26	33	40
BG	Trans- $\beta$ -carotene	k	0.0083 $\pm$ 0.0004	0.01580144 $\pm$ 0.00070011	0.02760271 $\pm$ 0.00100008	0.04790430 $\pm$ 0.00140013
		R <sup>2</sup>	0.979 <del><math>\pm</math>0.009</del>	0.948900 <del><math>\pm</math>0.041</del>	0.990984 <del><math>\pm</math>0.013</del>	0.987981 <del><math>\pm</math>0.014</del>
		p	<0.0001*	<0.0001*	<0.0001*	<0.0001*
BG	9-cis- $\beta$ -carotene	k	0.0071 $\pm$ 0.00050004	0.01250123 $\pm$ 0.00060010	0.02340230 $\pm$ 0.00100008	0.04290.0375 $\pm$ 0.00160010
		R <sup>2</sup>	0.968 <del><math>\pm</math>0.0190.951</del>	0.990 <del><math>\pm</math>0.0540.887</del>	0.984 <del><math>\pm</math>0.0140.980</del>	0.991 <del><math>\pm</math>0.0090.983</del>
		p	<0.0001*	<0.0001*	<0.0001*	<0.0001*
RPG	Trans- $\beta$ -carotene	k	0.00270020 $\pm$ 0.00110006	0.00390040 $\pm$ 0.00110006	0.00890086 $\pm$ 0.00130006	0.01840189 $\pm$ 0.00030011
		R <sup>2</sup>	0.634 <del><math>\pm</math>0.2820.589</del>	0.869 <del><math>\pm</math>0.0430.769</del>	0.896 <del><math>\pm</math>0.0790.939</del>	0.934 <del><math>\pm</math>0.0200.930</del>
		p	0.016*	<0.0001*	<0.0001*	<0.0001*
RPG	9-cis- $\beta$ -carotene	k	0.00300023 $\pm$ 0.00140008	0.00400038 $\pm$ 0.00100008	0.00850080 $\pm$ 0.00100005	0.0170 $\pm$ 0.00030010
		R <sup>2</sup>	0.679 <del><math>\pm</math>0.2260.516</del>	0.837 <del><math>\pm</math>0.0580.641</del>	0.910 <del><math>\pm</math>0.0610.937</del>	0.935 <del><math>\pm</math>0.0180.926</del>
		p	0.029*	0.0001*	<0.0001*	<0.0001*
RPG	Trans- $\alpha$ -carotene	k	0.0037 $\pm$ 0.0010	0.0049 $\pm$ 0.0009	0.0084 $\pm$ 0.0009	0.0132 $\pm$ 0.0006
		R <sup>2</sup>	0.761 <del><math>\pm</math>0.357</del>	0.883 <del><math>\pm</math>0.120</del>	0.911 <del><math>\pm</math>0.103</del>	0.918 <del><math>\pm</math>0.001</del>

<sup>a</sup> expressed in day<sup>-1</sup> at storage temperatures on a fresh weight basis and maintained at constant humidity ( $a_w = 0.5$  for NaBr)

R<sup>2</sup>: Coefficient of determination. Mean of triplicate determinations  $\pm$  standard deviation/error (linear regression; XLSTAT 2014). Standard error is standard deviation divided by the square root of the number of analyses.

k was the slope on the logarithmic carotenoid concentration (Y-axis) vs storage time (X-axis) graph and was obtained by linear regression.

First order equation:

$$\ln C = \ln C_0 - kt$$

Where: C: Carotenoid content of gari ( $\mu\text{g.g}^{-1}$ ) at storage time t; C<sub>0</sub>: Carotenoid content of food ( $\mu\text{g.g}^{-1}$ ) at initial time (before storage); t: storage time (day); k: degradation constant rate (day<sup>-1</sup>).

\* indicate a significant correlation at  $p < 0.05$  (linear regression; XLSTAT 2014).

Formatted Table

Formatted: Not Highlight

Formatted: Superscript

**Table 3.** Parameters of the Arrhenius<sup>a</sup> and Eyring<sup>b</sup> models for the carotenoids degradation in gari on a fresh weight basis between 19-40°C and maintained at constant humidity ( $a_w = 0.5$  for NaBr) in gari made from yellow cassava (BG) (A) and from white cassava with palm oil (RPG)

Carotenoid	Parameter	Type of gari		p
		BG	RPG	
Trans- $\beta$ -carotene	$\ln k_\infty$ (days <sup>-1</sup> )	20.1±0.9	27.1±1.2	0.039*
	$E_a$ (kJ.mol <sup>-1</sup> )	60.4±2.1	81.0±3.1	
	$R^2$	0.998	0.997	
9-cis- $\beta$ -carotene	$\ln k_\infty$ (days <sup>-1</sup> )	20.2±0.6	24.2±2.1	0.034*
	$E_a$ (kJ.mol <sup>-1</sup> )	61.2±1.6	73.7±5.2	
	$R^2$	0.999	0.988	

<sup>a</sup>Arrhenius model

$$k = k_\infty e^{-\frac{E_a}{RT}}$$

Where: T : temperature (K); k: degradation rate constant at T (day<sup>-1</sup>);  $k_\infty$ : value of k at T =  $\infty$  (day<sup>-1</sup>);  $E_a$ : Activation energy (kJ.mol<sup>-1</sup>); R: gas constant = 8.314 J · K<sup>-1</sup> · mol<sup>-1</sup>

<sup>b</sup>Eyring model

$$k = \frac{k_B}{h} T \cdot e^{-\frac{\Delta H^\ddagger - T\Delta S^\ddagger}{RT}}$$

Where:  $k_B$ : Boltzmann constant = 1.381 · 10<sup>-23</sup> J.K<sup>-1</sup>; h: Planck constant = 6.626 · 10<sup>-34</sup> J.s;  $\Delta H^\ddagger$ : activation enthalpy (kJ.mol<sup>-1</sup>);  $\Delta S^\ddagger$ : activation entropy (J.mol<sup>-1</sup>.K<sup>-1</sup>)


$R^2$ : Coefficient of determination.


Mean of triplicate determinations ± standard deviation error (linear regression: XLSTAT 2014).

Yellow cassava 01/1371 (YCBC); White cassava IITA 3303 with added palm oil (PWCRPG).

\* indicate a significant difference between BG and RPG at p<0.05 (One Way ANOVA T-test).

Formatted Table






**Fig. 1.** HPLC Chromatogram<sup>a</sup> of gari from TMS 01/1371 biofortified variety (BG) (A) **Fig. 1B.** and HPLC Chromatogram<sup>a</sup> of Gari-gari from IITA 3303 variety with palm oil (RPG) (B)<sup>a</sup> Identification of minor carotenoids (*i.e.* 5,6 and 5,8 epoxy-β-carotene) is tentative. Alpha-carotene was identified by

Formatted: Not Superscript/ Subscript

Formatted: Font: Not Bold







injection of a mixture of  $\alpha$  and  $\beta$  carotenes.

Formatted: Left

**Fig. 2.** Carotenoid content (on a fresh weight basis (FW)) of gari made from yellow biofortified cassava (BG) (A) or white cassava with red palm oil (RPG) (B) ( $\mu\text{g}\cdot\text{g}^{-1}$ ) and stored at four different temperatures (—) and constant humidity ( $a_w = 0.5$ ) for NaBr) over 80 days.  
Mean of triplicate determination

**Fig. 2.** Carotenoid content (on a fresh weight basis (FW)) of gari made from yellow biofortified cassava (BG) (A) or white cassava with red palm oil (RPG) (B) (expressed in  $\mu\text{g}\cdot\text{g}^{-1}$  trans- $\beta$ -carotene<sup>a</sup>) and stored at four different temperatures<sup>b</sup> (—) and constant humidity ( $a_w = 0.5$  for NaBr) over 80 days. Mean of triplicate determination.



<sup>a</sup> Concentration for all the carotenoids was determined from standard curve of pure trans- $\beta$ -carotene (Sigma, UK).  
<sup>b</sup>

Formatted: Width: 29.7 cm, Height: 21 cm

Formatted: Superscript

Formatted: Font: Not Bold

**Fig. 3.** Relationships between storage time, temperature, trans- $\beta$ -carotene true retention (TR) and vitamin A activity<sup>a</sup> of gari made from yellow cassava (BG) (A) and from white cassava with palm oil (RPG) (B) based on Arrhenius model predictions.


<sup>a</sup> Retinol Activity Equivalent (RAE) estimate-estimate was calculated for a bioconversion factor of 5:1 (La Frano et al., 2013)  $\text{RAE} = \{\text{all-trans-}\beta\text{-carotene content}\}$

A

/5 + minor  $\beta$ -carotene content/ 10} x unit (g).

Accepted Manuscript

Accepted Manuscript



**Highlights**

- Trans- $\beta$ -carotene degradation was modelled during storage of BG and RPG
- Trans and 9-cis  $\beta$ -carotene degradation fitted well the Arrhenius and Eyring models
- Initial trans- $\beta$ -carotene was twice as high in the BG compared to RPG
- Activation energy was lower (~20%) with BG compared to RPG
- Based on carotenoid stability, shelf life of BG was much shorter than RPG

Accepted Manuscript