



**Greenwich Academic Literature Archive (GALA)**  
– the University of Greenwich open access repository  
<http://gala.gre.ac.uk>

---

*Citation for published version:*

Bianco, Antonino, Thomas, Ewan, Bellafiore, Marianna, Martines, Francesco, Messina, Giuseppe, Battaglia, Giuseppe, Karsten, Bettina, Sahin, Fatma, Bielec, Grzegorz, Paoli, Antonio and Palma, Antonio (2015) Mediterranean diet and dietary supplementation as possible predicting variables of weight management: an update of the protein project. *Acta Medica Mediterranea*, 31:907. pp. 1256-1270. ISSN 0393-6384 (In Press)

*Publisher's version available at:*

---

**Please note that where the full text version provided on GALA is not the final published version, the version made available will be the most up-to-date full-text (post-print) version as provided by the author(s). Where possible, or if citing, it is recommended that the publisher's (definitive) version be consulted to ensure any subsequent changes to the text are noted.**

*Citation for this version held on GALA:*

Bianco, Antonino, Thomas, Ewan, Bellafiore, Marianna, Martines, Francesco, Messina, Giuseppe, Battaglia, Giuseppe, Karsten, Bettina, Sahin, Fatma, Bielec, Grzegorz, Paoli, Antonio and Palma, Antonio (2015) Mediterranean diet and dietary supplementation as possible predicting variables of weight management: an update of the protein project. London: Greenwich Academic Literature Archive.

Available at: <http://gala.gre.ac.uk/13996/>

---

Contact: [gala@gre.ac.uk](mailto:gala@gre.ac.uk)

## MEDITERRANEAN DIET AND DIETARY PROTEIN SUPPLEMENTATION AS POSSIBLE PREDICTING VARIABLES OF WEIGHT MANAGEMENT: AN UPDATE OF THE PROTEIN PROJECT

ANTONINO BIANCO<sup>1</sup>, EWAN THOMAS<sup>1</sup>, MARIANNA BELLAFFIORE<sup>1</sup>, FRANCESCO MARTINES<sup>2</sup>, GIUSEPPE MESSINA<sup>1</sup>, GIUSEPPE BATTAGLIA<sup>1</sup>, BETTINA KARSTEN<sup>3</sup>, FATMA NESE SAHIN<sup>4</sup>, GRZEGORZ BIELEC<sup>5</sup>, ANTONIO PAOLI<sup>6</sup>, ANTONIO PALMA<sup>1</sup>

<sup>1</sup>Sport and Exercise Research Unit, University of Palermo, Italy - <sup>2</sup>BioNeC Department, Audiology Section, University of Palermo - <sup>3</sup>Department of Life and Sports Sciences, University of Greenwich, United Kingdom - <sup>4</sup>Faculty of Sport Science, Ankara University, Turkey - <sup>5</sup>Department of Environmental Sciences, Warmia and Mazury University in Olsztyn, Poland - <sup>6</sup>Department of Biomedical Sciences, University of Padova, Italy

### ABSTRACT

**Introduction:** The Mediterranean diet is known to support the prevention of several chronic diseases and excessive weight gains. However, relevant investigations were performed in a clinical setting. The aim of this study was therefore to understand the effects of protein supplements intake on body mass index (BMI) in healthy active individuals following the Mediterranean diet.

**Materials and methods:** A face-to-face questionnaire was administered to 667 subjects, 627 living in a Mediterranean area (MD) and 40 in a non-Mediterranean area (NMD); questionnaire enquired dietary behaviors, including a comparison between dietary patterns and protein consumption between these two populations and within each population.

**Results:** Dietary patterns significantly varied between the MD and NMD populations ( $p < 0.001$ ), although, BMI was not significantly different. The major significant differences were found between the BMI of protein supplement users and non-protein supplement users within and between the two populations ( $p < 0.01$ ). No differences were found between the BMI of the protein supplement users of MD and NMD, while moderate differences were displayed between the non-supplement users of the two populations ( $p < 0.05$ ).

**Conclusions:** The Mediterranean diet does not appear to affect BMI. However, BMI significantly differed between protein supplement users and non-protein supplement users indicating a direct influence of such products on body weight and consequently BMI.

**Key words:** Protein project, questionnaire, supplements, dietary patterns, health.

Received January 30, 2014; Accepted March 30, 2015

### Introduction

The traditional Mediterranean diet (MD) has been defined as that dietary pattern adopted by the inhabitants of the Mediterranean area before the 1960s<sup>(1)</sup>. Such dietary regimen is characterized by a high consumption of vegetables, legumes, fruits, nuts, cereals, olive oil and fish with moderate consumption of dairy products and low intakes of meats<sup>(2)</sup>.

This particular nutritional pattern was first adopted due to the poor economic status of the Mediterranean population, as meat and poultry was too cost expensive unlike locally grown vegetables and fruits<sup>(3)</sup>. The mean macronutrient composition of this kind of diet has a moderate to high intake of lipids (30-40%), with a high prevalence of mono- and low saturated fats<sup>(4)</sup>. Carbohydrates (CHO) contribute between 40 and 55% to the caloric intake and mainly originate from complex CHOs, which show a

relatively low glycemic load. The remaining proportion is covered by protein intake<sup>(5)</sup>. Over the last decades the MD has been studied mainly under clinical and medical conditions and it was found to be effective in reducing: 1) morbidity and mortality of cardiovascular diseases, 2) peripheral artery diseases, 3) the occurrence of the metabolic syndrome and 4) obesity<sup>(6-8)</sup>. With particular regards to the effectiveness of such dietary regimen for weight loss, various authors such as Greco et al<sup>(9)</sup>, Grosso et al<sup>(10)</sup> and Bekkouche et al<sup>(11)</sup> stated that after a period of adherence to the MD, body weight (BW) and consequently Body Mass Index (BMI), significantly decreased. However, today there is inconsistent evidence about the application of these findings in healthy physically active individuals. Another factor that is known to influence BW is protein supplementation. A number of authors state a reduction in BW after the consumption of such nutritional components<sup>(12-15)</sup>.

Therefore the aim of the present study was to understand whether and to what extent the MD and protein supplement intake influence values of BMI in two populations of healthy and physically active subjects, one who adhered to a Mediterranean dietary pattern and one who followed a different dietary pattern.

## Materials and methods

### *Participants*

Permissions to conduct this survey were obtained from the managers of a representative number of commercial gyms located in Italy (Palermo and Pavia) and in Poland (Danzica). In Italy, suitable gyms were identified using a database of the CONI register (National Olympic Committee Register for Sport and Fitness Associations) Gyms in Poland were identified randomly through a sealed envelope method. To reduce heterogeneity and number of aerobic activities only gym attendees taking part in strength training (Gym, functional fitness, weightlifting, etc....) were included for this investigation. Gym users performing aerobic activities were consequently excluded. Based on these inclusion/exclusion criteria, 667 participants were retained; 627 from the MD<sup>(18-20)</sup> and 40 from NMD.

Subjects' ages ranged from 13 to 68 years (table 1). The gender-split resulted in 502 male and 165 female subjects out of which 189 male and 38 female subjects declared to use protein supplements.

### *Questionnaire procedure*

As in our previous works<sup>(18, 19, 21)</sup>, the face to face interview method was adopted<sup>(22)</sup> to evaluate the frequency consumption of protein supplements amongst participants, dietary behaviors and other related information. Easy understandable definitions of the supplements (common and commercial names of products or substances included within the definition of supplement: product intended to supplement the diet that contains one or more dietary ingredients<sup>(23)</sup>) and foods were provided to participants. The completion of the questionnaire indicated the agreement of the gym user to participate in the study. According to the Italian and Polish regulations, ethical approval was not required for this study. The same investigator for each area administered the questionnaire over a period of ten months.

### *Definition procedure*

According to Trichopoulou et al<sup>(4)</sup> the MD can be defined as a high prevalence of vegetables, fruits and nuts, legumes, and unprocessed cereals intake and a low intake of meat and meat products and dairy products (with the exception of cheese. Total intake of lipids can be high (around 40% of total energy intake, as in Greece), or moderate (around 30% of total energy intake, as in Italy). However, the overall ratio of beneficial mon-unsaturated to non-beneficial saturated lipids is high (due to the high mono unsaturated content of used olive oil). MD can also be expressed through the use of a pyramid such as that of Willet et al<sup>(24)</sup>. The non-Mediterranean diet presented in this study is typical for Central and Eastern Europe. It is characterized by consumption of sugar and saturated fatty acids as well as a low intake of fruits and vegetables<sup>(25-27)</sup>.

### *Data analysis*

Data analysis was performed using the EpiInfo software version 7.0 (CDC, Atlanta, GA, US) and the STATISTICA software 8.0 software for Windows (Tulsa, OK, US). The descriptive analysis was performed by calculating the means and standard deviations (SD). Frequency of weekly food intake was assessed and the chi-square test was performed in order to verify differences between dietary patterns of both populations; subsequently differences between BMI of MD and NMD groups were assessed through an unpaired t-test. The comparisons were made between groups and within groups, stratifying each sample for protein supple-

ment and non-protein supplement users. Statistical significance was set at a P value < 0.05.

## Results

### Participants

The anthropometric characteristics of participants are described in table 1.

	Age (years)	Height (Cm)	Weight (Kg)	BMI	Gender (M/F)
Mediterranean	28±9.9	174±9	72.5±12.6	23.9±3.1	480/147
Non-Mediterranean	29±7.1	175±9	74.1±16.1	24.1±3.5	22/18
<i>p</i>	Ns	Ns	Ns	Ns	n/a

**Table 1:** Anthropometric characteristics of participants. *Ns* = Not significant, *n/a* = Not applicable.

### Dietary patterns

Dietary patterns significantly varied between the analyzed populations. MD population declared a high consumption of MD key foods<sup>(3)</sup> (3 times per week or more) whereas in the NMD population an inverse trend was evident. Although, no differences in BMI are shown between the MD group and the NMD group, respectively.

Frequency per week	0	1	2	3	4	5	6	7
<b>Foods</b>								
<i>Bakery</i>	33.41	13.1	11.14	8.73	4.59	22.05	0.66	6.33
<i>Chicken</i>	3.28	16.38	27.07	27.07	11.35	11.14	0.87	2.84
<i>Cheese</i>	15.5	11.57	21.83	21.4	9.17	14.19	0.87	5.46
<i>Cold cuts</i>	23.58	20.09	19.87	15.07	8.08	9.17	1.53	2.62
<i>Fresh Fish</i>	11.57	34.5	31.88	13.97	5.46	1.97	0.66	0
<i>Legumes</i>	15.28	23.58	26.42	18.56	8.3	7.21	0	0.66
<i>Meat</i>	5.02	13.32	27.51	30.13	10.26	9.17	2.18	2.4
<i>Milk</i>	22.27	5.9	5.46	8.52	2.84	34.5	1.31	19.21
<i>Nuts</i>	63.54	15.5	9.83	4.59	1.09	3.93	0.22	1.31
<i>Canned Tuna</i>	30.13	26.64	21.62	10.7	4.59	4.8	0.44	1.09
<i>Eggs</i>	17.25	33.84	28.6	11.79	2.62	3.93	0	1.97
<i>Yogurt</i>	40.83	9.61	11.35	11.79	5.02	13.97	0.87	6.55

**Table 2:** Percentage of food consumption of the Mediterranean area (n=627).

### Protein consumption

A number of 201 subjects from the MD sample and 26 from the NMD sample declared to use protein supplements. The BMI of MD and NMD stratified for protein users was 24.34±2.99 and 25.36±3.48, respectively; the BMI of MD and

NMD stratified for non-protein users was 23.65±3.19 and 21.61±3.49, respectively. Differences are shown in table 5.

Frequency per week	0	1	2	3	4	5	6	7
<b>Foods</b>								
<i>Bakery</i>	15	5	7.5	15	5	15	7.5	30
<i>Chicken</i>	7.5	2.5	17.5	25	17.5	10	5	15
<i>Cheese</i>	7.5	2.5	17.5	22.5	22.5	12.26	1.89	7.55
<i>Cold cuts</i>	16.98	12.26	18.87	17.92	12.26	17.5	0	10
<i>Fresh Fish</i>	22.5	15	17.5	10	10	2.5	2.5	20
<i>Legumes</i>	10	35	40	5	2.5	2.5	0	5
<i>Meat</i>	12.5	28.5	18	10	10	5	2.5	12.5
<i>Milk</i>	22.5	2.5	10	15	2.5	20	2.5	25
<i>Nuts</i>	27.5	17.5	20	10	5	2.5	0	17.5
<i>Canned Tuna</i>	15	22.5	17.5	17.5	15	2.5	5	5
<i>Eggs</i>	2.5	2.5	15	7.5	7.5	20	5	40
<i>Yogurt</i>	17.5	10	15	10	5	12.5	5	25

**Table 3:** Percentage of food consumption of the Non-Mediterranean area (n=40).

Food	<i>p</i>	$\chi^2$
Bakery	<0.001	182.53
Chicken	<0.001	95.85
Cheese	<0.001	33.79
Cold cuts	<0.001	37.53
Fresh Fish	<0.001	38
Legumes	<0.001	59.91
Meat	<0.001	89.63
Milk	0.006	19.62
Nuts	<0.001	252.45
Canned Tuna	<0.001	99.34
Eggs	<0.001	858.61
Yogurt	<0.001	86.52

**Table 4:** Comparison of dietary patterns between the Mediterranean and Non-Mediterranean population.

## Discussion

The questionnaire was intended to retrieve data from a MD and NMD area in order to understand the differences between dietary patterns, with particular interest in MD, and how the intake of protein supplements affects the BMI.

Comparisons	<i>p</i>
BMI MD vs. BMI NMD	Ns
BMI pro vs. BMI No pro	< 0.001
BMI MD pro vs. BMI MD No pro	< 0.05
BMI NMD pro vs. BMI NMD No pro	< 0.001
BMI MD pro vs. BMI NMD pro	Ns
BMI MD No pro vs. BMI NMD No pro	<0.05

**Table 5:** Differences of BMI between groups.

*BMI = Body Mass Index; MD = Mediterranean Diet; NMD = Non-Mediterranean Diet; pro = protein supplement users; No pro = Non-protein supplement users; Ns = Not significant.*

Our manuscript presents many biases and limitations and we strongly believe that it can be considered only an interesting update of Protein Project, with stimulating inputs for professionals working on the field of sport nutrition and sport performance. Prior to jump to all positive evidences we want to highlight, in details, what we retain the main limits of this investigation. 1) Even though the comparison of food frequency has provided us with an opportunity to identify to identify a Mediterranean like pattern of dietary habits, people living in the Mediterranean area do not necessarily consume a Mediterranean diet. 2) No predictors of weight management and no determinants of health-related quality of life was studied. 3) The sample of participants coming from Poland cannot be considered statistically relevant to accomplish conclusions or interpretations.

The main outcomes highlight significant differences in BMI between the MD group those subjects that used and the NMD group (of both the MD and NMD areas) or the NMD group; whilst after a further stratification, significant differences in BMI were found between supplement and non-supplement users. This indicate that higher BMI values for the supplement users might be related to their quality of life (dietary behavior, kind of resistance training should be). Our results furthermore demonstrate that the use of other diets (mainly rich in proteins or zone diets) also appear to control and manage weight as outcome. Moreover, we did not identify a significant difference between the MD and the NMD groups, despite a significant difference in

the nutritional composition of the analyzed diets (table 4).

Other strategies can therefore be suggested to be as equally effective in weight loss and weight management as the MD. However, when compared for example to a low fat diet it is the MD diet which has been demonstrated to be more beneficial in reducing markers for cardiovascular diseases<sup>(28)</sup>. Mayneris-Perxachs et al.<sup>(29)</sup> recently demonstrated a change in lipid profiles in patients suffering with the metabolic syndrome after a one-year MD intervention, despite no changes in BW weight. The interesting role of the MD in weight management was also confirmed by Roccaldo et al.<sup>(30)</sup>.

The MD due to its composition has a low energy density and a relatively low glycemic load<sup>(31)</sup> compared to other dietary patterns. The high proportion of fruit and vegetables furthermore provide higher water content, which leads to an increase in saturation and lower calorie intake, thus facilitating a positive weight management. Other physiological explanations, which could elucidate the reason of the MD protecting mechanisms of weight, gain and/or weight loss can be suggested by plant-based foods that provide a large quantity of dietary fiber. These have been shown to increase satiety and satiation<sup>(32)</sup>. The effectiveness of the MD in weight management that arises from numerous studies may be due to the fact that these were performed in overweight and obese subjects with an intention of reducing the caloric intake to reduce the subjects BW<sup>(33, 34)</sup>.

A main concern in the present study is the difference in BMI values between protein and non-protein supplement users. Recent research, which investigated protein supplementation, in general demonstrated no weight gains or weight losses<sup>(12, 13, 35-37-39)</sup>. These however were longitudinal studies that investigated the long-term effects of such nutritional component consumption of such nutritional components. There are still uncertainties in scientific literature regarding direct comparisons of baseline anthropometric or metabolic parameters between supplement users and non-supplement users in healthy populations. Further studies can consequently be suggested, which in-depth investigate the difference between.

In conclusion, the MD compared to other dietary regimens does not seem to show any baseline difference regarding BMI values in healthy active individuals. Apart from dietary patterns, the intake of protein supplements seems to be a deter-

mining factor for a positive weight strategies. A direct comparison between protein and non-protein users showed higher BMI values in the MD group. These findings may be of interest for nutritionists involved in the treatment of underweight individuals or athletes.

## References

- 1) Trichopoulou A, Lagiou P. *Healthy traditional Mediterranean diet: an expression of culture, history, and lifestyle*. Nutr Rev. 1997 Nov; 55(11 Pt 1): 383-9.
- 2) Tapsell LC. *Foods and food components in the Mediterranean diet: supporting overall effects*. BMC Med. 2014; 12: 100.
- 3) Bach-Faig A, Berry EM, Lairon D, Reguant J, Trichopoulou A, et al. *Mediterranean diet pyramid today. Science and cultural updates*. Public Health Nutr. 2011 Dec; 14(12A): 2274-84.
- 4) Trichopoulou A, Martinez-Gonzalez MA, Tong TY, Forouhi NG, Khandelwal S, et al. *Definitions and potential health benefits of the Mediterranean diet: views from experts around the world*. BMC Med. 2014; 12(1): 112.
- 5) Karamanos B, Thanopoulou A, Angelico F, Assaad-Khalil S, Barbato A, et al. *Nutritional habits in the Mediterranean Basin. The macronutrient composition of diet and its relation with the traditional Mediterranean diet*. Multi-centre study of the Mediterranean Group for the Study of Diabetes (MGSD). Eur J Clin Nutr. 2002 Oct; 56(10): 983-91.
- 6) Ruiz-Canela M, Martinez-Gonzalez MA. *Lifestyle and dietary risk factors for peripheral artery disease*. Circ J. 2014; 78(3): 553-9.
- 7) Velazquez-Lopez L, Santiago-Diaz G, Nava-Hernandez J, Munoz-Torres AV, Medina-Bravo P, et al. *Mediterranean-style diet reduces metabolic syndrome components in obese children and adolescents with obesity*. BMC Pediatr. 2014; 14(1): 175.
- 8) Casas R, Sacanella E, Urpi-Sarda M, Chiva-Blanch G, Ros E, et al. *The effects of the mediterranean diet on biomarkers of vascular wall inflammation and plaque vulnerability in subjects with high risk for cardiovascular disease*. A randomized trial. PLoS One. 2014; 9(6): e100084.
- 9) Greco M, Chieffari E, Montalcini T, Accattato F, Costanzo FS, et al. *Early effects of a hypocaloric, Mediterranean diet on laboratory parameters in obese individuals*. Mediators Inflamm. 2014; 2014: 750860.
- 10) Grosso G, Pajak A, Mistretta A, Marventano S, Raciti T, et al. *Protective role of the Mediterranean diet on several cardiovascular risk factors: evidence from Sicily, southern Italy*. Nutr Metab Cardiovasc Dis. 2014 Apr; 24(4): 370-7.
- 11) Bekkouche L, Bouchenak M, Malaisse WJ, Yahia DA. *The Mediterranean diet adoption improves metabolic, oxidative, and inflammatory abnormalities in Algerian metabolic syndrome patients*. Horm Metab Res. 2014 Apr; 46(4): 274-82.
- 12) Miller PE, Alexander DD, Perez V. *Effects of whey protein and resistance exercise on body composition: a meta-analysis of randomized controlled trials*. J Am Coll Nutr. 2014; 33(2): 163-75.
- 13) Wu J, Dong J, Jiang X, Qin L. *Effects of soy protein supplement on overweight and obese population: meta-analysis of randomized controlled trials*. Wei Sheng Yan Jiu. 2013 Mar; 42(2): 185-9.
- 14) Soenen S, Plasqui G, Smeets AJ, Westerterp-Plantenga MS. *Protein intake induced an increase in exercise stimulated fat oxidation during stable body weight*. Physiol Behav. 2010 Dec 2; 101(5): 770-4.
- 15) Smigielski J, Jegier A, Hanke W, Bielecki W, Drygas W. *The effect of selected lifestyle factors and diet on mortality of men with documented physical fitness in the city of Lodz*. Int J Occup Med Environ Health. 2013 Aug; 26(4): 535-44.
- 16) Eisenberg ME, Wall M, Neumark-Sztainer D. *Muscle-enhancing behaviors among adolescent girls and boys*. Pediatrics. 2012 Dec; 130(6): 1019-26.
- 17) Arnberg K, Molgaard C, Michaelsen KF, Jensen SM, Trolle E, et al. *Skim milk, whey, and casein increase body weight and whey and casein increase the plasma C-peptide concentration in overweight adolescents*. J Nutr. 2012 Dec; 142(12): 2083-90.
- 18) Bianco A, Mammaia C, Thomas E, Bellafiore M, Battaglia G, et al. *Protein supplementation and dietary behaviours of resistance trained men and women attending commercial gyms: a comparative study between the city centre and the suburbs of Palermo, Italy*. J Int Soc Sports Nutr. 2014; 11: 30.
- 19) Bianco A, Mammaia C, Paoli A, Bellafiore M, Battaglia G, et al. *Protein supplementation in strength and conditioning adepts: knowledge, dietary behavior and practice in Palermo, Italy*. J Int Soc Sports Nutr. 2011; 8(1): 25.
- 20) Bianco A. *The use of protein supplementation among fitness center attendees*. The protein project. EJSS Journal. 2014; 2(1): 61-3.
- 21) Bianco A, Mammaia C, Thomas E, Ciulla F, Pupella U, et al. *Protein supplements consumption: a comparative study between the city centre and the suburbs of Palermo, Italy*. BMC Sports Sci Med Rehabil. 2014; 6:29.
- 22) Eldridge AL, Sheehan ET. *Food supplement use and related beliefs: Survey of community college students*. Journal of Nutrition Education. 1994 11//; 26(6): 259-65.
- 23) Administration USFaD. *What is a dietary supplement?* 2014; Available from: <http://www.fda.gov/aboutfda/transparency/basics/ucm195635.htm>.
- 24) Willett WC, Sacks F, Trichopoulou A, Drescher G, Ferro-Luzzi A, et al. *Mediterranean diet pyramid: a cultural model for healthy eating*. Am J Clin Nutr. 1995 Jun; 61(6 Suppl): 1402S-6S.
- 25) Boylan S, Welch A, Pikhart H, Malyutina S, Pajak A, et al. *Dietary habits in three Central and Eastern European countries: the HAPIEE study*. BMC Public Health. 2009; 9: 439.
- 26) El Ansari W, Stock C, Mikolajczyk RT. *Relationships between food consumption and living arrangements among university students in four European countries - a cross-sectional study*. Nutr J. 2012; 11: 28.
- 27) Smigielski J, Bielecki W, Drygas W. *Health and life style-related determinants of survival rate in the male residents of the city of Lodz*. Int J Occup Med Environ Health. 2013 Jun; 26(3): 337-48.

- 28) Serra-Majem L. *Efficacy of diets in weight loss regimens: is the Mediterranean diet appropriate?* Pol Arch Med Wewn. 2008 Dec; 118(12): 691-3.
- 29) Mayneris-Perxachs J, Sala-Vila A, Chisaguano M, Castellote AI, Estruch R, et al. *Effects of 1-year intervention with a mediterranean diet on plasma Fatty acid composition and metabolic syndrome in a population at high cardiovascular risk.* PLoS One. 2014; 9(3): e85202.
- 30) Roccaldo R, Censi L, D'Addezio L, Toti E, Martone D, et al. *Adherence to the Mediterranean diet in Italian school children (The ZOOM8 Study).* Int J Food Sci Nutr. 2014 Aug; 65(5): 621-8.
- 31) Willett WC, Leibel RL. *Dietary fat is not a major determinant of body fat.* Am J Med. 2002 Dec 30; 113 Suppl 9B: 47S-59S.
- 32) Schroder H. *Protective mechanisms of the Mediterranean diet in obesity and type 2 diabetes.* J Nutr Biochem. 2007 Mar; 18(3): 149-60.
- 33) Shai I, Schwarzfuchs D, Henkin Y, Shahar DR, Witkov S, et al. *Weight Loss with a Low-Carbohydrate, Mediterranean, or Low-Fat Diet.* New England Journal of Medicine. 2008; 359(3): 229-41.
- 34) Sacks FM, Bray GA, Carey VJ, Smith SR, Ryan DH, et al. *Comparison of weight-loss diets with different compositions of fat, protein, and carbohydrates.* N Engl J Med. 2009 Feb 26; 360(9): 859-73.
- 35) Antonio J, Peacock CA, Ellerbroek A, Fromhoff B, Silver T. *The effects of consuming a high protein diet (4.4 g/kg/d) on body composition in resistance-trained individuals.* J Int Soc Sports Nutr. 2014; 11: 19.
- 36) Shahar S, Kamaruddin NS, Badrasawi M, Sakian NI, Abd Manaf Z, et al. *Effectiveness of exercise and protein supplementation intervention on body composition, functional fitness, and oxidative stress among elderly Malays with sarcopenia.* Clin Interv Aging. 2013; 8: 1365-75.
- 37) Deibert P, Solleder F, Konig D, Vitolins MZ, Dickhuth HH, et al. *Soy protein based supplementation supports metabolic effects of resistance training in previously untrained middle aged males.* Aging Male. 2011 Dec; 14(4): 273-9.

#### Acknowledgments

We are grateful to Prof. Caterina Mammina (University of Palermo) and Prof. Anna Tagliabue (University of Pavia) for their invaluable help.

#### Corresponding author

FRANCESCO MARTINES  
BioNeC Department, Audiology Section  
University of Palermo  
(Italy)