

# Kent Academic Repository

## Full text document (pdf)

### Citation for published version

Dallimer, Martin and Tang, Zhiyao and Gaston, Kevin J. and Davies, Zoe G. (2016) The extent of shifts in vegetation phenology between rural and urban areas within a human-dominated region. *Ecology and Evolution*, 6 (7). pp. 1942-1953.

### DOI

<https://doi.org/10.1002/ece3.1990>

### Link to record in KAR

<http://kar.kent.ac.uk/55124/>

### Document Version

Publisher pdf

#### Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

#### Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

#### Enquiries

For any further enquiries regarding the licence status of this document, please contact:

[researchsupport@kent.ac.uk](mailto:researchsupport@kent.ac.uk)

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

## The extent of shifts in vegetation phenology between rural and urban areas within a human-dominated region

Martin Dallimer<sup>1,\*</sup>, Zhiyao Tang<sup>2,\*</sup>, Kevin J. Gaston<sup>3</sup> & Zoe G. Davies<sup>4</sup>

<sup>1</sup>Sustainability Research Institute, School of Earth and Environment, University of Leeds, Woodhouse Lane, Leeds LS2 9JT, UK

<sup>2</sup>Department of Ecology, College of Urban and Environmental Sciences, Peking University, Beijing 100871, China

<sup>3</sup>Environment and Sustainability Institute, University of Exeter, Penryn, Cornwall TR10 9EZ, UK

<sup>4</sup>Durrell Institute of Conservation and Ecology (DICE), School of Anthropology and Conservation, University of Kent, Canterbury, Kent CT2 7NR, UK

### Keywords

Enhanced Vegetation Index, green infrastructure, greenspace, growing season, urban ecology, urban heat island, urbanization.

### Correspondence

Martin Dallimer, Sustainability Research Institute, School of Earth and Environment, University of Leeds, Woodhouse Lane, Leeds LS2 9JT, UK.

Tel: +44 113 343 3036;

Fax: +44 113 343 5259;

E-mail: m.dallimer@leeds.ac.uk

### Funding Information

Seventh Framework Programme, (Grant/Award Number: 'Marie Curie Fellowship no. 273547') Engineering and Physical Sciences Research Council, (Grant / Award Number: 'EP/ F007388/1')

Received: 8 April 2015; Revised: 22

December 2015; Accepted: 5 January 2016

*Ecology and Evolution* 2016; 6(7): 1942–1953

doi: 10.1002/ece3.1990

\*These two authors contributed equally to this article.

## Introduction

Urbanization and climate change are two of the major environmental challenges facing the world today. Determining the broad swathe of consequences of climate change on species, communities and ecosystems have been the leading focus of research in recent decades (e.g., Bunn 2009). One of the most well-documented impacts has been the extension of the growing season for

## Abstract

Urbanization is one of the major environmental challenges facing the world today. One of its particularly pressing effects is alterations to local and regional climate through, for example, the Urban Heat Island. Such changes in conditions are likely to have an impact on the phenology of urban vegetation, which will have knock-on implications for the role that urban green infrastructure can play in delivering multiple ecosystem services. Here, in a human-dominated region, we undertake an explicit comparison of vegetation phenology between urban and rural zones. Using satellite-derived MODIS-EVI data from the first decade of the 20th century, we extract metrics of vegetation phenology (date of start of growing season, date of end of growing season, and length of season) for Britain's 15 largest cities and their rural surrounds. On average, urban areas experienced a growing season 8.8 days longer than surrounding rural zones. As would be expected, there was a significant decline in growing season length with latitude (by 3.4 and 2.4 days/degree latitude in rural and urban areas respectively). Although there is considerable variability in how phenology in urban and rural areas differs across our study cities, we found no evidence that built urban form influences the start, end, or length of the growing season. However, the difference in the length of the growing season between rural and urban areas was significantly negatively associated with the mean disposable household income for a city. Vegetation in urban areas deliver many ecosystem services such as temperature mitigation, pollution removal, carbon uptake and storage, the provision of amenity value for humans and habitat for biodiversity. Given the rapid pace of urbanization and ongoing climate change, understanding how vegetation phenology will alter in the future is important if we wish to be able to manage urban greenspaces effectively.

plants in temperate regions (e.g., Menzel and Fabian 1999; Noormets 2009). During the latter half of the 20th century, at mid- and high latitudes, a shift towards an earlier onset of spring and later autumn has been widely observed (Steltzer and Post 2009 and references therein). The magnitude of the phenomenon has been shown to be closely associated with changes in mean temperature (White et al. 1997; Fitter and Fitter 2002; Penuelas et al. 2002; White and Nemani 2003; Badeck et al. 2004; Chuine

*et al.* 2004; Piao *et al.* 2006; Jeong *et al.* 2011; Cong *et al.* 2013). However, other factors such as water availability, precipitation, photoperiod length, nitrogen deposition, and CO<sub>2</sub> concentrations are also known to influence vegetation phenology (Badeck *et al.* 2004; Korner and Basler 2010; Jeong *et al.* 2011; Cong *et al.* 2013).

Globally, the pace of land conversion to urbanized areas is rapid (Seto *et al.* 2012), with over half of the human population now living in towns and cities (United Nations 2013). Through the conversion of natural land surfaces to the built form, such as buildings, roads, and other sealed surfaces, urbanization radically alters many aspects of an ecosystem, including water availability, species composition, and soil properties (Gaston *et al.* 2010). One outcome of urbanization is an alteration of local and regional climate via the Urban Heat Island (UHI) effect, modifications to wind flow and turbulence, as well as shifts in patterns of cloud formation and precipitation (for reviews Gaston *et al.* 2010; Seto and Shepherd 2009). Temperature increases associated with UHI can be as great as several degrees Celsius, even in mid- and high latitude cities (e.g., Kershaw *et al.* 2010). Given the link between phenology and temperature, we would therefore expect seasonal patterns of vegetation growth and die-back to differ between urban centers and surrounding non-built-up areas. While temperature shifts associated with urbanization are undoubtedly important, ground-based observations indicate that the influence of urbanization on various aspects of the growing season is mediated by species and community composition, soil moisture, and topology (Fisher *et al.* 2007; Gazal *et al.* 2008; Hwang *et al.* 2011; Jochner *et al.* 2012), all of which can be substantially altered by urbanization.

Shifts in vegetation phenology can have a profound impact on ecosystem function, altering water, carbon, and energy balances, and affecting interspecific interactions and productivity (Schwartz 1998; White *et al.* 1999; Menzel 2000; Parmesan and Yohe 2003). This is particularly pertinent in urban areas where vegetation is key to the delivery of many important ecosystem services, including temperature mitigation (Susca *et al.* 2011; Park *et al.* 2012; Myint *et al.* 2013), pollution reduction (Manes *et al.* 2012; Pugh *et al.* 2012), carbon storage (Davies *et al.* 2011), recreational opportunities for human residents (EEA 2009), and habitat for biodiversity (Chace and Walsh 2006; McKinney 2008; Dallimer *et al.* 2012). Indeed, green infrastructure is increasingly being recognized as a critical component of urban areas, improving the quality of the environment for inhabitants (Gaston 2010; Niemela *et al.* 2010; Keniger *et al.* 2013).

Here, we undertake an explicit comparison of vegetation phenology between urban and adjacent rural zones. Using satellite-derived data, we examine variation in the

growing season across the 15 largest cities in Britain. Our aim (cf. White *et al.* 2002) is to do this without assessing the longer-term temporal trends associated with climate change. We therefore restrict our analyses to the first decade of the 21st century, as temperatures (Cane 2010; Wang *et al.* 2011) and phenological signals in vegetation (e.g., Jeong *et al.* 2011; Piao *et al.* 2011; Wu and Liu 2013) remained relatively static during this time.

We purposefully carry out our analyses in a human-dominated region. Some 80% of the UK population already lives in towns and cities, a proportion that the rest of the world is predicted to approach by 2050 (United Nations 2013). Further, we include all land covers in our dataset. Much of the existing literature has compared urban areas with nearby forested/natural landscapes (e.g., White *et al.* 1997; Zhang *et al.* 2004; Wu and Liu 2013), thereby excluding, for example, agricultural land, where management influences phenology. Some urban–rural differences in growing season that have been ascribed to urbanization might, therefore, be confounded by the radically different type, scale and management of vegetation occurring in urban compared to forested/natural areas. Our study cities are surrounded by a mosaic of land covers, such as patches of woodland and shrub, grass and croplands, which, although different from land covers in urban areas, encompass many of the types of heterogeneous vegetation structure and cover (e.g., woodland, shrub patches, amenity grassland) that are typically found in British cities. We test the following statements: (i) the growing season will be longer in urban compared to neighboring rural areas across the major cities of Britain, and (ii) any declines in growing season length that are associated with latitude will be notably less in urban areas.

## Materials and Methods

### Data acquisition and processing

Satellite-derived vegetation indices, related to the fraction of photosynthetically active radiation absorbed by plants, are widely used as a surrogate for vegetation activity (Huete *et al.* 2002; Sims *et al.* 2006; White *et al.* 2009). In this study, we use Moderate Resolution Imaging Spectroradiometer Enhanced Vegetation Index (MODIS-EVI) data to examine phenological trends within Britain's 15 largest cities (Fig. 1; Appendix S1 and Table S1). The dataset has a spatial resolution of 250 × 250 m, a temporal resolution of 16 days (16-day composite period (Justice *et al.* 2002), and we extracted data spanning a 9-year period in the UK from February 2000 (when MODIS-EVI data first became available) to December 2009. The MODIS-EVI data were downloaded from the Global Land Cover Facility (<http://glcf.umd.edu/>).




**Figure 1.** The location of the 15 study cities (Latitudes 50.72 to 55.95N) in Britain, northwest Europe. Darker green shading indicates increasing proportion of each city that was recorded as greenspace.

In order to investigate how the growing season within cities might be altered as a result of urbanization, comparative baseline data were required for adjacent rural zones. The urban extent of each city was delimited according to the 2006 Ordnance Survey definition using a Geographic Information System (GIS). Surrounding rural zones were defined as all non-built-up land uses (determined using Landsat TM; Appendix S2) lying in a buffer between 2 and 5 km around the urban boundary. Based on these urban and rural zones, within the GIS, we calculated the biweekly mean EVI for each study city and its associated rural area by averaging across all EVI pixels contained within the zones. The start of the growing season (SOS) each year was defined as the first day that the EVI increased above the annual mean EVI in the spring. The end of the growing season (EOS) was taken as the day the EVI decreased below the annual mean EVI in the autumn (Zhou *et al.* 2001; Suzuki *et al.* 2003). Finally, the length of the growing season (LOS) was the difference between the SOS and EOS. As the proportion of vegetation cover within each city and its adjacent rural area differs, a

specific annual mean EVI threshold was applied to each zone (Fig. 2). This approach is equivalent to, but simpler than, the phenology fitting curve used in other studies (Zhou *et al.* 2001; Suzuki *et al.* 2003).

In some years, peaks in EVI above the annual mean EVI threshold occurred during winter. Such peaks were characterized by a single period of high EVI followed by an immediate reduction and are likely to be due to factors such as cloud, atmosphere, and solar zenith angle. As it is not possible for plants in temperate regions to complete their growth phase in such a short period of time during the winter, we excluded EVI peaks from our definitions of SOS and EOS. We did this by discounting peaks in EVI that occurred prior to day 70 (11 March) or after day 315 (11 November); dates that were chosen based on our experience of the study region and inspecting the annual form of EVI. A further potential complication is the theoretical situation where low points in EVI could occur in the rural zone where that area is cultivated and harvested. Although low points did occur, EVI never dropped below the mean annual EVI prior to the autumn “green down.” We defined EVI thresholds using both mean and median EVI. There were no substantive differences in the outputs, so here we report results based on the mean only.

The choice of method used to determine the SOS and EOS can lead to considerable variation and limit the comparability of different studies. For example, satellite-derived data tend to deliver SOS dates earlier and EOS later than ground-based observations (White *et al.* 2009; Zhu *et al.* 2012), with heterogeneity in vegetation cover as one explanation for why this occurs (Badeck *et al.* 2004). Equally, the exact methodology used to estimate the SOS from the satellite data itself will lead to different dates being estimated. White *et al.* (2009) compared 10 methods for calculating the SOS, from empirical approaches to mathematical models. They found that no method could claim to be automatically superior to the others. Similarly, Cong *et al.* (2013) found that although there were significant differences between techniques in their estimates of the SOS and EOS, all were internally consistent and able to reveal the same patterns within the data. We therefore opted for a transparent and straightforward method for ascertaining the SOS and EOS (as outlined above). Finally, several methods can also be applied to interpolate between the EVI data points, such as linear, quadratic, cubic, and polynomial interpolation. The linear method might estimate an earlier SOS and later EOS, given a concave shape of EVI pattern, but a later SOS and earlier EOS, given a convex shape of EVI pattern (see Fig. 2 B and C). The selection of methods depends on the shape of the temporal changes of EVI, how many dates with EVI data are available, and the



**Figure 2.** (A) An example of the annual pattern in biweekly Moderate Resolution Imaging Spectroradiometer Enhanced Vegetation Index (MODIS-EVI) data, averaged across all years, for the urban extent of the city of Birmingham (closed circles; dashed line, annual mean EVI) and its surrounding rural zone (open circles; dotted line, annual mean EVI). SOS and EOS indicate the start and end of the growing season, respectively. To interpolate the estimated date of EOS and SOS from the 16-day interval of the EVI data, we assumed that EVI increased in spring, and decreased in autumn linearly within the 16 days of the interval. Therefore, the estimated SOS date (when the mean EVI intersected with observed EVI) is calculated as follows:  $SOS = day_1 + (mean\ EVI - EVI_1)/(EVI_2 - EVI_1) \times 16$ . Where  $day_1$  and  $day_2$  are the neighboring dates of the EVI values. Similarly, we calculated the EOS as following:  $EOS = day_3 + (mean\ EVI - EVI_4)/(EVI_3 - EVI_4) \times 16$ . (B) and (C) illustrate in more detail how linear interpolation between biweekly data points was used.

datasets and consistent with our desire to apply a straightforward and consistent method.

In this study, we wished to understand the impact of urbanization as a whole on vegetation phenology. This could be mediated through many possible drivers (e.g., temperature, water availability, nitrogen deposition, vegetation community composition, variation between native and non-native plant species in responses to photoperiod and/or temperature). We therefore did not attempt to disentangle their relative importance and instead examined whether differences in the SOS, EOS, and LOS between each city and its neighboring rural zone were associated with some key characteristics of urban form (the proportion of greenspace, dwelling density, extent of the urban area, distance to the nearest major urban area, and disposable household income; Table S3). Following previous studies, we predicted that the difference in SOS, EOS, and LOS between each city and its neighboring rural zone would be negatively associated with the proportion of greenspace in a city (cf. temperature mitigation associated with vegetation in cities; Susca et al. 2011; Park et al. 2012; Myint et al. 2013). We anticipated a similar association with disposable income, given that socio-demographics are often associated with many aspects of vegetation structure and coverage in cities (e.g., Hope et al. 2003; Luck et al. 2009). In contrast, we predicted that the difference in SOS, EOS, and LOS between each city and its neighboring rural zone would be positively associated with dwelling density (a metric of how intensively built-up an urban area is) and the extent of the urban area. Finally, cities can raise temperatures in a broad swathe of rural land around them (Zhang et al. 2004; Elmore et al. 2012). Rural areas close to several large cities are therefore likely to experience higher temperatures than more isolated areas, reducing the phenological differences between a city and its surrounding rural zone. We therefore predicted that differences in

purpose of the phenological phase estimation. When comparing the temporal and spatial patterns of phenological phases, it is important to select a method that is comparable among different times and different sites. We therefore selected linear interpolation, which is not only simple and reliable, but compatible among different

phenology would be lower when major urban areas were closer together.

### Statistical analyses


Linear regression was used to explore temporal trends for each of the growing season characteristics in the study cities and their neighboring rural zones which might be attributable to large scale changes in, for example, climate. As no relationships between the SOS, EOS, or LOS and time were apparent, either at the individual city level or for all cities combined (Table S2), mean values for these variables across the 10 year period were used in all further analyses. Across all the cities, paired *t*-tests were used to assess whether urbanization resulted in a significant difference in any of the growing season characteristics between urban and rural zones. At the individual city level, differences between urban areas and adjacent rural zones for the SOS, EOS, and LOS were investigated using Wilcoxon signed-rank tests, as the data were not consistently normal.

To determine whether SOS, EOS, and LOS varied with latitude, linear regression was applied (Table S3). We hypothesized that, if latitudinal trends were present, they would be less pronounced (i.e.,  $\beta$  lower) within the cities, and tested for this by including an interaction term between latitude and zone (rural or urban). If this term were to be significant, then this would be evidence that  $\beta$ s, and therefore latitudinal trends, differed between zones. Finally, we assessed the strength of any association between urban form and the difference in SOS, EOS, and LOS between urban and rural areas with partial Spearman's rank correlations, which allowed us to account for the likely influence of latitude. Our sample size ( $N = 15$ ) precluded us from undertaking more complex multivariate analyses.

### Results

When considering individual cities, no uniform pattern emerged in relation to the SOS in urban versus rural zones (Fig. 3A; Table S4); the SOS was significantly earlier in three urban centers, compared to adjacent rural areas, but significantly delayed in another two. In contrast, the EOS was consistently later in all urban areas, significantly so for four cities (Fig. 3B; Table S4). The LOS was extended in 13 cities, five of which were significant (Fig. 3C; Table S4). In all cases, differences were modest compared to the temporal resolution of the MODIS data.

Combining the data from across all 15 British cities, the LOS was significantly longer in urban versus rural areas by 8.8 days (192 days and 183 days, respectively). The mean SOS occurred on 3 April in urban areas compared to



**Figure 3.** Differences in the (A) start of the growing season (SOS), (B) end of the growing season (EOS) and (C) length of growing season (LOS), for each study city (latitude in brackets) when compared to its surrounding rural area. \* indicates a significant difference ( $P < 0.05$ ) between the urban and rural zones (Table S4).

4 April in adjacent rural zones, due to an average, non-significant, advance of 0.8 days (Table 1). Similarly, on average, the EOS was 8.0 days later in urban areas than rural zones, with a mean date of 12 October as opposed to 4 October.

**Table 1.** Paired *t*-tests assessing differences in the growing season characteristics between urban and rural zones across 15 British study cities: start of season, SOS; end of season, EOS; length of season, LOS.

Growing season variable	Zone	Mean	SE	df	<i>t</i>	<i>P</i>
SOS	Urban	93.36	1.23	24.721	0.376	0.710
	Rural	94.18	1.80			
EOS	Urban	285.25	1.87	24.336	-2.357	0.027
	Rural	277.30	2.81			
LOS	Urban	191.88	1.89	27.448	-3.036	0.005
	Rural	183.12	2.18			

Although the SOS was delayed with increasing latitude (1.9 and 0.9 days per degree of latitude in rural and urban areas, respectively), these trends were not significant (Fig. 4A; Table 3). Similarly, there was a non-significant advance in EOS, of 1.5 days in both rural and urban areas with each degree of latitude (Fig. 4B; Table 3). However, the LOS decreased significantly, shortening by 3.4 days and 2.4 days per degree of latitude in rural and urban zones respectively (Fig. 4C; Table 3). Across Britain, from south to north, this equates to a 17.6-day reduction in the LOS in the rural zones surrounding the study cities, and a 12.5-day decline in LOS within urban areas (Fig. 4C). For all three growing season characteristics, the latitudinal trends were weaker (but not significantly so) in urban than rural areas, but interaction terms, and therefore, the differences between the slopes were not significant (Table 2).

Shifts in the growing season characteristics were modest and, after accounting for the influence of latitude in a partial correlation, were not significantly associated with the aspects of built urban form examined (percentage of greenspace, dwelling density, extent of the urban area, distance to nearest major urban area). However, LOS was negatively associated with disposable household income (Table 3).


## Discussion

Across Britain's 15 largest cities, urbanization has extended the growing season by an average of 8.8 days within urban areas, compared to rural surroundings, by prolonging the end of the season (Table 1) (cf. Elmore *et al.* 2012; Garonna *et al.* 2014). However, this figure masks considerable variation at an individual city level, with the differences in the length of the growing season ranging between 3.3 days shorter in the urban versus rural areas of Bournemouth, compared to 13.1 days longer for Leicester (Fig. 3C). Much of the variability can be explained by the geographic location of the cities

(Table 1), with the growing season estimated as being 17.6 and 12.5 days longer in rural and urban areas, respectively, for the lowest compared to the highest latitude (Fig. 4C). Although the modest compared to the relatively coarse temporal resolution of MODIS data, as long as a consistent methodology is used, vegetation phenology variables estimated from satellite data are robust (Cong *et al.* 2013), we can thus conclude that trends detected in our study are sound.

Although we detected no uniform pattern for the start of the growing season in urban areas to be earlier than rural zones, the end of the growing season was always delayed. This suggests that the length of the season in urban zones might be more strongly determined by autumn, rather than spring, vegetation phenology. The precise timing of the end of the growing season is inherently more difficult to measure than the start due to the gradual "green down" observed in temperate areas (cf. a rapid "green-up" at the onset of warmer temperatures in the spring), which perhaps is part of the reason why the importance of changes in autumn phenology in driving growing season length remains understudied (Garonna *et al.* 2014). Nevertheless, our results reflect those for Europe as a whole (Garonna *et al.* 2014) as well as those along gradients of urbanization in mid-Atlantic forests in North America (Elmore *et al.* 2012). Autumn phenology is strongly influenced by soil moisture and hydrology (Garonna *et al.* 2014; Yang *et al.* 2015), so the patterns we observed might be at least partly driven by changes to these that occurs in cities (Elmore *et al.* 2012). Differences in species composition between urban and rural zones might also be important (Elmore *et al.* 2012).

Climate change is thought to be the major driver in the advance and extension of growing seasons in temperate latitudes (e.g., Schwartz 1998; Menzel and Fabian 1999). Nonetheless, ambient climate can also be altered by land-use changes associated with urbanization (e.g., Imhoff *et al.* 2010; Li *et al.* 2011). For instance, Kershaw *et al.* (2010) modeled UHI for the largest cities in the UK and concluded that average spring temperatures were between 0.2 and 1.9°C warmer (for Leicester and Newcastle, respectively) than the surrounding rural landscape. One explanation for the variation in UHI is the differences in urban form (e.g., percentage of greenspace, dwelling density, extent of the urban area) between cities. We might, therefore, also expect variation in city-wide attributes of urban form to be associated with vegetation phenology. However, after accounting for latitude, we did not uncover any significant associations. Including measures of the UHI for each city in our analyses might have increased our understanding of the intracity variability, but our overall finding is consistent with that of Zhang



**Figure 4.** The relationship between latitude and the (A) start of the growing season (SOS), (B) end of the growing season (EOS), and (C) length of growing season (LOS), for each of the 15 British study cities (open circles) and their surrounding rural areas (closed squares). Lines (dashed, cities; solid, rural areas) indicate a significant relationship ( $P < 0.05$ ) with latitude.

*et al.* (2004), who also failed to detect a relationship between phenological differences between adjacent urban and rural zones and city size.

The vegetation phenological variability observed in our study at an individual city level is likely to be attributable to local-scale vegetation and topographic features that



characterize each urban area and directly influence the extent of the UHI effect experienced. For example, the specific characteristics of buildings can alter temperature patterns in urban landscapes (Myint *et al.* 2013), while trees and shrubs mitigate UHI (Susca *et al.* 2011; Park *et al.* 2012; Myint *et al.* 2013). Even small areas of vegetation can reduce UHI effects in their immediate vicinity (Oliveira *et al.* 2011), as can water bodies, such as rivers (Hathway and Sharples 2012). Furthermore, vegetation phenology can be modified by many factors, including disease, competition, soil condition, nutrient and water availability, and weather patterns (Menzel 2000). Similarly, the composition of vegetation communities differs between rural and urban zones, and across the latitudinal

range of our study. Given that the responses of individual species to climate change are enormously varied (for a summary, see Korner and Basler 2010), this might confound our ability to uncover consistent phenological patterns with remote sensed data.

Interestingly, a smaller difference in LOS was associated with higher city-scale mean disposable household income. In the desert city of Phoenix, USA, higher neighborhood household incomes were associated with cooler temperatures (Jenerette *et al.* 2011). If this relationship was also apparent in our region, this would translate to a smaller difference in vegetation phenology between urban and rural settings. Indeed, income and socio-economic status is often associated with vegetation structure, type, species richness, and community composition in urban areas (e.g., Hope *et al.* 2003; Luck *et al.* 2009). This therefore provides a plausible mechanism through which this association could be accounted for.

The spatial extent over which cities alter local climate can extend beyond the urban boundary. For example, in eastern North America, the climate influence of cities extended up to 10 km into the adjacent rural zone (Zhang *et al.* 2004). Similarly, Elmore *et al.* (2012) found that the influence of urban land use could be detected up to 32 km from large cities. Differences in vegetation phenology between urban and rural zones are therefore likely to be mediated by the extent to which rural areas are within a “climate shadow” of other nearby cities. Rural areas close to many cities might experience elevated temperatures compared to more isolated zones. In our study, differences in phenology between urban and rural zones were not associated with the distance between large cities. One explanation for the lack of an association could be that in our study region, the high human population density in Britain means that the majority of the rural landscape is impacted by urban “climate shadows” to some extent, and future work could focus on the size and intensity of these effects. In addition, we deliberately carried out our analyses in a human-dominated region and included all land covers in our dataset. Much of the existing literature has compared urban areas with nearby

**Table 2.** Linear regression models of the relationships between each growing season characteristics and latitude, for the urban and rural zones associated with Britain’s largest 15 cities: start of season, SOS; end of season, EOS; length of season, LOS.

Growing season variable	Zone	$\beta^1$	SE	P
SOS ( $R^2 = 0.13$ )	Urban	0.94	0.72	0.213
	Rural	1.89	0.99	0.078
	Latitude $\times$ Zone Interaction <sup>2</sup>	−0.95	1.22	0.443
EOS ( $R^2 = 0.14$ )	Urban	−1.46	1.09	0.203
	Rural	−1.48	1.70	0.399
	Latitude $\times$ Zone Interaction <sup>2</sup>	0.02	2.06	0.991
LOS ( $R^2 = 0.50$ )	Urban	−2.39	0.97	0.028
	Rural	−3.37	0.98	0.005
	Latitude $\times$ Zone Interaction <sup>2</sup>	0.97	1.38	0.487

<sup>1</sup>General form of the regression equation  $y = \alpha + \beta_1 (\text{Latitude}) + \beta_2 (\text{Zone}) + \beta_3 (\text{Latitude} \times \text{Zone})$ , where zone is a dummy with the value 0 for rural and 1 for urban. Intercept ( $\alpha$ ) is not reported.  $\beta$  represent the slope of the relationship between the growing season characteristic and latitude. The interaction terms are the difference between those slopes.

<sup>2</sup>If significant, this interaction term would indicate that latitudinal trends differed between urban and rural zones.

**Table 3.** Partial Spearman’s rank correlations between the advance/delay (days) in each growing season characteristic in urban areas, compared to rural zones and key aspects of, while accounting for the likely influence of latitude across the 15 study cities: start of season, SOS; end of season, EOS; length of season, LOS.

Growing season variable	Greenspace (%)	Dwelling density (no./ha)	Urban extent (ha)	Distance to nearest major city (km)	Household disposable income (GB£)
SOS	−0.08	−0.40	0.01	−0.03	0.18
EOS	−0.39	0.02	−0.03	−0.23	−0.48
LOS	−0.49	0.35	−0.23	−0.31	−0.71**

Significance levels (\* $P < 0.05$ ; \*\* $P < 0.01$ ) adjusted to correct for multiple tests using the Holm–Bonferroni method.

forested/natural landscapes (e.g., White *et al.* 1997; Zhang *et al.* 2004; Wu and Liu 2013). Some urban–rural differences in growing season that have been ascribed to urbanization might, therefore, be confounded by the different type, scale and management of vegetation occurring in urban compared to forested/natural areas. In our region, the study cities are surrounded by a mosaic of land covers, such as patches of woodland, scrub, grasslands, and arable crop, which, although different from land covers in urban areas, encompass many of the types of heterogeneous vegetation structure and cover (e.g., woodland, scrub patches, amenity grassland; Davies *et al.* 2011) that are typically found in British cities.

## Limitations and Conclusions

Solely relying on satellite-derived measures of phenology is likely to have restricted our ability to detect consistent signals for how urbanization might impact the growing season. For example, EVI response curves for rural areas were often characterized by a peak in growth in May–June, followed by a decline in July and a lower plateau in August–September. This is likely to reflect that large areas are covered by agricultural crops and grasses that ripen and are harvested/cut for silage in July–August. This pattern is likely to be less apparent in urban areas where frequent mowing and irrigation might keep vegetation greener for longer, leading to an apparent delay in the end of the growing season. In contrast, water shortages for street trees and pollution might both act in the opposite direction. In spring, increased temperatures in urban areas will generally mean that plants start growing earlier than they otherwise would. However, vegetation composition in urban areas is often very different from rural landscapes (e.g., see Dallimer *et al.* 2012) and urban vegetation can include many non-native species whose phenology might be more temperature sensitive than native species, increasing the chances of observing an earlier start to the growing season. Disentangling the relative importance of these effects, among others, is extremely challenging and a topic for future research.

Vegetation is a key component of urban areas, delivering many ecosystem services such as temperature mitigation, pollution removal, carbon uptake, and storage, the provision of amenity value for humans and habitat for biodiversity. Given the rapid pace of urbanization (United Nations 2013) and ongoing climate change (IPCC 2013), understanding how vegetation phenology will alter in the future is important if we wish to be able to manage urban greenspaces effectively. The impacts of an extended growing season on vegetation communities are likely to be complex, not least because individual species and functional groups will respond

differently (Korner and Basler 2010; Liang *et al.* 2011), with factors such as winter chilling, photoperiod, and temperature constraining them in varied and interacting ways. Nonetheless, the presence of “green” for longer in temperate urban areas could act to reduce local temperatures, not only through direct transpiration effects and shading (thereby reducing the UHI (e.g., Schwartz 1996)), but also indirectly by mitigating warming via carbon storage and sequestration (Penuelas *et al.* 2009). A longer growing season might also be beneficial for urban and peri-urban agriculture and food production, especially as the influence of a city on climate can extend outside the built-up area.

## Acknowledgments

ZYT was a China Scholar Council fellow. MD was supported by an EU-FP7 Marie Curie Fellowship (no. 273547). We thank the anonymous reviewers who encouraged us to include more speculative elements in the Discussion regarding possible mechanisms behind vegetation phenology difference between urban and rural areas.

## Conflict of Interest

None declared.

## References

- Badeck, F. W., A. Bondeau, K. Bottcher, D. Doktor, W. Lucht, J. Schaber, *et al.* 2004. Responses of spring phenology to climate change. *New Phytol.* 162:295–309.
- Bunn, I. D. 2009. The United Nations and climate change: legal and policy developments. Pp. 61–72 *in* Sustainability 2009: The Next Horizon Conference. American Institute of Physics, Melbourne, FL.
- Cane, M. A. 2010. Climate science: decadal predictions in demand. *Nat. Geosci.* 3:231–232.
- Chace, J. F., and J. J. Walsh. 2006. Urban effects on native avifauna: a review. *Landsc Urban Plan* 74:46–69.
- Chuine, I., P. Yiou, N. Viovy, B. Seguin, V. Daux, and E. L. Ladurie. 2004. Historical phenology: grape ripening as a past climate indicator. *Nature* 432:289–290.
- Cong, N., T. Wang, H. Nan, Y. Ma, X. Wang, R. B. Myneni, *et al.* 2013. Changes in satellite-derived spring vegetation green-up date and its linkage to climate in China from 1982 to 2010: a multimethod analysis. *Glob. Chang. Biol.* 19:881–891.
- Dallimer, M., J. R. Rouquette, A. M. J. Skinner, P. R. Armsworth, L. M. Maltby, P. H. Warren, *et al.* 2012. Contrasting patterns in species richness of birds, butterflies and plants along riparian corridors in an urban landscape. *Divers. Distrib.* 18:742–753.
- Davies, Z. G., J. L. Edmondson, A. Heinemeyer, J. R. Leake, and K. J. Gaston. 2011. Mapping an urban ecosystem

- service: quantifying above-ground carbon storage at a city-wide scale. *J. Appl. Ecol.* 48:1125–1134.
- EEA. 2009. Ensuring quality of life in Europe's cities and towns. European Environment Agency, Copenhagen.
- Elmore, A. J., S. M. Guinn, B. J. Minsley, and A. D. Richardson. 2012. Landscape controls on the timing of spring, autumn, and growing season length in mid-Atlantic forests. *Glob. Chang. Biol.* 18:656–674.
- Fitter, A. H., and R. S. R. Fitter. 2002. Rapid changes in flowering time in British plants. *Science* 296:1689–1691.
- Fisher, J. I., A. D. Richardson, and J. F. Mustard. 2007. Phenology model from surface meteorology does not capture satellite-based greenup estimations. *Glob. Chang. Biol.* 13:707–721.
- Garonna, I., R. de Jong, A. J. W. de Wit, C. A. Múcher, B. Schmid, and M. E. Schaepman. 2014. Strong contribution of autumn phenology to changes in satellite-derived growing season length estimates across Europe (1982–2011). *Glob. Chang. Biol.* 20:3457–3470.
- Gaston, K. J. 2010. Urban ecology. Pp. 1–9 *in* K. J. Gaston, ed. *Urban ecology*. Cambridge Univ. Press, Cambridge.
- Gaston, K. J., Z. G. Davies, and J. L. Edmondson. 2010. Urban environments and ecosystem functions. Pp. 35–52 *in* K. J. Gaston, ed. *Urban ecology*. Cambridge Univ. Press, Cambridge.
- Gazal, R., M. A. White, R. Gillies, E. Rodemaker, E. Sparrow, and L. Gordon. 2008. GLOBE students, teachers, and scientists demonstrate variable differences between urban and rural leaf phenology. *Glob. Chang. Biol.* 14:1568–1580.
- Hathway, E. A., and S. Sharples. 2012. The interaction of rivers and urban form in mitigating the Urban Heat Island effect: a UK case study. *Build. Environ.* 58:14–22.
- Huete, A., K. Didan, T. Miura, E. P. Rodriguez, X. Gao, and L. G. Ferreira. 2002. Overview of the radiometric and biophysical performance of the MODIS vegetation indices. *Remote Sens. Environ.* 83:195–213.
- Hope, D., C. Gries, W. X. Zhu, W. F. Fagan, C. L. Redman, N. B. Grimm, et al. 2003. Socioeconomics drive urban plant diversity. *Proc. Natl Acad. Sci. USA* 100:8788–8792. doi:10.1073/pnas.1537557100
- Hwang, T., C. Song, J. Vose, and L. Band. 2011. Topography-mediated controls on local vegetation phenology estimated from MODIS vegetation index. *Landsc. Ecol.* 26:541–556. doi:10.1007/s10980-011-9580-8
- Imhoff, M. L., P. Zhang, R. E. Wolfe, and L. Bounoua. 2010. Remote sensing of the urban heat island effect across biomes in the continental USA. *Remote Sens. Environ.* 114:504–513.
- IPCC. 2013. *Climate Change 2013: the physical science basis*. *In* T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, P. M. Midgley, eds. *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK and New York.
- Jeong, S.-J., C.-H. Ho, H.-J. Gim, and M. E. Brown. 2011. Phenology shifts at start vs. end of growing season in temperate vegetation over the Northern Hemisphere for the period 1982–2008. *Glob. Chang. Biol.* 17:2385–2399.
- Jenerette, G. D., S. L. Harlan, W. L. Stefanov, and C. A. Martin. 2011. Ecosystem services and urban heat riskscape moderation: water, green spaces, and social inequality in Phoenix, USA. *Ecol. Appl.* 21:2637–2651.
- Jochner, S. C., T. H. Sparks, N. Estrella, and A. Menzel. 2012. The influence of altitude and urbanisation on trends and mean dates in phenology (1980–2009). *Int. J. Biometeorol.* 56:387–394.
- Justice, C. O., J. R. G. Townshend, E. F. Vermote, E. Masuoka, R. E. Wolfe, N. Saleous, et al. 2002. An overview of MODIS Land data processing and product status. *Remote Sens. Environ.* 83:3–15.
- Keniger, L. E., K. J. Gaston, K. N. Irvine, and R. A. Fuller. 2013. What are the benefits of interacting with nature? *Int. J. Environ. Res. Public Health* 10:913–935.
- Kershaw, T., M. Sanderson, D. Coley, and M. Eames. 2010. Estimation of the urban heat island for UK climate change projections. *Build. Serv. Eng. Res. Technol.* 31:251–263.
- Korner, C., and D. Basler. 2010. Phenology under global warming. *Science* 327:1461–1462.
- Li, J., C. Song, L. Cao, F. Zhu, X. Meng, and J. Wu. 2011. Impacts of landscape structure on surface urban heat islands: a case study of Shanghai, China. *Remote Sens. Environ.* 115:3249–3263.
- Liang, L. A., M. D. Schwartz, and S. L. Fei. 2011. Validating satellite phenology through intensive ground observation and landscape scaling in a mixed seasonal forest. *Remote Sens. Environ.* 115:143–157.
- Luck, G. W., L. T. Smallbone, and R. O'Brien. 2009. Socioeconomics and vegetation change in urban ecosystems: patterns in space and time. *Ecosystems* 12:604–620.
- Manes, F., G. Incerti, E. Salvatori, M. Vitale, C. Ricotta, and R. Costanza. 2012. Urban ecosystem services: tree diversity and stability of tropospheric ozone removal. *Ecol. Appl.* 22:349–360.
- McKinney, M. L. 2008. Effects of urbanization on species richness: a review of plants and animals. *Urban Ecosyst* 11:161–176.
- Menzel, A. 2000. Trends in phenological phases in Europe between 1951 and 1996. *Int. J. Biometeorol.* 44:76–81.
- Menzel, A., and P. Fabian. 1999. Growing season extended in Europe. *Nature* 397:659–659.
- Myint, S. W., E. A. Wentz, A. J. Brazel, and D. A. Quattrochi. 2013. The impact of distinct anthropogenic and vegetation features on urban warming. *Landsc. Ecol.* 28:959–978.
- Niemela, J., S.-R. Saarela, T. Soderman, L. Kopperoinen, V. Yli-Pelkonen, S. Vare, et al. 2010. Using the ecosystem services approach for better planning and conservation of

- urban green spaces: a Finland case study. *Biodivers. Conserv.* 19:3225–3243.
- Noormets, A. 2009. Phenology of ecosystem processes: applications in global change research. Springer, New York.
- Oliveira, S., H. Andrade, and T. Vaz. 2011. The cooling effect of green spaces as a contribution to the mitigation of urban heat: a case study in Lisbon. *Build. Environ.* 46:2186–2194.
- Park, M., A. Hagishima, J. Tanimoto, and K. Narita. 2012. Effect of urban vegetation on outdoor thermal environment: field measurement at a scale model site. *Build. Environ.* 56:38–46.
- Parmesan, C., and G. Yohe. 2003. A globally coherent fingerprint of climate change impacts across natural systems. *Nature* 421:37–42.
- Penuelas, J., I. Filella, and P. Comas. 2002. Changed plant and animal life cycles from 1952 to 2000 in the Mediterranean region. *Glob. Chang. Biol.* 8:531–544.
- Penuelas, J., T. Rutishauser, and I. Filella. 2009. Phenology feedbacks on climate change. *Science* 324:887–888.
- Piao, S. L., J. Y. Fang, L. M. Zhou, P. Ciais, and B. Zhu. 2006. Variations in satellite-derived phenology in China's temperate vegetation. *Glob. Chang. Biol.* 12:672–685.
- Piao, S. L., X. H. Wang, P. Ciais, B. Zhu, T. Wang, and J. Liu. 2011. Changes in satellite-derived vegetation growth trend in temperate and boreal Eurasia from 1982 to 2006. *Glob. Chang. Biol.* 17:3228–3239.
- Pugh, T. A. M., A. R. MacKenzie, J. D. Whyatt, and C. N. Hewitt. 2012. Effectiveness of green infrastructure for improvement of air quality in urban street canyons. *Environ. Sci. Technol.* 46:7692–7699.
- Schwartz, M. D. 1996. Examining the spring discontinuity in daily temperature ranges. *J. Clim.* 9:803–808.
- Schwartz, M. D. 1998. Green-wave phenology. *Nature* 394:839–840.
- Seto, K. C., and J. M. Shepherd. 2009. Global urban land-use trends and climate impacts. *Curr. Opin. Environ. Sustain.* 1:89–95.
- Seto, K. C., B. Güneralp, and L. R. Hutyrá. 2012. Global forecasts of urban expansion to 2030 and direct impacts on biodiversity and carbon pools. *Proc. Natl Acad. Sci. USA* 109:16083–16088.
- Sims, D. A., A. F. Rahman, V. D. Cordova, B. Z. El-Masri, D. D. Baldocchi, L. B. Flanagan, et al. 2006. On the use of MODIS EVI to assess gross primary productivity of North American ecosystems. *J. Geophys. Res. Biogeosci.* 111:2005–2012.
- Steltzer, H., and E. Post. 2009. Seasons and life cycles. *Science* 324:886–887.
- Susca, T., S. R. Gaffin, and G. R. Dell'Osso. 2011. Positive effects of vegetation: urban heat island and green roofs. *Environ. Pollut.* 159:2119–2126.
- Suzuki, R., T. Nomaki, and T. Yasunari. 2003. West-east contrast of phenology and climate in northern Asia revealed using a remotely sensed vegetation index. *Int. J. Biometeorol.* 47:126–138.
- United Nations 2013. World urbanization prospects, the 2012 revision. United Nations, Department of Economic and Social Affairs, Population Division, New York.
- Wang, X. H., S. L. Piao, P. Ciais, J. S. Li, P. Friedlingstein, C. Koven, et al. 2011. Spring temperature change and its implication in the change of vegetation growth in North America from 1982 to 2006. *Proc. Natl Acad. Sci. USA* 108:1240–1245.
- White, M. A., and A. R. Nemani. 2003. Canopy duration has little influence on annual carbon storage in the deciduous broad leaf forest. *Glob. Chang. Biol.* 9:967–972.
- White, M. A., P. E. Thornton, and S. W. Running. 1997. A continental phenology model for monitoring vegetation responses to interannual climatic variability. *Global Biogeochem. Cycles* 11:217–234.
- White, M. A., S. W. Running, and P. E. Thornton. 1999. The impact of growing-season length variability on carbon assimilation and evapotranspiration over 88 years in the eastern US deciduous forest. *Int. J. Biometeorol.* 42:139–145.
- White, M. A., R. R. Nemani, P. E. Thornton, and S. W. Running. 2002. Satellite evidence of phenological differences between urbanized and rural areas of the eastern United States deciduous broadleaf forest. *Ecosystems* 5:260–273.
- White, M. A., K. M. de Beurs, K. Didan, D. W. Inouye, A. D. Richardson, O. P. Jensen, et al. 2009. Intercomparison, interpretation, and assessment of spring phenology in North America estimated from remote sensing for 1982–2006. *Glob. Chang. Biol.* 15:2335–2359.
- Wu, X. C., and H. Y. Liu. 2013. Consistent shifts in spring vegetation green-up date across temperate biomes in China, 1982–2006. *Glob. Chang. Biol.* 19:870–880.
- Yang, Y., H. Guan, M. Shen, W. Liang, and L. Jiang. 2015. Changes in autumn vegetation dormancy onset date and the climate controls across temperate ecosystems in China from 1982 to 2010. *Glob. Chang. Biol.* 21:652–665.
- Zhang, X. Y., M. A. Friedl, C. B. Schaaf, A. H. Strahler, and A. Schneider. 2004. The footprint of urban climates on vegetation phenology. *Geophys. Res. Lett.* 31:12.
- Zhou, L. M., C. J. Tucker, R. K. Kaufmann, D. Slayback, N. V. Shabanov, and R. B. Myneni. 2001. Variations in northern vegetation activity inferred from satellite data of vegetation index during 1981 to 1999. *J. Geophys. Res. Atmos.* 106:20069–20083.
- Zhu, W., H. Tian, X. Xu, Y. Pan, G. Chen, and W. Lin. 2012. Extension of the growing season due to delayed autumn over mid and high latitudes in North America during 1982–2006. *Glob. Ecol. Biogeogr.* 21:260–271.

## Supporting Information

Additional Supporting Information may be found in the online version of this article:

**Appendix S1.** Selection of study cities.

**Appendix S2.** Determining non-developed land-uses within the surrounding rural zones.

**Table S1.** Characteristics of Britain's largest 'Urban Areas', in descending order of size, from the Office for National Statistics (ONS 2005).

**Table S2.** Linear regression models used to explore changes in the start of the growing season (SOS), end of the growing season (EOS), and length of the growing sea-

son (LOS), between 2000 and 2009, in Britain's 15 largest cities (listed in order of ascending latitude).

**Table S3.** Location and urban form characteristics Britain's 15 largest cities (listed in order of ascending latitude).

**Table S4.** For each individual city (listed in order of ascending latitude), vegetation growing season median start and end Julian days, and length of growing season (SOS, EOS and LOS respectively).