

Kent Academic Repository

Full text document (pdf)

Citation for published version

Geeves, Michael A. (2016) The ATPase mechanism of myosin and actomyosin. *Biopolymers*, 105 . pp. 483-491. ISSN 0006-3525.

DOI

<https://doi.org/10.1002/bip.22853>

Link to record in KAR

<http://kar.kent.ac.uk/55320/>

Document Version

Author's Accepted Manuscript

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>



The ATPase mechanism of myosin and actomyosin.

Journal:	<i>Biopolymers</i>
Manuscript ID	Draft
Wiley - Manuscript type:	Invited Review
Date Submitted by the Author:	n/a
Complete List of Authors:	Geeves, Michael; University of Kent School of Biosciences, School of Biosciences
Keywords:	ATPase mechanism, Molecular motor, transition states

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The ATPase mechanism of myosin and actomyosin.

Michael A Geeves

School of Biosciences, University of Kent, Canterbury, UK

Michael A Geeves
School of Biosciences,
University of Kent,
Canterbury, CT1 7NJ
UK

m.a.geeves@kent.ac.uk

Tel +44 1227 827597

Abstract

Myosins are a large family of molecular motors that use the common P-loop, Switch 1 and Switch 2 nucleotide binding motifs to recognise ATP, to create a catalytic site than can efficiently hydrolyse ATP and to communicate the state of the nucleotide pocket to other allosteric binding sties on myosin. The energy is of ATP binding is used to do work against an external load. In this short review I will outline current thinking on the mechanism of ATP hydrolysis and how the energy of ATP hydrolysis is coupled to a series of protein conformational changes that allow a myosin, with the cytoskeleton track actin, to operate as a molecular motor of distinct types.

1
2
3 Myosin was identified more than 100 years ago in muscle where it is the protein that drives muscle
4 contraction using ATP as the energy source^{1,2}. In the that last 25 years knowledge of the myosin
5 family has grown explosively and now includes at least 35 different classes in the myosin superfamily
6 found across all eukaryotic cells³. Myosins are a large family of ATP driven motor proteins which
7 use the cytoskeleton protein actin as a track along which the myosin motor can move cargos or
8 generate and sense mechanical forces in the cytoskeleton^{4,5}. Mammalian cells will typically express
9 more than a dozen myosins from at least 5 different classes and they are involved in a wide range of
10 actin cytoskeleton based activities from organising the cytoskeleton, transporting material around
11 the cell, generating forces and sensing forces in the cytoskeleton^{6,7}.

12
13
14
15
16
17
18 Myosins are defined by the presence of a conserved globular motor domain of about 90 kDa (Figure
19 1). The domain is roughly pear or tear-drop in shape and is distinguished by a large cleft that spits
20 the broad end of the molecule and a single α -helix stabilised by light chains at the narrow end. The
21 actin binding sites span either side of the large cleft and closing of the cleft is required for tight
22 binding to actin. The binding sites for the ATP phosphate groups are near the bottom of the cleft
23 where the familiar *P-loop* and *switch-1* and *switch-2* loops (SW1 & SW2) are found and identify
24 myosins as part of the broader family of the P-Loop containing ATP and GTPases. ATP binding and
25 ADP release does not occur via this large cleft but via a pocket on the side of the molecule. The Pi
26 groups enter the pocket first with the nucleoside blocking the exit of hydrolysis product Pi via the
27 pocket.

28
29
30
31
32
33
34
35 The C-terminus of the motor domain, consists of a long α -helix stabilised by calmodulin or
36 calmodulin-like light chains (between 2 and 6 such light chains depending upon the isoform). This
37 light chain binding domain (LCBD) is also known as the lever arm because it swings through and
38 angle of $\sim 60^\circ$ during the ATP cycle thereby amplifying small domain movements within the motor
39 domain to produce 5-20 nm movements of the tip of the lever relative to the actin binding sites.

40
41
42
43
44 The C-terminus of the motor domain extends into a further domain known as the cargo binding
45 domain. This region is highly variable even within family groups and comprises a combination of
46 protein and lipid binding motifs (the cargo binding motifs) and/or dimerization domains that can also
47 allow assembly into higher order myosin filaments^{4,8}.

51 Myosin ATPase cycle

52
53
54 The normal physiological substrate of myosin is Mg·ATP but unusually myosin can be defined as a
55 Mg²⁺ inhibited ATPase, since in the complete absence of Mg²⁺ myosins have an unusually high
56 turnover rate (eg $\sim 10 \text{ s}^{-1}$ for muscle myosin 2) that is reduced to 0.05 s^{-1} when Mg²⁺ is present. This
57
58
59
60

1
2
3 is because the products of the ATPase reaction, Mg-ADP and Pi, are both tightly bound in the ATP
4 site and an interaction with actin is normally required to release the products. The outline of the
5 myosin ATPase (and actin.myosin) cycle has been known for 40 years (Bagshaw & Trentham 1976,
6 Geeves 1991) but some details remain elusive in particular the catalytic mechanism of hydrolysis has
7 been debated continually. The following outline is that appropriate for the muscle myosin II but the
8 outline varies little between this and other myosin forms that have been studied. In its basic form
9 the cycle begins when Mg-ATP binds myosin rapidly and almost irreversibly followed by rapid
10 reversible hydrolysis of ATP to the tightly bound Mg-ADP-Pi. The low free energy change for the
11 hydrolysis step and the slow release of ATP or Pi from this complex indicates the stability of the two
12 forms, with reactant and products both tightly bound in the myosin pocket. In the absence of actin
13 the bound Pi has an average lifetime of ~20 sec. Pi escape is followed by a more rapid release of
14 Mg-ADP. The release of ADP is linked to Mg²⁺ release but whether they are release together or
15 sequentially, Mg²⁺ escapes first then promoting ADP release, appears to vary between myosin
16 isoforms (^{9 10 11} and may be linked to the precise role of ADP release in different classes of myosin¹².
17 However, the rate of product release observed experimentally is often much faster than that
18 occurring physiologically as there are additional regulation events that slow Pi release even further
19 in the absence of available actin sites. These vary across the family, but for some types of myosin
20 folding of two myosin heads together (e.g. smooth muscle myosin II¹³) or of motor and cargo
21 domains together (myosin V¹⁴) or stabilising motor and filaments structures¹⁵ to further
22 trap/stabilise the M-ADP-Pi complex are common. This is one of the differences in the myosin
23 compared to many G-proteins (or even kinesin). M-ADP-Pi is the stable complex that sits and waits
24 for a signal to complete its biological role and that signal can come either via the actin (making actin
25 site available) or via the myosin – relieving inhibition of the motor domain. There are exceptions
26 (and across the 35 members of the myosin family there are plenty of exceptions to most rules) and
27 for some myosin involved in mechanical signalling the mechanically loaded A-M-ADP complex is a
28 long lived stable complex and the loss of load leads to rapid release of ADP^{16 17 18}.

29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46 The complexes of myosin with ATP and ADP.Pi are stable and therefore might be expected to
47 crystalize but the rapid reversibility of the hydrolysis step (which is accompanied by a large change
48 of conformation) makes this largely unsuccessful. However, crystals of myosin with putative
49 transition state analogues, such as ADP-AlF₄, ADP-BeF₃, and ADP-Vi have all been crystallised^{19 20}. In
50 addition to complexes with inhibitors such as blebbistatin bound to the motor domain have given
51 insights into the catalytic mechanism of ATP hydrolysis²¹ and how the conformational changes
52 accompanying ATP binding, hydrolysis and ordered product release couple the ATPase cycle to the
53 biological role of different myosins. These are discussed in the section below.
54
55
56
57
58
59
60

1
2
3 The key point of relevance to the biological role is that ATP binds to myosin – and the combination
4 of the binding energy and hydrolysis step primes the myosin into a conformation, which, while
5 stable, is ready to interact with actin to complete mechanical work or drive mechanical signals
6 through the cytoskeleton.
7
8

9 10 Actin activated myosin ATPase and the mechanical cycle

11
12 A simple cartoon outline of the major events and conformation changes in the actin-myosin ATPase
13 cycle is outlined in Figure 2 and described in detail in the legend. Here it can be seen that the role of
14 ATP is to bind to myosin, force apart the major cleft and thereby reduce the affinity of myosin for
15 actin by about 1000 fold, then hydrolyse ATP. These activities are linked at the simplest level with
16 the closing of the two switch elements onto the Pi of ATP. On ATP binding SW1 closes first and is
17 coupled with the opening of the major cleft and hence actin dissociation. SW2 closure follows and is
18 linked to the *recovery stroke* in which the *converter* domain and *lever* are repositioned into the *pre-*
19 *power* or *working-stroke* conformation. In addition to any structural constraint that require SW1 to
20 close before SW2 the time of events also ensures the ordering of events. SW1 closure and actin
21 dissociation takes about ~1 msec at saturating ATP for a fast muscle myosin and the hydrolysis step
22 occurs about 10 times slower in ~10 msec. The resulting complex is stable for ~20 sec unless it
23 encounters an actin binding site which results in the completion of the rest of the cycle in ~20 msec
24
25
26
27
28
29
30
31
32

33 The molecular details of the nucleotide-free actin·myosin complex, the detached M·ATP and the
34 post-recovery M·ADP·Pi conformations are well defined from crystal structures^{22,23} and high
35 resolution EM images²⁴ and such images have been used in molecular dynamic simulations generate
36 plausible models of the transitions between defined structural states^{25,26,27}. The transition between
37 the detached M·ADP·Pi state back to the A·M state remain more problematic and this, frustratingly,
38 is of course the business end of the cycle where myosin can generate force and movement. As set
39 out in the cartoon (Fig 2) actin binding is accompanied by Pi release, myosin cleft closure and force
40 generation but which order these events take place in remains hotly debated. These are drawn as
41 all occurring on a single step, step 3. But the order is important for understanding the way the
42 energy of the event is efficiently coupled to force and movement. There are several possible
43 scenarios but evidence is hard to generate because such structures are going to be short-lived,
44 possibly unstable and the mechanical forces involved are large at the molecular scale and may well
45 influence the conformation route taken between M·ADP·Pi and A·M. Most structural methods work
46 on protein in the absence of any external load and under such conditions all of the molecular events
47 consolidated in step 3 are instantaneously followed by transition via route 4a/5a to the stable A·M·D
48 complex that can easily be formed by addition of ADP to the stable A·M complex. The more
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 interesting route is that via step 4 and 5 where the load on the motor domain (and the external work
4 being done) slows the transition through the molecular events. The load-induced slowing of the
5 cycle leads to the classical force-velocity relationship observed when the recording forces and
6 movements generated by an ensemble of motor molecules working together against a load. That is
7 the rate of the ATPase cycle slows as the opposing load increases to a point where the ensemble of
8 motors continues to use ATP but can no longer move the load. A possible outline of events and
9 some of the alternatives are described below
10
11
12
13

14
15 Actin rebinding to M.ADP.Pi initiates the process of product release and force generation and the
16 current view favours the lower 50 kDa domain making the important hydrophobic inter face with
17 actin– in particular the contact of the activation loop (Dictyostelium Myosin II residues ⁵¹⁹GRQPP₅₂₃)
18 has been shown to have a role in promoting Pi release ²⁸. But does Pi release occur before or after
19 cleft closure and forming the complete actin myosin interface? Two views clash here: crystal
20 structures indicate the most straight forward route for Pi to be released from the pocket (since ADP
21 blocks Pi from escaping back the way it came in when attached to ATP) is through a *backdoor* which
22 allows a Pi route out via the cleft. When the cleft closes this route is no longer accessible the
23 backdoor is closed ²²²⁹. Recent mutational studies attempting to block this back door have
24 supported such an idea³⁰. The counter view is that other routes are possible and breathing motions
25 of the structure, SW1 opening – and mechanical loading across the domains may aid such alternate
26 exit routes ²³³¹. The argument against Pi release before cleft closure is that since Pi escape is linked
27 to force generation it does not seem to make sense to generate force at the actin-myosin interface
28 until the majority of the actin myosin interface is in place. A partially formed interface will be
29 weaker and more prone to mechanical failure. Recent fluorescent probe studies from the Thomas
30 lab ^{3233 34} indicate that in the absence of actin the conformation of the myosin is dynamic and actin
31 traps the post power stroke conformation and this precedes Pi release, a slower process. The
32 events surrounding step 3 therefore remain under-defined and until there is more experimental
33 evidence the differing viewpoints will remain strongly held by different experimental camps.
34
35
36
37
38
39
40
41
42
43
44
45

46
47 The one clear undisputed result is that once the myosin has completed its working stroke Pi release
48 is irreversible but while force is being generated –and the energy not dissipated, Pi can bind
49 reversibly into the site and the cycle is freely reversible back through steps 3 and 2 (see Fig 2) to
50 form tightly bound ATP. This is illustrated in Figure 2. As the Pi is released in step 3 the blue circle
51 rotates and generates a force against an external load attached to the tail (indicated by the bent tail)
52 while the tail remains bent Pi can rebind into the pocket, but as soon as the tail moves the load
53 forward and the tail straightens against the load, the energy has been dissipated and Pi can no
54
55
56
57
58
59
60

1
2
3 longer reverse the process. The domain rotates further in the absence of load, further preventing Pi
4 rebinding and allowing ADP release.
5

6
7 The role of ADP release is much better understood and occurs after the Pi release and force
8 generating events. The key role here is for ADP to remain in the pocket until the “working stroke”
9 has been completed. That is, it remains in the pocket as long any force generated by or
10 accompanying Pi release has been dissipated (step 4). This makes the efficiency of the system very
11 high and prevents futile burning of ATP. Any myosin designed to have very long-lived force holding
12 states have relatively slow rates of ADP release and or a strong load dependence of the rate
13 constant of ADP release¹²¹⁸³⁵. This aspect is discussed in more detail below where the adaptations of
14 myosin to different mechanical functions is outlined. In essence ADP remains in the pocket while
15 the energy from ATP breakdown can still be used. Once ADP is released, the high concentration of
16 cellular ATP ensures very rapid rebinding of ATP and dissociation from actin losing any mechanical
17 (force or strain sensing) contact with the actin cytoskeleton.
18
19
20
21
22
23
24

25 Catalytic Mechanism of Hydrolysis

26
27
28 The crystal structure of the ATP and ADP.Pi analogues revealed that the active site was not fully
29 formed until the SW1 and SW2 elements closed down on to the γ -Pi of ATP. The structures then
30 allow identification of residues likely to be involved in the catalytic mechanism. Despite several
31 different crystal structures of putative transition state analogues and extensive kinetic studies of the
32 hydrolysis step, two mechanism have remained possible: An associative (direct attack by water via a
33 penta-covalent γ -Pi) or a dissociative mechanism (attack of water on a stabilised meta-phosphate
34 intermediate) with opinion swinging in either direction over time. The most recent contribution to
35 the debate has been from a series of quantum mechanical/ molecular mechanics calculations
36 (QM/MM) of the hydrolysis pathway based on crystal structures of analogues of the M.ATP structure
37 (predominantly an ADP vanadate structure)and the transition states using data from a range of
38 different myosin types³⁶³⁷. A summary of this recent work view is presented here and the reader is
39 invited to explore a recent accessible summary for a detailed evaluation of the evidence³⁸.
40
41
42
43
44
45
46
47

48 In the following description residue numbers refer to those of Dictyostelium myosin II. The
49 phosphates of Mg^{2+} ATP are tightly bound through interactions with the three loops. α - & β - Pi
50 contact the P-loop (¹⁷⁹GESGAGKTEN₁₈₇ makes 8 H-bonds formed mainly through peptide backbone
51 contacts), while the γ -Pi are stabilised via SW1 (²³⁶SSR₂₃₈) and SW2 (⁴⁵⁵ISGF₄₅₈) again primarily via
52 peptide H-bonds (see Fig 3A). The closed position of SW1 and SW2 is further stabilised by salt bridge
53 between Glu459 and Arg238. The primary catalytic side chains are Glu-459, Ser-181 and Ser236/7
54
55
56
57
58
59
60

1
2
3 and involves 3 water molecules. There are three elements to the mechanism envisaged (outlined in
4 Fig 3B)

5
6
7 – breaking of the β - γ bond phosphate bond between the γ -phosphorus and the β - γ bridging
8 oxygen and stabilisation of the resulting meta-phosphate group. A wide range of interaction with the
9 peptide backbone of SW1, SW2 and the P-loop contribute to this stabilisation of the metaphosphate
10 (listed above) and results in almost zero change in energy between the ATP and metaphosphate
11 states.
12

13
14
15
16 - a combination of Glu459 and Ser237 polarize an attacking water molecule allowing formation of
17 the OH⁻ anion which can more easily attack the stabilised metaphosphate.
18

19
20 - creation of a low energy pathway to transfer the abstracted proton to the γ -phosphate group via
21 so called *proton wires*. The contribution of Glu459 to the catalytic mechanism and the role of
22 protein wires have only been appreciated in the most recent simulations.
23
24

25
26 The combination of the three elements generates a pathway with small intermediate energy levels
27 (+2.7 to -3 kcal/mol) allowing reversibility and relatively low energy barriers between states (8.7-
28 10.3 kcal/mol) allowing high rates of transition consistent with experimental observations.
29
30

31 Overview of the ATPase & mechanical cycles

32
33
34 With the above outline of the actin-myosin ATPase mechanism the role of ATP in the overall
35 mechanism can be considered. Two viewpoints are essential – the thermodynamic and the kinetic.
36 How the binding energies and the energy of ATP hydrolysis is used to drive the work cycle - and how
37 the timing of events contributes the efficiency of the cycle.
38
39

40
41 The energy associated with ATP binding is essential to first of all displace the actin from the myosin,
42 itself very tightly bound with an affinity of ~ 10 nM. This is associated with SW1 closing and opening
43 the myosin cleft. This is followed by the re-priming or recovery-stroke returning the myosin tail into
44 the *up* position ready to complete the next working-stroke. This is associated with SW2 closing.
45
46 These two major conformational changes are triggered by SW1 then SW2 closing on to the γ -Pi and
47 in doing so create the active site that can stabilise the metaphosphate and attacking OH⁻. The whole
48 process is accompanied by very small free energy changes allowing each of these events to be
49 readily reversible – except actin rebinding to the M-ATP form. As stated above the actin dissociation
50 event is faster than the recovery-stroke/hydrolysis step ensuring the recovery stroke does not occur
51 while bound to actin which could result in a futile mechanical cycle.
52
53
54
55
56
57
58
59
60

1
2
3 The timing of the hydrolysis step is important for another reason. This sets the time during which the
4 myosin is unable (or unlikely) to rebind to an actin site – once M·ADP·Pi is formed it can rebind an
5 actin site and complete the ATPase cycle. Thus the lifetime of the M·ATP is important to ensure the
6 myosin can diffuse to its next productive actin site (reattaching to the site just detached from would
7 not produce net forward movement) and this varies for different types of myosin. It is noticeable
8 that myosins that spend most of their ATP cycle time attached to actin can have very fast ATP
9 hydrolysis steps (³⁹). The lifetime of M·ATP can partially control the *duty ratio* – the fraction of time
10 a myosin remains tightly bound to actin during a single ATPase cycle.

11
12
13
14
15
16 Like the ATP hydrolysis, the Pi release step and force generation occur with little change in free
17 energy and the step remains freely reversible. Only if the free energy is dissipated through the
18 completion of the working stroke does the Pi step become essentially irreversible and the system is
19 committed to completion of the ATP hydrolysis. Exactly how the energy of the system is stored, in
20 for example, a bending or stretching of an elastic component remains to be defined. Current
21 thinking focusses on some combination of the lever-arm bending and to central β -sheet distortion.
22 One issue is to understand if such a mechanically distorted state appears different at the single
23 molecule level vs an ensemble of myosin motors. Certainly in a sophisticated structure such as the
24 muscle sarcomere any mechanical force generated by a single myosin will rapidly dissipate into
25 various compliant elements of the super structure.

26
27
28
29
30
31
32
33
34 The ADP release step varies for different type of actin-myosin motor and can occur with zero,
35 positive or negative free energy changes. Bloemink & Geeves (2011)¹² outlined how the free energy
36 of ADP release can be modified in different myosins to generate motors of different types. A fast
37 ADP release event with a significant negative free energy change is characteristic of a motor
38 generating fast movement even under load. A slower motor with high efficiency for both
39 movement and force-holding actions has a slower ADP release and the rate of ADP release is readily
40 inhibited by load on the motor. A myosin designed for generation tension over long periods and
41 sensing tension changes in the cytoskeleton has a small or zero free energy associated with ADP
42 release and effectively ADP release is blocked by any small load on the motor. Finally if ADP release
43 occurs with a positive free energy change then ADP is retained by the motor unless it receives
44 additional information. This is similar to many G-proteins that require an exchange factor to release
45 GDP, and is observed in a two-headed processive motor such as myosin V; the ADP on the attached
46 head that has completed its working stroke remains in the trapped in nucleotide pocket. If he
47 second head attaches to actin and goes through its force generation step then it pulls on the rear
48 head opening up the exit route for ADP allowing ADP to be exchanged for ATP, the head then
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 detaches and the ATPase cycle restarts. Such a process ensures the heads work alternately and
4 allows the motor to *walk* hand-over-hand along an actin filament.
5
6

7 This flexible ADP release mechanism allows quite small changes in the free energy of ADP release to
8 tune the myosin for different types of mechanical function. Of course optimising each myosin for its
9 own function involves more than just changing the free energy of ADP release. Other factors that
10 are altered between myosin isoforms include the length of the lever arm (varying from 2 to 6
11 calmodulin binding domains) which can alter the size of the step – and thus on simple lever arm
12 theory adjusting the distance moved per ATP hydrolysed and the force developed in each working
13 stroke⁴⁰⁴¹. The overall speed of the cycle also allows for different speeds of movement and different
14 response times to different mechanical or chemical signals. And the duty ratio – the fraction of each
15 ATPase cycle time the motor spends strongly attached to actin – can be altered to change between
16 efficient force holders/sensors where the long lived A·M·D complexes can maintain force with little
17 ATP turnover but by reducing the response time to changes in force and being very slow transport
18 systems³⁹¹².
19
20
21
22
23
24
25
26

27 There remain many molecular details we do not fully understand (such as the exact details of Pi
28 release and force generation) but we are beginning to understand how the *design* of the myosin
29 motor has allowed the development of a range of motors tuned for different roles within the cell.
30 For each motor the basic ATPase cycle remains the same; the mechanical cycle is largely the same
31 and the mechanism of catalysis remains the same. What differs is how the cycle is tweaked
32 to alter the cycle time, to alter where in each cycle myosin spends most of its time, and to alter how
33 the energy from ATP hydrolysis is coupled to generate different types of mechanical activity. These
34 we are beginning to understand. Coupled to this is the question of how myosin activities are
35 regulated and integrated with the rest of cellular responses - but that as they say is a different story.
36
37
38
39
40
41
42
43
44

45 **References**

- 46
47 (1) Kuhne. *Untersuchungen über das Protoplasma und die Contractilität.*; Leipzig, 1864.
48
49 (2) Englehardt, W. A.; Ljubimowa, M. N. *Nature* **1939**, *144*, 668–669.
50
51 (3) Odronitz, F.; Kollmar, M. *Genome Biol* **2007**, *8* (9), R196.
52
53 (4) Krendel, M.; Mooseker, M. S. *Physiology (Bethesda)*. **2005**, *20* (4), 239–251.
54
55 (5) Coluccio, L. M. *Myosins. A super family of Molecular Motors*; Springer, Netherlands, 2008;
56 Vol. 7.
57
58 (6) Bement, W. M. *Proc. Natl. Acad. Sci.* **1994**, *91* (24), 11767c – 11767.
59
60

- 1
2
3 (7) Hartmann, M.; Spudich, J. A. *J Cell Sci* **125**, 1627–1632.
4
5 (8) Korn, E. D. *Proc. Natl. Acad. Sci. U. S. A.* **2000**, *97* (23), 12559–12564.
6
7 (9) Bagshaw, C. R.; Trentham, D. R. *Biochem. J.* **1974**, *141* (2), 331–349.
8
9 (10) Fujita-Becker, S.; Dürrwang, U.; Erent, M.; Clark, R. J.; Geeves, M. A.; Manstein, D. J. *J. Biol. Chem.* **2005**, *280* (7), 6064–6071.
10
11 (11) Hannemann, D. E.; Cao, W.; Olivares, A. O.; Robblee, J. P.; De La Cruz, E. M. *Biochemistry* **2005**, *44* (24), 8826–8840.
12
13 (12) Bloemink, M. J.; Geeves, M. A. *Seminars in Cell and Developmental Biology*. 2011, pp 961–967.
14
15 (13) Wendt, T.; Taylor, D.; Trybus, K. M.; Taylor, K. *Proc Natl Acad Sci U S A* **2001**, *98* (8), 4361–4366.
16
17 (14) Kremmentsov, D. N.; Kremmentsova, E. B.; Trybus, K. M. *J. Cell Biol.* **2004**, *164* (6), 877–886.
18
19 (15) Woodhead, J. L.; Zhao, F.; Craig, R.; Egelman, E. H.; Alamo, L.; Padrón, R. *Nature* **2005**, *436* (7054), 1195–1199.
20
21 (16) Veigel, C.; Molloy, J. E.; Schmitz, S.; Kendrick-Jones, J. *Nat. Cell Biol.* **2003**, *5* (11), 980–986.
22
23 (17) Greenberg, M. J.; Lin, T.; Goldman, Y. E.; Shuman, H.; Ostap, E. M. *Proceedings of the National Academy of Sciences*. 2012, pp E2433–E2440.
24
25 (18) Greenberg, M. J.; Shuman, H.; Ostap, E. M. *Biophys. J.* **2014**, *107* (12), L41–L44.
26
27 (19) Fisher, A. J.; Smith, C. A.; Thoden, J. B.; Smith, R.; Sutoh, K.; Holden, H. M.; Rayment, I. *Biochemistry* **1995**, *34* (28), 8960–8972.
28
29 (20) Sweeney, H. L.; Houdusse, A. *Annu. Rev. Biophys.* **2010**, *39*, 539–557.
30
31 (21) Allingham, J. S.; Smith, R.; Rayment, I. *Nat. Struct. Mol. Biol.* **2005**, *12* (4), 378–379.
32
33 (22) Sweeney, H. L.; Houdusse, A. *Annu Rev Biophys* **2010**, *39*, 539–557.
34
35 (23) Geeves, M. A.; Holmes, K. C. *Adv Protein Chem* **2005**, *71*, 161–193.
36
37 (24) Behrmann, E.; Mueller, M.; Penczek, P. A.; Mannherz, H. G.; Manstein, D. J.; Raunser, S. *Cell* **2012**, *150* (2), 327–338.
38
39 (25) Kühner, S.; Fischer, S. *Proc. Natl. Acad. Sci. U. S. A.* **2011**, *108* (19), 7793–7798.
40
41 (26) Fischer, S.; Windshügel, B.; Horak, D.; Holmes, K. C.; Smith, J. C. *Proc. Natl. Acad. Sci. U. S. A.* **2005**, *102* (19), 6873–6878.
42
43 (27) Cecchini, M.; Alexeev, Y.; Karplus, M. *Structure* **2010**, *18* (4), 458–470.
44
45 (28) Várkuti, B. H.; Yang, Z.; Kintszes, B.; Erdélyi, P.; Bárdos-Nagy, I.; Kovács, A. L.; Hári, P.; Kellermayer, M.; Vellai, T.; Málnási-Csizmadia, A. *Nat. Struct. Mol. Biol.* **2012**, *19* (3), 299–306.
46
47 (29) Preller, M.; Holmes, K. C. *Cytoskeleton* **2013**, *70* (10), 651–660.
48
49 (30) Llinas, P.; Isabet, T.; Song, L.; Ropars, V.; Zong, B.; Benisty, H.; Sirigu, S.; Morris, C.; Kikuti, C.;
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 Safer, D.; Sweeney, H. L.; Houdusse, A. *Dev. Cell* **2015**.
4
5 (31) Kull, F. J.; Endow, S. A. *J. Cell Sci.* **2013**, *126* (Pt 1), 9–19.
6
7 (32) Trivedi, D. V.; Muretta, J. M.; Swenson, A. M.; Davis, J. P.; Thomas, D. D.; Yengo, C. M. *Proc.*
8 *Natl. Acad. Sci.* **2015**, *112* (47), 201517566.
9
10 (33) Muretta, J. M.; Rohde, J. A.; Johnsrud, D. O.; Cornea, S.; Thomas, D. D. *Proc. Natl. Acad. Sci.*
11 **2015**, *2015*, 201514859.
12
13 (34) Muretta, J. M.; Petersen, K. J.; Thomas, D. D. *Proc. Natl. Acad. Sci. U. S. A.* **2013**, *110* (18),
14 7211–7216.
15
16 (35) Veigel, C.; Schmitz, S.; Wang, F.; Sellers, J. R. *Nat. Cell Biol.* **2005**, *7* (9), 861–869.
17
18 (36) Kiani, F. A.; Fischer, S. *Proc Natl Acad Sci U S A* **2014**, *111* (29), E2947–E2956.
19
20 (37) Grigorenko, B. L.; Kaliman, I. A.; Nemukhin, A. V. *J. Mol. Graph. Model.* **2011**, *31*, 1–4.
21
22 (38) Kiani, F. A.; Fischer, S. *Curr. Opin. Struct. Biol.* **2015**, *31*, 115–123.
23
24 (39) De La Cruz, E. M.; Ostap, E. M. *Curr Opin Cell Biol* **2004**, *16* (1), 61–67.
25
26 (40) Anson, M.; Geeves, M. A.; Kurzawa, S. E.; Manstein, D. J. *EMBO J.* **1996**, *15* (22), 6069–6074.
27
28 (41) Uyeda, T. Q.; Abramson, P. D.; Spudich, J. a. *Proc. Natl. Acad. Sci. U. S. A.* **1996**, *93* (April),
29 4459–4464.
30
31 (42) Rayment, I.; Holden, H. M.; Whittaker, M.; Yohn, C. B.; Lorenz, M.; Holmes, K. C.; Milligan, R.
32 a. *Science (New York, N.Y.)*. 1993, pp 58–65.
33
34
35 -----
36

37 *Banga, I. 1942. The phosphatase activity of myosin. Stud. Inst. Med. Chem. Univ. Szeged.*
38 *I:27–36.*
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Fig 1 Structure of the myosin motor domain
4
5

6 The post-rigor structure of the myosin motor domain ⁴². The structure of the myosin cross-bridge
7 shown as a ribbon diagram in the orientation it would take on binding to actin viewed from the
8 pointed (-) end of the actin filament. The N-terminus is shown green and the nucleotide binding P-
9 loop and adjoining helix are shown yellow; the upper 50K is red; the lower 50K domain is grey. Note
10 the cleft separating the upper and lower 50K domains. The lower 50K domain appears to be the
11 primary actin-binding site. The N-terminal boundary of the upper 50K domain comprises the
12 disordered loop 1 (between the points marked A and B). The upper and lower 50K domains are also
13 connected by a disordered loop (loop 2 between C and D). The C-terminal long helix (dark blue)
14 carries two calmodulin-like light chains and joins onto the thick filament. The relay helix and
15 converter domain are shown. In this conformation of the cross-bridge (post-rigor state) the lever
16 arm is in the post-power stroke position or "DOWN", as in the rigor state. The colouring corresponds
17 with sub-domain boundaries. For clarity, the proximal end of the relay helix is shown light blue
18 although it is actually part of the lower 50K domain. The distal end (beyond the kink) is firmly
19 attached to the converter domain. In the post-rigor structure the relay helix is straight (no kink).
20 Reproduced from Geeves & Holmes 2005 *Advances in Protein Chemistry* **71**, 161-193 (2005)²³
21
22
23
24
25
26
27
28
29
30
31
32

33 Fig 2 The ATP driven actin-myosin mechanical cycle
34

35 The myosin cross-bridge is shown as consisting of 3 major components. 1. The central core of the
36 myosin head is shown in red (filled circle) with the lower 50k domain projecting up to make contact
37 with one actin monomer (grey circles). The lower 50k domain is one half of the major cleft that splits
38 the actin binding site. The P-loop that binds to the γ -Pi of ATP is part of the central core. This red
39 segment is drawn as a fixed reference point during the cycle. 2. The upper 50 K domain with SW-1
40 (switch 1) is shown in yellow. This is drawn as two parts; the upper jaw of the major cleft that splits
41 the actin binding site and a yellow ring with SW-1 projecting into the center. 3. A blue ring
42 represents the relay loop with SW-2 projecting to the center while the converter domain and the
43 lever arm (shown with 2 light chains attached) is projected out from the ring. The nucleotide pocket
44 is shown as lying on top of the central core (red) bounded by two circles one part of the upper 50 K
45 domain (yellow) the other part of the relay/converter domain (blue). Each ring has a segment
46 missing to represent the entrance to the nucleotide pocket.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 A·M at the top left represents the rigor actin-myosin complex with the upper 50kDa domain cleft
4 closed to allow both sides of the cleft to make contact with actin. The nucleotide pocket entrance is
5 open with both SW1 & 2 open.
6

7
8 Step 1 ATP binds to myosin and myosin dissociates from actin. In detail ATP binds into the nucleotide
9 pocket and the yellow segment rotates to close SW-1 onto the ATP in doing so the major cleft opens
10 destroying the actin binding site leading to dissociation from actin.
11

12
13 Step 2. The recovery stroke and ATP-hydrolysis. The blue segment rotates to bring SW-2 into contact
14 with ATP thus rotating the converter/light-chain binding domain to complete the recovery stroke or
15 re-priming of the motor while detached from actin. Only after both SW1 and SW2 are closed is ATP
16 hydrolysed to form the stable M·ADP·Pi complex.
17

18
19 Step 3 Actin rebinding and the power stroke. After hydrolysis the lower 50kDa part of the cleft
20 rebinds to actin, the cleft closes and both upper and lower 50kDa domains bind actin, this involves a
21 rotation of the yellow segment which triggers both Pi release and the rotation of the blue segment
22 generating a force – represented as a distortion of the converter domain. The exact order of cleft
23 closure, Pi release and power stroke remains under debate. The location of the “elasticity” within
24 the cross bridge is not defined.
25
26

27
28 Step 4 “Sliding”. Provided the force generated in step 3 is large enough the load is moved 5-10 nm
29 by the cross bridge working-stroke. This is shown as a relative displacement of the end of the lever
30 relative to the actin.
31

32
33 Step 5 Opening of the nucleotide pocket. Once the strain in the converter is dissipated by the sliding
34 movement, the blue segment rotates further to open the nucleotide pocket and allows ADP to
35 escape in step 6. An efficient motor requires that ADP release be limited until the cross-bridge has
36 completed its movement or working stroke.
37
38

39
40 Step 4a/5a. The strain holding cross-bridge. If the force generated in step 3 is too small to move the
41 load no sliding takes place. The further movement of the blue segment against the load is strongly
42 inhibited (~5-100 fold for different myosins). The degree to which this rotation is inhibited in a load
43 bearing myosin defines the load sensitivity of the myosin. This is a function of the size of the
44 rotation, the length of the lever arm and the stiffness of the “spring” in the structure.
45
46
47
48

49
50 Reproduced from: Bloemink & Geeves (2011) Seminars in Cell and Developmental Biology 22, 961-
51 967¹²
52

1
2
3 Fig 3 A Stabilisation of the 3 phosphates in the myosin nucleotide pocket.

4
5 The ATP binding site in the catalytically active conformation, i.e. switch 1 and switch 2 closed and the
6 stabilizing salt bridge formed between Arg 238 (of SW1) and Glu 459 (of SW2). ATP has been
7 modelled into the 1VOM crystal structure (Dyctyostelium myosin II with Mg^{2+} .ADP and vanadate in
8 the active site). Only the three phosphates are depicted. Dotted lines show hydrogen bonds shorter
9 than 2.8 Å between the heavy atoms. The Mg^{2+} coordination sphere is represented with thick grey
10 lines. Taken from Kiani & Fischer PNAS 2014 11, E2947-E2956³⁶.

11
12
13
14
15
16
17
18 Fig 3B The catalysis of the ATP hydrolysis step and the role of Glu459.

19
20 Reactions on the left: during the breaking of water W_a , a proton is first transferred onto Glu459
21 before being transferred to the final γ -Pi. Reaction on the right: the proton is transferred onto the γ -
22 Pi without transiting Glu459. a) ATP reactant state **R**. Arrows with black heads show covalent
23 rearrangements. b & c) Metaphosphate intermediate **m**. d) Stable intermediate **e** with protonated
24 Glu459. E) Product precursor **g**, with proton on γ -Pi. Arrows with white heads indicate
25 rearrangements of the H-bond network. f) Final ADP·Pi product **P**.

26
27
28
29
30 Numbers in parentheses indicate the energy (kcal/mol, relative to the reaction state R) of the stable
31 intermediates. Numbers on the large arrows between boxes indicate the energy barriers between
32 states. Taken from Kiani & Fischer Current Opinion in Structural Biology Volume 31, 1-140³⁸




Figure 1
159x111mm (220 x 220 DPI)




Figure 2
264x295mm (96 x 96 DPI)




Figure 3A
121x105mm (96 x 96 DPI)




Figure 3B
188x213mm (96 x 96 DPI)