

Kent Academic Repository

Full text document (pdf)

Citation for published version

Matechou, Eleni and McCrea, Rachel S. and Morgan, Byron J. T. and Nash, Darryn and Griffiths, Richard A. (2016) Open models for removal data. *Annals of Applied Statistics*. ISSN 1932-6157.

DOI

<https://doi.org/10.1214/16-AOAS949>

Link to record in KAR

<http://kar.kent.ac.uk/55734/>

Document Version

Publisher pdf

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

OPEN MODELS FOR REMOVAL DATA

BY ELENI MATECHOU, RACHEL S. MCCREA¹, BYRON J. T. MORGAN,
DARRYN J. NASH AND RICHARD A. GRIFFITHS

University of Kent

Individuals of protected species, such as amphibians and reptiles, often need to be removed from sites before development commences. Usually, the population is considered to be closed. All individuals are assumed to (i) be present and available for detection at the start of the study period and (ii) remain at the site until the end of the study, unless they are detected. However, the assumption of population closure is not always valid. We present new removal models which allow for population renewal through birth and/or immigration, and population depletion through sampling as well as through death/emigration. When appropriate, productivity may be estimated and a Bayesian approach allows the estimation of the probability of total population depletion. We demonstrate the performance of the models using data on common lizards, *Zootoca vivipara*, and great crested newts, *Triturus cristatus*.

1. Introduction. When protected species occur on a site scheduled for development, there may be a legal requirement for them to be relocated to an alternative site before the development can proceed [Germano et al. (2015)]. In the UK, such relocations are often required for amphibians and reptiles. These relocations are typically achieved by conducting regular surveys during which detected animals are relocated to a suitable alternative habitat. We refer to collection and removal as translocation. The duration of the translocation is determined by the predicted size of the community and/or constituent species present. Current guidelines (developed 18 years ago) state that translocations should last between 60 and 120 “suitable” days [HGBI (1998)]. After the prescribed period has elapsed, a minimum of five trap-free days is required before the translocation is considered to be complete. Although it is acknowledged that some individuals may be left behind, this is considered to be a “reasonable effort” in line with current UK legislation. More recent guidance is less prescriptive, but suggests survey effort may need to last for 1–3 years [Natural England/Defra (2015)].

Our proposed methods were motivated by removal sampling data of the kind illustrated in Tables 1 and 2. The species surveyed are respectively common lizards,

Received December 2015; revised May 2016.

¹Supported in part by NERC Fellowship Grant NE/J018473/1.

Key words and phrases. Common lizard, depletion, great crested newts, RJMCMC, stopover model.

TABLE 1

Removal sampling: Counts of common lizards removed in 2007. A dash indicates that no sampling took place on that day. Temp denotes the maximum daily temperature (°C) recorded on each visit to the study site. Here, the number of removed individuals, $D = 213$

Date	Count	Temp	Date	Count	Temp	Date	Count	Temp	Date	Count	Temp
24-May	0	20	02-Jul	2	15	10-Aug	7	19	18-Sep	6	16
25-May	–	–	03-Jul	7	17	11-Aug	–	–	19-Sep	6	16
26-May	–	–	04-Jul	3	16	12-Aug	–	–	20-Sep	6	14
27-May	–	–	05-Jul	1	15	13-Aug	12	18	21-Sep	4	14
28-May	–	–	06-Jul	1	15	14-Aug	8	20	22-Sep	–	–
29-May	–	–	07-Jul	–	–	15-Aug	1	18	23-Sep	4	14
30-May	–	–	08-Jul	–	–	16-Aug	10	14	24-Sep	1	14
31-May	–	–	09-Jul	1	16	17-Aug	9	16	25-Sep	5	11
01-Jun	0	22	10-Jul	–	–	18-Aug	–	–	26-Sep	5	12
02-Jun	–	–	11-Jul	–	–	19-Aug	–	–	27-Sep	0	12
03-Jun	–	–	12-Jul	2	17	20-Aug	6	17	28-Sep	2	12
04-Jun	–	–	13-Jul	–	–	21-Aug	–	–	29-Sep	–	–
05-Jun	1	21	14-Jul	–	–	22-Aug	–	–	30-Sep	1	14
06-Jun	0	21	15-Jul	–	–	23-Aug	–	–	01-Oct	3	16
07-Jun	2	21	16-Jul	0	18	24-Aug	–	–	02-Oct	0	16
08-Jun	0	23	17-Jul	0	16	25-Aug	–	–	03-Oct	1	16
09-Jun	–	–	18-Jul	3	18	26-Aug	–	–	04-Oct	0	16
10-Jun	–	–	19-Jul	–	–	27-Aug	–	–	05-Oct	0	16
11-Jun	1	26	20-Jul	0	13	28-Aug	–	–	06-Oct	0	16
12-Jun	1	20	21-Jul	–	–	29-Aug	–	–	07-Oct	–	–
13-Jun	–	–	22-Jul	–	–	30-Aug	–	–	08-Oct	0	16
14-Jun	–	–	23-Jul	1	18	31-Aug	–	–	09-Oct	1	14
15-Jun	0	13	24-Jul	–	–	01-Sep	–	–	10-Oct	1	15
16-Jun	–	–	25-Jul	0	16	02-Sep	9	18	11-Oct	1	15
17-Jun	–	–	26-Jul	–	–	03-Sep	–	–	12-Oct	0	16
18-Jun	–	–	27-Jul	–	–	04-Sep	–	–	13-Oct	1	16
19-Jun	–	–	28-Jul	–	–	05-Sep	1	18	14-Oct	–	–
20-Jun	1	18	29-Jul	–	–	06-Sep	8	17	15-Oct	0	16
21-Jun	2	19	30-Jul	–	–	07-Sep	2	18	16-Oct	0	12
22-Jun	3	16	31-Jul	–	–	08-Sep	–	–	17-Oct	0	15
23-Jun	–	–	01-Aug	1	18	09-Sep	11	18	18-Oct	0	15
24-Jun	–	–	02-Aug	0	15	10-Sep	7	18	19-Oct	0	13
25-Jun	2	15	03-Aug	1	15	11-Sep	9	18	20-Oct	–	–
26-Jun	–	–	04-Aug	–	–	12-Sep	1	18	21-Oct	–	–
27-Jun	2	15	05-Aug	–	–	13-Sep	1	17	22-Oct	0	12
28-Jun	2	17	06-Aug	1	16	14-Sep	5	18	23-Oct	0	13
29-Jun	5	18	07-Aug	1	18	15-Sep	–	–	24-Oct	0	9
30-Jun	–	–	08-Aug	4	20	16-Sep	–	–	25-Oct	0	19
01-Jul	–	–	09-Aug	3	20	17-Sep	4	12			

TABLE 2

Removal sampling: Counts of male great crested newts removed in 2010. A dash indicates that no sampling took place on that day. Temp denotes the minimum daily air temperature (°C) recorded on each visit to the study site. Here, the number of removed individuals, $D = 741$. The nine missing covariate values were imputed using the average of the previous and next observed values

Date	Count	Temp	Date	Count	Temp	Date	Count	Temp
18-Mar	65	7	13-Apr	0	2.8	09-May	1	7.1
19-Mar	115	5.2	14-Apr	0	6.5	10-May	0	9
20-Mar	185	12.4	15-Apr	0	1.1	11-May	0	-0.7
21-Mar	49	6.5	16-Apr	0	-	12-May	0	2.2
22-Mar	64	3.6	17-Apr	0	3.5	13-May	0	13.2
23-Mar	9	2.3	18-Apr	-	-	14-May	1	0.6
24-Mar	38	5.2	19-Apr	-	-	15-May	1	6.9
25-Mar	102	12.4	20-Apr	-	-	16-May	0	1.4
26-Mar	16	5.2	21-Apr	0	-1	17-May	4	1.4
27-Mar	7	2.7	22-Apr	0	-3.2	18-May	0	-
28-Mar	5	3.1	23-Apr	0	-4.2	19-May	1	2.2
29-Mar	8	10.8	24-Apr	0	-4	20-May	1	10.8
30-Mar	22	-	25-Apr	0	6	21-May	0	9.3
31-Mar	0	3.1	26-Apr	16	4.2	22-May	-	-
01-Apr	2	-	27-Apr	2	4	23-May	-	-
02-Apr	-	-	28-Apr	1	7	24-May	1	-
03-Apr	-	-	29-Apr	3	4.3	25-May	0	-
04-Apr	-	-	30-Apr	3	7.7	26-May	0	18
05-Apr	-	-	01-May	4	5.1	27-May	0	10.3
06-Apr	3	-	02-May	1	8.2	28-May	0	-
07-Apr	3	3.7	03-May	0	4.2	29-May	0	15
08-Apr	2	2.1	04-May	0	2.5	30-May	0	11.5
09-Apr	0	1.3	05-May	1	2	31-May	0	-
10-Apr	4	1.8	06-May	1	4.9	01-Jun	0	5.1
11-Apr	0	3.2	07-May	0	3	02-Jun	0	3.9
12-Apr	0	6.1	08-May	0	3	03-Jun	0	3.9

Zootoca vivipara, and great crested newts, *Triturus cristatus*. Both of these are protected species which are frequently removed from the path of development, giving rise to data of the form illustrated.

The original model for removal sampling dates back to Moran (1951) and Zippin (1956). The model is the same as model M_b in the Otis notation for closed population models [McCrea and Morgan (2014), Chapter 3], which allows for behavioural response to trapping with recapture probability constrained to zero. Under this model, expected numbers of individuals decrease geometrically over time as a consequence of the assumptions of closure and constant capture probability. We refer to this as the geometric model. Recent work on removal sampling includes Bohrmann and Christman (2013), who discuss sampling design, and Dorazio, Jelks and Jordan (2005), who present a hierarchical model for when

the sample site can be considered to be made up of several distinct subsites distributed spatially, with independent counts taken at each site. In addition, [Dorazio, Jelks and Jordan \(2005\)](#) allow for heterogeneity in capture using a beta-binomial distribution. We note in passing that such an extension to the basic removal model corresponds to models for digit preference in fecundability studies when the total number of individuals in the study is unknown; see [Ridout and Morgan \(1991\)](#). [Dorazio et al. \(2008\)](#) propose a Bayesian nonparametric model for dealing with unobserved sources of heterogeneity and [Ruiz and Laplanche \(2010\)](#) account for the effect of individual variability on population size estimation from removal data.

Existing removal models assume population closure for the duration of the study. However, it is often the case when sampling amphibians and reptiles that the study period encompasses the reproduction period, and hence the population is renewed by the emergence of newborn individuals, violating the assumption of closure. This population renewal is suggested by the data set of common lizards (Table 1) where it can be seen that the number of individuals detected peaks in August, three months after the study commenced. The ecology of the species suggests that this is the result of a renewal of the population due to birth [[Avery \(1975\)](#)]. Motivated by this and similar data sets, we develop a model that allows for a group or groups, of unknown size, of individuals to emerge/arrive at an unknown time while accounting for individuals that may die or permanently emigrate during the study period. In addition, the model estimates the number of undetected individuals, that is, the number of individuals remaining at the site at the end of the study. We note that the number of undetected individuals is the demographic parameter of interest in removal studies. We fit the model using a Bayesian approach to obtain the posterior distribution of the number of undetected individuals. An attractive feature of the Bayesian approach is that it is straightforward to estimate the probability that no animals remain (i.e., that the population is totally depleted). This is analogous to the use of Bayesian methods to estimate the probabilities of bird species being endangered [see [Brooks et al. \(2008\)](#)].

Additionally, when the study takes place at an unfenced breeding site and commences before the start of the breeding season, individuals can migrate into or out of the breeding site, thus violating the assumption of population closure. Populations of great crested newts are a typical example because they are known to migrate to breeding ponds in a staggered manner and then depart at the end of the breeding season. In this case, the number of renewal groups is unknown. We therefore further extend the model to allow for an unknown number of renewal groups and we use a reversible jump [RJ, [Green \(1995\)](#)] MCMC algorithm to fit the model and estimate the renewal pattern and the number of renewal groups.

The model we present responds to a practical ecological need, as data of the type we analyse are often encountered. It provides a completely general modelling framework which extends existing simple models to include features such as survival and recruitment. Conclusions resulting from using rules of thumb or inappropriate model fitting, with no allowance for features such as renewal or variable

detection, may be used to conform to current legislative guidelines, therefore it is critical that new statistical approaches are developed so that deleterious effects on protected species can be prevented.

We introduce the model and the parameters in Section 2. We present each motivating example and the results of model fitting in Sections 3 and 4, respectively. Further extensions to the model are discussed in Section 5. Convergence diagnostics, RJMCMC details and simulation results are provided in the supplemental article [Matechou et al. (2016)].

2. Model. We assume that there are T sampling occasions taking place at times $\tau_t, \in \mathbf{R}, t = 1, \dots, T$ and individuals detected at an occasion are permanently removed from the study site. The data set of a single species, \mathbf{n} , is a vector of length T with entry $n_t, t = 1, \dots, T$ denoting the number of individuals of that species removed at occasion t . The total number of individuals of a species detected and removed from the site is denoted by $D = \sum_{t=1}^T n_t$, and the unknown number of individuals that will remain at the site when the study ends by N with $N \geq 0$. The primary aim of sampling is population depletion, and hence the main objective of modelling is to estimate N or its posterior distribution.

Each individual in the population belongs to one of $T + 2$ categories. If an individual belongs to category $t, t = 1, \dots, T$, then it was removed at sampling occasion t . The N individuals that were never detected and not removed belong to category $T + 1$. The remainder of the individuals, M , left the study site either through death or emigration before being detected and before the study ended.

The probability, γ_t , of an individual belonging to category t is a function of the following parameters:

- $\beta_{t-1}, t = 1, \dots, T$: entry parameters. The proportion of individuals that became available for detection for the first time at sampling occasion t .

The proportion of individuals that became available for detection at least once during the study is $\Omega = \sum_{t=1}^T \beta_{t-1}$, while the complement of $\Omega, 1 - \Omega$, is equal to the proportion of individuals that arrived after the end of the study and therefore never became available for detection. It is anticipated that Ω will be close to 1 by study design.

- ϕ_t : survival probability. The probability an individual present at sampling occasion t remains until occasion $t + 1$. The parameter ϕ is used to account for natural mortality and emigration, in addition to the removal process for detected individuals, and therefore denotes apparent survival as it is referred to in the capture-recapture literature.
- p_{t-b} : detection probability. The probability of detecting an individual that is present at sampling occasion t and first became available for detection at occasion b , hence having avoided detection $t - b$ times, at occasions $b, \dots, (t - 1)$. The parameter p_0 denotes the probability of being detected upon arrival.

Therefore, the probability γ_t can be expressed as

$$\gamma_t = \begin{cases} \sum_{b=1}^t \left[\beta_{b-1} \left\{ \prod_{k=b}^{t-1} \phi_k \right\} p_{t-b} \right], & t = 1, \dots, T, \\ \sum_{b=1}^T \left[\beta_{b-1} \left\{ \prod_{k=b}^{T-1} \phi_k \right\} \left(1 - \sum_{k=b}^T p_{k-b} \right) \right] + (1 - \Omega), & t = T + 1, \\ 1 - \sum_{t=1}^{T+1} \gamma_t, & t = T + 2, \end{cases}$$

where the empty product $\prod_{k=t}^{t-1} \phi_k$ is taken to be unity.

If we denote the vector of model parameters by θ , the data are described by the multinomial distribution,

$$(2.1) \quad \mathcal{P}(\mathbf{n}|\theta) = \frac{(N + M + D)!}{(\prod_{t=1}^T n_t!) N! M!} \left\{ \prod_{t=1}^T \gamma_t^{n_t} \right\} (\gamma_{T+1})^N \left(1 - \sum_{t=1}^{T+1} \gamma_t \right)^M.$$

Alternatively, conditioning on the event of surviving until detection (or until the end of the study for undetected individuals), we can work in terms of the conditional probabilities,

$$(2.2) \quad \mathcal{P}_c(\mathbf{n}|\theta) = \frac{(N + D)!}{(\prod_{t=1}^T n_t!) N!} \left\{ \prod_{t=1}^T \left(\frac{\gamma_t}{\sum_{t=1}^{T+1} \gamma_t} \right)^{n_t} \right\} \left(\frac{\gamma_{T+1}}{\sum_{t=1}^{T+1} \gamma_t} \right)^N.$$

Instead of using equation (2.1), we shall follow the conditional approach and use equation (2.2), which is not a function of M , for two reasons: (1) M is of no ecological interest in this case as the aim of sampling is to estimate N and the probability that it is zero, and (2) using the unconditional approach of equation (2.1) results in poor mixing in the MCMC algorithm to be used, especially for parameter M . If M is of interest, then the mixing of the chain can be improved by using more sophisticated proposal distributions that account for the correlations between M and $\{\phi_t\}$ and running the algorithm for longer.

We assume that individuals arriving or emerging into the population do so in groups, called renewal groups, and we model the arrival pattern of these groups using the probability density function (p.d.f.) of a normal distribution. Mixtures of normal distributions have also been used by [Matechou et al. \(2014\)](#) to model emergence of butterfly broods and by [Matechou et al. \(2015\)](#) to model arrival of flocks of birds at stopover sites.

For the data set of common lizards we allow for one renewal group of unknown size, which corresponds to newborn individuals. We assume that the proportion of individuals in this renewal group is equal to π_2 , with $\pi_1 = 1 - \pi_2$ denoting the proportion of individuals already present at the start of the study. We further assume that the arrival times of members of the renewal group can be described by

a normal p.d.f. with mean μ and variance σ^2 so that the proportion of individuals with arrival time in the interval $(\tau_{b-1}, \tau_b]$ is equal to

$$\beta_{b-1} = \pi_2 P(\tau_{b-1} < X \leq \tau_b)$$

for $b \in \{2, \dots, T\}$ while $\beta_0 = \pi_1 + \pi_2 P(X \leq \tau_1)$, where $X \sim N(\mu, \sigma^2)$.

For the data set of great crested newts, we allow for an unknown number, G , of renewal groups, each of unknown size and unknown arrival/emergence pattern, and each modelled by a normal p.d.f. as explained in the previous paragraph. In this case, π_g, μ_g and $\sigma_g^2, g = 1, \dots, G$, are respectively population fractions, mean arrival times and variance of arrival times of the G renewal groups, with $\sum_{g=1}^G \pi_g = 1$. The proportion of individuals with arrival time in the interval $(\tau_{b-1}, \tau_b]$ is now given as

$$\beta_{b-1} = \begin{cases} \sum_{g=1}^G \pi_g P(\tau_{b-1} < X_g \leq \tau_b), & b = 2, \dots, T, \\ \sum_{g=1}^G \pi_g P(X_g \leq \tau_1), & b = 1, \end{cases}$$

where $X_g \sim N(\mu_g, \sigma_g^2)$.

We adopt a Bayesian approach for inference and fit the model by using an MCMC algorithm. In the case of an unknown number of renewal groups, we use an RJMCMC algorithm that moves between models with different numbers of groups. The two detailed applications that follow are designed to take into account known features of the studied animals.

3. Common lizards. To facilitate the redevelopment of a large area of brown-field land in Cumbria, a region in the north of England, a programme of reptile collection and removal was undertaken in 2007. Three areas of land that supported reptiles were enclosed by purpose-built reptile exclusion fencing. Within each of the resulting compartments, artificial refuges (0.5 m² sheets of felt and metal) were placed at a density of approximately 50 ha⁻¹ within a suitable habitat. These refuges were checked once per day in the mornings only during suitable weather conditions. Common lizards were captured by hand and moved to an ex situ receptor site. The resulting removal data and daily records of temperature (maximum temperature in °C) are presented in Table 1.

Profile likelihood plots for N obtained from fitting the geometric model (which corresponds to the case of $\beta_0 = 1, \beta_{t-1} = 0, \forall t > 1$, and $\phi_t = 1, \forall t$) to the data show [supplemental article Matechou et al. (2016), Figure S1(a)] that the maximum-likelihood estimate obtained for N is equal to 903 individuals. This estimate for N is over four times the sample size and the likelihood surface is practically flat over a large range of values for N . The curvature around the maximum likelihood estimate for N is low, resulting in profile likelihood intervals that extend

beyond values of N greater than 10,000, 50 times the sample size. The wide profile likelihood intervals are due to the fact that the assumption of population closure is violated because it is evident that a large proportion of the common lizard population actually arrived/emerged long after the start of the study. The result suggests that when there is renewal of the population during a removal study, then a different model from the geometric is needed. For comparison, we also show the profile log-likelihood plot for N obtained from fitting the model presented in Section 2, which allows for one renewal group to demonstrate that the likelihood surface is no longer flat in this case [supplemental article, (Matechou et al. 2016), Figure S1(b)]. Note that we constrain $\phi = 1$ for this illustration.

Based on the literature concerned with the effect of temperature on the detection probability of reptiles [Sewell et al. (2012)], we logistically regress detection probability on maximum temperature and its square, and hence set

$$\log\left(\frac{\eta_t}{1 - \eta_t}\right) = \alpha_0 + \alpha_1 x_t + \alpha_2 x_t^2,$$

with x_t the maximum recorded temperature on sampling occasion t and $\alpha_0, \alpha_1, \alpha_2$ the regression coefficients, which gives the following expression for p_{t-b} :

$$(3.1) \quad p_{t-b} = \eta_t \left\{ \prod_{k=b}^{t-1} (1 - \eta_k) \right\}^{\mathbb{I}(t > b)},$$

where the indicator variable $\mathbb{I}(t > b)$ is equal to 1 if $t > b$ and 0 otherwise.

We chose a Unif[0, 1000] prior for the population size and a Dirichlet(1, 1) for the proportion of N in each group, π_1, π_2 . We set the prior for the mean arrival time of the renewal group to be Normal with a mean which corresponds to the end of July – beginning of August, based on the ecology of the species [Avery (1975)] and a standard deviation of roughly one month (30 days). Finally, we chose a half-normal prior for the standard deviation of arrival times of the renewal group with parameter 0.1, a $N(0, 1)$ prior for the coefficients of the covariates in the logistic regression model for detection probability, p , and a Unif(0, 1) prior for ϕ .

We provide the results of convergence diagnostics in Section 1.2 of the supplemental article [Matechou et al. (2016)]. The posterior distribution and cumulative posterior distribution for N are shown in Figure 1. Although the probability that the population has actually been depleted, which corresponds to the probability that $N = 0$, is lower than 10%, most of the posterior distribution mass is concentrated on values that are close to 0. In fact, the model estimates that, with a 95% probability, there were at most eight common lizards at the site after the end of the study. From the observed data, one might have expected that it would be likely that the population was totally depleted at the end of the sampling due to the observed trailing zeros; however, as noted above, the probability of total depletion is actually less than 0.1.

Because in this application the renewal group is thought to consist of juvenile common lizards which are born in the current breeding season, by comparing the




FIG. 1. Common lizard data. (a) Posterior distribution, and (b) cumulative posterior distribution of N . Also shown in (b) is the value of N which corresponds to the 95% quantile of the posterior distribution.

estimated proportion of individuals in each emergence group to the estimated total population, we can also extract information on the number of juveniles per adult (i.e., the rate of productivity). The posterior mean for π_2/π_1 is equal to 3.16 with a 95% posterior credible interval (CI) of (2.20, 4.43). Avery (1975) reports that female common lizards each produce between 3 and 11 juveniles per breeding season, with an average of 7.74. Therefore, assuming an equal sex-ratio of individuals in our study, our estimate of productivity is in close agreement with this previous finding.




FIG. 2. *Common lizard data: assessment of model fit. The vertical bars show the width of the 95% quantile interval of values generated from 1000 randomly sampled parts of the chain, the circles show the mean simulated values and the stars the observed values. The gaps in the data correspond to days when sampling was not conducted.*

The mean arrival time of the juvenile renewal group has a posterior mean of 77.89 [95% posterior CI = (76.34, 80.47)], which corresponds to the beginning of August, as would be expected given the northern location of this study site in England [Van Damme, Bauwens and Verheyen (1990)].

The posterior mean of apparent survival probability is 0.997 [95% posterior CI = (0.989, 0.999)], suggesting that the reduction in the size of the population by the end of the study is mostly due to sampling instead of emigration or death.

The fit of the model is assessed by generating data from randomly chosen parts of the chain and comparing them to the true data set in Figure 2, where it can be seen that the model provides a good description of the increase in the number of individuals detected due to population renewal, and the actual numbers of common lizards detected are mostly encompassed by the 95% quantile intervals of the simulated values. The model achieves a good fit to the data despite the sparseness of the data and the fact that no samples were collected during the days when emergence of juveniles peaked.

Finally, the posterior distribution of detection probability as a quadratic function of maximum temperature is summarised in Figure 3. We find that detection probability is at its maximum in mild weather when maximum temperature is roughly equal to 14°C, while it decreases considerably as maximum temperature decreases or increases. Our conclusion regarding the relationship between detection probability and temperature is in agreement with literature on the subject [Gent and




FIG. 3. *Common lizard data. Summaries of the posterior distribution of detection probability as a function of maximum daily recorded temperature. A logistic-quadratic function has been fitted.*

Gibson (1998), Joppa et al. (2009), Reading (1997)]. The large width of the posterior CI at low temperatures is due to the fact that there were very few surveys conducted under these conditions compared to higher temperatures.

4. Male great crested newts. An extensive removal study was conducted at a site in the east of England in 2010 prior to a substantial commercial development project. The areas where great crested newts were most likely to be found were fenced off to intercept the animals coming in from their hibernation areas to the ponds which were in the middle of the site. The trapping consisted mainly of collecting animals from pitfall traps, and this was supplemented by two evenings of sampling using torchlight. The data we consider here consist of removals of adult male individuals. Minimum air temperature was also recorded for most sampling occasions.

An RJMCMC algorithm was used to fit a model with an unknown number of renewal groups assuming constant survival probability, ϕ , and detection probability varying with minimum air temperature at sampling occasion t , x_t , such that

$$\log\left(\frac{\eta_t}{1 - \eta_t}\right) = \alpha_0 + \alpha_1 x_t.$$

Missing values for the covariate were imputed using the average of minimum temperatures recorded on the two adjacent sampling occasions.

We used a vague $\text{Unif}\{1, \dots, 20\}$ for the number of renewal groups, G , a $\text{Unif}[0, 1000]$ prior for N , a Dirichlet with all concentration parameters equal to 1 for π_g , $\forall g$, a $\text{Unif}(1, T)$ prior for μ_g to reflect our expectation that individuals are arriving during the study and not before or after, a half-normal with parameter

0.1 for $\sigma_g \forall g$, a $N(0, 1)$ for the coefficients of the logistic regression model for p and a $\text{Unif}(0, 1)$ prior for ϕ .

We present details on the RJMCMC algorithm and the convergence diagnostics used in Section 2 of the supplemental article [Matechou et al. (2016)].


The posterior distribution for G , the number of arrival groups, is mostly (87%) concentrated on the values 7 and 8 [supplemental article, (Matechou et al. 2016), Figure S5(a)]. The posterior distribution for N is mostly (80%) concentrated on 0 [supplemental article, (Matechou et al. 2016), Figure S5(b)], and we estimate that there is a probability of 5% that 2 or more individuals were at the site after the study ended. The low estimated number of undetected individuals is unsurprising, as not only do the data have a large number of trailing zeros at the end of the study, but the estimated mean arrival times of the renewal groups are primarily in the first half of the study (see Figure 4).

Posterior summaries of the probability of detection as a function of minimum air temperature are presented in Figure 5 where it can be seen that, as expected [Sewell, Beebee and Griffiths (2010)], newts are more likely to fall into the traps as the temperature increases. Although the slope of the logistic curve is steep, the 95% posterior CIs are fairly wide. The large width of the posterior CIs is possibly due to a moderate effect of minimum air temperature on detection probability, but it could also be an artefact of the increased uncertainty due to the use of RJMCMC.


The posterior mode for survival probability is around 0.1, with the posterior mean equal to 0.19 (95% posterior CI: 0, 0.6 and 95% HPD interval: 0, 0.5) because the posterior density has a long right tail. This low apparent survival probability in comparison to the common lizard apparent survival probability is likely due to study design. Within the newt study, the traps are outside the ponds, and so newts are able to more easily leave the study area without falling into the traps.

The fit of the model is assessed graphically in Figure 4(b), where it is seen that all of the observed counts lie within the boundaries of the 95% quantile intervals of the counts simulated by the model using randomly chosen iterations of the algorithm.

5. Possible extensions. The effect of unmodelled individual heterogeneity in either detection or survival probability on the estimation of population size has been well documented in the literature [McCrea and Morgan (2014), Chapter 3, and references therein]. Our simulations presented in Section 3 of the supplemental article [Matechou et al. (2016)] suggest that, in the case of removal data, such as those considered in this paper, if there exists heterogeneity in p which the model does not account for, then the posterior for N tends to be concentrated to the left of the true value for N . If N is small, as in the case studies considered in Sections 3 and 4, then the resulting bias is also small. However, when N is large and the population is far from being depleted, then the bias in N can be substantial.



(a)



(b)

FIG. 4. Male great crested newt data. (a) Estimated entry parameters obtained at each iteration of the RJMCMC algorithm, gray lines, together with the mean of all iterations, shown by the black line. The values on the x-axis correspond to sampling occasions. (b) Assessment of model fit. The observed counts, stars, lie within the 95% quantile intervals of counts simulated from the model, bars, using randomly selected iterations of the algorithm.

If individuals are expected to exhibit variation in their detectability, then the probability of detection can be modelled as a random variable with a beta-geometric distribution with respective mean and shape parameters η and θ [see Morgan (2009), Section 2.3], which gives

$$(5.1) \quad p_{t-b} = \eta \frac{\prod_{k=b}^{t-1} \{1 - \eta + (k - b)\theta\}}{\prod_{k=b}^t \{1 + (k - b)\theta\}}.$$




FIG. 5. Male great crested newt data. Summaries of the posterior distribution of detection probability as a function of minimum daily recorded temperature. A logistic function has been fitted.

Note that in the case of common detection probabilities between individuals (i.e., when $\theta = 0$) p_{t-b} simplifies to a standard geometric model with probability of success η , and

$$p_{t-b} = \eta \prod_{k=b}^{t-1} (1 - \eta) = \eta(1 - \eta)^{t-b}.$$

Similarly, if it is anticipated that detection probability depends on covariates, such as environmental factors, as was the case for the applications in Sections 3 and 4, then η can be modelled as a function of these using a logistic regression model. If $\mathbf{x}_1, \dots, \mathbf{x}_q$ are q environmental covariates and $\boldsymbol{\alpha}$ is a set of coefficients of length $q + 1$, then

$$\log\left(\frac{\eta_t}{1 - \eta_t}\right) = \alpha_0 + \alpha_1 x_{1t} + \dots + \alpha_q x_{qt},$$

which, for example, in the case $\theta = 0$ gives

$$p_{t-b} = \eta_t \prod_{k=b}^{t-1} (1 - \eta_k).$$

It is easy to show that when appropriate the model can naturally accommodate both heterogeneity and environmental covariates, as long as the covariate(s) are discrete or discretised so that the values for all years are not all different. In this

discrete case, the recapture probabilities are suitable products of terms of the form illustrated in expression (5.1).

An alternative approach for modelling heterogeneity, both in detection and survival probabilities, is provided by [Pledger \(2000\)](#) via the use of finite mixture models. Finite mixture models have been used extensively in the capture-recapture literature for both open and closed populations, and they provide an effective way to model heterogeneity.

Our presented model assumes that there is no temporary emigration, and hence once an individual has been removed or has departed from the study it does not return. However, amphibians and reptiles have activity patterns that vary according to weather conditions and may become unavailable for detection for a period of time. This issue of temporary unavailability for capture is the equivalent of temporary emigration in the capture-recapture literature; see, for example, [Barker \(1997\)](#), [Kendall, Nichols and Hines \(1997\)](#) and [Kendall and Nichols \(2002\)](#). Although for the data sets considered in this paper we estimated that the population was practically depleted before the expected start of the hibernation period (i.e., end of October [[Beebee and Griffiths \(2000\)](#)]), it is possible that some individuals evaded detection by becoming less active as winter approached. We are currently working on extensions of the models presented in this paper that use a multi-state approach to allow for temporary emigration of individuals, developing methodology from standard capture-recapture literature [[Pradel \(2005\)](#)].

Removal data alone may be insufficient to estimate all parameters of interest; however, models for removal data have survival and detection parameters in common with capture-recapture models. Therefore, it may be possible to perform an integrated analysis—simultaneously modelling removal data and independent capture-recapture data [see [Besbeas et al. \(2002\)](#)]. This integrated analysis would result in generally improved precision of parameter estimates and may overcome challenges such as near-singularity and parameter redundancy of models [[Cole and McCrea \(2016\)](#)].

6. Discussion. Translocations of protected species from the path of development are widespread in the UK and globally. However, such actions are expensive, often poorly designed and monitored, and undetected animals may comprise a significant proportion of the population left behind at the development site [[Germano et al. \(2015\)](#), [Lewis et al. \(2014\)](#)]. The models we propose here provide a basis for determining the effectiveness of such translocations, and thereby improve policy and guidance for such actions.

We have proposed a new model for the case of removal data when the assumption of population closure is violated because of: (a) individuals emerging/arriving sometime after the start of the study, either through birth and/or immigration, and renewing the population and (b) individuals departing from the study site before being detected through death and/or emigration.

The model allows for either a fixed number of known renewal groups or an unknown number, and estimates their sizes as well as the means and variances of arrival times of the individuals in these groups. The model of this paper responds to a practical need, as the data described are commonly encountered.

We provided two applications:

(i) a data set of common lizards collected during a period that encompassed the reproduction period, and hence there was one renewal group which consisted of juveniles that emerged in late summer. We estimated the probability of population depletion as less than 10%. The ratio of juveniles to adults in the population was also derived as a by-product of the analysis.

(ii) a data set of male great crested newts collected during the time when individuals are migrating to ponds to breed, where we used an RJMCMC algorithm to account for and estimate the number of renewal groups. In this case, the probability of population depletion was estimated at around 80%.

We have also incorporated environmental covariates in the estimation of detection probabilities and, although we have not undertaken a detailed model-selection exercise for our choice of covariates, we have shown using simulation that the model provides a good description of the data. The functional form of the covariate regression used for each application was guided by ecological knowledge. Our results suggest that detection probability of common lizards is lower when the maximum daily temperature is too low or too high, with the optimum being roughly at 14°C. We found that the detection probability of great crested newts varied with minimum air temperature, and as minimum air temperature increased, so did detection probability which is in-line with previous findings relating weather conditions to detectability of these animals [see Sewell, Beebee and Griffiths (2010)].

It is important to note that, while the new model incorporating both arrivals and departures provides good descriptions of the two data sets that we analyse in the paper, if capture probability is high, then there could be difficulties in estimating ϕ : if individuals are removed soon after arrival, then retention due to survival cannot be observed. The model has performed well in both case studies considered, but care should be taken when fitting complex models to sparse data. The diagnosis of possible parameter redundancy of the new model, along with how it might be overcome by combining additional information, is an area of current research.

A great advantage of adopting a Bayesian approach for model fitting is that we can evaluate the probability that more than n individuals remain at the study site for any n . In further work, we plan to investigate the posterior predictive distribution after each sample is collected to use this as a guide for calculating the number of samples still required before total population depletion is achieved.

7. Acknowledgements. We thank Herpetologic Ltd. for providing the great crested newt removal data, particularly Jon Cranfield for helpful information about the sampling protocol. We also thank Cristiano Villa for the discussion about the

choice of priors for the parameters. The comments of the Associate Editor and three referees greatly improved the paper.

SUPPLEMENTARY MATERIAL

Supplement to “Open models for removal data” (DOI: [10.1214/16-AOAS949SUPP](https://doi.org/10.1214/16-AOAS949SUPP); .pdf). We provide results of convergence diagnostics, details of the RJMCMC algorithm employed and simulation results.

REFERENCES

- AVERY, R. A. (1975). Clutch size and reproductive effort in the lizard *Lacerta vivipara* Jacquin. *Oecologia* **19** 165–170.
- BARKER, R. J. (1997). Joint modeling of live recapture, tag-resight and tag-recovery data. *Biometrics* **53** 666–677.
- BEEBEE, T. J. C. and GRIFFITHS, R. A. (2000). *Amphibians and Reptiles*. HarperCollins, London.
- BESBEAS, P., FREEMAN, S. N., MORGAN, B. J. T. and CATCHPOLE, E. A. (2002). Integrating mark-recapture-recovery and census data to estimate animal abundance and demographic parameters. *Biometrics* **58** 540–547. [MR1933532](#)
- BOHRMANN, T. F. and CHRISTMAN, M. C. (2013). Optimal allocation of sampling effort in depletion surveys. *J. Agric. Biol. Environ. Stat.* **18** 218–233. [MR3067276](#)
- BROOKS, S. P., FREEMAN, S. N., GREENWOOD, J. J. D., KING, R. and MAZZETTA, C. (2008). Quantifying conservation concern—Bayesian statistics, birds and the red lists. *Biological Conservation* **141** 1436–1441.
- COLE, D. J. and MCCREA, R. S. (2016). Parameter redundancy in discrete state-space and integrated models. *Biom. J.* To appear.
- DEFRA (2015). <https://www.gov.uk/guidance/reptiles-protection-surveys-and-licences>. Natural England/Defra.
- DORAZIO, R. M., JELKS, H. L. and JORDAN, F. (2005). Improving removal-based estimates of abundance by sampling a population of spatially distinct subpopulations. *Biometrics* **61** 1093–1101. [MR2216203](#)
- DORAZIO, R. M., MUKHERJEE, B., ZHANG, L., GHOSH, M., JELKS, H. L. and JORDAN, F. (2008). Modeling unobserved sources of heterogeneity in animal abundance using a Dirichlet process prior. *Biometrics* **64** 635–644. [MR2432438](#)
- GENT, T. and GIBSON, S. E. (1998). *Herpetofauna Workers’ Manual*. Joint Nature Conservation Committee, Peterborough.
- GERMANO, J. M., FIELD, K. J., GRIFFITHS, R. A., CLULOW, S., FOSTER, J., HARDING, G. and SWAISGOOD, R. R. (2015). Mitigation-driven translocations: Are we moving wildlife in the right direction? *Frontiers in Ecology and the Environment* **13** 100–105.
- GREEN, P. J. (1995). Reversible jump Markov chain Monte Carlo computation and Bayesian model determination. *Biometrika* **82** 711–732. [MR1380810](#)
- HGBI (1998). *Herpetofauna Groups of Britain and Ireland—Evaluating Local Mitigation/translocation Programmes: Maintaining Best Practice and Lawful Standards*. HGBI Advisory Notes for Amphibian and Reptile groups. Preprint.
- JOPPA, L. N., WILLIAMS, K. R., TEMPLE, S. A. and CASPER, G. S. (2009). Environmental factors affecting sampling success of artificial cover objects. *Herpetological Conservation and Biology* **5** 143–148.
- KENDALL, W. L. and NICHOLS, J. D. (2002). Estimating state-transition probabilities for unobservable states using capture-recapture/resighting data. *Ecology* **83** 3276–3284.

- KENDALL, W. L., NICHOLS, J. D. and HINES, J. E. (1997). Estimating temporary emigration using capture-recapture data with Pollock's robust design. *Ecology* **78** 563–578.
- LEWIS, B., GRIFFITHS, R. A., WILKINSON, J. W. and ARNELL, A. (2014). Examining the Fate of Local Great Crested Newt Populations Following Licensed Developments. Report WM031, London: Department for Environment, Food and Rural Affairs.
- MATECHOU, E., DENNIS, E. B., FREEMAN, S. N. and BRERETON, T. (2014). Monitoring abundance and phenology in (multivoltine) butterfly species: A novel mixture model. *Journal of Applied Ecology* **51** 766–775.
- MATECHOU, E., NICHOLLS, G., MORGAN, B. J. T., COLLAZO, J. A. and LYONS, J. E. (2015). Bayesian analysis of Jolly–Seber type models; incorporating heterogeneity in arrival and departure. *Environ. Ecol. Stat.* To appear. DOI:10.1007/s10651-016-0352-0.
- MATECHOU, E., MCCREA, R. S., MORGAN, B. J. T., NASH, J. D. and GRIFFITHS, R. A. (2016). Supplement to “Open models for removal data.” DOI:10.1214/16-AOAS949SUPP.
- MCCREA, R. S. and MORGAN, B. J. T. (2014). *Analysis of Capture–Recapture Data*. Chapman & Hall/CRC, Boca Raton, FL. MR3330977
- MORAN, P. A. P. (1951). A mathematical theory of animal trapping. *Biometrika* **38** 307–311.
- MORGAN, B. J. T. (2009). *Applied Stochastic Modelling*, 2nd ed. CRC Press, Boca Raton, FL. MR2468192
- PLEDGER, S. (2000). Unified maximum likelihood estimates for closed capture-recapture models using mixtures. *Biometrics* **56** 434–442.
- PRADEL, R. (2005). Multievent: An extension of multistate capture-recapture models to uncertain states. *Biometrics* **61** 442–447. MR2140915
- READING, C. J. (1997). A proposed standard method for surveying reptiles on dry lowland heath. *Journal of Applied Ecology* **34** 1057–1069.
- RIDOUT, M. S. and MORGAN, B. J. T. (1991). Modelling digit preference in fecundability studies. *Biometrics* **47** 1423–1434.
- RUIZ, P. and LAPLANCHE, C. A. (2010). A hierarchical model to estimate the abundance and biomass of salmonids by using removal sampling and biometric data from multiple locations. *Canadian Journal of Fisheries and Aquatic Sciences* **67** 2032–2044.
- SEWELL, D., BEEBEE, T. J. C. and GRIFFITHS, R. A. (2010). Optimising biodiversity assessments by volunteers: The application of occupancy modelling to large-scale amphibian surveys. *Biological Conservation* **143** 2102–2110.
- SEWELL, D., GUILLERA-ARROITA, G., GRIFFITHS, R. A. and BEEBEE, T. J. C. (2012). When is a species declining? Optimizing survey effort to detect population changes in reptiles. *PLOS ONE* **7** e43387.
- VAN DAMME, R., BAUWENS, D. and VERHEYEN, R. F. (1990). Evolutionary rigidity of thermal physiology—The case of the cool temperate lizard, *Lacerta vivipara*. *Oikos* **57** 61–67.
- ZIPPIN, C. (1956). An evaluation of the removal method of estimating animal populations. *Biometrics* **12** 163–189.

E. MATECHOU
 R. S. MCCREA
 B. J. T. MORGAN
 SCHOOL OF MATHEMATICS, STATISTICS
 AND ACTUARIAL SCIENCE
 UNIVERSITY OF KENT
 CORNWALLIS MATHS BUILDING
 CANTERBURY
 KENT CT2 7NF
 UNITED KINGDOM
 E-MAIL: E.Matechou@kent.ac.uk
 R.S.McCrea@kent.ac.uk
 B.J.T.Morgan@kent.ac.uk

D. J. NASH
 R. A. GRIFFITHS
 INSTITUTE OF CONSERVATION AND ECOLOGY
 SCHOOL OF ANTHROPOLOGY AND CONSERVATION
 UNIVERSITY OF KENT
 DURRELL MARLOWE BUILDING
 CANTERBURY, CT2 7NR
 UNITED KINGDOM
 E-MAIL: djn8@kent.ac.uk
 R.A.Griffiths@kent.ac.uk