

DOCTOR OF PHILOSOPHY

Evaluation of cultivation, legume undersowing and nitrogen interventions on wheat development

Rial Lovera, Karen E.

Award date:
2015

Awarding institution:
Coventry University
Royal Agricultural University

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of this thesis for personal non-commercial research or study
- This thesis cannot be reproduced or quoted extensively from without first obtaining permission from the copyright holder(s)
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Evaluation of cultivation, legume
undersowing and nitrogen interventions
on wheat development**

By

Karen E. Rial Lovera

*A thesis submitted in partial fulfilment of the University's requirements
for the Degree of Doctor of Philosophy*

September 2015

Coventry University

In association with the Royal Agricultural University

Declaration

I declare that this research is the result of my own work and except where stated and referenced otherwise, all the written work and investigations are my own. This work has not been accepted or submitted for an academic award elsewhere.

Karen E Rial Lovera

Dedication

To my beloved parents Juan and Rosa, who without their sacrifice, love and encouragement I would not have achieved so much.

This item has been removed due to 3rd Party Copyright. The unabridged version of the thesis can be viewed in the Lanchester Library Coventry University.

Anonymous.

Acknowledgement

I would like to express my gratitude to my supervisors Nicola Cannon, Paul Davies and John Conway for their valuable advice, support and guidance in the shaping of both myself and my professional abilities throughout the course of my PhD study.

My sincere thanks and gratitude to all the technical staff who helped to make this project possible, Tony Norris for his valuable assistance in field operations, Sally Rice, Sylvia de Oliveria and Darren Hopkins for their assistance in the lab and for all those encouraging conversations at time of hard work. To the library team, especially to Theano Manoli for provide me with many interlibrary loans and for so enriching and cheering conversations, and all those many laughs.

I would also like to extend my gratitude to Roger Payne for his valuable advice and for guidance with statistics.

The financial support from the Royal Agricultural University's Research Committee for carrying out this research is also greatly acknowledged.

To my colleagues in the 'PhD Office', Sara, Harriet, Vijaya, Kemi, Adeola, Matthew and all the new 'PhDs', many thanks for all those many conversations which made for a very stimulating and friendly work environment.

I would also like to gratefully thank everybody who has provided help and support to me throughout every stage of my life, especially to my parents, all new and old friends, and my Venezuelan and Spanish family.

Lastly but not the least, my thanks to Marco Forno for his help, belief in me and most of all for his patience.

Abstract

Agriculture is facing increasing pressures to produce food that meets specific market and/or nutrition requirements, while using inputs in such a way that can ensure economic and environmental goals more efficiently. Two field experiments were conducted in 2013 and 2014 at the Royal Agricultural University's Harnhill' Manor Farm, Cirencester, UK to evaluate the influence of selected cultivation techniques, N fertilisation and undersowing legumes on spring wheat growth and development. To explore, in particular, the yield components contributing to grain yield and quality, as well as weed pressure influences together with changes in soil mineral N (SMN) content. Cultivation techniques included conventional tillage (CT), high intensity non-inversion tillage (HINiT) and low intensity non-inversion tillage (LINiT); mineral N fertilisation rates of 0, 70, 140 and 210 kg N ha⁻¹ and two undersown legume species, black medic and white clover, plus no undersowing treatment. The performance of the management practices was strongly influenced by the weather. In 2013, under dry weather conditions, LINiT seems to be a suitable alternative to CT, while N fertilisation did not encourage greater grain yield. In 2014, CT appears to be a more reliable practice, while the application of up to 140 kg N ha⁻¹ seemed to be enough to increase grain yield. Dry weather conditions at the time of broadcasting did not allow the undersowing species to be fully established, resulting in no effects on weed control and crop growth. In 2013, the initial poor plant establishment and slow crop growth under LINiT was compensated for by the soils ability to retain moisture, and thereby reducing crop water stress during the dry periods. This finally resulted in statistically similar grain yield to CT. In 2014 when water was not a limiting factor, poor plant establishment and crop growth, low SMN content and high weed pressure under LINiT resulted in lower grain yield than CT. In both years, HINiT resulted in low SMN content and high weed pressure resulting in poor grain yield. Across experiments, HINiT and LINiT saved energy-use and production costs, but CT could be more energy-use efficient and have high economic return if higher grain yield is assured. N fertilisation significantly promoted wheat growth, although under dry conditions with higher residual soil N, the N fertilisation did not increase yield. Under low SMN level applying up to 140 kg N ha⁻¹ increased grain yield produced, but N fertilisation is energy consuming and its use does not always ensure a higher economic return.

Table of Content

Dedication.....	i
Acknowledgement	ii
Abstract.....	iii
Table of Content	iv
List of Tables	ix
List of Figures.....	xiii
List of Plates	xix
Abbreviations	xx
CHAPTER ONE - Introduction.....	1
1.1. Background.....	1
1.2. Problem statement and study aim.....	2
1.3. Objectives of the study	4
1.4. Thesis structure.....	4
CHAPTER TWO - Literature review	6
2.1. Cultivation techniques	6
2.1.1. Conventional tillage.....	6
2.1.2. Non-inversion tillage	7
2.2. Cultivation techniques and soil properties.....	9
2.2.1. Soil physical properties	9
2.2.2. Soil chemical properties	11
2.3. Cultivation techniques and soil nitrogen availability	11
2.4. Cultivation techniques and weed suppression	12
2.4.1. Cultivation techniques and herbicide-use.....	13
2.5. Cultivation techniques and diseases	14
2.6. Cultivation techniques and crop yield	15
2.7. Cultivation techniques and grain protein.....	17
2.8. Nitrogen fertilisation and plant growth	17
2.8.1. Nitrogen-use and nitrogen-use efficiency.....	19
2.8.2. Nitrogen fertilisation and grain protein	20
2.8.3. Nitrogen fertilisation and weeds.....	21

2.9.	Intercropping	22
2.9.1.	Undersowing.....	23
2.9.2.	Cereal-legume intercropping	24
CHAPTER THREE - Materials and methods		27
3.1.	Experimental site	27
3.1.1.	Meteorological conditions	28
3.2.	Experimental design and treatments.....	30
3.2.1.	Details of experimental treatments	32
3.3.	Aboveground assessments.....	40
3.3.1.	Developmental stages	40
3.3.2.	Assessments.....	40
3.4.	Soil assessments	43
3.4.1.	Soil chemical analysis	43
3.4.2.	Soil physical analysis	47
3.5.	Statistical analysis	48
CHAPTER FOUR - Core experiment I – spring wheat 2013		49
4.1.	Introduction	49
4.2.	Material and methods	50
4.2.1.	Experimental site	50
4.2.2.	Experimental design and treatment structure	51
4.2.3.	Meteorological conditions	51
4.2.4.	Assessments.....	52
4.2.5.	Statistical analysis	53
4.3.	Results	54
4.3.1.	Establishment	54
4.3.2.	Tillers and total shoot production.....	55
4.3.3.	Mid-season wheat biomass and nitrogen uptake	57
4.3.4.	Mid-season legume biomass and nitrogen accumulation	60
4.3.5.	Mid-season total weed biomass and nitrogen accumulation	61
4.3.6.	Weed species composition.....	66
4.3.7.	Plant height and ears number.....	71
4.3.8.	Disease scoring.....	71

4.3.9.	Final biological harvest	72
4.3.10.	Soil mineral nitrogen	79
4.3.11.	Soil moisture (gravimetric) content	84
4.3.12.	Soil pH.....	86
4.3.13.	Soil penetration resistance	86
4.4.	Discussion.....	88
4.4.1.	Plant establishment, tillers and total shoots number.....	88
4.4.2.	Mid-season plant biomass and nitrogen uptake.....	90
4.4.3.	Plant height and ears number.....	95
4.4.4.	Ears, straw and total wheat biomass	96
4.4.5.	TGW, grains per ear, final grain yield and harvest index.....	97
4.4.6.	Wheat N yield, grain protein and N efficiencies	99
4.4.7.	Non-wheat biomass and N yield at harvest	102
4.4.8.	Soil moisture content	103
4.4.9.	Soil mineral nitrogen	105
4.4.10.	Soil pH.....	107
4.4.11.	Soil penetration resistance	108
4.5.	Conclusions	108
CHAPTER FIVE - Core experiment II - spring wheat 2014		112
5.1.	Introduction	112
5.2.	Materials and methods.....	113
5.2.1.	Experimental site	113
5.2.2.	Experimental design and treatment structure	113
5.2.3.	Meteorological conditions	113
5.2.4.	Assessments.....	114
5.2.5.	Statistical analysis	115
5.3.	Results	116
5.3.1.	Overwinter growth assessment.....	116
5.3.2.	Plant establishment, tiller numbers and total shoots.....	120
5.3.3.	Mid-season wheat biomass and nitrogen uptake	120
5.3.4.	Mid-season weed and legume biomass and nitrogen uptake.....	124
5.3.5.	Weed species composition.....	132

5.3.6.	Plant height and ears number.....	135
5.3.7.	Diseases scoring	137
5.3.8.	Final biological harvest	138
5.3.9.	Soil mineral nitrogen	144
5.3.10.	Soil moisture (gravimetric) content.....	149
5.3.11.	Soil pH.....	152
5.3.12.	Soil penetration resistance	152
5.4.	Discussion.....	155
5.4.1.	Overwinter growth assessment.....	155
5.4.2.	Soil penetration resistance	156
5.4.3.	Plant establishment, tillers and total shoot production	157
5.4.4.	Mid-season plant biomass and nitrogen uptake.....	158
5.4.5.	Plant height and ears number.....	163
5.4.6.	Ear, straw and total wheat biomass	164
5.4.7.	TGW, grains per ears, final grain yield and harvest index	165
5.4.8.	Wheat N yield, grain protein and N efficiencies	166
5.4.9.	Non-wheat biomass and N yield at harvest	168
5.4.10.	Soil moisture content	169
5.4.11.	Soil mineral nitrogen	170
5.5.	Conclusions	172
CHAPTER SIX - Effect of weather on spring wheat performance following different cultivation regimes		174
6.1.	Introduction	174
6.2.	Material and methods	175
6.3.	Results and discussion.....	177
6.4.	Conclusions	182
CHAPTER SEVEN - Economic and energy-use evaluation of spring wheat production under different cultivation techniques, nitrogen fertilisation rates and undersowing ..		184
7.1.	Introduction	184
7.2.	Material and methods	185
7.2.1.	Energy considerations.....	185
7.2.2.	Economic analysis	187
7.3.	Results and discussion.....	189

7.3.1.	Energy considerations.....	189
7.3.2.	Economic analysis	197
7.4.	Conclusion	200
CHAPTER EIGHT - General discussion and conclusions.....		201
8.1.	Introduction	201
8.2.	Cultivation techniques	202
8.2.1.	Wheat performance.....	202
8.2.2.	Weed pressure.....	204
8.2.3.	Soil mineral nitrogen	206
8.2.4.	Energy consumption and economic impact.....	207
8.3.	Nitrogen fertilisation	208
8.3.1.	Wheat performance.....	208
8.3.2.	Weed pressure.....	210
8.3.3.	Soil mineral nitrogen	211
8.3.4.	Energy consumption and economic impact.....	211
8.4.	Legume undersowing	212
8.5.	Interactions	212
8.6.	Concluding remarks.....	212
8.7.	Implications	214
8.8.	Future work	215
References		217
Appendices		275
Appendix 1. Illustrated keys used for disease assessment		276
Appendix 2. Weed species biomass across core experiments.....		279
Appendix 3. Data set for meta-analysis.....		281
Appendix 4. Energy balance.....		282
Appendix 5. Publications prepared during this investigation.....		284

List of Tables

Table 2. 1. Some detrimental effects of conventional tillage	7
Table 2. 2. Some reported non-inversion tillage effects.....	8
Table 2. 3. Some reported cereal diseases favoured by different cultivation systems ...	15
Table 2. 4. Detrimental effects of non-inversion tillage on final grain yield	16
Table 2. 5. Nitrogen fertilisation effects on cereal plant growth and development	18
Table 2. 6. Adverse and beneficial effects of legume intercropping on cereal productivity.....	25
Table 3. 1. Initial physiochemical properties of the experimental site.....	27
Table 3. 2. Cropping systems used in the site from 2007 to 2012	28
Table 3. 3. Experimental wheat crop details	32
Table 3. 4. Nitrogen treatments application dates with corresponding spring wheat growth stages (GS)	37
Table 3. 5. Dates and growth stages for above ground crop assessment.....	40
Table 4. 1. Diary of 2013 field operations.....	51
Table 4. 2. Above ground assessments dates for spring wheat 2013	53
Table 4. 3. Soil assessment dates for spring wheat 2013	53
Table 4. 4. Effect of cultivation techniques and nitrogen fertilisation on spring wheat establishment	54
Table 4. 5. Effect of cultivation techniques and nitrogen fertilisation on spring wheat tillers and total shoots production.....	56
Table 4. 6. Effect of cultivation techniques and nitrogen fertilisation on spring wheat biomass and N uptake (May 2013).....	57
Table 4. 7. Effect of cultivation techniques and nitrogen fertilisation on spring wheat biomass and N uptake (July 2013)	60
Table 4. 8. Effect of cultivation techniques and nitrogen fertilisation on legume biomass and N uptake (May and July 2013)	61

Table 4. 9. Effect of cultivation techniques and nitrogen fertilisation on weed biomass and N uptake (May and July 2013)	62
Table 4. 10. Effect of cultivation techniques and nitrogen fertilisation on weed species biomass (July 2013).....	66
Table 4. 11. Effect of cultivation techniques and nitrogen fertilisation on <i>Stellaria media</i> L. and <i>Lolium perenne</i> L. biomass	68
Table 4. 12. Cultivation techniques and nitrogen fertilisation effects on wheat height and wheat ear number.....	71
Table 4. 13. Effect of cultivation techniques and nitrogen fertilisation on spring wheat ears, straw and total wheat biomass.....	72
Table 4. 14. Effect of cultivation techniques and nitrogen fertilisation on TGW, grains per ear, final grain yield and harvest index	73
Table 4. 15. Effect of cultivation techniques and nitrogen fertilisation on spring wheat N uptake, grain protein and N harvest index	75
Table 4. 16. Effect of cultivation techniques, nitrogen fertilisation and undersowing on spring wheat N-efficiency parameters	76
Table 4. 17. Effect of cultivation techniques and nitrogen fertilisation on weed and legume biomass and N uptake (Harvest 2013).....	77
Table 4. 18. Effect of nitrogen fertilisation on soil pH (June 2013).....	86
Table 4. 19. Key outcomes for the cultivation techniques effects during 2013 cropping season	109
Table 4. 20. Key outcomes for the nitrogen fertilisation effects during 2013 cropping season	111
Table 5. 1. Diary of 2014 field operations.....	113
Table 5. 2. Above ground assessments for spring wheat 2014	115
Table 5. 3. Soil assessments for spring wheat 2014	115
Table 5. 4. Weed and legume biomass and N uptake (Overwinter 2013).....	116
Table 5. 5. Effect of cultivation techniques on establishment, tiller number and total shoot of spring wheat.....	120

Table 5. 6. Effect of cultivation techniques and nitrogen fertilisation on spring wheat biomass and N uptake (May 2014).....	121
Table 5. 7. Effect of cultivation techniques and nitrogen fertilisation on spring wheat biomass and N uptake (July and August 2014).....	123
Table 5. 8. Effect of cultivation techniques and nitrogen fertilisation on weed and legume biomass and N uptake (May 2014).....	124
Table 5. 9. Effect of cultivation techniques and nitrogen fertilisation on weed and legume biomass and N uptake (July 2014).....	126
Table 5. 10. Effect of cultivation techniques and nitrogen fertilisation on weeds and legumes biomass and N uptake (August 2014).....	129
Table 5. 11. Effect of cultivation techniques, nitrogen fertilisation and undersowing on <i>Sinapis arvensis</i> L. and <i>Stellaria media</i> L. biomass (July 2014).....	133
Table 5. 12. Cultivation techniques and nitrogen fertilisation effect on spring wheat height and ears number.....	136
Table 5. 13. Cultivation techniques and nitrogen fertilisation effect on ears, straw and total wheat biomass	138
Table 5. 14. Cultivation techniques and nitrogen fertilisation effect on spring wheat TGW, grains per ear, final grain yield and harvest index.....	140
Table 5. 15. Cultivation techniques and nitrogen fertilisation effect on total wheat and grain N uptake, grain protein content, and N harvest index	142
Table 5. 16. Cultivation techniques and nitrogen fertilisation effect on N- efficiency parameters.....	143
Table 5. 17. Cultivation techniques, nitrogen fertilisation and undersowing effects on weed and legume biomass and N uptake (Harvest 2014).....	144
Table 5. 18. Key outcomes for the cultivation techniques effects during 2014 cropping season	172
Table 5. 19. Key outcomes for the nitrogen fertilisation effects during 2014 cropping season	173
Table 6. 1. Cultivation techniques treatments terminology used by Vijaya Bhaskar <i>et al.</i> (2013b) and their equivalents in the present study	175
Table 6. 2. Details of the spring wheat cropping seasons.....	176

Table 6. 3. Monthly air temperature (°C) for the study period (2012-2014) and for the long-term records (2002-2012). Royal Agricultural University meteorological station, (NGR SP 42 004 011).....	177
Table 6. 4. Monthly and cumulative rainfall (mm) for the study period (2012-2014) and the long-term records (2002-2012). Royal Agricultural University meteorological station, (NGR SP 42 004 011).....	178
Table 7. 1. Energy equivalent of inputs and outputs	187
Table 7. 2. Seeds cost, contractor costs and grain price considered for all core experiments.....	188
Table 7. 3. Analysis of energy indices for spring wheat production in 2013.....	192
Table 7. 4. Analysis of energy indices for spring wheat production in 2014.....	195
Table 7. 5. Economic analysis for spring wheat production in 2013	198
Table 7. 6. Economic analysis for spring wheat production in 2014	199
Table 8. 1. Key findings for contrasting cultivation techniques for all core experiments	203
Table 8. 2. Trends in weed growth between cultivation techniques across core experiments.....	205
Table 8. 3. Key findings for different nitrogen fertilisation treatments for all core experiments.....	209
Table 8. 4. Trends in weed growth under different nitrogen fertilisation rates across core experiments.....	210

List of Figures

Figure 3. 1. Mean precipitation and air temperature during 2013 and 2014 cropping seasons in comparison with the 10-year average. Royal Agricultural University meteorological station (NGR SP 42 004 011).....	29
Figure 3. 2. Field trial design.....	31
Figure 4. 1. Mean air temperatures and precipitation during the 2013 experimental period in comparison with the 10-year average. Royal Agricultural University meteorological station (NGR SP 42 004 011).....	52
Figure 4. 2. Effect of cultivation techniques and nitrogen fertilisation interaction on spring wheat establishment.....	55
Figure 4. 3. Effect of cultivation techniques and nitrogen fertilisation interaction on spring wheat total shoots production	56
Figure 4. 4. Effect of cultivation techniques and nitrogen fertilisation interaction on spring wheat biomass (May 2013)	58
Figure 4. 5. Effect of cultivation techniques and nitrogen fertilisation interaction on spring wheat N uptake (May 2013)	59
Figure 4. 6. Effect of cultivation techniques and undersowing interaction on spring wheat N uptake (May 2013)	59
Figure 4. 7. Effect of cultivation techniques and nitrogen fertilisation interaction on weed biomass (July 2013)	63
Figure 4. 8. Effect of cultivation techniques and undersowing interaction on weed biomass (July 2013).....	63
Figure 4. 9. Effect of nitrogen fertilisation and undersowing interaction on weed biomass (July 2013).....	64
Figure 4. 10. Effect of cultivation techniques and nitrogen fertilisation on weed N uptake (July 2013)	65
Figure 4. 11. Effect of cultivation techniques and undersowing on weed N uptake (July 2013).....	65
Figure 4. 12. Effect of cultivation techniques and nitrogen fertilisation on broadleaved weeds biomass (July 2013).....	67

Figure 4. 13. Effect of cultivation techniques and nitrogen fertilisation interaction on <i>Stellaria media</i> L. biomass (July 2013).....	68
Figure 4. 14. Effect of cultivation techniques and nitrogen fertilisation interaction on <i>Lolium perenne</i> L. biomass (July 2013)	69
Figure 4. 15. Effect of cultivation techniques and undersowing interaction on <i>Lolium perenne</i> L. biomass (July 2013)	70
Figure 4. 16. Effect of cultivation techniques and undersowing interaction on <i>Avena fatua</i> L. biomass (July 2013)	70
Figure 4. 17. Effect of cultivation techniques and undersowing interaction on spring wheat TGW.....	74
Figure 4. 18. Effect of cultivation techniques and nitrogen fertilisation interaction on legumes biomass (Harvest 2013).....	77
Figure 4. 19. Effect of cultivation techniques and nitrogen fertilisation interaction on total legumes N uptake (Harvest 2013)	78
Figure 4. 20. Effect of undersowing and nitrogen fertilisation interaction on total legumes N uptake (Harvest 2013)	78
Figure 4. 21. Soil mineral nitrogen (kg N ha ⁻¹) under three cultivation techniques.....	79
Figure 4. 22. Soil mineral nitrogen (kg N ha ⁻¹) under four nitrogen fertilisation treatments	80
Figure 4. 23. Effect of cultivation techniques and undersowing interaction on soil mineral nitrogen (kg N ha ⁻¹) (June 2013)	81
Figure 4. 24. Effect of nitrogen fertilisation and undersowing interaction on soil mineral nitrogen (kg N ha ⁻¹) (June 2013)	81
Figure 4. 25. Effect of cultivation techniques, nitrogen fertilisation and undersowing interaction on soil mineral nitrogen (kg N ha ⁻¹) (June 2013)	82
Figure 4. 26. Effect of nitrogen fertilisation and undersowing interaction on soil mineral nitrogen (kg N ha ⁻¹) (July 2013)	83
Figure 4. 27. Soil mineral nitrogen and wheat N uptake in 2013 season (kg N ha ⁻¹) under three cultivation techniques	83
Figure 4. 28. Soil mineral nitrogen and wheat N uptake in 2013 season (kg N ha ⁻¹) under four nitrogen fertilisation rates	84

Figure 4. 29. Effect of cultivation techniques and nitrogen fertiliser treatments on soil gravimetric moisture content (%) with mean precipitation during the 2013 experimental period in comparison to the 10-year average.....	85
Figure 4. 30. Effect of cultivation techniques and soil depth interaction on soil penetration resistance (kPa) (February 2013) with cultivation techniques soil disturbance depth.....	87
Figure 4. 31. Effect of cultivation techniques and soil depth interaction on soil penetration resistance (kPa) (mean values 2013) with cultivation techniques soil disturbance depth.....	88
Figure 5. 1. Mean air temperatures and precipitation during 2014 experimental period in comparison with the 10-year average. Royal Agricultural University meteorological station (NGR SP 42 004 011).....	114
Figure 5. 2. Cultivation techniques and nitrogen fertilisation interaction effect on overwintering weed N uptake.....	117
Figure 5. 3. Cultivation techniques, nitrogen fertilisation and undersowing interaction effect on overwintering weed N uptake.....	118
Figure 5. 4. Cultivation techniques and nitrogen fertilisation effect on overwinter legume biomass	119
Figure 5. 5. Cultivation techniques and nitrogen fertilisation effect on overwinter legume N uptake.....	119
Figure 5. 6. Cultivation techniques and undersowing interaction effect on wheat N uptake (May 2014)	122
Figure 5. 7. Cultivation techniques and undersowing interaction effect on wheat N uptake (July 2014)	123
Figure 5. 8. Cultivation techniques and nitrogen fertilisation interaction effect on legume biomass (May 2014)	125
Figure 5. 9. Cultivation techniques and nitrogen fertilisation interaction effect on legume N uptake (May 2014).....	125
Figure 5. 10. Cultivation techniques and nitrogen fertilisation interaction effect on weed biomass (July 2014).....	127
Figure 5. 11. Cultivation techniques and nitrogen fertilisation interaction effect on broadleaf weed biomass (July 2014)	128

Figure 5. 12. Cultivation techniques and nitrogen fertilisation interaction effect on weed N uptake (July 2014)	128
Figure 5. 13. Cultivation techniques and nitrogen fertilisation interaction effect on weed biomass (August 2014).....	130
Figure 5. 14. Cultivation techniques and nitrogen fertilisation interaction effect on weed N uptake (August 2014)	131
Figure 5. 15. Cultivation techniques and nitrogen fertilisation interaction effect on legume biomass (August 2014)	131
Figure 5. 16. Cultivation techniques and nitrogen fertilisation interaction effect on legume N uptake (August 2014).....	132
Figure 5. 17. Cultivation techniques and nitrogen fertilisation interaction effect on <i>Sinapsis arvensis</i> L. biomass (July 2014).....	134
Figure 5. 18. Cultivation techniques, nitrogen fertilisation and undersowing interaction effect on <i>Sinapsis arvensis</i> L. biomass (July 2014)	135
Figure 5. 19. Cultivation techniques and nitrogen fertilisation interaction effect on spring wheat height.....	136
Figure 5. 20. Cultivation techniques and nitrogen fertilisation interaction effect on spring wheat ears number	137
Figure 5. 21. Cultivation techniques and undersowing interaction effect on spring wheat straw biomass	139
Figure 5. 22. Cultivation techniques and nitrogen fertilisation interaction effect on spring wheat grains per ear	140
Figure 5. 23. Cultivation techniques and undersowing interaction effect on spring wheat harvest index.....	141
Figure 5. 24. Soil mineral nitrogen (kg N ha ⁻¹) under three cultivation techniques.....	145
Figure 5. 25. Soil mineral nitrogen (kg N ha ⁻¹) under four nitrogen fertilisation treatments	145
Figure 5. 26. Cultivation techniques and nitrogen fertilisation interaction effect on soil mineral nitrogen content (July 2014).....	146
Figure 5. 27. Cultivation techniques and undersowing interaction effect soil mineral nitrogen content (August 2014).....	146

Figure 5. 28. Effect of cultivation techniques, nitrogen fertilisation and undersowing interaction on soil mineral nitrogen (kg N ha ⁻¹) (August 2014)	147
Figure 5. 29. Soil mineral nitrogen (kg N ha ⁻¹) and total wheat N uptake under three cultivation techniques at each month of assessment	148
Figure 5. 30. Soil mineral nitrogen (kg N ha ⁻¹) and total wheat N uptake under four N fertilisation rates at each month of assessment.....	149
Figure 5. 31. Effect of cultivation techniques and nitrogen fertilisation on soil gravimetric moisture content (%) with mean precipitation during the 2014 experimental period in comparison to the 10-year average.....	150
Figure 5. 32. Cultivation techniques and nitrogen fertilisation interaction effect on soil moisture content (%) (June 2014)	151
Figure 5. 33. Cultivation techniques and undersowing interaction effect on soil moisture content (%) (August 2014)	151
Figure 5. 34. Effect of cultivation techniques and soil depth interaction on soil penetration resistance (kPa) (March 2014) with cultivation techniques soil disturbance depth	153
Figure 5. 35. Effect of cultivation techniques and soil depth interaction on soil penetration resistance (kPa) (June 2014) with cultivation techniques soil disturbance depth	154
Figure 5. 36. Effect of cultivation techniques and soil depth interaction on soil penetration resistance (kPa) (August 2014) with cultivation techniques soil disturbance depth	155
Figure 6. 1. Sixty years of long-term cropping season and off-season rainfall (mm). Royal Agricultural University meteorological station (NGR SP 42 004 011)	179
Figure 6. 2. Weighed mean differences in crop grain yield in HINiT compared with CT as affected by growing season rainfall (<300 mm; 300-500 mm; >500 mm).....	180
Figure 6. 3. Weighed mean differences in crop grain yield in LINiT compared with CT as affected by growing season rainfall (<300 mm; 300-500 mm; >500 mm).....	181
Figure 7. 1. Amount and energy inputs assigned to various agricultural management practices for all core experiments across the cropping seasons.....	186
Figure 7. 2. Energy consumption for all core experiments (MJ ha ⁻¹).....	190

Figure 7. 3. Direct and indirect input energy, and renewable and non-renewable energy for all core experiments	191
Figure 7. 4. Nitrogen fertilisation and undersowing treatment interaction effect on energy-use efficiency (2013)	194
Figure 7. 5. Nitrogen fertilisation and undersowing treatment interaction effect on energy productivity (2013)	194
Figure 7. 6. Cultivation techniques and nitrogen fertilisation treatment interaction effect on energy productivity (2014)	196
Figure 7. 7. Cultivation techniques and nitrogen fertilisation treatment interaction effect on energy-use efficiency (2014)	197
Figure 8. 1. Chapters interlinking different agricultural management practices, weather conditions, and energy and economic considerations for spring wheat production.....	202

List of Plates

Plate 3. 1. White mustard over existing soil cover. Winter 2012	28
Plate 3. 2. Trial design after cultivation techniques, nitrogen application and undersowing treatments.....	30
Plate 3. 3. Kuhn power harrow combination seed drill	33
Plate 3. 4. Seedbed conditions after contrasting cultivation systems	33
Plate 3. 5. Simba Xpress with a Simba ST bar fitted ahead	34
Plate 3. 6. Vaderstad Rapid-A system disc combined with seed drill	34
Plate 3. 7. Eco-dyn integrated seed drill	35
Plate 3. 8. Fertiliser sprayer	37
Plate 5. 1. Overwinter soil cover	112

Abbreviations

ADAS	Agricultural Division of Advisory Service
ASAE	American Society of Agriculture Engineers
BM	black medic
°C	degree Celsius (centigrade)
cm	centimetre
CT	Conventional tillage
DEFRA	Department for Environment, Food and Rural Affairs
DM	dry matter
FAO	Food and Agriculture Organization of the United Nations
g	grams
GS	growth stages
h	hour
ha	hectare
HGCA	Home-Grown Cereals Authority
HINiT	High intensity non-inversion tillage
IPCC	Intergovernmental Panel on Climate Change
kg	kilogram
kPa	kilo Pascal
l	litre
LINiT	Low intensity non-inversion tillage
LSD	Fisher's Protected Least Significant Difference
M	Molar
MAFF	Ministry of Agriculture, Fisheries and Food
mg	milligrams
min	minute
ml	millilitre

mm	millimetre
MPa	Mega Pascal
N	Nitrogen
N0	No nitrogen application
N70	70 kg N ha ⁻¹
N140	140 kg N ha ⁻¹
N210	210 kg N ha ⁻¹
NABIM	National Association of British and Irish Millers
NIAB	National Institute of Agricultural Botany
NO ₃ -N	Nitrate nitrogen
NH ₄ -N	Ammonium nitrate
Nus	No undersowing
P	Phosphorus
ppm	parts per million
SED	Standard error of the difference
SMN	Soil mineral nitrogen
SOM	Soil organic matter
t	tonnes
TGW	Thousand grain weight
UK	United Kingdom
UKMO	United Kingdom Met Office
UN	United Nations
WC	white clover
§	section symbol

CHAPTER ONE

Introduction

1.1. Background

Since the nineteen seventies the global population has doubled and, according to projections of the United Nations Population Division, today's population (7.3 billions) is likely to reach over nine billions by 2050 and perhaps 10.9 billion by 2100 (United Nations, 2013, 2014). Most recent estimates by the Food and Agriculture Organization of the United Nations (FAO) and the Government Office for Science (GO-Science) suggest that feeding a world population of ten billion people will demand raising overall food production by 60% from 2005-2007 levels (FAO, 2013a; GO-Science, 2011). However, a growing population also means that more land is needed to build infrastructures; to produce bio-energy crops and biodiversity protection; further reducing the expansion for agriculture land (FAO, 2013a). To meet these demands, cereal production will have to grow (FAO, 2009; Hawkesford *et al.*, 2013; Semenov *et al.*, 2014) but cereal crop yield growth will probably be at slower rate than in the past, with a predicted slowdown to 0.7% per annum (4.3 tonne ha⁻¹ by 2050) (FAO, 2009). Increased productivity is expected to be mainly from increases of yields and cropping intensity, and to a lesser extent from land expansion through sustainable intensification (Alexandratos & Bruinsma, 2012; Conway, 1998; Conway & Waage, 2010).

Global food production will also be challenged by unpredictable weather events, as result of changes to the climate due to the rise in greenhouse gases (GHGs) emissions, in some part from the effect of denitrification of nitrate fertilisers (Reynolds & Borlaug, 2006a,b; HGCA, 2012; Jenkinson, 2010). In fact, projections suggest that by 2030 the release of N₂O, as a GHG, will increase by 35 to 60% due to increases in N fertilisation and manure production (FAO, 2003). In order to avoid major climate changes, global emission of GHGs must reportedly be reduced by at least 50-60% (IPPC, 2007). Smith *et al.* (2007) suggested that by 2030 the potential mitigation options in agriculture will be about 89% from soil carbon sequestration, 9% from mitigation of methane and 2% from mitigation of soil N₂O emissions. The considerable challenge for plant and crop scientists' for the next century is to increase crop yields while reducing the use of

fertilisers and fossil fuel (Hamilton, 2009), and increasing resources-use efficiency to ensure greater sustainability (Hawkesford *et al.*, 2013). To overcome this challenge only a multidisciplinary approach is likely to success (Parry & Hawkesford, 2012).

During the nineteen sixties, wheat yield increases were reached due to the exploitation of semi-dwarf wheat varieties (Reynolds & Borlaug, 2006b). The dwarfing genes led to more grains per m² and greater harvest index (HI), also increasing the amounts of N fertiliser used globally, which could be used without causing lodging (Hawkesford *et al.*, 2013; HGCA, 2012; Reynolds & Borlaug, 2006b). As a result, wheat production became more intensified, resulting in increased food production for humankind through the end of the 20th century (Borlaug, 1971). However, global food security was never completely realised, particularly in developing countries (Blaustein, 2008; Reynolds & Borlaug, 2006a).

Regardless of the substantial increases in wheat yield in the last decades, further increase is still required to fulfil the demand of a growing population (Hawkesford *et al.*, 2013) (e.g. 50% in the next few decades; Reynolds *et al.*, 2012). However, global wheat yields seem to be reaching a plateau in many countries, which could relate to HI already reaching its maximum (Brown, 2012), e.g. 0.50 for spring wheat (Reynold *et al.*, 2009). In the UK, increases in wheat grain yield have been achieved due to genetic improvements, but at a rate of less than 0.1 tonnes ha⁻¹ per year (Hawkesford *et al.*, 2013). In order to accomplish the increasing yield needed, several national and international programmes have being created (e.g. 20:20 Wheat; G20-led Wheat Initiative; Wheat Yield Consortium). Further improvements will also require increases in total crop biomass and this will require further improvement in efficient use of the available resources (Long *et al.*, 2006; Parry & Hawkesford 2010; Reynolds *et al.*, 2012).

1.2.Problem statement and study aim

Farmers are facing increasing pressures to produce wheat that meets economic yield targets; specific nutrition and market requirements (Mercier & Hyberg 1995), while

making use of inputs in such a way that can ensure economic and environmental goals efficiently (DEFRA, 2011, 2014b; Hawkesford *et al.*, 2013).

As the awareness of combined ecological and economic costs of maintaining yields in intensive-input systems has grown, several studies have been concerned with the transition from conventional to organic agriculture systems (Clark *et al.*, 1999; Gomiero *et al.*, 2011; Hass *et al.*, 2001; Liu *et al.*, 2007; Pimentel *et al.*, 2005). However, organic production often results in lower yields, as a consequence often of greater weed and disease pressures and increasing soil nutrition deficiency (Moakes & Lampkin, 2011; Pridham & Entz, 2008; Ryan *et al.*, 2004). In the UK the classified organic land is decreasing, according to the National Survey produced by the UK Government Department for Environment, Food and Rural Affairs (DEFRA, 2014). In spite of a steady growth of the organic land since 2006, a decrease in the fully converted and in-conversion organic land was recorded from 2009 to 2013 by 11% and 80%, respectively (DEFRA, 2014). Such a decline in organic land is related in part to the reduction of the financial support to the organic sector by Government policies, as agri-environment payments in the UK are the lowest in the European Union accordingly to the Soil Association (2011).

Whilst accepting that organic farming is providing for a specialist niche food product market, the lower organic yields compared with conventional production systems together with a decrease in organic certified land, have made necessary a different approach to better combine cultural practices and the use of chemical and external inputs, so as to increase crop yield and making a better use of resources.

The overall aim of the investigation presented in this thesis is to study how the interaction between cultivation techniques, nitrogen fertilisation and legume-undersowing influence the ability of spring wheat to grow and develop. Further investigations consider treatment repercussions on the yield components that contribute to yield and grain quality. Towards increasing understanding of factors which influence crop development, essential soil processes, plant disease development and weed diversity.

1.3.Objectives of the study

The purpose of this study was to evaluate how a changing agronomy could support a more integrated crop management approach, by including combinations of external inputs and cultural practices. In order to also assess relationships between these management practices for both, increasing yields and a more profitable production cropping system.

The specific objectives of the study were:

1. To estimate the productivity of spring wheat intercropped with black medic (*Medicago lupulina* L.) and white clover (*Trifolium repens* L.) compared with sole wheat crop.
2. To evaluate the potential of black medic and white clover when undersown with spring wheat for weed suppression.
3. To determine the benefits that different soil cultivation techniques disturbance (conventional tillage, high intensity non-inversion tillage and low intensity non-inversion tillage) have on weed pressure and crop productivity.
4. To evaluate the effect of nitrogen nutrition on crop productivity and weed infestation.
5. To investigate nitrogen availability within all the interactions among components for spring wheat production.
6. To investigate weather patterns, in different growing seasons, affecting spring wheat production under the field site conditions.
7. To determine the profitability and energy consumption of the modified management practices for spring wheat production.

1.4.Thesis structure

Chapter 1 - Introduction. Brief overview of the justification of this study and outlining the aim and objectives examined.

Chapter 2 - Literature review. Review of the agricultural management practices, such as cultivation techniques, nitrogen fertilisation and undersowing, and how these determine

crop growth and development. This section also highlight challenges related to the intrinsic combination of factors affecting crop production.

Chapter 3 - Materials and methods. Reports the methodologies and techniques adopted in this study.

Chapter 4 - Core experiment – 2013. Influence of contrasting cultivation techniques, nitrogen fertilisation and legume undersowing in spring wheat. Reports outcomes of the core experiment during 2013 cropping season, and also provides a critical discussion of the influence of the management systems studied and the main conclusions.

Chapter 5 - Core experiment – 2014. Influence of contrasting cultivation techniques, nitrogen fertilisation and legume undersowing in spring wheat. Provides the outcomes of the core experiment performed during 2014 cropping season with critical discussion and main conclusions.

Chapter 6 - Effects of weather conditions on spring wheat performance following different cultivation techniques regimes. A meta-analysis of the effect of rainfall affecting spring wheat yield under contrasting cultivation techniques.

Chapter 7 - Economics and energy considerations for contrasting cultivation techniques, nitrogen fertilisation and undersowing in spring wheat. Illustrates the economic and energy consideration of the adoption of the agricultural management practices adopted.

Chapter 8 - General discussion and conclusions. A critical discussion of agricultural practices selected in this study is presented highlighting strengths and limitations. This section reaches the conclusions, and suggests implications of the study, with possibilities for further research.

CHAPTER TWO

Literature review

2.1.Cultivation techniques

Soil cultivation is considered one of the most important practices in agriculture due to its effect on the soil physical, chemical and biological properties (Wild, 1988). Cultivation techniques aim to prepare a suitable soil environment for seed emergence and plant growth. Ahn & Hintze (1990) define tillage as any mechanical or manual cultivation operations which create physical loosening of the soil, and according to Lal (1979) soil tillage is the alteration of the soil properties in order to modify soil conditions for crop production.

To obtain a more suitable medium for crop growth and development, soil manipulation by cultivation techniques can influence soil compaction, aeration and erosion; crop residues distribution into the soil; weeds and diseases suppression (Gajri *et al.*, 2002) and soil N mineralisation (López-Bellido *et al.*, 2005; Silgram & Shepherd, 1999). Additionally, energy and labour costs for crop production can also vary depending on the cultivation technique adopted (Ozpinar & Cay, 2005).

In the UK, tillage systems can be categorised into two main classes: inversion tillage system also referred as conventional tillage, and non-inversion tillage also broadly known as conservation tillage (Davies & Finney, 2002).

2.1.1. Conventional tillage

Conventional tillage (ploughing) involves primary and secondary cultivation operations to prepare the seedbed (Gajri *et al.*, 2002). Primary cultivation is the main operation consisting on the inversion of the soil by the use of mouldboard ploughs. Those used currently are mostly reversible and consist of a coulter frame with a series of mouldboards, forward rake points, vertical plates and tail pieces attached to the rigid plough frame (Soffe, 2003). Depending on soil type and cultivation speed, the mouldboards ploughs working depth is around 20 to 25 cm depth (Brassington, 1986).

This primary operation is often followed by secondary cultivations which creates a smooth seedbed by the use of a power harrow generally in conjunction with a seed drill (Bell, 1996; Soffe, 2003). A power harrow consists of almost vertical pair of tines, each one attached to a gear, which drives or is driven by adjacent gears, resulting in contra-rotating sets of neighbouring tines (Soffe, 2003). The harrows move rapidly across the soil surface leaving a level and smooth seedbed (Brassington, 1986). The combination of primary and secondary cultivations provide a regular soil surface which allow good seed-soil contact (Braunack & Dexter, 1989), and bury crop residues which can also disrupt weed, pest and diseases life cycles, giving better crop germination and growth conditions (Gajri *et al*, 2002).

However, several detrimental effects of continuous conventional tillage practices have been reported (Table 2.1). Despite these reported negative effects, the use of the plough is still justified by many farmers to loosen crusted and compacted soils, towards optimising yields, but in some cases its use may just be for ease of drilling (Morris *et al*, 2010).

Table 2. 1. Some detrimental effects of conventional tillage

Effect	Reference
Increase in soil erosion	El Titi (2003); Lal <i>et al.</i> (2007); Larson & Osborne (1982)
Increase in organic matter oxidation	Lal <i>et al.</i> (2007); Mitchell <i>et al.</i> (2004); Six <i>et al.</i> (2000)
Lower work rates	Akbarnia <i>et al.</i> (2010); Ozpinar & Çay (2005)
Increase in cost and energy-use	Akbarnia <i>et al.</i> (2010); Arvidsson (2010); Ozpinar & Çay (2005)

2.1.2. Non-inversion tillage

Non-inversion tillage systems have now become more widely used alternatives to conventional tillage on different soil types (IPCC 2014a, Kassam *et al.*, 2009), including heavy clays (Cannell & Hawes, 1994; Holland, 2004), on around 15% of the arable land in Europe (Jones *et al.*, 2006). In the UK, conservation tillage systems involve the use of tines and disc harrows without inverting the soil, but incorporating

much of the crop residues in the top-soil layers although maintaining a proportion on the soil surface (Carter *et al.*, 2003; Peigné *et al.*, 2007; Soane *et al.*, 2012). Tines can be found in different shapes and angles, from straight to curve and either fixed or moving with crumblers attached to mounted sections or front boards (Christian, 1994). Tines are designed to remove any compacted layer by lifting and shattering the soil and to breakdown residues (Morris *et al.*, 2010), and are often followed by disc harrows at a shallow depth of around 12 to 15 cm, depending upon soil type and cultivation speed (Soffe, 2003). Disc harrows comprise two or four adjustable axles with concave discs mounted along its length (SMI, 2003). Axles are angled for forward motion. The front axle discs cut and throw soil outwards, while rear axle discs throw soil inward (Soffe, 2003). Press wheels are usually attached at the rear of the cultivator in order to level and firm the soil surface prior to seed drilling (Morris *et al.*, 2010). It is often required for drilling, the use of a high output cultivation drill combined with tines and discs ahead of the seed coulter in order to facilitate seed depth and emergence (Bell, 1996).

Non-inversion tillage creates seedbed and soil environment conditions that allow seed germination with less soil movement and inversion (Cannell, 1994). Table 2.2 summarises reported effects of non-inversion tillage.

Table 2. 2. Some reported non-inversion tillage effects

Effect	Reference
Reduction of soil erosion	Carter (1991); Hussain <i>et al.</i> (1999); López <i>et al.</i> (1996)
Improved soil moisture retention	Carter (1991); Hussain <i>et al.</i> (1999); Sharma <i>et al.</i> (2011)
Reduction of organic matter oxidation	Morris <i>et al.</i> (2010); Soane <i>et al.</i> (2012)
Increase in soil microbial activity	Mitchell <i>et al.</i> (2004); Morris <i>et al.</i> (2010); Soane <i>et al.</i> (2012)
Reduction of costs and energy-use	Jones <i>et al.</i> (2006); Knight (2004); Sánchez-Girón <i>et al.</i> (2004); Triplett & Dick (2008)
Higher work rates	Jones <i>et al.</i> (2006); Triplett & Dick (2008)

2.2.Cultivation techniques and soil properties

Variations of tillage intensity commonly alter the soil physical, chemical and biological properties resulting in changes of the soil functional quality, as widely reported (Aziz *et al.*, 2009; Celik *et al.*, 2011; Ding *et al.*, 2011).

2.2.1. Soil physical properties

Around 50% of the crop root mass grows in the seedbed created by the tillage operations (Finney & Knight, 1973). Tillage effects on soil physical properties are highly important therefore, to the essential crop growth and development. Differences in tillage intensity can temporally change soil physical conditions (Rasmussen, 1999; Tebrügge & Düring, 1999). If the soil is too hard, too dry, lacks of oxygen or its temperature is far from optimal, it can limit root growth (Bengough *et al.*, 2006; Kaspar & Bland, 1992). However, these changes depend on soil type, climate conditions and the type and extent of previous tillage systems adopted (Rhoton, 2000).

Some of the most important soil physical properties of the seedbed created by the cultivation techniques practices are soil bulk density and soil penetration resistance (Cassel, 1982). Soil bulk density influences soil water and air dynamics and also how crop roots grow (Unger & Cassel, 1991). The immediate tillage effect is loosening the soil which then decreases the bulk density. However, this effect is not permanent and it can be reduced or even reversed with subsequent events such as continuous soil disturbance or increasing rainfall conditions. Among cultivation techniques, conventional tillage reportedly presents lower soil bulk density compared with non-inversion tillage systems, especially in the upper 0-15 cm soil layer (Kaspar *et al.*, 1992; Munkholm *et al.*, 2003; Stokes *et al.*, 1992). However, several other studies reported higher bulk density under conventional tillage compared with non-inversion tillage systems (Griffith *et al.*, 1977; Lal, 1979; Sharma *et al.*, 2011). This is mainly related to the high presence of residues under non-inversion systems which reduce the bulk density, particularly in relatively compacted soils (Ghuman & Sur, 2001). Such variable results highlight the widely contrasting impact differences of tillage and their dependence on environmental conditions and soil type. Moreover, alterations of bulk

density by tillage systems tend to be maximised after initial cultivation practices are undertaken, but might possibly decrease with time or even become insignificant at the end of the cropping season (López *et al.*, 1996; Pelegrin *et al.*, 1990).

Cultivation techniques can also affect the soil total pore space arrangement which is inversely correlated with bulk density (Carter & Ball, 1993; Guérif *et al.*, 2001). Reductions in total pore space can indicate increases in soil penetration resistance. This is especially relevant for crop development as it can affect the crop root growth (Cassel, 1982). Atwell (1993) reported that a penetration resistance greater than 2MPa can potentially reduce root growth. However, a penetration resistance of 1.5MPa has been used as a reference to assess tillage practices impact on soil loosening (Carter, 1988). Soil penetration resistance reportedly increases when increasing soil depth, mainly due to increases in soil particles friction (Bradford, 1986; Campbell & O'Sullivan, 1991; Grant & Lafond, 1993). Differences between cultivation techniques on penetration resistance have been widely studied. Non-inversion tillage reportedly presents higher penetration resistance in the upper 15cm soil layer than conventional tillage (Aikins & Afuakwa, 2012; Grant & Lafond, 1993; Munkholm *et al.*, 2003; López *et al.*, 1996). This has also been commonly observed particularly in soils with poor structure and low soil organic matter (Hill, 1990). However, cultivation effects on soil penetration resistance is often reported to be temporal and dissipating after several years as the soil recover its former state, as reported by several studies (Campbell & Henshall, 1991; Franzluebbbers *et al.*, 1995; Martinez *et al.*, 2008). Penetration resistance also decreases when soil moisture content increases, as water absorption weakens connections between soil particles (Marshall & Holmes, 1988).

Soil moisture content reportedly increases with increasing soil residues cover. Such residues enhance soil structure improving infiltration and protecting the soil from evaporation and runoff (Allmaras *et al.*, 1977; López *et al.*, 1996). Greater presence of plant residues on the soil surface under non-inversion tillage, therefore, results in higher soil moisture compared with conventional tillage, as widely reported (De Vita *et al.*, 2007; López *et al.*, 1996; Sharma *et al.*, 2011). However, differences in soil moisture content between tillage practices are also highly dependent in weather pattern particularly rainfall, as reported by De Vita *et al.* (2007).

2.2.2. Soil chemical properties

Changes in tillage intensity can also potentially influence soil reaction (pH), organic matter stratification and nutrient distribution (Staley, 1999; White, 1990), although this also depends on environmental conditions and soil type (López-Fando & Pardo, 2009; Thomas *et al.*, 2007). Several authors (Hickman, 2007; Houx III *et al.*, 2011; Vijaya Bhaskar *et al.*, 2013a) reported decreases in soil pH with less tillage intensity. This soil acidification under non-inversion tillage systems is related to the breakdown of crop residues on the soil surface, which possibly result in organic acids accumulation, causing lower soil pH (Blevins *et al.*, 1977). Additionally, without soil inversion, variations on soil pH are slow, being influence by slow movement of carbonic acid (H₂CO₃) through the soil profile (López-Pando & Pardo, 2009).

2.3.Cultivation techniques and soil nitrogen availability

Soil movement created by the cultivation techniques reportedly results in temporary increase of the availability of soil mineral nitrogen (SMN). This occurs by modifications of the soil environment, such as water content and temperature, and increasing growth and activity of soil fauna promoting organic matter oxidation (Silgram & Shepherd, 1999; Wild, 1988). However, the SMN availability created by tillage operations varies with operations timing and weather conditions during and after cultivation performance, as widely reported (El Titi, 2003; Kapusta *et al.*, 1996; Radford *et al.*, 1992). Additionally, Silgram & Shepherd (1999) reported that such tillage effects are often noticeable just for a few weeks. The presence and nature of crop residues incorporated or left on the soil surface also affects the availability of SMN. Crop residues with high C/N ratio cannot provide enough N to the microbial population, promoting rapid N-immobilization and limiting its availability to the subsequent crop (El Titi, 2003; Silgram & Shepherd, 1999).

Non-inversion tillage systems reportedly can result in lower SMN than conventional tillage due to transitory N limitations (Al-Khasi *et al.*, 2005; Braim *et al.*, 1992; McConkey *et al.*, 2002; Silgram & Shepherd, 1999). This is perhaps the result of lower organic matter oxidation in spring, autumn and winter but increased mineralisation in

summer (Blevins & Frye, 1993; Riley *et al.*, 1994; Schomberg *et al.*, 1994). Additionally, lower SMN has often been related to higher N-immobilization and nitrification of the crop residues (López-Bellido *et al.*, 2013; Soane *et al.*, 2012). However, increases in SMN availability by increasing tillage intensity do not necessarily support greater crop productivity (Greenwood, 1982), due to the increasing risk of leaching which can result in nitrates being less available during high crop demand (Silgram & Shepherd, 1999).

2.4.Cultivation techniques and weed suppression

Weeds can provide food and habitats for a range of beneficial organisms which can also benefit the crop if weed population is low (Cussans, 1968; Storkey, 2006). Above critical thresholds, weeds can compete and reduce the main crop yield and quality, which make weeds a major factor affecting cereal production (Froud-William *et al.*, 1983b; Marshall *et al.*, 2003). Weed occurrence is significantly influenced by the cultivated crop species, crop rotation, tillage practices, timing and type of weed management, and primarily by environmental factors, such as weather conditions, location and season of the year (Derksen *et al.*, 1995; Shrestha *et al.*, 2002; Tuesca *et al.*, 2001).

Tillage practices often modify weed abundance and species composition in crops (Ball & Miller, 1993; Froud-Williams *et al.*, 1981; Hakansson, 2003). Tillage affects the weed populations by changing the seed distribution both vertically and horizontally, affecting the seeds viability, emergence and seedling survival. Also by dismembering vegetative structure of perennial weeds, and thereby stimulating bud growth and depleting their food reserves (Clements *et al.*, 1996a; Streit *et al.*, 2002; Swanton *et al.*, 2000). Inverting the soil reportedly buries most seed initially present in the soil surface, but it can also relocate buried seeds back to the topsoil (Colbach *et al.*, 2006; Hakansson, 2003). Without soil inversion, weed seeds are maintained on the surface and distributed less down the soil profile due to less soil movement (Ball, 1992; Froud-Williams *et al.*, 1983a).

Among weed species, it has been widely reported that increasing tillage intensity increases broadleaf weed species frequency but decreases grass weeds (Froud-Williams *et al.*, 1983b; Tuesca & Puricelli, 2007; Tuesca *et al.*, 2001). Grass weed species are also highly susceptible to mechanical disturbance, which restricts their presence under conventional tillage. Limited soil disturbance under non-inversion tillage systems can result commonly in a greater incidence of grass weeds (Hakansson, 2003). Additionally, non-inversion tillage systems have also been related to increases in grass weeds, such as *Poa* spp and *Alopecurus myosuroides* (Froud-Williams *et al.*, 1983b), which are able to germinate on a soil surface covered by residues that can maintain moisture (Mester & Buhler, 1991; Tuesca & Puricelli, 2007). Generally, broadleaf weed seeds have greater longevity and marked dormancy; with annual inversion of the soil bringing to the soil surface dormant buried seed allowing their germination (Froud-Williams *et al.*, 1983b). Some broadleaf species, such as *Chenopodium album* L., have also been linked more to conventional tillage due to the accumulation of residues on the soil surface under non-inversion systems creating more shady conditions, reducing those species ability to germinate (Teasdale, 1993).

Tillage systems also affect weed population by altering soil temperature and moisture required by many weed species to break dormancy (Thompson *et al.*, 1977). Temperatures above or below the optimum range for germination, can possibly decrease seeds germination whilst, some species germinate better under alternating rather than constant temperatures (Vincent & Roberts, 1977).

In terms of weed species diversity, soil disturbance created by cultivation often prevents one species becoming dominant within the community and can, therefore, increase diversity (Cardina *et al.*, 2002; Clark *et al.*, 2007; Sousa, 1984).

2.4.1. Cultivation techniques and herbicide-use

Historically, the inclusion of herbicide applications reduced the importance of cultivation techniques as a major mean of weed control (Froud-Williams *et al.*, 1983b). However, an increasing number of weeds showing resistant to a wide range of herbicide active ingredients have been identified in the UK, particularly black grass (*Alopecurus*

myosuroides), wild oat (*Avena fatua* L.) and Italian ryegrass (*Lolium perenne* L.) (Davies & Finney, 2002). Herbicide resistant weeds are increasing interest in the complementary use of cultivation techniques and herbicide applications as more integrated weed control strategy (Finch *et al.*, 2014). For instance, the use of pre-emergence herbicides can exert beneficial weed control under non-inversion tillage systems with most of the weeds remaining in the soil surface before the establishment of the crop (Calado *et al.*, 2010). In addition, several authors report that herbicide application can reduce possible differences between tillage for weed suppression (Derksen *et al.*, 1995; Vijaya Bhaskar *et al.*, 2014b).

However, herbicide's effectiveness for weed control can also reportedly be reduced by residues left on the soil surface (Buhler, 1995). Sadeghi *et al.* (1998) found 70% of herbicide interception by crop residues, reducing its subsequent suppression effect on weeds. Additionally, the herbicide type and application time can also influence the efficacy of controlling weeds (Derksen *et al.*, 1995; Soane *et al.*, 2012; Streit *et al.*, 2002). For example, Anken *et al.* (2004) reported that the broad-spectrum systemic herbicide such as glyphosate reduces its effectiveness under low temperatures and also frequent rainfall condition after application. Underlying cultivation techniques influences on weed control are important in selecting an effective herbicide, with the associate cost also affecting the crop enterprise profitability (Sayili *et al.*, 2006).

2.5.Cultivation techniques and diseases

The effects of tillage on plant disease can vary depending on crop type, biology of the pathogen, soil type and prevailing environmental conditions (Bailey & Duczek, 1996; Conway, 1996). Cultivation techniques modify the soil environment affecting pathogens, but can also change the distribution of crop residues which for many pathogens are primary inoculum source (Jenkyn *et al.*, 2004). Under non-inversion tillage, higher residues can provide substrate for residue borne pathogens but, also modifies soil temperature and moisture encouraging diseases (Bockus & Shroyer, 1998; Sutton & Vyn, 1990; Watkins & Boosalis, 1994). If the contaminated residues are destroyed and buried by ploughing the inoculum is also destroyed, but if residues are

left undisturbed the pathogen survives resulting on the disease development (Sumner *et al.*, 1981). This is the case of some cereal diseases presented in Table 2.3, which are favoured by non-inversion tillage, while others occur more frequently under conventional tillage (Table 2.3).

Table 2. 3. Some reported cereal diseases favoured by different cultivation systems

Non-inversion tillage	Reference
<i>Gaeumannomyces graminis</i> var. <i>tritici</i> (take-all of wheat)	Ennaifar <i>et al.</i> (2005); Sumner <i>et al.</i> (1981)
<i>Pyrenophora tritici repentis</i> (tan spot of wheat)	Sumner <i>et al.</i> (1981)
<i>Mycosphaerella graminicola</i> (leaf blotch)	Vijaya Bhaskar <i>et al.</i> (2014a)
<i>Rhizoctonia solani</i> (rhizoctonia stunt)	HGCA (2008a); Sumner <i>et al.</i> (1981)
Conventional	
<i>Bipolaris sorokiniana</i> (common root rot)	Bailey & Duczek (1996); Conway (1996)

2.6.Cultivation techniques and crop yield

Crop yield is affected by interactions of the crop with such factors as the growing environment, soil type and tillage intensity (Rasmussen, 1999), although interactions are not always consistent nor predictable (Jones *et al.*, 2006). Cultivation techniques effects on the soil environment affect crop growth and development and also final crop yield. Several studies report crop yields under conventional tillage are either higher or comparable to those from non-inversion tillage (Gruber *et al.*, 2012; Rasmussen, 1999; Soane *et al.*, 2012; Vijaya Bhaskar *et al.*, 2013b). Table 2.4 summarise reported detrimental effect of non-inversion tillage resulting in lower yield compared with conventional tillage. Mixing and/or incorporation of the residue in the soil allows better seed-soil contact, increasing germination and crop establishment and resulting in better final yield under conventional tillage (Table 2.4)

Table 2. 4. Detrimental effects of non-inversion tillage on final grain yield

Effect	Reference
Poor crop establishment	Hemmat & Taki (2001); Vijaya Bhaskar <i>et al.</i> (2013b); Wilkins <i>et al.</i> (1989)
Higher weed and disease pressure	Gruber <i>et al.</i> (2012); McConkey <i>et al.</i> (1996); Vijaya Bhaskar <i>et al.</i> (2013b)
Higher soil penetration resistance	Cassel <i>et al.</i> (1995); Rasmussen (1999); Vyn & Raimbault (1993)
N immobilization	Chen <i>et al.</i> (2007); López- Bellido <i>et al.</i> (2013); Wang <i>et al.</i> (2012); Wang <i>et al.</i> (2015)

The effect of non-inversion tillage on crop yield is also highly variable depending on weather conditions, as reported by several authors (e.g. De Vita *et al.*, 2007; López-Bellido *et al.*, 1998; Ordoñez-Fernández *et al.*, 2007). Such variations are partially associated with modifications in soil moisture content, e.g. greater water storage during dry periods (Bonfil *et al.*, 1999). Carr *et al.* (2006) found almost 40% greater yield under non-inversion than conventional tillage, attributed to the greater soil moisture holding capacity in alleviating crop water stress.

Soil type plays a very important role on the performance of the cultivation techniques and their effect on the final crop yield. Knight (2004) reported that non-inversion tillage yielded lower than conventional tillage in two of three years under clay soil while, non-inversion tillage resulted in the greatest yield in three years on a light chalkland soil. Kumar *et al.* (2013) also reported greater yield under non-inversion compared with conventional tillage in a sandy loam soil, attributed to better soil moisture content under non-inversion treatment. In contrast, Munkholm *et al.* (2003) concluded that yield decreased under non-inversion tillage on light-textured soils mainly due to greater soil compaction in a moist climate. Repeated tillage systems effects on yield are sometimes contradictory and depend on site-specific factors, such as soil type, environmental conditions and previous management history (Arvidsson *et al.*, 2013; Morris *et al.*, 2010; Rasmussen, 1999).

2.7. Cultivation techniques and grain protein

For the wheat industry, grain protein is one of the main factors being sought (Wall *et al.*, 1979), with bread making potential largely determined by the quantity and quality of the grain protein (Hruskova & Famera, 2003). Several factor interactions are reported to influence grain quality, including cultivar, soil type, grain storage conditions, N availability and environmental conditions (Blumenthal *et al.*, 1991; Borghi *et al.*, 1997; De Vita *et al.*, 2007; Gooding & Davies, 1997).

Cultivation techniques effects on grain protein are mainly through modifications of soil moisture and soil nitrate content (López-Bellido *et al.*, 1998). It has been stated that an excess of soil moisture can lead to a decrease in grain protein content (Robinson *et al.*, 1979), while water stress increases protein content (Rao *et al.*, 1993; Terman *et al.*, 1996). Conventional tillage can result in higher protein content compared with non-inversion tillage systems, mainly due to higher N availability (De Vita *et al.*, 2007; López-Bellido *et al.*, 1998). In contrast, others studies reported no significant differences between tillage treatments on grain protein content (Bassett *et al.*, 1989; Cox & Shelton, 1992).

2.8. Nitrogen fertilisation and plant growth

A high amount of N is required by crops as it is an essential constituent of chlorophyll, and a major component of proteins and enzymes which catalyse essential reactions for the crop life (Blevins, 1989). Wheat development can be influenced by N applications mainly in four difference ways presented in Table 2.5.

These N fertilisation influences are interrelated even though they might not always be beneficial. For instance, it has been reported that increases in crop aboveground biomass might lead to increases in ears number per plant with high grain number. This can result in high grain yield while harvest index (the ratio of harvested grain to the total aboveground dry matter) stays unchanged (Cabrera-Bosque *et al.*, 2009; Cossani *et al.*, 2009).

Table 2. 5. Nitrogen fertilisation effects on cereal plant growth and development

Effect	Reference
Increase in the size and duration of canopy growth	Gooding & Davies (1997); Ottensson <i>et al.</i> (2008)
Affects individual plants within the crop stand, e.g. determining tillers survival which will sustain ears	Greenwood (1982); HGCA (2008b); Ottensson <i>et al.</i> (2008); Sarandon & Gianibelli (1990)
Produces heavy stems potentially increasing losses by lodging and disease attacks	Hay & Walker (1989); HGCA (2008b); Gerba <i>et al.</i> (2013)
Determines grain quantity and quality	Gooding & Davies (1997); Greenwood (1982); Smith <i>et al.</i> (1990)

N uptake by a crop and its effect on annual yields are often variable even on the same site and when the N available fulfils the crop demand. This considerable variability is due to the vegetative growth relative to grain yield and how it responds to water and nutrient availability, crop management and climate conditions (Ferguson, 1967; Hay & Walker, 1989; López-Bellido, *et al.*, 1998). For instance, Pearman *et al.* (1977) observed a lack of N effect on grain yield attributed to lower autumn precipitations which reduced any potential N leaching leaving high level of residual N in the soil. This was also reported by several other studies (Barracough *et al.*, 1989; Cabrera-Bosque *et al.*, 2009; López-Bellido *et al.*, 1998) while others have also reported decreases in yields with increasing N application (Cossani *et al.*, 2009; Fois *et al.*, 2009; Gooding & Davies, 1997). However, N application often increases grain yield (Abad *et al.*, 2005; Brennan *et al.*, 2014; Halvorson *et al.*, 2001; Raun *et al.*, 2010).

Regarding grain qualities, it has been suggested that N application improves the specific grain weight (Pushman & Bingham, 1975). However, several studies report negative effect of N fertiliser on specific grain weight mainly by increases in the grains number per ear (López-Bellido *et al.*, 2000; Rasmussen *et al.*, 1997; Smith & Davies, 1990).

2.8.1. Nitrogen-use and nitrogen-use efficiency

In the UK, wheat crops generally need large amount of available N in order to produce high grain yields and high protein content (Barraclough *et al.*, 2010; Greenwood, 1982). In order to optimise resources use, it is essential to understand the complex inter-relationship between N availability, uptake and utilisation during crop growth (Barraclough *et al.*, 2010).

N-use efficiency (NUE) in cereal crops is defined as grain dry matter yield per unit of available N from the soil and/or fertiliser (yield efficiency) (Moll *et al.*, 1982; Le Gouis *et al.*, 2000),

$$\text{NUE (kg kg}^{-1}\text{)} = \frac{\text{Grain yield}}{\text{N available}} = \text{NUpE} \times \text{NUtE}$$

NUE can be further divided into two primary components: (1) N uptake efficiency (NUpE); the efficiency with which the plant absorbs N from the soil, and (2) N utilisation efficiency (NUtE); the efficiency with which the absorbed N is used to produce grain dry matter (Moll *et al.*, 1982; Le Gouis *et al.*, 2000),

$$\text{NUpE} = \frac{\text{Total plant N uptake}}{\text{N available}} \quad \text{NUtE} = \frac{\text{Grain yield}}{\text{Total plant N uptake}}$$

Therefore, improvements in either NUpE or NUtE could improve NUE.

In crops generally, including cereals, the full recovery of N fertilisers is never achieved mainly due to the dependence on several factors such as soil type, climate conditions, crop variety, SMN availability, and the nature of the inorganic fertiliser applied (Burger & Jackson, 2004; Schulten & Schnitzer, 1998). In the case of wheat, Ladha *et al.* (2005) reported that the average recovery of N fertiliser is around 54%. In a study on winter wheat grown in eastern England by Powlson *et al.* (1992), wheat recovered an average of 68% of the N fertiliser applied, while 18% was retained in the soil and 14% lost by leaching and de-nitrification. The efficiency in the use of the N absorbed by the plant to produce grain yield is highly dependent on genotypic variations but also by the response to the environmental conditions (Hirel *et al.*, 2007).

The efficiency in the partitioning of aboveground biomass per unit of N uptake to the grain (NUtE) is an important process influencing yield quantity and grain quality (Le Gouis *et al.*, 2000; Ortiz-Monasterio *et al.*, 1997; Simpson *et al.*, 1983). From the same N uptake, crops with higher NUtE will produce higher yields or the same yield with a lower N uptake.

The N contained in the aboveground crop can count as much as 50 to 80% of the grain N content at harvest (Cox *et al.*, 1985; HGCA, 2008b; Sarandon & Caldiz, 1990; Xu *et al.*, 2005) depending on the variety and environmental conditions. Palta *et al.* (1993) reported increases in N remobilisation efficiency from the vegetative organs to the grain in conditions of water stress during grain filling period, as the plant needs to make better use of its N accumulated at anthesis. Conversely, N remobilization during grain filling can be reduced by foliar diseases (Dimmock & Gooding, 2002) and high temperatures (Heitholt *et al.*, 1990) due to acceleration of senescence of the vegetative parts (Gooding & Davies, 1997).

In other words, NUE can sometimes decrease with N application (Campbell *et al.*, 1977; Clark *et al.*, 1990; Sieling *et al.*, 1998) due to wrong application practices such as high amounts and/or wrong timing. If the amount of N applied is too high, the plant might not be able to take-up all decreasing NUpE, if so the N uptake may be utilised less efficiently (NUtE), and hence less NUE. In addition, NUE highly depends upon the response to water availability during the growing season (Hatfield & Prueger, 2004; Semenov *et al.*, 2007). During lower water availability, NUE decrease is mainly due to less N available for the crop. At high water availability, in contrast, the N available could be lost by leaching.

2.8.2. Nitrogen fertilisation and grain protein

Grain protein is often considered the most important singular criterion in defining grain quality specifically for the bread making industry. However, grain protein content is determined by genotype (Johnson *et al.*, 1985; Stoddard & Marshall, 1990) and influenced by growing environmental conditions (Blackman & Payne, 1987; Rao *et al.*,

1993). N availability and distribution is one of the major determining factors of the grain protein concentration (Fowler & Brydon 1989; Hirel *et al.*, 2007; Hunter & Stanford, 1973; Olson *et al.*, 1976). N application can also improve grain protein, commonly showing a linear response in a wide range of growing environments (Gao *et al.*, 2012; López-Bellido *et al.*, 1998; López-Bellido *et al.*, 2001). However, weather conditions throughout the growing season play an important role highlighting any possible positive or negative effects of N fertilisation on grain protein content. Applying N with appropriate soil moisture content mainly leads to increases in yield with limited effect on grain protein (Campbell *et al.*, 1977; Clark *et al.*, 1990; Smith and Gooding, 1999), whereas water stress reduces crop yield, while it may increase grain protein (Campbell *et al.*, 1977; Terman *et al.*, 1996). For instance, Kosmolak & Crowle (1980) reported an increase of 1% of the protein content with an application of 26 kg N ha⁻¹ to a crop yielding 2.5 t ha⁻¹; whereas Penny *et al.* (1978) reported the same increase of protein when only 46 kg N ha⁻¹ was applied but to a crop yielding 5 t ha⁻¹, with this variation mainly attributed to water availability across the growing season.

Increasing soil moisture can potentially reduce grain protein content, particularly if this occurs prior the grain filling due to the dilution of early N reserves by increases in vegetative growth (Smith & Gooding, 1999). Conversely, rainfall events during the summer can have positive effects on grain N particularly due to wetting / drying cycles of the soil affecting N mineralisation, beside the increase of rainfall at this point possibly increases diseases levels which reduces yields and therefore increasing the grain protein (Farrant, 1972; Smith *et al.*, 1990). In other words, the effect of the interaction between N application and predominant weather conditions on the protein content can also depend on the crop developmental stage.

2.8.3. Nitrogen fertilisation and weeds

Competition between plants is highly dependent on many factors including availability of nutrient, especially N (Sweeney *et al.*, 2008). Weeds can be directly influenced by N fertilisation, but also indirectly by increasing crop competitiveness against weeds. Crop ability to suppress weeds is increased by N application, mainly by promoting faster

growth which increases the competition for resources, resulting in the reduction of weed species number and biomass (Grundy *et al.*, 1993; Jørnsgård *et al.*, 1996). Conversely, weed growth can respond positively to N fertilisation possibly due to differential NUE compared with the crop (Di Tomaso, 1995; Sheibani & Ghadiri, 2012). The response to N fertilisation also greatly differs among different weed species (Iqbal & Wright, 1997). In a long-term experiment, Moss *et al.* (2004) reported that *Stellaria media* L. was highly favoured by N-rich conditions while other species such as *Medicago lupulina* L. and *Equisetum arvense* L. were highly disadvantaged. Iqbal & Wright (1997) also reported that the relative competitive abilities of *Sinapis arvensis* L. were greater than a wheat crop when 120 kg N ha⁻¹ was applied, while *Phalaris minor* Retz. was less competitive under the same conditions. Jørnsgård *et al.* (1996) observed that *Lamium* spp and *Veronica* spp had lower N optima than wheat.

2.9. Intercropping

Liebman & Dyck (1993) defined intercropping as spatial diversification of cropping systems by growing two or more crop simultaneously. This practice has been known to improve soil fertility, especially in the case of leguminous crops cropped with non-legumes crops (Fujita *et al.*, 1992; Shafi *et al.*, 2007; Thorsted *et al.*, 2006). The ability of legumes to fix atmospheric N represents a valuable source of organic N by utilising their nodulated-roots and residues (Anil *et al.*, 1998; Bakht *et al.*, 2009; Giller, 2001; Giller & Wilson, 1991; Kumar & Goh, 2002). However, the amount of N fixed can vary with the legume species, environmental conditions and crop management (Hamdi, 1995).

Intercropping is a practice widely used in developing countries as a way of increasing crop production per land area when capital investments are limited (Dakora 1996; Francis, 1986; Machado, 2009). However, interest in the use of intercropping in developed countries has also increased as a potential way to maintain or increase crop production while reducing fertilisers and pesticide use (Horwith, 1985; Machado, 2009).

2.9.1. Undersowing

In the UK, intercropping is mainly in the form of undersowing (Hartl, 1989) which consists of growing two or more crops sharing the same area for a part of their life cycle (Duncan & Schapaugh, 1997; Wallace *et al.*, 1996). For instance, perennial legumes can be sown either with winter or spring sown cereal crop in the spring in order to develop a subsequent ley to avoid a period of bare ground (Hartl, 1989). Legumes add potential benefits to the intercropping system such as improvement of soil fertility by fixation and N release (Badaruddin & Meyer, 1990; Hauggaard-Nielsen *et al.*, 2001).

Other potential benefits of undersowing is the suppression of weed, pest and diseases although it can also vary depending on the species grown, biomass production, time of sowing, harvest management and prevalent environmental conditions (Badaruddin & Meyer, 1990; Hartwig & Ammon, 2002). Weed suppression by the intercrop is mainly by the reducing the available space for weeds to germinate and grow, and by enhancing competition for resources such as light and nutrients (Anil *et al.*, 1998; Liebman, 1986; Liebman & Dyck, 1993). Banik *et al.* (2006) and Hauggaard-Nielsen *et al.* (2003) reported greater suppression of weeds under intercropping over monocrops, mainly due to higher interspecific competition for resources and complementarity between intercrop species in improving their competitive abilities against weeds. Additionally, reduction in the incidence of pests and diseases by intercropping compared with monocropping has been reported (Hiltbrunner *et al.*, 2002; Teasdale, 1996; Theunissen, 1997; Vilich-Meller, 1992).

Undersowing can also affect soil quality by covering the soil and reducing N leaching, while its biomass adds organic matter (Duda *et al.*, 2003; Hartwig & Ammon, 2002). This addition of organic matter by the undersown species can improve the soil structure and can also reduce soil compaction reducing soil deterioration (Bristow & Horton, 1996; Teasdale & Mohler, 1993). However, undersowing can lead to competition between the crops growing together (Vandermeer, 1989) which can result in a decrease in growth, development (Crawley, 1997) and final yield of the main crop (Clements & Williams, 1967).

2.9.1.1. Undersowing effects on yield

Interactions among plant species inside the intercropping system occur during the growth process as the plants exploit the same resources (Vandermeer, 1989), which generally creates competition (Andersen *et al.*, 2004; Hauggaard-Nielsen *et al.*, 2001). This interspecific competition for resources commonly results in intercrop yields intermediate to that of the sole crop (Hauggaard-Nielsen *et al.*, 2001; Thorsted *et al.*, 2006). Competition for resources can be reduced by manipulating the initial advantage of one of the crops inside the intercropping system through delaying understorey crop sowing, or by increasing the seed rate of the main crop (Andersen *et al.*, 2007). Charles (1958) reported, for instance, yield reduction in a cereal crop when the understorey crop was sown at the same time as the main crop. No reduction was observed when the understorey was sown when the main crop was already well established.

Nevertheless, in some cases the intercrop can enhance the productivity of the system (Fukai & Trenbath, 1993; Vandermeer, 1989). The cereal yield advantage of undersowing has been reported to be in part due to vigorous growth of the undersown species, which suppress weeds without affecting the main crop. In the case of undersown legume by enhancing the main crop N uptake (Brennan & Smith, 2005; Hauggaard-Nielsen *et al.*, 2008; Zhang & Li, 2003).

2.9.2. Cereal-legume intercropping

Growing a cereal crop with a legume is the most common type of intercropping (Francis, 1989). However, it is important to consider that the species should have a synergistic effect with each other and no antagonistic interactions. When using legumes it is important to acknowledge the time at which the leguminous plants are releasing the previously fixed N, and when the cereal crop is capable of utilising (Charles, 1958). Generally, legume swards have to be ploughed in order to release N and then followed by a cereal crop to take up the resultant N. Species such as white clover (*Trifolium repens* L.) recovers and spreads rapidly following suppression due to its prostrate and stoloniferous growth, being suitable for this practice (Jones, 1992). However, negative

effects have been observed when incorporating more than one crop in a cropping system (Pridham &ENZ, 2008). Table 2.6 shows detrimental effects of intercropping resulting in lower cereal yield. Conversely, several studies reported potential benefits of the incorporation of legumes in a continuous arable cropping system without compromising the cereal grain yield (Table 2.6).

Table 2. 6. Adverse and beneficial effects of legume intercropping on cereal productivity

Detrimental effects	Reference
Competition for resources between crops	Thorsted <i>et al.</i> (2006); Jones & Clements (1993)
More land to produce the same yield than monoculture (greater land ratio)	Lithourgidis <i>et al.</i> (2011); Ofori & Stern, (1987); Reynolds <i>et al.</i> (1994)
Beneficial effects	
Suppression of weeds	Haymes & Lee (1994); Thiessen-Martens <i>et al.</i> (2005); Walker <i>et al.</i> (2011).
Increase N availability to main crop	Känkänen <i>et al.</i> (2001); Thiessen-Martens <i>et al.</i> (2005); Walker <i>et al.</i> (2011).

In cropping systems with external input such as N fertiliser, legume-cereal intercrops reduce their advantages over cereal only crops (Ghaley *et al.*, 2005; Jensen, 1996; Ofori & Stern, 1987). This may relate to the increase of competitiveness of wheat adversely affecting the legume intercrop (Gooding & Davies, 1997), and the negative responses of legumes to N fertilisation (Moss *et al.*, 2004).

In summary, the success of cereal-legumes intercrops depends on the development and maintenance of the desirable balance of the components (Clements *et al.*, 1994b).

2.9.2.1. Undersowing effect on grain protein

The response of cereal grain protein when intercropped with legume species is highly variable (Berry *et al.*, 2002; Jones & Clements, 1993). Several authors reported greater grain protein content of the intercropped cereal primarily due to the complementary use of resources by the intercropped species (Hauggaard-Nielsen *et al.*, 2001; Jensen 1996;

Lauk & Lauk, 2008). Although this is thought to be related to the lower competitiveness of the legume crop for soil N and competition for resources such as light, thus limiting total cereal biomass production (Gooding *et al.*, 2007; Hauggaard-Nielsen *et al.*, 2006).

CHAPTER THREE

Materials and methods

The methods and techniques used in the present study were consistent between experiments; and are included in this chapter to avoid repetition. When methodologies differ or additional information is required, this will be given in the appropriate chapter.

3.1. Experimental site

Field experiments were initiated in March 2013 (Core experiment I) and March 2014 (Core experiment II). The field experiments were conducted on Evesham soil series, with characteristics shown in Table 3.1., at the Royal Agricultural University's Harnhill' Manor Farm (NGR SP 075 006), near Cirencester, UK, situated at 51° 42'N latitude, 01° 59'W longitude, at an altitude of 132 m above sea level. Soil texture was clay with a soil pH around neutral and initial soil mineral nitrogen (SMN) Index of 0 (Table 3.1).

Table 3. 1. Initial physiochemical properties of the experimental site

Parameters	Values	Parameters	Values	Index
pH	6.90	SMN (kg ha ⁻¹)	25.33	0
% Sand	22.59	P (mg l ⁻¹)	8.00	1
% Silt	37.48	K (mg l ⁻¹)	208.67	2
% Clay	39.93	Mg (mg l ⁻¹)	105.27	3

Following spring wheat harvest on 22 August 2012, broad-spectrum systemic herbicide – glyphosate [*N*-(phosphonomethyl)glycine], was applied at a rate of 4 l ha⁻¹ across the entire experimental site (2.7 ha). After herbicide application, white mustard (*Sinapsis alba* L. cv. Tilney) was broadcasted over existing soil cover to grow over the winter 2012 (Plate 3.1). White mustard was used as a break crop to help to reduce infections of take-all disease (*Gaeumannomyces graminis* var. *tritici*) (HGCA, 2006; Vijaya Bhaskar, 2014). The cropping system history is summarized in Table 3.2.

Plate 3. 1. White mustard over existing soil cover. Winter 2012



Table 3. 2. Cropping systems used in the site from 2007 to 2012

Year	Cropping system
2007 – 2010	White clover (<i>Trifolium repens</i> L.) (WC) – Perennial ryegrass (<i>Lolium perenne</i> L.) ley
2010 – 2011	Organic winter wheat (<i>Triticum aestivum</i> L.) cv Clare, bi-cropped with black medic (<i>Medicago lupulina</i> L.) cv Virgo Pajbjerg (BM) and WC cv Aberpearl.
2011 – 2012	Organic spring wheat cv Paragon bi-cropped with BM and WC

Previous research on the site during 2010 - 2012 evaluated three contrasting cultivation techniques, including conventional plough-based tillage and two non-inversion tillage systems which differ on the degree of soil disturbance and residue incorporation (30% or >50% of residue cover) (Vijaya Bhaskar, 2014). Similar treatments were then used in the present study (See § 3.2.1).

3.1.1. Meteorological conditions


During the 2013 cropping season, the maximum and minimum air temperature was recorded in July of 19.0 °C and March of 3.1 °C while, the maximum and minimum rainfall documented was in March of 76.8 mm and July of 31.3 mm (Figure 3.1). The

2013 season experienced lower air temperatures compared with the long-term average. The 2013 spring period faced higher precipitations, particularly during March and May, while precipitations at summer time were lower compared to the seasonal average.

In the 2014 cropping season, the maximum and minimum air temperature were recorded in July of 18.0 °C and March of 7.17 °C, whereas the maximum and minimum rainfall documented were in May of 97.3 mm and March of 39.5 mm. 2014 experienced higher air temperatures across the season with higher precipitations during the spring period and lower precipitations recorded at summer time compared to long-time average (Figure 3.1).

The 2013 growing season experienced lower air temperature compared to the 2014 cropping season except during July and August when temperatures were higher in 2013 than in 2014 season. The 2013 season experienced lower precipitations compared with 2014 season except in March when precipitations were higher in 2013 (Figure 3.1).

Figure 3. 1. Mean precipitation and air temperature during 2013 and 2014 cropping seasons in comparison with the 10-year average. Royal Agricultural University meteorological station (NGR SP 42 004 011)



3.2. Experimental design and treatments

For both cropping seasons, field trials followed a split-split plot randomized block design. The selected field of 2.7 ha was divided into three separate blocks (90 m x 100 m). Each block was divided into three fully randomized main plots (cultivation treatments) of 30 m x 100 m. The main plots were divided into four fully randomized sub-plots (N treatments) (7.5 m x 100 m) and these were divided into three fully randomized split-sub plots (undersowing treatments) (30 m x 33.3 m) (Plate 3.2 & Figure 3.2). The treatment structure was as follows:

Spring wheat (3 block) x cultivation technique (9 main plot) x N fertilisation (36 sub plot) x undersowing (108 split-sub plot)

Plate 3. 2. Trial design after cultivation techniques, nitrogen application and undersowing treatments





Figure 3. 2. Field trial design



Before land preparation was initiated (20 March 2013 and 24 March 2014), foliar contact herbicide, glyphosate, was sprayed at 2 l ha⁻¹. Previous research on adjacent organic land reported considerable high weed pressure, particularly of grass weeds (Vijaya Bhaskar, 2014). Application of herbicide was made to reduce weed competition towards improving future crop yields.

The crop structure for both cropping seasons is detailed in Table 3.3.

Table 3.3. Experimental wheat crop details

Variety	Spring wheat cv Paragon
Sowing date	10 April 2013 / 18 April 2014
Seed rate	200 kg ha ⁻¹
Thousand grain weigh	41.6 g
Seeds m ⁻²	480
Harvest date	27 August 2013 / 31 August 2014

Twenty days after sowing spring wheat in 2013, the experimental area was treated with triple superphosphate fertiliser (TSP) [Ca(H₂PO₄)₂H₂O] applied at a rate of 75 kg P₂O₅ ha⁻¹. Phosphate fertiliser was based on initial background level of soil P (Table 3.1) (DEFRA, 2010).

After 2013 harvest the field was left with surface residues over the winter and then 2014 core experiment was established.

3.2.1. Details of experimental treatments

3.2.1.1. Cultivation techniques

Three cultivation treatments (main plots) were examined in this study, two non-inversion tillage treatments and one conventional tillage treatment. More detailed information for each treatment is as follows:

- Conventional Tillage (CT); involved one pass of a five furrow Kverneland reversible plough, to a working depth of 20 cm, and then furrow pressed. Following, a Kuhn power harrow combination seed drill was used at a working depth of 8 cm (Plate 3.3). The percentage of soil cover by crop residues after drilling was typically assumed as 0% (Plate 3.4).

Plate 3. 3. Kuhn power harrow combination seed drill



Plate 3. 4. Seedbed conditions after contrasting cultivation systems



- High Intensity – Non-inversion Tillage (HINiT); consisted of two passes of a mounted Simba X-press with a Simba ST bar fitted ahead (Plate 3.5) at a working depth of 25 cm and 12 cm. A Vaderstad Rapid-A system disc in combination with seed drill (Plate 3.6) was employed to a working depth of 8 cm. The percentage of soil cover by crop residues after drilling was typically assumed as 30% (Plate 3.4).

Plate 3. 5. Simba Xpress with a Simba ST bar fitted ahead



Plate 3. 6. Vaderstad Rapid-A system disc combined with seed drill



- Low intensity – Non-inversion Tillage (LINiT); involved one pass of a mounted Simba X-press with a Simba ST bar fitted ahead at a working depth of 25 cm and 12 cm. A heavy planter Eco-dyn integrated seed drill followed at a working depth

of 26 cm to drill the crop (Plate 3.7). The percentage of soil cover by crop residues after drilling was typically assumed as >50% (Plate 3.4).

Plate 3. 7. Eco-dyn integrated seed drill



All the treatments had a uniform planting depth of 2 cm and coulter row spacing of 12.5 cm for 200 kg ha⁻¹ volume of seeds. Crop residues were a mix of straw and herbicide-killed weeds and legumes from the previous cropping season (Plate 3.4).

Varietal choice

Spring wheat was selected due to increasing recognition of its high premium opportunities. In addition, unlike winter wheat, spring wheat has shorter growing season and present fewer tillers allowing studying more comprehensively agricultural practice-induced effects on yield and yield components. The spring wheat variety was selected with particular attention to its potential bread-making quality and disease resistance. The oldest spring wheat in Group 1 (bread-making/milling) of the HGCA Recommended list – cv Paragon was selected, as it has a very good disease resistance and relatively long but stiff straw (HGCA, 2013). Paragon produces grains with very high protein content, and it has been reported to produce quality grain even under untreated trial conditions (NABIM, 2013).

Sowing date

Grain yield and other characteristics of wheat can be influenced by variations in sowing date (Hayward, 1990). Variable weather patterns across seasons can affect the selection of a suitable date for sowing. In the present study, the sowing date was adopted on the basis of pre-sowing cultivations and weather conditions, particularly prevailing precipitation.

Seed rate

Seeding rate can vary depending on sowing date, crop variety, soil type and weather conditions (Finch *et al.*, 2014). Spring wheat varieties with lower production of tillers than winter wheat (Wibberley, 1984) should be complemented with high plant density to compensate towards securing a more optimum plant population. In this study, due to late sowing and in order to compare different cultivation techniques, a seeding rate of ≥ 400 seed m^{-2} was chosen.

Drilling depth

To ensure good seed distribution, drilling aims to be sufficiently deep enough. It can, however, be challenging to control due to its high dependency on seedbed preparation influenced by the contrasting pre-sowing cultivation techniques. A poor consolidated seedbed can, for example, result in deep seed placement while if the seedbed is too dense seeds may not be adequately covered resulting in losses due to pest damage (Atkinson, 2008). Additionally, the presence of residues in the non-inversion tillage systems may interfere with drills and generate both, an uneven sowing depth and seeding row spacing (Siemens *et al.*, 2004). Sowing depth recommendations for wheat are usually between 4 and 2 cm depth, depending upon soil type and conditions (HGCA, 2008b). For all the experiments, a uniform sowing depth of 2 cm and coulter row spacing of 12.5 cm were used for all the combination drills. Even though, the sowing depth and row space were kept almost uniform, the drills performance may vary depending on the pre-sowing cultivation techniques used.

3.2.1.2. Nitrogen treatments

Four mineral N fertiliser treatments (sub-plots) were used in the present study,

- N0 – No N fertilisation
- N70 – 70 kg N ha⁻¹
- N140 – 140 kg N ha⁻¹
- N210 – 210 kg N ha⁻¹

N fertiliser was applied as ammonium nitrate solution, NH₄NO₃ (34.5% N), sprayed using a 6 m width sprayer, Case IH GEM 3000sp (Plate 3.8). Tramlines were used as a form of traffic control. At all application rates, half of the treatments dose was applied at seedling growth and the remainder was applied at tillering stage. Further details are listed in Table 3.4.

Plate 3. 8. Fertiliser sprayer



Table 3. 4. Nitrogen treatments application dates with corresponding spring wheat growth stages (GS)

Year	Timing	Date	On/after GS
2013	1 st	30 April	13
	2 nd	23 May	21
2014	1 st	10 May	13
	2 nd	30 May	21

Nitrogen fertiliser type

Several studies have compared different types of inorganic N fertilisers, reporting no differences in yield response between fertilisers type (Christensen & Meintz, 1982; Garrido-Lestache *et al.*, 2004), suggesting that the amount of N applied and timing are more relevant, irrespectively of product type. Ammonium nitrate (NH₄NO₃) is one of the most common inorganic sources of N, containing 35% of N and with very rapidly available nitrate (Finch *et al.*, 2014).

Nitrogen doses

Soil N supply (SNS) is defined as the soil mineral N (SMN) in soil plus the estimate of mineralisable N. The highest N dose used in this study, 210 kg N ha⁻¹, was selected following recommendations by DEFRA (2010), based on the SNS status assessed before the establishment of the field trial (February 2013). Subsequent treatments rates were based on 2/3 (140 kg N ha¹) and 1/3 (70 kg N ha¹) of the highest treatment dose respectively.

Timing and splitting of nitrogen applications

N fertiliser timing is one of the main factors influencing yields and grain quality (Borghi *et al.*, 1997; López-Bellido *et al.*, 1998). However, it has been argued in the case of spring wheat crops, for a most efficient utilization, N application timing is generally less critical due to the rapid crop growth and development (Gooding & Davies, 1997). Regardless of timing, splitting of N fertiliser doses have also been reported to potentially improve wheat N-use efficiency (López-Bellido *et al.*, 2005; Mahler *et al.*, 1994) although, the proportions of the split should be determined based on initial soil fertility status. In the present study, N applications were split by half, being applied at the seedling stage (on/or after GS13) and at tillering (on/or after GS21). Splitting applications were based on recommendations for the higheest doses, N210 and N140 (DEFRA, 2010), whereas the lowest dose (N70) was split in order to maintain the same conditions for all applications.

3.2.1.3. Undersowing

For the undersowing treatments (split-sub plot), two legumes species were compared, white clover (*Trifolium repens*) cv Aberpearl at 7 kg ha⁻¹ (WC) and black medic (*Medicago lupulina*) cv Virgo Pajbjerg sown at 8 kg ha⁻¹ (BM). Legumes were hand broadcasted into the established spring wheat stand or not undersown (Nus), on 7 May 2013 and 12 May 2014, on/or after GS13.

Undersown legume species choice

Legumes species were selected based on their ability to grow into the emerged crop; their ability to compete against weeds, and potentially less aggressive competition towards the main crop (Döring *et al.*, 2013; Rosenfeld *et al.*, 2011; Vijaya Bhaskar *et al.*, 2013c).

White clover (WC) is a perennial legume with a slow establishment but considerable production of biomass (Döring *et al.*, 2013) which made it particularly suitable for undersowing, and it also offers good weed control (Rosenfeld *et al.*, 2011). WC ability to fix atmospheric N results in a widely variable contribution of N which is estimated of about 250 kg ha⁻¹ of N per year (Smýkal *et al.*, 2015). This legume species also exhibits great winter hardiness and persistence, attributable to the formation of a complex network of stolons (Jones, 1992; Smýkal *et al.*, 2015). A great number of WC varieties have been developed as a result of a large number of breeding programs (Döring *et al.*, 2013) and grouped into small, medium and large leaved types (Rosenfeld *et al.*, 2011). WC cv Aberpearl is a small leaved variety, best suitable for undersowing due to less aggressive growth against the main wheat crop.

Black medic (BM) is a short-lived annual/biannual specie (Clapham *et al.*, 1987) slow to establish, but with a fast develop (Hartmann *et al.*, 2009) and great biomass production allowing good weed suppression (Döring *et al.*, 2013). BM cv Pajbjerg is the only variety currently use in the UK (Rosenfeld *et al.*, 2011).

3.3. Aboveground assessments

3.3.1. Developmental stages

For all sampling assessments, growth stages (GS) were based on the decimal code devised by Zadoks *et al.*, (1974). Date of maturity was judged when grain moisture content had reached 15%. Details of dates and developmental stage for each plant assessment are given in Table 3.5.

Table 3. 5. Dates and growth stages for above ground crop assessment

Assessments	Approximate wheat growth stage
Wheat establishment and tillers number	On/or before GS31
Wheat shoot numbers	On/or after GS31
Growth assessments (wheat, weeds & legumes – BM & WC – biomass)	On/or before GS31; on/or after GS61; on/or after GS71
Plant height	On/or after GS71
Wheat ears number	On/or after GS71
Diseases assessment	On/or after GS71
Biological harvest	On/or after GS87

3.3.2. Assessments

Plant establishment

Plant establishment was determined by counting the number of plants inside a 0.25 m² random quadrat for each split-sub plot with ten replications. The assessments were performed twice a week following drilling to monitor the emergence of the crop, and were continued until no further plants emerged.

Tillers number and shoots number

Wheat tillers number and total wheat shoot number (main stem and tillers) were assessed using a 0.25 m² random quadrat with ten replications per each split-sub plot.

Growth assessments

In 2013 two destructive harvests were assessed by growth analyses, while in 2014 three destructive harvests were sampled. Samples of all the aboveground plant material were collected from random areas in order to avoid bias. Sampling was conducted with three replications and avoiding the previously disturbed area. All the aboveground plant material inside a 0.25 m² random quadrat was recovered using scissors or secateurs. All cut material was placed as quickly as possible inside labelled plastic bags and sealed. In the laboratory, wheat, legumes (WC & BM) and weed species were separated and fresh weights recorded. Dry weight (DM) yield was recorded after drying samples at 105°C overnight.

The non-undersowing (Nus) treatment was not completely free of legumes due to some small natural regeneration, by stolons (WC) and adventitious bud in roots (BM), in spite of pre-cultivation herbicide applications. Legumes were, therefore, separated from weed even though they were considered legumes weeds in these plots rather than deliberately undersown legumes.

Plant height and wheat ears number

Height of the main wheat shoot was measured from the tip of the ear to the nearest centimetre from the ground level, at fifteen randomly selected fertile shoots per split-sub plot. Measurements were taken using a rising disk apparatus, a rectangular (30 x 50 cm) expanded polystyrene weighing 50 g with a hole centrally bored (4 cm diameter) into which a graduated wooden rod was inserted. The scale was positioned so that the top of the disc recorded zero when the base of the disc was at soil level.

The number of wheat ears was assessed using a 0.25 m² random quadrat with ten replications per each split-sub plot.

Disease assessment

The incidence of septoria leaf blotch of wheat (*Zymoseptoria tritici* Desm. Quaedvlieg & Crous), and leaf rust of wheat (*Puccinia triticina* Erikss.) severity percentages were evaluated on 10 flag leaf samples per each split-sub plot, using an illustrated key (James, 1971). Take-all (*Gaeumannomyces graminis* var *tritici*) severity percentages

was evaluated in 10 plants per each split-sub plot, using an illustrated key (MAFF, 1976). Illustrated keys are given in Appendix 1.

Final biological harvest

The trial was hand harvested following the same protocol used for growth assessments, using 0.25 m² random quadrats, with three replications at each split-sub plot. All the plant material was separated into wheat (separated by ears and straw), legumes (BM & WC) and weeds, and their fresh weight assessed. Ears were cut off at the peduncle and number recorded. All samples were dried at 105°C overnight and DM recorded. Ears were threshed by hand and the amount of grain was weighed to obtain total grain weights and grain yield, which was corrected to 15% grain moisture content. Thousand grain weight (TGW) was recorded after using an automatic feeder and counter (Farmtec, Scunthorpe). Harvest index (HI) was determined as the ratio of grain weights to the total wheat aboveground biomass (Donald & Hambling, 1976).

Plant nitrogen content

In order to determine the N content, all the plant samples were course milled and then sub sampled and further micro-milled (0.5 mm sieve) (Cyclotec 1093 Sample Mill) to obtain a fine sample with a narrow particle size distribution. A sub sample of 25 mg (\pm 0.05 mg) of the ground material plus 50 mg of tungsten oxide were placed into aluminium foils and weighed on a five place analytical balance. Encapsulated samples were then analysed on an Elemental Cube CNS auto analyser (Elementar Analysensysteme GmbH). Grain protein content was obtained by multiplying grain N% by 5.7 (Osborne, 1907). Total grain N uptake, total wheat N uptake (total grain N uptake plus total straw N uptake), total legume and weeds N uptake, and the N harvest index were calculated using the following formulas (Fageria *et al.*, 2008; Moll *et al.*, 1982),

$$\text{Total grain N uptake (kg ha}^{-1}\text{)} = \left(\frac{\text{Grain yield (t ha}^{-1}\text{)}}{100} \times \text{grain N\%} \right) \times 1000$$

$$\text{Total plant N uptake (kg ha}^{-1}\text{)} = \left(\frac{\text{DM (t ha}^{-1}\text{)}}{100} \times \text{N\%} \right) \times 1000$$

$$\text{Nitrogen harvest index (\%)} = \left(\frac{\text{N\% in grains}}{\text{N\% in grains} + \text{N\% in straw}} \right) \times 100$$

Nitrogen-use efficiency parameters

The following N-efficiency parameters were calculated for each treatment:

- N-use efficiency (NUE; kg kg⁻¹) as the ratio of grain yield to N supply.
- N uptake efficiency (NUpE; kg kg⁻¹) as the ratio of total aboveground plant N uptake to N supply.
- N utilisation efficiency (NUtE; kg kg⁻¹) as the ratio of grain yield to total aboveground plant N uptake.
- N harvest index (NHI; %) as the ratio of grain N uptake to total aboveground plant N uptake.

The amount of N supply is the sum of SMN (ammonium plus nitrate) at sowing, mineralised N and N fertiliser applied. Mineralised N was estimated as the total plant N uptake at harvest plus mineral N left in soil profile after wheat harvest in control plots (N0) minus initial mineral N presented before wheat seeding (Huggins & Pan, 1993; Miao *et al.*, 2015). N in the roots was not considered in the calculations, due to practical experimental difficulties and sometimes low biomass. The terminology for N efficiency parameters follows Delogu *et al.* (1998), Huggins & Pan (1993), López-Bellido *et al.* (2005); Moll *et al.* (1982), Pierce & Rice (1988) and Sowers *et al.* (1994).

3.4. Soil assessments

3.4.1. Soil chemical analysis

Field sampling

Soils were sampled to a depth of 25 cm at fifteen randomly points following a W pattern within each split-sub plot using a Dutch auger. Samples were hand-crumbled

and hand-mixed to form representative samples. Samples were placed in sealed and labelled bags, and rapidly transferred to the laboratory. Initial processing and analyses occurred within 12 hours of sampling. The following analyses were then conducted:

Soil mineral nitrogen

Soil available mineral N (ammonium and nitrate) (SMN) was determined by a potassium sulphate (K₂SO₄) extraction method (Faithfull, 2002). Soils were sampled to a 25 cm depth, following the field sampling procedure previously described. Assessments were initiated from March (2013 and 2014) and continued throughout the cropping seasons at 4-week intervals and finished in August 2013 and 2014. Analyses were conducted on the same day of sampling.

In the laboratory, samples were passed through a 6.7 mm mesh sieve and any plant material or visible stone were removed. From each sample, three analytical replications of 25 g (\pm 0.02 g) were taken and transferred onto a labelled extraction bottle adding 100 ml of 0.5 M K₂SO₄ solution. In each extraction batch, 2 blanks were included. The extraction bottles were shaken vigorously using a shaker unit (Gerhardt, Germany) for 30 mins and allowed to stand for 15 min. The soil solutions were then filtered through Whatman No 40 filter papers onto a 60 ml labelled sample bottles, discarding the first couple of drops. Samples bottles containing the collected extracts were store frozen in preparation for analysis of NH₄ and NO₃ on a FiAstarTM 5000 Analyser (DK) based on flow injection analysis and colorimetric methods.

To convert SMN (mg l⁻¹) to an area basis (kg ha⁻¹), the soil bulk density is required and calculated as Unkovick *et al.* (2008),

$$\text{SMN} \left(\frac{\text{mg}}{\text{kg}} \right) = \frac{\text{extraction volume (ml)} \times \text{total extract mineral N (NO}_3^-, \text{NH}_4^+) \left(\frac{\text{mg}}{\text{l}} \right)}{\frac{\text{Soil dry weight (g)}}{1000}}$$

$$\text{SMN} \left(\frac{\text{kg N}}{\text{ha}} \right) = \text{SMN} \left(\frac{\text{mg}}{\text{kg}} \right) \times \text{bulk density} \times \text{depth factor}$$

Soil moisture content

Soil dry matter was assessed by weighing 50 g (± 0.05 g) of fresh soil samples from each split-sub plot and oven dried at 105°C overnight and the weights retaken. The soil moisture (gravimetric) percentage was obtained using the following formula (Brady & Weil, 1999),

$$\text{Soil gravimetric moisture(\%)} = \frac{\text{soil fresh weight} - \text{soil dry weight}}{\text{Soil dry weight}} \times 100$$

Assessments were initiated from March (2013 and 2014) and continued throughout the cropping seasons at 4-week intervals and finished in August 2013 and 2014.

Soil pH

Determination of soil pH was conducted by weighing three analytical replicas (per sample) of 20 g of sieved (≤ 2 mm) air-dry soil into shaking bottles and adding 50 ml of deionised water and shaken vigorously for 15 min, using a shaker unit (Gerhardt, Germany) and allowed to stand. A pH electrode was immersed in the solution, swirling a couple of times allowing the pH to stabilize before taking readings (Faithfull, 2002). Before pH measurements, calibration of the pH meter (Omega Engineering, USA) was performed according to manufacturer's instructions using buffers of pH 7.0 and 4.0 to cover the pH range of the soil samples.

Soil potassium and magnesium

Soil potassium (K) and magnesium (Mg) were assessed before the establishment of the experimental study (March 2013). K and Mg were determined by ammonium nitrate extraction (Faithfull, 2002). From each split-sub plot, three analytical samples of 10 g (± 0.05 g) of sieved (≤ 2 mm) air-dry soil was transferred into a 150 ml shaking bottles and 50 ml of N ammonium nitrate dispensed. Bottles were shaken vigorously using a shaker unit (Gerhardt, Germany) for 30 mins and allowed to stand for 15 mins. Solutions were subsequently filtered through Whatman No 2 filter papers, discarding the first couple of drops. The concentration of K in the extraction samples and in six working standards (0, 1, 2, 3, 4 and 5 $\mu\text{g K ml}^{-1}$) and two blanks, were determined using a flame photometer.

From the standard graph, the $\mu\text{g K ml}^{-1}$ equivalent in the samples were determined and blank value subtracted and difference multiplied by five (initial extraction ratio), resulting in the number of mg l^{-1} extractable K in the air-dry soil samples.

To determine Mg concentration, sub-samples of the ammonium nitrate extraction were used. From the extracted solutions, 5 ml was pipetted into a 100 ml volumetric flask adding 1 ml buffer and diluted to 100 ml's with deionised water. The concentration of Mg in the solutions and in six working standards (0, 0.2, 0.4, 0.6, 0.8 and 1 $\mu\text{g Mg ml}^{-1}$) and two blanks were determined by using an atomic absorption spectrophotometer (Thermo Scientific Inc., USA).

Soil phosphorus

Soil phosphorus (P) content was measured before the establishment of the experimental study (March 2013) by the Olsen Method (Olsen *et al.*, 1954). From each split-sub plot, three analytical samples of 5 g (± 0.05 g) of sieved (≤ 2 mm) air-dry soil were weighed and transferred into 150 ml shaking bottles. A teaspoon of powdered charcoal and 100 ml of sodium bicarbonate (NaHCO_3) reagent, at pH 8.5, were added to the bottles and then shaken vigorously using a shaker unit (Gerhardt, Germany) for 30 mins and allowed to stand for 15 mins. Solutions were filtered through Whatman No 2 filter papers, discarding the first few drops of filtrate. From the extractions, 5 ml was pipetted into a 100 ml conical flask slowly adding 1 ml of 1.5 M sulphuric acid. When frothing ceased from releasing carbon dioxide, 20 ml of ammonium molybdate (1.2% m/v)/ascorbic acid solution was added and allowed to stand for 30 mins. Working standard solutions of 0.25, 0.5, 1, 2, 3, 4, 5 and 6 $\mu\text{g P ml}^{-1}$ were used in order to obtain the equivalent $\mu\text{g P ml}^{-1}$ of the samples and 2 blanks were used. Finally, the absorbance of the samples, standards and blanks were measured using a spectrophotometer (Cecil Instruments Lt., UK) at 880 nm wave-lengths. P was assessed before the establishment of the experimental study (March 2013).

3.4.2. Soil physical analysis

Soil texture

Soil texture was determined following the Bouyoucos Hydrometer Method (Bouyoucos, 1962) before the establishment of the experimental study (March 2013). From each main plot, three analytical samples of 50 g (± 0.01 g) of sieved (≤ 2 mm) air-dry soil were placed into 250 ml shaking bottles. 100 ml of Calgon solution was added and shaken for 400 mins. The solutions were transferred into a 1000 ml cylinder and diluted to 1000 ml's using deionised water. The top of the cylinder were sealed with parafilm and inverted 20 times; placed on the bench and timed immediately with a stop watch. A hydrometer was inserted into each cylinder without disturbing the solution approximately 20 sec prior to a reading being taken; then removed and rinsed immediately. Readings were taken at 40 secs, 4 mins, 37 mins and 2 hours. Readings gave the density in g l^{-1} . To correct the readings for temperature and density, readings were calibrated against the hydrometer in the Calgon-water control solution and subtracted from all the readings. The percentage of sand, silt and clay fractions was plotted on the triangular texture chart to determine texture class (MAFF, 1988).

Soil bulk density

Soil bulk density was determined following the ISO 11272:1998 method (ISO 11272:2014) for non-gravelly soils. Bulk density was only measured before the cultivations treatments performance (2013), in order to obtain initial values. Soil samples were taken from each main plot with three replications at a depth of 30 cm with 5 cm intervals. Bulk density was determined on undisturbed soil samples using a steel sampler cylinder of 358.36 cm^3 which was driven perpendicular, without deflection and compaction, into the soil surface. Samplers were removed carefully in order to prevent any loss of soil. Using a flat-bladed knife, the excess of soil from the sample holder was removed leaving the bottom of the sample holder flat and even with edges of the holder. Samplers were placed in a labelled plastic bag and sealed. In the laboratory the weight of the soil samples at a 5 cm interval were recorded to calculate soil water content. Samples were placed in an oven at 105°C until constant mass was reached (minimum 48 hours) and then the dry weight was recorded. Using the oven dry weight and the

volume of the sample holder, the soil bulk density was calculated using the following formula,

$$\text{Soil bulk density (g cm}^{-3}\text{)} = \frac{\text{Soil dry mass (g)}}{\text{Volume of the sample (cm}^{-3}\text{)}}$$

Soil bulk density value was then used to convert SMN (mg l^{-1}) to an area basis (kg ha^{-1}), see § 3.4.1. Soil chemical analysis - soil mineral nitrogen.

Soil penetration resistance

Soil penetration resistance was measured at 5, 10, 15 and 30 cm soil depth using a hand cone penetrometer with base area of 3.33 cm^2 , 60° included angle and 80 cm driving shaft (Model 06.01.SA, Eijkelkamp Agrisearch Equipment, The Netherlands), following standard procedures (ASAE, 1994). Assessments were performed at 10 random positions from each main plot before and after cultivations, and harvest. The device was pushed perpendicular into the soil and the resistance (in N, Newton) and appropriate depth were recorded. The cone resistance was estimated by the ratio of the manometer reading (N) to the base area (cm^2) and then transformed to mega Pascal (MPa) units.

3.5. Statistical analysis

All the data collected were analysed using the split-split plot analysis of variance (ANOVA) model in Genstat (15th Edition VSN International Ltd, Hemel Hempstead, UK). Uniformity and residuals of all the data sets were verified before reporting results. The ANOVA results are reported quoting treatment means, residual degrees of freedom (df), standard error of difference (SED) or Fisher's Protected Least Significant Differences (LSD) and the *P*-value at significant level of $P < 0.05$. When necessary, correlation and regression analyses were also applied.

CHAPTER FOUR

Core experiment I – spring wheat 2013

4.1. Introduction

Cultivation techniques significantly influence the soil environment affecting crop germination, growth and development (Arvidsson *et al.*, 2013; Gajri *et al.*, 2002; Morris *et al.*, 2010; Silgram & Sheperd, 1999). In a clay soil, the use of plough and power harrows in conventional tillage (CT) can break the soil structure and clods, while mixing and incorporating plant residues creating a smooth and level soil surface (Bell, 1996; Soffe, 2003). Increasing soil movement can also intensify organic matter mineralisation, increasing N availability for the crop after cultivation operations (Silgram & Shepherd, 1999; Wild, 1988). However, continued use of CT operations reportedly leads to negative effects (see Table 2.1), as reported by others (e.g. Mitchell *et al.*, 2004; Six *et al.*, 2000)

In order to reduce potential negative effects by CT, non-inversion tillage has been widely used and increasingly adopted (Cannell & Hawes, 1994; Holland 2004; Jones *et al.*, 2006). These tillage systems create a seedbed using discs and tines without soil inversion, leaving a substantial part of the plant residues on the soil surface and/or mixed within topsoil layers (Soffe, 2003). Non-inversion tillage systems also reportedly increase organic matter in the very surface layer (Carter, 1991) and can reduce costs (Morris *et al.*, 2010; Soane *et al.*, 2012). However, detrimental effects of non-inversion tillage can sometimes result in lower crop yield compared with CT (see Table 2.4), as reported by several studies (López-Bellido *et al.*, 1998; Vijaya Bhaskar *et al.*, 2013b).

Mineral N application effects on crop production have been widely studied due to its influences on crop growth and development (Cossani *et al.*, 2009; López-Bellido, *et al.*, 1998; Otteson *et al.*, 2008) (see Table 2.5). For grain production, N fertilisation often increases yield when increasing N rates until this response is reduced with over-supply, while grain protein generally gives a linear response to N increases in various growing environments (López-Bellido *et al.*, 1998; López-Bellido *et al.*, 2001). However, several authors reported that weather conditions play an important role in highlighting

negative or positive effects of N fertilisation supply on grain protein and its negative correlation with grain yield (Campbell *et al.*, 1977; Smith & Gooding, 1999; Terman *et al.*, 1996). N fertilisation can also potentially benefit weeds, which then compete for resources with the crop (Di Tomaso, 1995; Moss *et al.*, 2004; Sheibani & Ghadiri, 2012). Although, it can also indirectly control weed pressure by encouraging crop growth biomass and increasing crop competitiveness (Grundy *et al.*, 1993; Jørnsgård *et al.*, 1996).

Legumes intercropped in UK are commonly utilised in the form of undersowing. Undersowing legume into the wheat crop stand in spring (Hartl, 1989) can potentially reduce competition between undersown and main crops (Charles, 1958). Undersowing covers the ground after the main crop harvest, allowing understorey species to possibly reduce weed presence by competing for resources (Liebman & Dyck, 1993; Thiessen-Martens *et al.*, 2001). Additionally, undersowing can suppress pests and diseases (Hiltbrunner *et al.*, 2002) but such effects depend on the undersown species, sowing time, harvest management and environmental conditions (Hartwig & Ammon, 2002). Using undersowing species with vigorous growth can potentially control weeds without negatively affecting the main crop and allowing sometimes greater final yields (Brennan & Smith, 2005; Haymes & Lee, 1994; Hauggaard-Nielsen *et al.*, 2008).

A field experiment was established in order to evaluate the effect of contrasting cultivation techniques, increasing mineral N fertilisation rates and undersowing legumes on growth, development and final yield of spring wheat.

4.2. Material and methods

4.2.1. Experimental site

The field experiment was performed at the Royal Agricultural University's Harnhill' Manor Farm, Cirencester, UK (NGR SP 075 006). The area was previously managed by either conventional tillage or by two inversion tillage systems distinguished by soil movement intensity and soil surface coverage (30% or >30%) (Vijaya Bhaskar, 2014).

Sinapsis alba cv Tilney was subsequently broadcasted in order to control take-all disease (*Gaeumannomyces graminis* var *tritici*) incidence in the field (HGCA, 2006).

4.2.2. Experimental design and treatment structure

The study was conducted from March 2013 to August 2013 on a field previously cropped with organic spring wheat cv Paragon. The experimental design and treatment structure were previously described in Chapter 3, Material and Methods, § 3.1. Before cultivation techniques operations in March 2013, 2 l ha⁻¹ of non-selective contact herbicide, a.i. glyphosate (Round-up) was applied across the entire experimental site. The 2013 dates for each field operation are reported in Table 4.1.


Table 4. 1. Diary of 2013 field operations

Field operation	Approximate date
Herbicide application	03 March 2013
Land preparation	20 March 2013
Spring wheat sowing	10 April 2013
Nitrogen applications	30 April / 23 May 2013
Undersowing	7 May 2013
Harvest	27 August 2013

4.2.3. Meteorological conditions

During the 2013 cropping season, maximum of 18.95°C was recorded in July while minimum temperature of 3.05 °C was recorded in March. Maximum and minimum rainfall documented were 76.8 mm in March and 31.3 mm in July. The 2013 growing season experienced lower air temperatures compared with the long-term average. The spring experienced higher precipitations, particularly in March and May, while rainfall in summer time was much lower, compared with the long-time average (Figure 4.1).

Figure 4. 1. Mean air temperatures and precipitation during the 2013 experimental period in comparison with the 10-year average. Royal Agricultural University meteorological station (NGR SP 42 004 011)



4.2.4. Assessments

4.2.4.1. Above ground assessments

Above ground assessment were previously described in Chapter 3, Material and Methods, § 3.3. Further details are presented in Table 4.2.

Table 4. 2. Above ground assessments dates for spring wheat 2013

Assessments	Approximate date
Wheat establishment	17 May 2013
Wheat tiller number	20 May 2013
Wheat shoot numbers	10 June 2013
Growth assessments (wheat, weeds and legumes – BM and WC – biomass)	24 May 2013 01 July 2013
Plant height	26 July 2013
Wheat ears number	27 July 2013
Diseases assessment	29 July 2013
Biological harvest	27 August 2013

4.2.4.2. Soil assessments

Soil assessments were previously described in Chapter 3, Material and Methods, § 3.4. Further details are given in Table 4.3.

Table 4. 3. Soil assessment dates for spring wheat 2013

Assessments	Approximate date
Soil mineral nitrogen, moisture (gravimetric) content	Monthly. March 2013 – August 2013
Soil pH	20 April, 10 June, 15 August 2013
Soil penetration resistance	29 February, 20 April, 10 June, 15 August 2013

4.2.5. Statistical analysis

Statistical analysis and reporting results were previously described in Chapter 3, Material and Methods, §3.5. The severity of diseases was transformed using log-normal transformation ($\log(x+1)$) ($x + 1$; x = percentage of leaf infected) in Genstat (15th Edition VSN International Ltd, Hemel Hempstead, UK), to reduce heterogeneity of variance.

4.3.Results

4.3.1. Establishment

There was a significant cultivation techniques effect on wheat establishment ($P<0.001$) with an overall mean establishment of 77%. CT resulted in greater plant establishment (96%) than HINiT (85%), followed by LINiT (53%) (Table 4.4). N fertilisation and undersowing treatments did not show significant effects on crop establishment.


Table 4. 4. Effect of cultivation techniques and nitrogen fertilisation on spring wheat establishment

	Establishment (number m⁻²)
CT	461.10c
HINiT	406.30b
LINiT	252.40a
<i>SED (4 df)</i>	10.86
<i>P</i>	***
N0	356.50a
N70	365.60a
N140	378.90a
N210	392.00a
<i>SED (18 df)</i>	18.44
<i>P</i>	<i>ns</i>

Values followed by same letter, do not differ significantly at $P<0.05$. ***= $P<0.001$, *ns*= no significant

Due to interactions, LINiT resulted in significant higher ($P<0.05$) crop establishment when N210 was applied (Figure 4.2). CT resulted in greater establishment than LINiT in spite of the N dose applied, while differences between CT and HINiT were only significant with N140 application.

Figure 4. 2. Effect of cultivation techniques and nitrogen fertilisation interaction on spring wheat establishment



Error bars representing average LSD ($P<0.05$)

4.3.2. Tillers and total shoot production

Tillers production and total shoot number were significantly affected ($P<0.01$) by cultivation techniques and N fertilisation treatments. A higher numbers of tillers were produced under CT and HINiT compared with LINiT, while the highest total shoots number was under CT (Table 4.5). Increases in N rate significantly increased ($P<0.01$) tillers and total shoots number, with the higher tillers and shoots number with N210, compared specifically with the unfertilised treatment. There was no significant effect of undersowing on tillers and total shoot production.


Additionally, the production of total shoots was significantly affected ($P<0.01$) by cultivation techniques and N fertilisation interaction, providing greater production under CT with 140 kg N ha⁻¹ was applied (Figure 4.3). On the other hand, HINiT resulted in greater shoot number when either N70 or N210 were applied, while LINiT only increased shoots number with N210 (Figure 4.3).

Table 4. 5. Effect of cultivation techniques and nitrogen fertilisation on spring wheat tillers and total shoots production

	Tillers (number m⁻²)	Total shoot (number m⁻²)
CT	1103.60b	895.60c
HINiT	961.70b	674.80b
LINiT	630.00a	612.50a
<i>SED (4 df)</i>	69.7	17.1
<i>P</i>	**	***
N0	721.60a	625.00a
N70	875.60ab	703.10ab
N140	932.70bc	753.00bc
N210	1063.90c	829.40c
<i>SED (18 df)</i>	79.7	44.8
<i>P</i>	**	**

Values followed by same letter, do not differ significantly at $P < 0.05$. ** = $P < 0.01$, and *** = $P < 0.001$

Figure 4. 3. Effect of cultivation techniques and nitrogen fertilisation interaction on spring wheat total shoots production



4.3.3. Mid-season wheat biomass and nitrogen uptake

May assessment

During May assessment, CT and HINiT produced a significantly greater wheat DM ($P<0.001$) and total wheat N uptake ($P<0.01$) than LINiT (Table 4.6). N fertilisation significantly increased wheat biomass production ($P<0.05$) and total wheat N uptake ($P<0.01$), compared with unfertilised treatment.


Table 4. 6. Effect of cultivation techniques and nitrogen fertilisation on spring wheat biomass and N uptake (May 2013)

	Wheat DM (t ha ⁻¹) May	Total wheat N uptake (kg N ha ⁻¹) May
CT	0.502b	20.53b
HINiT	0.456b	18.65b
LINiT	0.186a	8.83a
<i>SED (4 df)</i>	<i>0.034</i>	<i>1.51</i>
<i>P</i>	***	**
N0	0.341a	13.09a
N70	0.389b	15.87b
N140	0.389b	17.02b
N210	0.407b	18.03b
<i>SED (18 df)</i>	<i>0.023</i>	<i>1.08</i>
<i>P</i>	*	**

Values followed by same letter, do not differ significantly at $P<0.05$. *= $P<0.05$; ***= $P<0.01$; ***= $P<0.001$, and *ns*= no significant

Due to significant interactions ($P<0.05$), LINiT resulted in a greater wheat DM with 210 kg N ha⁻¹, compared specifically with N70 and N0 (Figure 4.4). CT and HINiT produced higher wheat biomass compared with LINiT, with up to 140 kg N ha⁻¹. However, under N210, CT resulted in higher wheat DM than HINiT, and then LINiT.


Figure 4. 4. Effect of cultivation techniques and nitrogen fertilisation interaction on spring wheat biomass (May 2013)



Error bars representing average LSD ($P < 0.05$)


A significant interaction between cultivation techniques and N fertilisation treatments was also observed on wheat N uptake (Figure 4.5). LINiT resulted in greater wheat N uptake when N210 was applied, compared specifically with N70 and N0. CT increased N uptake when either N140 or N210 were applied than specifically with N0. However, no significant effect was found between N rates under HINiT.

Figure 4. 5. Effect of cultivation techniques and nitrogen fertilisation interaction on spring wheat N uptake (May 2013)



Additionally, due to significant interaction, LINiT resulted in lower N uptake than CT and HINiT, regardless of the undersowing treatment (Figure 4.6).

Figure 4. 6. Effect of cultivation techniques and undersowing interaction on spring wheat N uptake (May 2013)



July assessment

At the July assessment, wheat DM and total wheat N uptake were not significantly affected by cultivation techniques (Table 7). On the contrary, N fertilisation treatments significantly increased ($P<0.001$) wheat DM, compared with unfertilised condition, while N uptake significantly increased ($P<0.001$) with increasing N rate application (Table 4.7). No significant effects of undersowing or treatment interactions were observed on wheat biomass and N uptake at the July assessment.

Table 4. 7. Effect of cultivation techniques and nitrogen fertilisation on spring wheat biomass and N uptake (July 2013)

	Wheat DM (t ha ⁻¹) July	Total wheat N uptake (kg N ha ⁻¹) July
CT	6.93a	125.40a
HINiT	7.56a	128.40a
LINiT	6.66a	143.20a
<i>SED (4 df)</i>	0.58	11.27
<i>P</i>	<i>ns</i>	<i>ns</i>
N0	5.25a	70.90a
N70	7.19b	125.90b
N140	7.82b	154.50c
N210	7.94b	178.10d
<i>SED (18 df)</i>	0.45	11.12
<i>P</i>	***	***

Values followed by same letter, do not differ significantly at $P<0.05$. ***= $P<0.001$; and *ns*= no significant

4.3.4. Mid-season legume biomass and nitrogen accumulation

Cultivation techniques treatments showed no significant effect on legume DM and N uptake during both growth assessments (Table 4.8). N fertilisation treatments only affected legume biomass in July, resulting in a significant reduction ($P<0.05$) of DM when N was applied, regardless of the rate. Moreover, legume N uptake was not significantly affected by N fertilisation at any assessment time.

No significant effects of undersowing or treatment interactions were observed on legume biomass and N uptake.

Table 4. 8. Effect of cultivation techniques and nitrogen fertilisation on legume biomass and N uptake (May and July 2013)

	Legume DM (t ha ⁻¹)	Legume N uptake (kg N ha ⁻¹)	Legume DM (t ha ⁻¹)	Legume N uptake (kg N ha ⁻¹)
	May	May	July	July
CT	0.0001a	0.001a	0.0580a	1.343a
HINiT	0.0010a	0.0362a	0.0196a	0.390a
LINiT	0.0004a	0.0153a	0.0144a	0.334a
<i>SED (4 df)</i>	<i>0.0005</i>	<i>0.0196</i>	<i>0.0245</i>	<i>0.6259</i>
<i>P</i>	<i>ns</i>	<i>ns</i>	<i>ns</i>	<i>ns</i>
N0	0.0002a	0.0087a	0.0809b	1.608a
N70	0.0004a	0.0130a	0.0216a	0.535a
N140	0.0002a	0.0076a	0.0103a	0.291a
N210	0.0011a	0.0394a	0.0099a	0.322a
<i>SED (18 df)</i>	<i>0.0005</i>	<i>0.0186</i>	<i>0.0242</i>	<i>0.5142</i>
<i>P</i>	<i>ns</i>	<i>ns</i>	*	<i>ns</i>

Values followed by same letter, do not differ significantly at $P < 0.05$. * = $P < 0.05$; and *ns* = no significant

4.3.5. Mid-season total weed biomass and nitrogen accumulation

May assessment

HINiT significantly resulted ($P < 0.05$) in a greater total weed biomass than CT and LINiT, at the May assessment (Table 4.9). No significant effect of cultivation techniques was observed on total weed N uptake. N fertilisation, undersowing or any treatment interaction did not significantly affect weed DM and total weed N uptake (Table 4.9).

July assessment

Among cultivation techniques, HINiT significantly increased ($P < 0.05$) total weed DM, particularly when compared with CT (Table 9). Weed N uptake was significantly lower ($P < 0.05$) under CT when compared with HINiT and LINiT (Table 4.9). N fertiliser application significantly increased ($P < 0.001$) weed DM and N uptake regardless of the

rate used compared with N0. Undersowing resulted in a non-significant effect on total weed DM or weed N yield at July assessment.


Table 4. 9. Effect of cultivation techniques and nitrogen fertilisation on weed biomass and N uptake (May and July 2013)

	Total weed DM (t ha ⁻¹)	Total weed N uptake (kg N ha ⁻¹)	Total weed DM (t ha ⁻¹)	Total weed N uptake (kg N ha ⁻¹)
	May	May	July	July
CT	0.0289a	1.11a	0.8600a	22.75a
HINiT	0.0540b	1.95a	1.9980b	44.07b
LINiT	0.0184a	0.83a	1.4560ab	38.40b
<i>SED (4 df)</i>	<i>0.0084</i>	<i>0.33</i>	<i>0.2583</i>	<i>5.09</i>
<i>P</i>	*	<i>ns</i>	*	*
N0	0.0364a	1.38a	0.9100a	13.19a
N70	0.0410a	1.55a	1.6320b	36.96b
N140	0.0303a	1.19a	1.7530b	44.98b
N210	0.0275a	1.08a	1.4570b	45.16b
<i>SED (18 df)</i>	<i>0.0063</i>	<i>0.24</i>	<i>0.1554</i>	<i>4.25</i>
<i>P</i>	<i>ns</i>	<i>ns</i>	***	***

Values followed by same letter, do not differ significantly at $P < 0.05$. * = $P < 0.05$; and *ns* = no significant


Due to significant interactions ($P < 0.001$), HINiT increased weed biomass when N70 or N140 were applied, specifically when compared with N0 (Figure 4.7). LINiT increased weed DM with N140 and N210 applications compared with N0. No significant differences were observed under CT between N rates.

Figure 4. 7. Effect of cultivation techniques and nitrogen fertilisation interaction on weed biomass (July 2013)




Due to significant interactions ($P<0.001$), BM and WC increased weed DM under non-inversion tillage treatments compared with CT. However, under Nus, HINiT resulted in higher weed DM than CT and LINiT (Figure 4.8).

Figure 4. 8. Effect of cultivation techniques and undersowing interaction on weed biomass (July 2013)



Additionally, WC increased weeds DM under fertilised conditions regardless of the rate (Figure 4.9). Under Nus, weed DM was higher at N140, compared with N0. BM significantly increased weed DM when N70 and N140 were applied, compared specifically with the unfertilised treatment.


Figure 4. 9. Effect of nitrogen fertilisation and undersowing interaction on weed biomass (July 2013)



Error bars representing average LSD ($P < 0.05$)


Cultivation techniques significantly interacted ($P < 0.05$) with N fertilisation, resulting in higher weed N uptake under HINiT and LINiT than CT when either N140 or N210 were applied (Figure 4.10). No significant differences were observed between cultivation techniques at N0.

Figure 4. 10. Effect of cultivation techniques and nitrogen fertilisation on weed N uptake (July 2013)



Cultivation techniques also significantly interacted ($P < 0.01$) with undersowing, with BM and WC resulting in a significant lower weed N uptake under CT than HINiT and LINiT, while Nus resulted in lower weed N uptake following either CT or LINiT than HINiT (Figure 4.11).

Figure 4. 11. Effect of cultivation techniques and undersowing on weed N uptake (July 2013)



4.3.6. Weed species composition

Irrespective of management treatments, a total of 39 weed species were recorded during the 2013 cropping season. At both growth assessment times, weeds were separated by species and grouped into broadleaf and grass weeds. In May, no grass weeds were observed, and total weed DM was mainly broadleaved weeds (Table 4.9).

At the July assessment, HINiT significantly increased ($P<0.05$) broadleaf weed DM compared with CT and LINiT (Table 4.10). Moreover, a significant increase ($P<0.001$) of broadleaf weed DM was observed with applications of 70 and 140 kg N ha⁻¹. Grass weed were not significantly affected by any treatment or interactions at this assessment time.

Table 4. 10. Effect of cultivation techniques and nitrogen fertilisation on weed species biomass (July 2013)


	Broadleaf weed	Grass weed
	DM (t ha ⁻¹)	DM (t ha ⁻¹)
	July	July
CT	0.772a	0.089a
HINiT	1.673b	0.325a
LINiT	0.949a	0.507a
<i>SED (4 df)</i>	<i>0.183</i>	<i>0.15</i>
<i>P</i>	*	<i>ns</i>
N0	0.778a	0.131a
N70	1.409b	0.223a
N140	1.318b	0.436a
N210	1.019a	0.438a
<i>SED (18 df)</i>	<i>0.124</i>	<i>0.133</i>
<i>P</i>	***	<i>ns</i>

Values followed by same letter, do not differ significantly at $P<0.05$. *= $P<0.05$; ***= $P<0.001$ and *ns*= no significant

No significant undersowing treatments effect was observed for either broadleaved weeds or grass weeds at any assessment time.

Furthermore, HINiT significantly increased ($P<0.05$) broadleaf weeds when N70 and N140 was applied, compared with CT and LINiT (Figure 4.12).

Figure 4. 12. Effect of cultivation techniques and nitrogen fertilisation on broadleaved weeds biomass (July 2013)



Among weed species, five grasses and 33 broadleaved weed species were recorded. Weed species recorded in May were also present in July. To prevent over-counting, weed biomass by species was analysed only from the July assessment. The list of all weed species recorded can be found in Appendix 2.

The dominant broadleaf weed species were *Stellaria media* L., *Fallopia convolvulus* L. and *Sinapsis arvensis* L., accounting for 29.8%, 12.1% and 9.7% of the total weed biomass recorded. Dominant grass weeds were *Avena fatua* L. and *Lolium perenne* L. constituting 11.9% and 8.1% of the total weed biomass.

Within all the species only *Stellaria media*, *Lolium perenne* and *Avena fatua* were significantly affected by management treatments. Other species were not significantly affected by any treatment, or occurred too infrequently to permit treatment effects to be appropriately tested.

Stellaria media DM significantly increased ($P<0.05$) under HINiT compared with CT and LINiT, while its biomass significantly increased ($P<0.05$) with N70 and N140 (Table 4.11).


Table 4. 11. Effect of cultivation techniques and nitrogen fertilisation on *Stellaria media* L. and *Lolium perenne* L. biomass

	<i>Stellaria media</i> L. DM (t ha ⁻¹)	<i>Lolium perenne</i> L. DM (t ha ⁻¹)
CT	0.2251a	0.0155a
HINiT	0.8544b	0.0318a
LINiT	0.1969a	0.3010b
<i>SED (4 df)</i>	0.1327	0.0436
<i>P</i>	*	**
N0	0.2422a	0.0717a
N70	0.5396b	0.0916a
N140	0.5254b	0.1865b
N210	0.3947ab	0.1146ab
<i>SED (18 df)</i>	0.0994	0.036
<i>P</i>	*	*

Values followed by same letter, do not differ significantly at $P < 0.05$; * = $P < 0.05$; ** = $P < 0.01$; and *ns* = no significant


Furthermore, due to interactions, HINiT significantly increased ($P < 0.05$) *Stellaria media* biomass with either N70 or N140 applications compared with N0 and N210 (Figure 4.13).

Figure 4. 13. Effect of cultivation techniques and nitrogen fertilisation interaction on *Stellaria media* L. biomass (July 2013)




Lolium perenne DM was greater under LINiT ($P<0.01$) compared to CT and HINiT, and N140 significantly increased ($P<0.05$) its DM compared specifically with N0 (Table 4.11). Under LINiT, *Lolium perenne* biomass was significantly higher ($P<0.01$) when N140 was applied compared with any other N rate (Figure 4.14). Additionally, under fertilised conditions LINiT resulted in higher *Lolium perenne* DM compared with CT and HINiT, while non-significant effects were observed between cultivation techniques at N0.

Figure 4. 14. Effect of cultivation techniques and nitrogen fertilisation interaction on *Lolium perenne* L. biomass (July 2013)




Furthermore, under LINiT, *Lolium perenne* DM increased when BM was undersown compared with WC and Nus (Figure 4.15).

Figure 4. 15. Effect of cultivation techniques and undersowing interaction on *Lolium perenne* L. biomass (July 2013)



Avena fatua DM was greater ($P < 0.01$) under HINiT compared with CT and LINiT at Nus (Figure 4.16). Higher *Avena fatua* DM was also found under HINiT when BM was undersown or Nus compared with WC.

Figure 4. 16. Effect of cultivation techniques and undersowing interaction on *Avena fatua* L. biomass (July 2013)



4.3.7. Plant height and ears number

The different cultivation techniques gave no significant effect on wheat height (Table 4.12). However, significantly ($P<0.001$) taller plants were recorded with 210 kg N ha⁻¹, when specifically compared with N0 and N70. There was no significant effect of undersowing and no treatment interactions on wheat height.

Wheat ear number per m² was significantly increased ($P<0.01$) under CT and HINiT compared with LINiT (Table 4.12). Nevertheless, there were no significant effects of N fertilisation rates, undersowing and any treatments interactions on total wheat ears number.

Table 4. 12. Cultivation techniques and nitrogen fertilisation effects on wheat height and wheat ear number

	Plant height (cm)	Ears (number m ⁻²)
CT	95.52a	596.10b
HINiT	96.72a	617.60b
LINiT	94.66a	484.00a
<i>SED (4 df)</i>	1.38	25.10
<i>P</i>	<i>ns</i>	**
N0	87.59a	597.00a
N70	96.31b	506.50a
N140	98.47bc	581.10a
N210	99.36c	579.10a
<i>SED (18 df)</i>	1.37	49.53
<i>P</i>	***	<i>ns</i>

Values followed by same letter, do not differ significantly at $P<0.05$; **= $P<0.01$; ***= $P<0.001$; and *ns*= no significant

4.3.8. Disease scoring

Assessments showed low incidence of leaf blotch (*Zymoseptoria tritici* Desm. Quaedvlieg & Crous) and take-all (*Gaeumannomyces graminis* var *tritici*). No significant effect of any treatments or treatments interaction was observed, and no results are therefore presented.

4.3.9. Final biological harvest

Ear, straw and total wheat biomass

Cultivation techniques did not significantly affect ear, straw or total wheat DM (Table 4.13). However, N fertilisation treatments only significantly affected ($P<0.01$) straw DM, resulting in an increase in biomass production when N210 was applied, particularly when compared with N0. There was no significant effect of undersowing or any treatment interaction on ear, straw biomass and total wheat DM.

Table 4. 13. Effect of cultivation techniques and nitrogen fertilisation on spring wheat ears, straw and total wheat biomass

	Ears DM (t ha⁻¹)	Straw DM (t ha⁻¹)	Total wheat DM (t ha⁻¹)
CT	7.13a	5.22a	13.33a
HINiT	7.17a	5.40a	13.73a
LINiT	7.59a	4.93a	13.38a
<i>SED (4 df)</i>	0.28	0.25	0.763
<i>P</i>	<i>ns</i>	<i>ns</i>	<i>ns</i>
N0	6.88a	4.50a	12.28a
N70	6.98a	4.98ab	13.26a
N140	7.89a	5.45bc	14.11a
N210	7.44a	5.81c	14.29a
<i>SED (18 df)</i>	0.63	0.34	0.877
<i>P</i>	<i>ns</i>	**	<i>ns</i>

Values followed by same letter, do not differ significantly at $P<0.05$; **= $P<0.01$, and *ns*= no significant

TGW, grains per ear and grain yield

Cultivation techniques did not significantly affect thousand grain weight (TGW) (Table 4.14). Increasing N fertilisation rates significantly reduced ($P<0.001$) TGW, with 29.31 g (LSD 2.07) observed when N210 was applied, compared with 35.52 g under N0. Additionally, CT significantly interacted ($P<0.01$) with Nus resulting in lower TGW compared with non-inversion tillage treatments (Figure 4.17).

The number of grains per ear was significantly higher under LINiT ($P<0.01$) than with CT and HINiT (Table 4.14). In addition, the N fertilisation, regardless of the N rate, also significantly increased ($P<0.05$) grain number per ear compared with N0 (Table 4.14). Yet undersowing, or any other treatment interactions, did not significantly affect number of grains per ears.

Overall, final spring wheat grain yield mean obtained in 2013 cropping season was 5.65 t ha⁻¹. Cultivation techniques significantly affected ($P<0.05$) grain yield with LINiT resulting in a higher yield than HINiT while showing statistically comparable yield with CT (Table 4.14). No significant effect was observed by N fertilisation, undersowing or any treatments interactions on final grain yield.


Table 4. 14. Effect of cultivation techniques and nitrogen fertilisation on TGW, grains per ear, final grain yield and harvest index

	TGW (g)	Grain per ears	Grain yield (t ha⁻¹)	HI (%)
CT	31.94a	30.63a	5.58ab	45.77ab
HINiT	32.40a	29.29a	5.30a	43.81a
LINiT	32.79a	37.96b	6.08b	48.61b
<i>SED (4 df)</i>	0.75	0.97	0.21	1.094
<i>P</i>	<i>ns</i>	**	*	*
N0	35.52c	28.59a	5.61a	50.33c
N70	33.03b	33.73b	5.50a	46.03b
N140	31.65b	33.70b	5.86a	45.21ab
N210	29.31a	34.48b	5.63a	42.69a
<i>SED (18 df)</i>	0.99	1.76	0.44	1.479
<i>P</i>	***	*	<i>ns</i>	***

Values followed by same letter, do not differ significantly at $P<0.05$; *= $P<0.05$; ** = $P<0.01$, and *ns*= no significant

Harvest index (HI) was significantly affected ($P<0.05$) by cultivation techniques treatments, resulting in greater HI under LINiT, compared specifically with HINiT (Table 4.14). Increasing N fertilisation rate significantly decreased ($P<0.001$) HI, with N0 resulting in the highest HI (Table 4.14).

Figure 4. 17. Effect of cultivation techniques and undersowing interaction on spring wheat



Wheat nitrogen yield

The average total wheat N uptake across treatments was 150.5 kg ha⁻¹. Cultivation techniques did not significantly affect wheat N uptake (Table 4.15). N fertilisation treatments significantly affected ($P < 0.001$) wheat N uptake resulting in lower uptake with N0 and N70, compared with N140 and N210 (Table 4.15).

Total grain N uptake was significantly higher ($P < 0.05$) with LINiT compared with CT and HINiT (Table 4.15). Furthermore, grain N uptake was significantly lower ($P < 0.001$) under N0 compared specifically with N210.

Regardless of management treatments, the average grain protein obtained was 12.28%. LINiT significantly increased ($P < 0.001$) grain protein content (13.02%) followed by CT (12.11%) and then by HINiT (11.71%) (LSD 0.33) (Table 4.15). Increases of N rates resulted in a highly significant effect ($P < 0.001$) on grain protein with N210 producing 14.46% of grain protein.

N harvest index (NHI) was not significantly affected by cultivation treatments. However, application of 210 kg N ha⁻¹ resulted in a significant lower ($P < 0.001$) NHI compared particularly with unfertilised conditions (Table 4.15).

Table 4. 15. Effect of cultivation techniques and nitrogen fertilisation on spring wheat N uptake, grain protein and N harvest index

	Total wheat N uptake (kg ha⁻¹)	Total grain N uptake (kg ha⁻¹)	Grain protein (%)	NHI (%)
CT	144.70a	118.30a	12.11b	82.07a
HINiT	140.0a	113.90a	11.71a	80.77a
LINiT	166.7a	137.90b	13.02c	82.48a
<i>SED (4 df)</i>	<i>8.16</i>	<i>6.61</i>	<i>0.12</i>	<i>0.64</i>
<i>P</i>	<i>ns</i>	<i>*</i>	<i>***</i>	<i>ns</i>
N0	120.60a	102.6a	10.12a	84.56c
N70	137.7a	114.0ab	11.87b	82.44bc
N140	163.6b	135.6bc	12.68b	82.13b
N210	180.0b	141.3c	14.46c	77.95a
<i>SED (18 df)</i>	<i>12.26</i>	<i>10.67</i>	<i>0.46</i>	<i>1.11</i>
<i>P</i>	<i>***</i>	<i>***</i>	<i>***</i>	<i>***</i>

Values followed by same letter, do not differ significantly at $P<0.05$; $*$ = $P<0.05$; $***$ = $P<0.001$; and ns = no significant

There were no significant effects of undersowing or any treatments interactions on total wheat and grain N uptake, grain protein and NHI.

Wheat N efficiency

Regardless of the treatments, N-use efficiency (NUE) was on average 30 kg kg⁻¹. Cultivation treatments did not exert a significant effect on NUE and N uptake efficiency (NUpE). However, N utilisation efficiency (NUtE) was significantly higher ($P<0.05$) under CT and HINiT than with LINiT (Table 4.16).

N fertilisation treatments significantly affected ($P<0.001$) NUE, NUpE and NUtE (Table 4.16). NUE decreased when increasing N rates and a similar trend was observed for NUpE although no significant differences were found between N140 and N210. NUtE decreased with increasing N rates but no significant differences were observed between N70 and N140 (Table 4.16).

Undersowing treatments significantly affected ($P<0.05$) NUE, resulting in greater NUE under Nus specifically when compared with BM, although, no significant effects were

observed on NUpE or NUtE (Table 4.16). No significant differences were found between any treatment interactions on N-use efficiency parameters.

Table 4. 16. Effect of cultivation techniques, nitrogen fertilisation and undersowing on spring wheat N-efficiency parameters

	NUE (kg kg⁻¹)	NUpE (kg kg⁻¹)	NUtE (kg kg⁻¹)
CT	31.67a	0.761a	40.30b
HINiT	30.43a	0.702a	40.84b
LINiT	27.82a	0.728a	37.02a
<i>SED (4 df)</i>	<i>1.545</i>	<i>0.044</i>	<i>0.846</i>
<i>P</i>	<i>ns</i>	<i>ns</i>	<i>*</i>
N0	49.08d	0.999c	48.94c
N70	30.07c	0.744b	40.29b
N140	23.48b	0.626a	37.30b
N210	17.26a	0.553a	31.02a
<i>SED (18 df)</i>	<i>2.289</i>	<i>0.042</i>	<i>1.995</i>
<i>P</i>	<i>***</i>	<i>***</i>	<i>***</i>
BM	28.41a	0.709a	38.76a
Nus	31.72b	0.759a	40.04a
WC	29.78ab	0.724a	39.37a
<i>SED (44 df)</i>	<i>1.202</i>	<i>0.029</i>	<i>0.61</i>
<i>P</i>	<i>*</i>	<i>ns</i>	<i>ns</i>

Values followed by same letter, do not differ significantly at $P \leq 0.05$; * = $P < 0.05$; *** = $P < 0.001$; and *ns* = no significant

Weed and legume biomass and nitrogen yield

Total weed biomass and N uptake were not significantly affected by cultivation techniques treatments (Table 4.17). Although, no significant effects of N fertilisation were observed on weed DM, weed N uptake was significantly lower ($P < 0.05$) with N0 than with N140 and N210. No significant effect of undersowing or any treatment interaction on weed DM or N uptake was observed.

CT resulted in significant higher ($P < 0.01$) legume DM and N uptake than HINiT and LINiT (Table 4.17). N0 resulted in significantly higher ($P < 0.001$) legume biomass and N uptake compared with N fertilised treatments.


Table 4. 17. Effect of cultivation techniques and nitrogen fertilisation on weed and legume biomass and N uptake (Harvest 2013)

	Total weed DM (t ha ⁻¹)	Total weed N uptake (kg ha ⁻¹)	Legume DM (t ha ⁻¹)	Total legume N uptake (kg ha ⁻¹)
CT	0.874a	15.76a	0.112b	3.300b
HINiT	1.217a	20.06a	0.042a	1.381a
LINiT	1.329a	25.05a	0.004a	0.117a
<i>SED (4 df)</i>	0.2656	5.82	0.0154	0.466
<i>P</i>	<i>ns</i>	<i>ns</i>	**	**
N0	0.934a	13.60a	0.177b	5.399b
N70	1.291a	20.23ab	0.015a	0.453a
N140	1.301a	24.46b	0.014a	0.417a
N210	1.033a	24.46b	0.004a	0.129a
<i>SED (18 df)</i>	0.2164	3.741	0.017	0.59
<i>P</i>	<i>ns</i>	*	***	***

Values followed by same letter, do not differ significantly at $P < 0.05$; * = $P < 0.05$; ** = $P < 0.01$, *** = $P < 0.001$; and *ns* = no significant


Due to significant interactions ($P < 0.001$), at N0, CT significantly increased the legume DM (Figure 4.18) and N uptake (Figure 4.19) compared with HINiT and then by LINiT. No significant effects were observed between any other N rate and cultivation techniques interaction.

Figure 4. 18. Effect of cultivation techniques and nitrogen fertilisation interaction on legumes biomass (Harvest 2013)




Error bars representing average LSD ($P < 0.05$)

Figure 4. 19. Effect of cultivation techniques and nitrogen fertilisation interaction on total legumes N uptake (Harvest 2013)



Significantly higher ($P < 0.05$) legume N uptake was observed with N0 under Nus, compared with BM and WC (Figure 4.20).


Figure 4. 20. Effect of undersowing and nitrogen fertilisation interaction on total legumes N uptake (Harvest 2013)



Total non-wheat (legumes plus weeds) DM and N uptake were not significantly affected by cultivation techniques, N fertilisation rates, undersowing or any treatment interaction, and the results are not, therefore, presented.


4.3.10. Soil mineral nitrogen

Throughout the assessments, cultivation technique effects on SMN were only evident in May, resulting in a significantly higher ($P<0.001$) SMN under CT (146.9 kg N ha⁻¹) and LINiT (141.5 kg N ha⁻¹) compared with HINiT (114.6 kg N ha⁻¹) (LSD 9.31) (Figure 4.21).




N fertilisation treatments resulted in a strong significant effect ($P<0.001$) on SMN content throughout all the assessments times (Figure 4.22). In May, higher SMN contents were observed under high N rates, where N210 resulted on 226.5 kg N ha⁻¹ while N0 showed 15.60 kg N ha⁻¹ (LSD 30.81). In June and July, lower SMN content was recorded when either N0 or N70 were applied, compared with N140 and N210. At the end of the season, August, N210 showed the highest SMN content (43.81 kg N ha⁻¹).

Figure 4. 22. Soil mineral nitrogen (kg N ha⁻¹) under four nitrogen fertilisation treatments




There was not a significant effect of undersowing treatments on SMN content throughout the assessments. However, at June assessment, due to interactions, WC and Nus treatments resulted in higher SMN content under LINiT than CT and HINiT (Figure 4.23). No significant differences were found when BM was undersown under any tillage treatment.

Figure 4. 23. Effect of cultivation techniques and undersowing interaction on soil mineral nitrogen (kg N ha⁻¹) (June 2013)




Under WC, the highest SMN was observed with N210 (Figure 4.24). Under Nus, the highest SMN was recorded with either N140 or N210. In the case of BM, SMN content increased when increasing N rate although no differences were observed between N70 and N0.

Figure 4. 24. Effect of nitrogen fertilisation and undersowing interaction on soil mineral nitrogen (kg N ha⁻¹) (June 2013)




At the June assessment, there was a significant cultivation \times N fertilisation \times undersowing interaction affecting SMN content (Figure 4.25). When N140 was applied under Nus, LINiT resulted in higher SMN content than CT and HINiT. Likewise, at N210 with WC, LINiT resulted in higher SMN content than CT and HINiT. Under N210 with undersown BM, CT resulted in higher SMN compared with HINiT and statistically similar to LINiT. In addition, LINiT resulted in higher SMN content than HINiT under Nus at N210. No significant differences were observed between treatments interactions under N0 or with N70 rate.

Figure 4. 25. Effect of cultivation techniques, nitrogen fertilisation and undersowing interaction on soil mineral nitrogen (kg N ha^{-1}) (June 2013)




At the July assessment, under WC, the highest SMN content was obtained with N210. Under BM and Nus, the highest SMN was with either when N210 or N140 were applied (Figure 4.26).

Figure 4. 26. Effect of nitrogen fertilisation and undersowing interaction on soil mineral nitrogen (kg N ha⁻¹) (July 2013)




The relationship between SMN content and wheat N uptake at mid-season growth assessments and harvest time among treatments are plotted in Figure 4.27 and 4.28.

Figure 4. 27. Soil mineral nitrogen and wheat N uptake in 2013 season (kg N ha⁻¹) under three cultivation techniques



LSD at $P < 0.05$ at each month of assessment; * = $P < 0.05$; ** = $P < 0.01$; *** = $P < 0.001$; and ns = no significant

Figure 4. 28. Soil mineral nitrogen and wheat N uptake in 2013 season (kg N ha⁻¹) under four nitrogen fertilisation rates



SMN (LSD)	30.81***	38.39***	29.02***	16.82***
Wheat N uptake (LSD)	2.27**	-	23.35***	28.41***

LSD at $P < 0.05$ at each month of assessment; ** = $P < 0.01$ and *** = $P < 0.001$


4.3.11. Soil moisture (gravimetric) content

Across assessments, the cultivation treatments influence on soil moisture content was variable with significant differences found only in March, July and August (Figure 4.29). In March, soil moisture content was significantly lower ($P < 0.01$) under CT compared with HINiT and LINiT (Figure 31). In July, CT resulted in a significant ($P < 0.05$) lower moisture content than HINiT, and LINiT. Furthermore, in August the moisture content was significantly lower ($P < 0.05$) under CT than LINiT.

Variable soil moisture contents were observed across the cropping season under different N fertilisation treatments, except in August when no significant effect was observed (Figure 4.29). In May, the soil moisture was higher when under N70, compared specifically with N140. In June, the N0 and N70 showed the highest soil moisture content. In July, the N210 presented significantly higher moisture content.

Throughout the assessments times, there were no significant effects of undersowing treatments, or any treatment interaction on soil moisture content.

Figure 4. 29. Effect of cultivation techniques and nitrogen fertiliser treatments on soil gravimetric moisture content (%) with mean precipitation during the 2013 experimental period in comparison to the 10-year average



Error bars representing LSD ($P < 0.05$) at each month of assessment

4.3.12. Soil pH

Irrespective of the management treatments and assessment times, soil pH was 7.02 (mean as a result of Repeated Measurements ANOVA). However, soil pH was only significantly affected ($P<0.01$) by N fertilisation at the June assessment with higher pH under unfertilised plots compared with fertilised ones, regardless of the N rate applied (Table 4.18). No significant effect of cultivation techniques, N fertilisation, undersowing or any treatment interaction was observed at any other assessment time.

Table 4. 18. Effect of nitrogen fertilisation on soil pH (June 2013)


	<u>pH</u>
N0	7.225b
N70	6.979a
N140	6.877a
N210	6.976a
<i>SED (18 df)</i>	<i>0.100</i>
<i>P</i>	<i>**</i>

Values followed by same letter, do not differ significantly at $P<0.05$; **= $P<0.01$

4.3.13. Soil penetration resistance


Before cultivation operations in February 2013, soil penetration resistance was evaluated in order to further test the previous year effect under contrasting tillage systems and to obtain initial values for the present study. Results showed a significant ($P<0.05$) cultivation techniques \times soil depth interaction (Figure 4.30). At this assessment time, at 15 cm depth, CT resulted in lower soil penetration resistance compared specifically with LINiT. At 30 cm depth, HINiT resulted in lower penetration resistance compare with CT. Penetration resistance significantly increased ($P<0.001$) with depth, with 5 cm and 10 cm soil layers showing lower resistance than 15 cm and 30 cm soil layers.

Figure 4. 30. Effect of cultivation techniques and soil depth interaction on soil penetration resistance (kPa) (February 2013) with cultivation techniques soil disturbance depth



After cultivation operations (April 2013), at any assessment time, soil penetration resistance was not significantly affected by cultivation techniques treatments. The mean values were therefore reported in Figure 4.31. However, at 30 cm depth significantly higher ($P < 0.001$) penetration resistance was found compared with any above soil layer.

Figure 4. 31. Effect of cultivation techniques and soil depth interaction on soil penetration resistance (kPa) (mean values 2013) with cultivation techniques soil disturbance depth



4.4. Discussion

4.4.1. Plant establishment, tillers and total shoots number

Cultivation techniques effect

Overall plant establishment was 77% of the seed sown (480 seeds m^{-2}), regardless of the management treatments. Plant establishment and early crop growth were affected by modifications of the seedbed conditions created by cultivation techniques operations, as reported by Strudley *et al.* (2008). The greater soil disturbance generated by CT in this clay soil created a level and even seedbed and incorporated plant residues into the soil. This allowed better seed-soil contact resulting in significantly greater plant establishment. More limited soil disturbance and greater plant residues cover on the soil surface under non-inversion tillage systems produced a coarser and highly variable seedbed conditions resulting in lower plant establishment, as also reported by other

studies (Känkänen *et al.*, 2011; Pietola & Tanni, 2003; Rieger *et al.*, 2008). Maximum reduction of soil disturbance and greater presence of residues under LINiT resulted in lower plant density compared with HINiT and CT. The presence of plant residues on the soil surface can also interfere with drill operations causing an uneven seedling depth and seed-soil contact under non-inversion tillage, as others report (Siemens *et al.*, 2004; Wilkins *et al.*, 1989). Additionally, high plant residues soil cover, particularly wheat straw, under non-inversion tillage treatments can release phytotoxic substances negatively affecting crop establishment and growth, as widely reported (Alam, 1990; Elliot *et al.*, 1976; Lovett & Jessop, 1982; Rice, 1984).

Variations of the seedbed conditions that affected plant establishment also appeared to affect wheat tillers production. Differences in tiller number per unit area between HINiT and CT were not as large as those observed for plant establishment, as also reported by Wade *et al.* (2006). Plants under HINiT compensated for lower plant density, at least partially, by tillering. Maximum reduction of tillage intensity and greater soil residues cover, under LINiT, can slow soil warming and drying resulting in a cold soil environment (Børrensen & Njos, 1990; Shinnars *et al.*, 1994), and consequently affected tillers production and shoot number. Better soil conditions under CT increased plant number and increased the early performance of the crop, represented by increasing tillers and total shoot number.

Nitrogen fertilisation effect

Tiller formation and total shoots were positively correlated with N supply as also reported by Power & Alessi (1978) and Rieger *et al.* (2008). An increasing number of tillers were produced by increasing N rate up to 210 kg N ha⁻¹ and resulting in larger number of shoots, like reported by Weisz *et al.* (2001). N application at the wheat tillering stage has provided additional N to the crop promoting additional tiller development, as reported by other authors (Otterson *et al.*, 2008; Power & Alessi, 1978; Sarandon & Gianibelli, 1990; Weisz *et al.*, 2001).

Interaction effect

Even though N fertilisation rates did not significantly affect crop establishment, CT resulted in greater number of plants per unit area when 140 kg N ha⁻¹ was applied. This

suggests N application could have encouraged a more rapid and stronger seedling growth promoting final establishment (Power & Alessi, 1978). Less soil movement intensity without inversion and the higher presence of plant residue cover under LINiT possibly resulted in initial N-immobilisation. Applying up to 210 kg N ha⁻¹ supported more plants establishment, therefore, and also increased shoot production under LINiT. HINiT appears to increase N availability reducing the need for extra N supply, as greater shoot number was observed with up to 70 kg N ha⁻¹. Nevertheless, maximum tillage intensity by soil inversion under CT resulted in a higher number of shoots when N was applied up to 140 kg N ha⁻¹ suggesting N fertilisation also encouraged higher tillers survival. This was also observed by Ottenson *et al.* (2008) reporting that shoots numbers can be increased with N fertiliser.

4.4.2. Mid-season plant biomass and nitrogen uptake

4.4.2.1. Wheat

Wheat biomass varied throughout the mid-season assessments among cultivation techniques and N fertilisation rates.

Cultivation techniques effect

Treatments that affected plant establishment and tiller production had a significant effect on wheat DM at the May assessment. Higher crop biomass production under CT seems to be related with greater plant population and tillers produced. Under HINiT, in spite of lower wheat establishment compared with CT, it seems that compensatory effect of tillering may result on similar biomass compared with ploughed soils, as also reported by Rieger *et al.* (2008). Variations in seedbed conditions and high rainfall conditions observed in May resulted in slow initial wheat growth under LINiT, as Mehdi *et al.* (1999) report. The greater presence of plant residues under LINiT, could have reduced soil temperature in the top layers, as the seedbed remains moist for longer slowing wheat growth as suggested by Cannell (1985). Acharya & Sharma (1994) also reported that a reduction in tillage intensity increases seedbed variability, resulting in less mid-season cereal biomass compared with CT.

Total wheat N uptake was, to a large extent, determined by cultivation techniques that significantly affected crop growth and biomass accumulation, as also reported by Gastal & Lemaire (2002). Initial growth assessment showed that CT and HINiT had significantly greater wheat N uptake and, therefore, greater wheat biomass than LINiT.

In spite of the increasing number of shoots by increasing tillage intensity, wheat biomass at the July assessment seemed to be more affected by environmental conditions. The ability of the soil under contrasting tillage intensity to maintain moisture at times of low rainfall (in and before July) was more likely to create variations in wheat growth across assessments, as also reported by others (Mehdi *et al.*, 1999; Lafond *et al.*, 2006). Non-inversion tillage, and specifically LINiT improved soil moisture content during dry weather conditions, reducing wheat stress and supporting crop growth. This possibly compensated for poor early growth under non-inversion tillage, at least partially, as differences between cultivation techniques in wheat DM at the July assessment were not significant. It seems that early crop growth was most likely to be affected by an uneven seedbed condition affecting plant establishment, rather than moisture availability as observed later in the season.

Nitrogen fertilisation effect

N supply is widely reported to substantially increase crop biomass (Campbell *et al.*, 1977; López-Bellido *et al.*, 2005; Pearman *et al.*, 1977). This was also observed in the present study throughout all growth assessments. Increases in wheat biomass before anthesis was mainly related by increases in tiller and shoot production, and by increasing leaf expansion, like also reported by Pearman *et al.* (1977) and Gastal & Lemaire (2002). In May, crop growth was slower and the canopy smaller, and its N demand was also expected to be low. Increasing N rates up to 70 kg N ha⁻¹ seems to be enough to have increased wheat biomass at this stage. However, in July, crop canopy was bigger possibly leading to a higher N demand. This suggests that N uptake was based on the soil N availability to meet such high crop N demand, as also reported by Justes *et al.* (1994) and Power & Alessi (1978).

Interaction effect

During the first growth assessment, it seems that N availability was a limiting factor for the crop for early growth under LINiT, as higher wheat DM and N uptake was obtained with up to 210 kg N ha⁻¹. Reducing soil disturbance under LINiT possibly reduced plant-available N by immobilization, increasing the need for additional N supply in order to meet crop requirements, like also reported by Radford *et al.* (1992). Additionally in May decreases in N uptake under LINiT compared to HINiT and CT in all undersowing treatments were observed. This suggests a more relevant effect of the cultivation techniques on the crop N uptake than the undersowing.

4.4.2.2. Legumes

Establishment and biomass of undersown legume species were quite low throughout growth assessments. Before and until a week after legume broadcast, higher temperatures and absence of rainfall resulted in low soil moisture content which could possibly have affected germination, establishment and biomass production. It could be assumed that the lack of undersowing effects in a large range of treatment assessments was an indirect effect of low legume establishment rather than a direct effect of the undersowing *per se*.

Nitrogen effect

Legume persistence has often been reported incompatible with N fertilisation (Soussana & Arregui, 1995). Undersown species, *Trifolium repens* L. (WC) and *Medicago lupulina* L. (BM) exhibited smaller biomass, slow growth rate and increased leaf and root longevity resulting in poor competitive ability under N-rich conditions (Döring *et al.*, 2013). These patterns resulted in lower legume biomass at the July assessment under fertilised conditions, as also reported by Moss *et al.* (2004) and Döring *et al.* (2013). Additionally, N fertilisation can increase wheat competitiveness resulting in an indirect adverse effect on legume undersown (Gooding & Davies, 1997).

4.4.2.3. Weeds

Most of the weeds species identified are commonly report in spring wheat production (HGCA, 2010) and their presence was influenced by the agricultural managements adopted and time of assessment as reported by Menalled *et al.* (2001). Total weed biomass constituted between 8.1% and 16.9% of the total plant aboveground biomass at the May and July assessment, respectively.

Cultivation techniques effect

Grass weeds appear to be a major challenge for cereal production, specifically when non-inversion tillage systems are adopted (Vijaya Bhaskar *et al.*, 2014). To overcome this challenge, a broad spectrum pre-cultivation herbicide glyphosate was applied, irrespective of treatments. Although, the approach employed in this study for the herbicide application did not allow us to test its specific impact on weed dynamics, it is possible to speculate on its relative effect on weed occurrence. Some studies (e.g. Mavunganidze *et al.*, 2014) report that a broad-spectrum herbicide as glyphosate controls both grass and broadleaf weed species. However, in the present study it appears that by applying herbicide, grass weeds seemed to be restricted and were less relevant compared with broadleaf weed species, throughout all of the mid-season assessments, like also reported by Ewald & Aebischer (2000) and Marshall & Nowakowski (1996).

As grass weed species prevalence was low, broadleaf weed species mainly accounted for the differences between tillage treatments on the total weed DM. This was observed as HINiT resulted in higher total weed and broadleaf weeds DM than LINiT and CT. Others authors (e.g. Clements *et al.*, 1996a; Swanton *et al.*, 2000) reported lower broadleaf weed species under CT compared with non-inversion tillage practices, like observed in this study particularly with HINiT. Delayed sowing due to increased rainfall (March 2013) could have allowed the emergence, after herbicide application, of weeds retained in the soil under non-inversion tillage. This condition combined with increases in soil disturbance intensity is the possible reason for high biomass of short-lived annual broadleaf weeds under HINiT. Higher presence of soil residues cover under LINiT created shadowing, reducing germination of some broadleaf species after herbicide application, as reported by Teasdale *et al.* (1991). For CT, weeds that escaped foliar

contact herbicide are likely to grow but the subsequent soil inversion is thought to have reduced broadleaf weed presence. This situation also provides a head-start for the primary crop, such that it can effectively compete with later emerging weeds, as other studies (Mahn, 1984; Menalled *et al.*, 2001; Wicks *et al.*, 1988) reporting lower weed biomass under CT also in spite of herbicide-use.

Nitrogen fertilisation effect

Weed DM at the May assessment was not significantly affected by N fertilisation. This was probably a result of the N fertilisation promoting a good start to the crop increasing its competitiveness against weeds, as the crop biomass and N uptake were higher, like also reported by Moss *et al.* (2004) and Davis & Liebman (2001). However, later in the season (in July), total weed DM increase their response to N fertilisation and broadleaf species DM increased especially with 70 and 140 kg N ha⁻¹ applications. This was also reported by several authors (Blackshaw *et al.*, 2003; Jørnsgård *et al.*, 1996; Małecka & Blecharczyk, 2008; Moss *et al.*, 2004) suggesting the weed N response depends on their differential competitiveness to uptake N.

Interaction effect

At the July assessment, HINiT resulted in a greater total weed and broadleaved weeds biomass with either N70 or N140 rate. Such effect is perhaps related to the increase of *Stellaria media* L. biomass, as the major weed species recorded. The increase of *Stellaria media* under HINiT is probably the result of seed retention in the soil and to the subsequent soil disturbance. Additionally, *Stellaria media* seems to be able to grow and reproduce under N-rich conditions and severe competition (Moss *et al.*, 2004). These traits make *Stellaria media* the major weed species in this study high N conditions and one of the commonest species of intensively grown cereals in the UK (Moss *et al.*, 2004).

Non-inversion tillage effect on total grass weed DM was not evident. However, seedbed conditions under LINiT were more advantageous for *Lolium perenne* L. to grow. This grass specie is susceptible to soil disturbance and specifically mechanical soil inversion, as also reported by others (e.g. Froud-Williams *et al.*, 1983b; Hakansson, 2003; Tuesca & Puricelli, 2007). Additionally, the fast growth behaviour of *Lolium perenne* requires

high N supply (Daepf *et al.*, 2001), resulting in greater biomass under fertilised conditions, which could explain its greater biomass under LINiT combined with high N applications.

BM interacted with HINiT resulting in greater *Avena fatua* biomass and BM \times LINiT interaction increased *Lolium perenne* biomass. These two weed species contributed to 20% of the total weed DM. Suggesting their role, to a large extent, for the total weed DM increases under BM and WC interaction with non-inversion tillage treatments. Slow growth of the undersown legume species allowed weed species with faster growth to establish, such as *Lolium perenne*, *Avena fatua*, and *Stellaria media* under fertilised N conditions, as also reported by Moss *et al.* (2004). Whereas under Nus, N70 and N140 application there was increased weed biomass, confirming positive weeds growth response to N supply.

4.4.3. Plant height and ears number

Cultivation techniques effect

Plant height reflects prevailing growing conditions and is affected by several factors, such as crop variety, soil conditions, weather patterns, and also by agricultural management performed (Malhi *et al.*, 2007). Overall, the 2013 cropping season underwent uncommon and contrasting weather conditions compared to the long-term seasonal average. It appears that cultivation techniques effects on the early crop performance were overshadowed, or compensated for, by dry weather later in the season. Greater crop residue cover, particularly under LINiT, appears to limit soil water loss (Martinez *et al.*, 2008) possibly promoting later plant growth under lower rainfall conditions, as also reported by Radford *et al.* (1992) and Guy & Lauver (2007). This perhaps compensated for slow early crop growth resulting in statistically similar plant height between tillage systems, as also observed on wheat biomass later in the season.

Agricultural management effects on plant establishment, and tiller production and survival affected ear production per unit area in the present study, like also reported by Mc-Master *et al.* (1994). The reduction of tillage intensity and greater soil coverage

under LINiT created higher seedbed variability affecting shoot survival and resulting in fewer ears compared with CT and HINiT, as also reported by Boomsma *et al.* (2010).

Nitrogen fertilisation effect

N application promoted vigorous plant growth, as previously discussed, which seems to also result in taller plants compared with unfertilised conditions, agreeing with Liu *et al.* (2013) findings. No significant effect of N fertilisation on ear number has been related, in some cases elsewhere, to the incidence of diseases, particularly *Fusarium* spp. (Pearman *et al.*, 1977; Rieger *et al.*, 2008). However, in this study, *Fusarium* spp. was not observed. Consequently, it seems ear number may merely be an expression of greater competition for available resources between an increasing shoot number, under fertilised conditions, as also reported by Power & Alessi (1978) and Pearman *et al.*, (1977).

4.4.4. Ears, straw and total wheat biomass

Cultivation techniques effect

Weather conditions greatly affect plant growth, specifically those developments occurring during the dry period observed at grain-filling stage. Initial crop growth and development was highly encouraged under CT with more favourable seedbed conditions and lower weeds occurrence compared to non-inversion tillage systems, as mentioned by Mehdi *et al.* (1999). However, the ability to save soil moisture under non-inversion tillage systems perhaps reduced the initial differences observed between tillage systems. This may have resulted in similar ear, straw and total wheat DM, agreeing with Sainju *et al.* (2012). Saving, or perhaps increasing, soil moisture may be vital for crop production specifically under contrasting weather patterns or in months with scarcity of precipitations (Hansen *et al.*, 2011).

Nitrogen fertilisation effect

One of the main effects of mineral N fertilisation is to increase the size and duration of cereal canopy growth (Gooding & Davies, 1997). This was evident as N applications

increased wheat DM since the early growth assessments, and continued to increase straw DM at harvest especially with 210 kg N ha⁻¹ rate, agreeing with Pearman *et al.* (1977). It seems that the lack of N significant effect on ear production was also translated to the total ear DM. As ear biomass is mainly defined by grains number and their weight, it is also possible that the N fertilisation influence on TGW and grains per ear counteracted resulting in a similar ear DM. This agrees with López-Bellido *et al.* (2000) and Rasmussen *et al.* (1997) reporting that raising N rate increased grains per ear while decreasing seed weight.

4.4.5. TGW, grains per ear, final grain yield and harvest index

Cultivation techniques effect

Greater ear production under CT and HINiT possibly created more competition for resources, resulting in lower number of grains produced compared with LINiT. This agrees with Pollard *et al.*, (1981) reporting that less grains at harvest under ploughed plots were a result of more ears competing for resources. Fewer ear numbers under LINiT resulted in more grains per ear.

Thousand grain weight (TGW) is reportedly to be genetically determined (Mogensen *et al.*, 1985). However, several authors agree that the expression of TGW seems to be influenced by the cropping environment (so called GxE interaction). Therefore under stress situations, grains often appeared smaller and with less weight (De Vita *et al.*, 2007; López-Bellido *et al.*, 1998). Treatments that resulted in higher grains number per ear could possibly have caused a shrinking effect of the grains, while lower grains number increased their weigh (López-Bellido *et al.*, 1998). This may have resulted in compensation between cultivation treatments on TGW.

Final grain yield is mainly determined by number of ears per unit area, grains per ear and TGW (HGCA, 2008b). Each of these yield components are related to growing conditions at different growth stages, although, each of them can in some part compensate for developments at earlier phases (HGCA, 2008b), as previously

discussed. This also emphasises the difficulties of relating final grain yield to the individual yield components, as previously highlighted by Gooding & Davies (1997).

Final grain yield under LINiT was higher than HINiT, and statistically similar to that of CT. Higher number of grains under LINiT could possibly have made the final grain yield under this treatment comparable with that of CT. Moreover, grain yield was also related to the crop N uptake, with LINiT resulting in high total wheat N uptake and yield. This may perhaps have been due to the soil being wet and cool at the beginning of the cropping season, leading to N losses and possibly high immobilisation under LINiT. Later in the season, dry and warm conditions combined with high soil moisture availability under LINiT, leading to increased N mineralisation. This allowed more N available for the plant, supporting the final yield under LINiT. Such results were also observed by Fox & Bandel (1986) reporting that comparable yield under reduced tillage with CT is the result of differences in N availability. In contrast, the lower grains number per ear and wheat N uptake under HINiT resulted in a lower final grain yield. These observations clearly suggest that for the 2013 cropping season, N and moisture availability, and number of grains per ear, have largely determined differences between cultivation techniques treatments on the final yield, as suggested by other studies (Brennan *et al.*, 2014; De Vita *et al.*, 2007; Fox & Bandel, 1986; Hansen *et al.*, 2011).

As reported by Zhang *et al.* (2012), the present study showed that wheat yield can be increased without increases in DM production if the number of grains increases. This was observed in the harvest index (HI), which is the ratio of aboveground DM partitioning to grain yield, which was higher under LINiT than HINiT, and statistically similar to that of CT.

Nitrogen fertilisation effect

Increasing N supply increased grains number mainly by promoting nutrient availability, as also reported by Alijani *et al.* (2012) and Ferrise *et al.* (2010). Additionally, the higher availability of N might perhaps have reduced floret mortality resulting in higher number of grain with increasing N fertilisation rates, as suggested by Ferrante *et al.*, (2010). However, increases in grains per ear by N application appear to promote shortening of the grains reducing their weight, presented as lower TGW, agreeing with

several authors (Campbell *et al.*, 1977; López-Bellido *et al.*, 2000; Pearman *et al.*, 1977; Rasmussen *et al.*, 1997). Ferguson (1967) also suggested that grains per ear and TGW in practice are often negative, being influenced by intra and inter-plant competition for resources available.

Grain yield depends on ears number, grains per ear and TGW (Campbell *et al.*, 1977; HGCA, 2008b), and it seems these yield components compensated resulting in similar grain yield between N treatments. Additionally, wheat yield response to N fertilisation is also influenced by factors such as growing environment conditions, soil type and cultivar (López-Bellido *et al.*, 2012). In this study dry weather conditions and the high clay content of the soil could have generated a significant accumulation of N in the soil profile. This caused additional soil available N for the crop, perhaps, contributing to high yield under unfertilised conditions. A lack of wheat yield response to N fertiliser application occurring overall. Similarly, several authors also reported no response of crops to N fertiliser attributable to high reserve of SMN (Abad *et al.*, 2005; Corbeels *et al.*, 1998; Johnson & Mattern, 1987; Miao *et al.*, 2015; López-Bellido *et al.*, 1996). Increases in straw DM by N applications seem to have not supported final grain yield, resulting in a lower HI with increasing N rates, as also reported by Borghi (2000) and Pearman *et al.* (1977).

4.4.6. Wheat N yield, grain protein and N efficiencies

Cultivation techniques effect

Higher soil moisture and N availability under LINiT appears to encourage later crop growth increasing grain N uptake compared with CT and HINiT, although this was not observed in total wheat N uptake, and as reported by Fox & Bandel (1986). N availability does not always increase total plant N uptake as the wheat ability to capture N depends on various factors, especially synchronisation of soil NO₃⁻ with crop demand (Halvorson *et al.*, 2001; Liao *et al.*, 2004).

Wheat grain protein is greatly dependent on genotype (Johnson *et al.*, 1985; Stobard & Marshall, 1990) but it is also influenced by the predominant growing environment (Rao

et al., 1993). Generally, there is a negative relationship between grain protein and final grain yield, reportedly due to energy constraints and N dilution effects (Terman *et al.*, 1996; McNeal *et al.*, 1982; Löffler *et al.*, 1985; Pearman *et al.*, 1978). However, this study like Kramer (1979) and Johnson & Mattern (1987), showed that for the same phenotype, grain yield and protein correlation is not always negative. There was no evidence of dilution effect of N assimilated resulting from high grain yield under LINiT, as grain protein content was also higher under LINiT than CT, and by HINiT. This effect perhaps is related to the increased N availability under LINiT, as also reported by López-Bellido *et al.* (2001). Gao *et al.* (2012) also suggested that increases in grain protein content can be ascribed to an increase in soil N supply, due to improvements in soil moisture content.

Although tillage systems significantly affected SMN, it had no significant effect on the crop N-use efficiency (NUE), agreeing with Brennan *et al.* (2014) and Giacomini *et al.* (2010). NUE can be partitioned into N capture by roots (uptake efficiency, NUpE) and its conversion to grain by shoots (utilisation efficiency, NUtE) (Moll *et al.*, 1982). Novoa & Loomis (1981) defines NUtE as the physiological efficiency of the N-use as it is the grain yield divided by the total crop aboveground N at maturity. Among cultivation techniques, LINiT resulted in lower NUtE compared with CT and HINiT. However, there were no observed relation between the crop total N uptake and the NUtE as also reported by Barraclough *et al.* (2010).

Nitrogen fertilisation effect

An increase in total wheat N uptake was evident with the application of 140 and 210 kg N ha⁻¹. However, grain N uptake was highly affected by increases in N rate resulting, particularly in higher grain N under N210 when compared with unfertilised conditions. These increases in total wheat and grain N uptake were also reported by Campbell *et al.* (1977) and Pearman *et al.* (1977). High crop N uptake under N-rich conditions can indicate the amount of unused N fertiliser present in a rooting zone at the time of crop requirement (Campbell *et al.*, 1977). In the present study, high-level of SMN content and high N uptake under high N rates treatments was not translated to higher yields while it was in grain quality, as also reported by Barraclough *et al.* (2014).

Reportedly, grain protein content is modified by the growing conditions, including interactions between management, genotype and environment (Gooding & Davies, 1997; Graybosch *et al.*, 1996; Zhu & Khan, 2001). In this study, grain protein content varied with N supplied with values from 10% in unfertilised treatment to 14% with 210 kg N ha⁻¹ applied. Similar results were also reported by other authors (Garrido-Lestache *et al.*, 2004; Godfrey *et al.*, 2010; López-Bellido *et al.*, 2001; Wieser & Seilmeier, 1998). Although, the highest grain protein was obtained with up to 210 kg N ha⁻¹, grain yield stayed unchanged by increasing N suggesting that higher N supply is needed in order to optimise grain protein rather than to maximise grain yield. This was also observed by Barraclough *et al.* (2014) evaluating several wheat varieties, including spring wheat cv Paragon in the UK, in reporting no correlation between grain yield and protein content at a given-N rate.

Crop N-efficiency parameters, NUE, NUpE, NUtE and NHI decreased under N fertiliser treatments. These results are mainly due to increases in aboveground N relative to grain yield reducing efficiency in the use of N, as also reported by Huggins & Pan (1993). Unfertilised treatment resulted in 52% higher NUE than the mean value for the N application treatments. This confirms that soil nitrate-N levels, under N0, were perhaps enough contribution to high grain yield as previously discussed and also reported by López-Bellido & López-Bellido (2001). Here, the low NUpE with increasing N rates can indicate higher available N than the crop demanded (Huggins & Pan, 2003). Furthermore, under high N rates, low NUtE and high protein suggest that once N is taken by the crop the physiological efficiencies decrease (Huggins *et al.*, 2010). Decrease in NHI under high N rates shows that high N uptake was not proportional to the final grain when increasing N supply. This agrees with López-Bellido & López-Bellido (2001), although it is contrary to Rozas *et al.* (1999) reporting greater N uptake resulted in greater yield thus increased NHI.

Undersowing effect

Low establishment and slow growth of the legume undersowed species resulted in a lack of significant differences between undersowing treatments, regardless of interactions, in almost every wheat development and production assessment, and in SMN and moisture content across the cropping season. However, NUE under Nus was

greater than with BM and similar to WC. However, cultivation techniques and N fertilisation exerted greater effects on N-efficiency parameters, making it more difficult to correlate such small undersowing effects on NUE.

4.4.7. Non-wheat biomass and N yield at harvest

4.4.7.1. Weeds

Application of broad-spectrum non-residual herbicide prior to cultivation operations appears to have reduced weed pressure regardless of the management treatments, as also reported by Derksen *et al.* (1995). However, weeds still occurred although their biomass seems to vary across the growing season. The dry weather conditions observed at the end of the cropping season and the enhancing wheat competition ability by increases in biomass, could have negatively affected weed development causing a decline in weed biomass at harvest (Jørnsgård *et al.*, 1996; Mas & Verdú *et al.*, 2003). This was more marked on the major weed species, *Stellaria media* whose seeds germinate at or close to the soil surface making this species more susceptible to drought (Bond *et al.*, 2007). Visual assessments corroborated weeds prevalence reduction during dry weather, affecting their biomass at harvest. Weed N uptake increased under fertilised treatments suggesting that weed N uptake was driven by differences in N supply, as also reported by Kamiji *et al.* (2014).

4.4.7.2. Legumes

Legume development patterns varied throughout the growing season. During mid-season growth assessments legume growth and development were poor, and do not appear to be related to any management treatment. However, at harvest time legume DM production increased, although still small reinforces that WC and BM have slower growth, as reported by other studies (Döring *et al.*, 2013; Wallace, 2001).

Cultivation techniques effect

Higher variability and weed pressure under non-inversion tillage slowed legume growth, resulting in lower legume DM and N uptake than under CT at harvest time.

Nitrogen fertilisation effect

Legume DM and N uptake were reduced when N was applied compared to unfertilised conditions. This reinforces the negative relation between legume DM production under N-rich conditions as reported by several authors (Moss *et al.*, 2004; Soussana & Arregui, 1995). N fertilisation effect on N uptake was related to legume DM production.

Interaction effect

Increasing tillage intensity favoured legume growth and N uptake under unfertilised conditions, possibly due to less variability of the seedbed and weed pressure under CT. Under unfertilised conditions, Nus resulted in higher N uptake compared with BM and WC species. Nus treatment was not completely free of legume probably due to some natural regeneration, by stolons (WC) and adventitious bud in roots (BM), in spite of pre-cultivation herbicide applications. Legumes were, therefore, separated from weeds even though they were legume weeds in these plots rather than deliberately undersown legumes. This could possibly have triggered differences on legume N uptake.

4.4.8. Soil moisture content

Soil moisture content followed a temporal pattern following rainfall events across the cropping season. This was particularly evident in March and May when rainfall and soil moisture content were higher followed by low moisture during summer months.

Cultivation techniques effect

In the present study, right after cultivation operations (in March 2013), increasing tillage intensity and soil inversion under CT reduced soil moisture. This was likely a consequence of breaking the soil water-related pores and increasing evaporation intensifying water loss (Reicosky *et al.*, 1999). In contrast, less tillage intensity and

without soil inversion together with greater plant residues cover under LINiT and HINiT resulted in higher soil moisture content. These results agree with several studies (Fabrizzi *et al.*, 2005; Fuentes *et al.*, 2003; Hatfield *et al.*, 2001; Lampurlanés *et al.*, 2001) reporting greater soil moisture under non-inversion tillage. High moisture content under non-inversion tillage treatments is attributable to several factors such as more water-related pores (Bescansa *et al.*, 2006). Additionally, plant residues protected the soil reducing evaporation and run-off, and in part decreasing soil temperature slowing soil drying, particularly at the surface (Baumhardt & Jones, 2002; Beyaert *et al.*, 2002; Shinnars *et al.*, 1994). These conditions were particularly more evident in July and August when crop development had possibly dried the soil as increased transpiration, and limited rainfall occurred highlighting the soil's ability under LINiT to potentially save soil moisture content. This has also been reported by several authors (Baumhardt & Jones, 2002; Bescansa *et al.*, 2006; Malhi *et al.*, 2007; Singh *et al.*, 1998).

The potential accumulation of soil moisture by non-inversion tillage systems during dry periods possibly encouraged wheat growth, and is most likely to be one of the possible explanations for the equivalent final grain yield between LINiT and CT. Since summers are expected to be drier in the UK as a result of climate change (Christensen & Christensen, 2007; Jenkins *et al.*, 2008, 2009) greater water availability observed under non-inversion tillage would seem to favour its use.

Nitrogen fertilisation effect

N fertilisation effect on plant growth towards maturity seems to indirectly affect soil moisture content from May to July. At this time, N supply promoted crop biomass production possibly increasing crop water uptake and reducing soil moisture levels. This agrees with Campbell *et al.* (1977) who reported a rise in plant water uptake under heavily N fertilised growing conditions. López-Bellido *et al.* (2007a) also related lower crop growth under unfertilised conditions resulting in higher soil moisture content levels. During crop production phase, specifically in July, higher soil moisture content was observed in the plots fertilised with high N rates. This was probably due to rapid biomass stimulation by N supply, covering the soil and reducing the surface susceptible to evaporation, and enhancing moisture storage, as also reported by several authors (Corbeels *et al.*, 1998; Hatfield *et al.*, 2001; López-Bellido *et al.*, 2007b). This occurred

despite the dry weather conditions and that during crop production phase ample N and water uptake, and transpiration were expected (HGCA, 2008b). Despite the variable soil moisture response under different N treatments at each month of the growing season, it seems that there is a general pattern suggesting that increasing N fertilisation will promote higher crop biomass potentially reducing soil moisture content.

4.4.9. Soil mineral nitrogen

Cultivation techniques effect

Interactions between biological processes and solute transfer in the soil result in inorganic N dynamic, which is influenced by weather conditions, soil type and cropping system (Oorts *et al.*, 2007). In the present study, contrasting soil disturbance and plant residues soil cover left by the cultivation techniques affected soil N mineralisation, as reported by other studies (Myrbeck *et al.*, 2012; López-Bellido & López-Bellido, 2001; Silgram & Shepherd, 1999). However, cultivation effects are often reported to be temporal and frequently to increase SMN content when increasing tillage intensity mainly by exposing organic matter to decomposition, as reported by Myrbeck *et al.* (2012) and Silgram & Shepherd (1999). In the present study no significant differences were observed after cultivation operations, however, as also reported by Fuentes *et al.* (2003) and Oorts *et al.* (2007). While no leaching measurements were conducted, it is suspected that lack of significant differences between tillage treatments after cultivations operations (March 2013) was related with high rainfall recorded at the end of March. This possibly increased the risk of leaching the N mineralised after cultivations operations, as reported by Oorts *et al.* (2007) comparing tillage systems in a clay textured soil.

Differences between cultivation techniques on SMN were then noticeable in May when higher levels of N was recorded compared with previous months following the N fertiliser applications. Plant N uptake is highly related to SMN levels (Brennan *et al.*, 2014). Differences in plant establishment under tillage treatments led to variations in plant N uptake, possibly resulting in variation of the residual SMN left in the soil. Lower plant populations under LINiT resulted in less N uptake probably leaving higher

residual N in the soil, as reported by several authors (Brennan *et al.*, 2014; Riley, 1998; Thomsen & Sørensen, 2006). On the contrary, despite increasing soil disturbance intensity under HINiT, higher plant number resulted in greater N uptake reducing SMN levels when compared specifically with LINiT. However, under CT, SMN level seems more related to a larger mineralisation rate at this time of the cropping season rather than being related with crop N uptake.

From June to August, towards wheat maturity, SMN level decreased when crop N uptake increased. These temporal variations of SMN were similar to those found by Fuentes *et al.* (2003) and Oorts *et al.* (2007) who reported that small seasonal differences in SMN between cultivation techniques were attributed to delayed N mineralisation. Conditions of low water availability limit soil microbiota slowing N mineralisation (Jenkinson *et al.*, 1987; Rasmussen *et al.*, 1998). Crop N uptake potentially depletes soil available N, therefore.

Nitrogen fertilisation effect

Application of mineral N fertiliser increased SMN levels, as also reported by other studies (Angás *et al.*, 2006; Giacomini *et al.*, 2010; Liebig *et al.*, 2002; Lu *et al.*, 2010; Zhao *et al.*, 2014). However, SMN decreased rapidly as the cropping season progressed, mainly due to increases in crop N uptake as the crop approaches to maturity (Fuentes *et al.*, 2003). The remaining soil N also depended on initial N levels (initial soil N plus N mineral fertilisation applied) and possible mineralisation. At initial crop growth stages, applying more N than the crop needed could lead to N accumulation in the soil (Angás *et al.*, 2006). This was observed at the initial assessments months, when large SMN content was observed with 210 kg N ha⁻¹ applied. While, under N₀ and N₇₀, a rapid depletion of SMN was observed across the experimental time due to higher plant uptake than the N supplied. This agrees with Zhao *et al.* (2014) who reported a low SMN content under low mineral N addition as the crop uptake was higher than the supply, whereas the opposite happened under N-rich conditions which exceeded crop uptake levels. Moreover, differences found between N rates treatments decreased with time as a result of a proportional balance between N supply and crop N consumption.

Interaction effect

Undersowing legume species, such as BM and WC, have been recommended for reducing N leaching and for increasing soil available N, even before their biomass incorporation into the soil (Döring *et al.*, 2013). These characteristics could possibly suggest a greater effect in SMN content, although this was only observed when undersowing interacted with other management treatments. At the June assessment, BM and WC increased SMN levels with N210 under LINiT compared with Nus, while under CT with N210, BM increased SMN compared to WC and Nus. SMN increases by legume undersown suggests, perhaps, a release of N by the legume species as reported by Bergkvist (2003). Känkänen *et al.* (2001) also reported increases in SMN when legumes were undersown under high levels of N fertilisation, compared with monocrops. However, this was not translated in greater crop N uptake contrary to results reported by Thorsted *et al.* (2006). Interactions between cultivation techniques and undersowing has also been reported to affect crop growth, biomass production, grain yield and weed population (Teasdale *et al.*, 1991) affecting soil residual N, therefore, SMN levels. However, in the present study, cultivation techniques and N fertilisation showed a more marked effect making it more difficult to determine undersowing effect irrespectively of treatments interactions

4.4.10. Soil pH

Soil pH was found to be significantly affected by the N applications at the June assessment. This is possibly due to N fertiliser increased the anion H^+ from the ammonium nitrification reducing soil pH (Magdoff *et al.*, 1997), as reported by Schroder *et al.* (2011) and Zhao *et al.* (2014). However, this effect of N fertilisation was only observed in June, suggesting that changes in pH are more highly influenced by environmental conditions and temporal variations as also reported by Spiegel *et al.* (2007). This perhaps also explain the lack of cultivation techniques effect on pH, contrary to Vijaya Bhaskar *et al.* (2013a) who reported higher soil pH under CT than reduced tillage systems.

4.4.11. Soil penetration resistance

Soil penetration resistance indicates how easily roots can penetrate the soil. High penetration resistance, therefore, can restrict root growth affecting crop production (Gregory, 1994). In the present study, penetration resistance at all soil depth layers was lower than 2MPa, which is the upper limit for uninterrupted root growth (Atwell, 1993). Before cultivation techniques operations, soil penetration resistance showed a depth gradient with increasing resistance from top soil layers (0 to 10 cm) to 15cm and then 30cm depth (despite soil moisture content not being directly assessed at different soil depths), as reported by several studies (Bradford, 1986; Campbell & O'Sullivan, 1991; Grant & Lafond, 1993; Martinez *et al.*, 2008). At this assessment time, the cultivation effects on penetration resistance indicate the legacy of the tillage operation previously used. At 5cm and 10cm soil layer, no significant differences were observed between tillage treatments suggesting a diminished effect of contrasting tillage treatment effect after a year from the initial operations, as also reported by Martinez *et al.* (2008). However, at 15cm soil layer, LINiT resulted in higher penetration resistance than CT. This is possibly the result of more intense soil movement created by CT equipment reducing soil compaction, compared with reduced tillage intensity under LINiT. Similar observations were also reported by Martinez *et al.* (2008) and Ozpinar & Çay (2005). Furthermore, soil inversion by ploughs increased soil compaction below plough-working depth (approx. 20 cm) increasing penetration resistance at 30 cm, compared specifically with HINiT, as also reported by Ardivissov *et al.* (2013). Nevertheless, significant differences between contrasting cultivation operations were reduced after their performance in March 2013. Across the 2013 growing season, penetration resistance was only related with depth resulting in lower resistance at top soil layers depth (0 -15cm) compared to 30cm, as observed by Grant & Lafond, (1993).

4.5. Conclusions

Key findings for the initial investigation on cultivation techniques are listed in Table 4.19. Initial investigation with spring wheat revealed the importance of seedbed conditions created by contrasting cultivation techniques in determining plant

establishment, growth and development. The success among cultivation techniques was initially demonstrated on plant establishment, and finally by reductions of negative effects under dry summer conditions.

Table 4. 19. Key outcomes for the cultivation techniques effects during 2013 cropping season

	CT	HINiT	LINiT
Tillage intensity	High	Intermediate	Low
Seedbed	Fine	Coarser	Much coarser
Seedbed evenness	Level / uniform	Variable / Not uniform	Highly variable / Not uniform
Plant establishment	High	Intermediate	Low
Tiller production	High	High	Low
Plant height	Statistically not significant		
Ears number	High	High	Low
TGW	Statistically not significant		
Number of grains per ear	Low	Low	High
Grain yield	¹ Comparable	Low	High
SMN	Low	Low	High
Moisture content	Low	Intermediate	High

¹Statistically comparable with LINiT and HINiT

The potential initial plant establishment and higher number of tillers under CT were overshadowed by competitive and compensatory effects occurring on ear number and grains per ear, but this did not reduce final yield. However, less soil tillage intensity and greater plant residues cover under LINiT resulted in greater seedbed variability causing a slow early crop growth and less plant establishment than CT and HINiT. Nevertheless, greater soil moisture content and soil N availability under LINiT encouraged later crop growth, inducing greater grain number per ear and resulting on comparable yield with CT. In spite of initial cultivation influences on weed DM at mid-season assessments, no-significant differences were observed by harvest time. This was the result of

vigorous crop growth possibly competing against weeds and also the prevalent dry weather conditions inducing natural decay of the weeds.

Consequently, under this study soil and weather conditions, CT performance was constant from high crop establishment until final yield. On the basis of yield, LINiT seems, however, to be an acceptable alternative to CT. This agrees with other authors reporting equivalent performance of non-inversion tillage compared with CT, under deficient rainfall conditions during the cropping season (Brennan *et al.*, 2014; De Vita *et al.*, 2007; López-Bellido *et al.*, 1996). The present study was encouraging, therefore, for the adoption of non-inversion tillage system as LINiT, for spring wheat production. However, with weather uncertainty further experimental study was necessary.

Regarding N fertilisation, key findings from the initial investigation are listed in Table 4.20. N fertilisation encouraged crop growth compared with unfertilised conditions. However, those differences between N rates were diminished, with no effects of N fertilisation treatment on final grain yield, mainly as a result of prevailing dry conditions and crop compensatory effects. Higher grain numbers by increasing N fertilisation supply was not supported at harvest, resulting in lower TGW creating a compensation effect and eclipsing N fertilisation influence on yield. SMN level in the soil under unfertilised condition appeared to be enough to encourage high yields as yields were similar than with N fertilised plots. In addition, N fertilisation also encouraged weed growth, increasing its biomass production although this seems to have not affected final wheat yield. In this study soil and prevalent weather conditions N fertilisation did not encourage yield gains, although further study is needed.

The scarcity of rainfall right after undersowing legume species were broadcasted resulted in poor establishment and slow growth, causing overall failure of the undersowing treatments. Even though some effects were observed, cultivation techniques and N fertilisation treatments effects were more evident making it more difficult to observe undersowing effects alone. Therefore, undersowing legume effects need to be investigated further to confirm that environment conditions in this season played a more relevant role than the undersown treatments *per se*.

Table 4. 20. Key outcomes for the nitrogen fertilisation effects during 2013 cropping season

	N0	N70	N140	N210
Tiller production	Low	¹ Comparable	² Comparable	High
Plant height	Low	Medium	³ Comparable	High
Ears number	Statistically not significant			
TGW	High	Medium	Medium	Low
Number of grains per ear	Low	High	High	High
Grain yield	Statistically not significant			
SMN	Highly low	Low	Medium	High

¹Statistically comparable with N0 and N140; ²Statistically comparable with N70 and N210;

³Statistically comparable with N70 and N210.

CHAPTER FIVE

Core experiment II - spring wheat 2014

5.1.Introduction

After spring wheat harvest on 28 August 2013, the soil was left with legume cover during the winter months and before drilling 2014 spring wheat. Soil cover was a combination of wheat straw, black medic, white clover and weeds (Plate 5.1). Due to the legume species ability to continue to grow after 2013 harvest, it was thought they could perhaps function as a catch crop accumulating N and limiting losses by denitrification (Döring *et al.*, 2013; Jones, 1992; Jensen, 1991). In addition, the legume species might develop a dense canopy which could potentially assist with controlling weed establishment and growth (Breland, 1996; Clements & Donaldson, 1997). The experiment was, therefore, repeated with spring wheat cv Paragon to reinforce the 2013 crop performance findings.

Plate 5. 1. Overwinter soil cover



5.2. Materials and methods

5.2.1. Experimental site

The field experiment was conducted at the Royal Agricultural University's Harnhill' Manor Farm, Cirencester, UK, (NGR SP 075 006). Before cultivation techniques operations in 2014, 2 l ha⁻¹ of a non-selective contact herbicide, a.i. glyphosate (Round-up), was applied throughout the experimental field.

5.2.2. Experimental design and treatment structure

The study was conducted from March 2014 to August 2014 on a field previously cropped with spring wheat cv Paragon. The experimental design and treatment structure were as previously described in Chapter 3, Material and Methods, § 3.1. The 2014 dates of each field operation are described in Table 5.1.

Table 5. 1. Diary of 2014 field operations



Field operation	Approximate date
Herbicide application	11 March 2014
Land preparation	24 March 2014
Spring wheat sowing	18 April 2014
Nitrogen applications	10 May / 30 May 2014
Undersowing	14 May 2014
Harvest	31 August 2014

5.2.3. Meteorological conditions

During the 2014 cropping season, maximum and minimum air temperatures were recorded in July (18.0 °C) and March (7.18 °C). Maximum and minimum rainfalls were recorded in May (97.3 mm) and March (39.5 mm). The 2014 growing season experienced lower air temperatures and the spring period experienced higher rainfall

conditions while rainfall in summer was lower compared to the long-term seasonal average (Figure 5.1).

Figure 5. 1. Mean air temperatures and precipitation during 2014 experimental period in comparison with the 10-year average. Royal Agricultural University meteorological station (NGR SP 42 004 011)



5.2.4. Assessments

5.2.4.1. Above ground assessments

Above ground assessments were previously described in Chapter 3, Material and Methods, § 3.3. Further details are presented in Table 5.2.

Table 5. 2. Above ground assessments for spring wheat 2014

Assessments	Approximate date
Overwinter legumes and weeds biomass	28 February 2014
Wheat establishment	07 May 2014
Wheat tillers number	28 May 2014
Wheat shoot numbers	16 June 2014
Growth assessments (wheat, weeds and legumes – BM and WC – biomass)	28 May 2014 / 01 July 2014 / 01 August 2014
Plant height	28 July 2014
Wheat ears number	28 July 2014
Diseases assessment	29 July 2014
Biological harvest	31 August 2014

5.2.4.2. Soil assessments

Soil assessments were previously described in Chapter 3, Material and Methods, § 3.4. Further details of the assessment dates are given in Table 5.3.

Table 5. 3. Soil assessments for spring wheat 2014

Assessments	Approximate date
Soil mineral nitrogen and moisture (gravimetric) content	Monthly. March 2014 – August 2014
Soil pH	March 2014
Penetration resistance	March, June and August 2014

5.2.5. Statistical analysis

Statistical analysis and reporting results are as previously described in Chapter 3, Material and Methods, §3.5. The severity of diseases was transformed using log-normal transformation ($\log(x+1)$) ($x + 1$; x = percentage of leave infected) in Genstat (15th Edition VSN International Ltd, Hemel Hempstead, UK), to reduce heterogeneity of variance.

5.3. Results

5.3.1. Overwinter growth assessment

Overwinter assessments showed significantly higher ($P<0.01$) weed biomass and N uptake under LINiT than CT and HINiT (Table 5.4). N fertilisation significantly affected ($P<0.001$) weed DM, resulting in higher weed DM with N140 and N210 than with N0 and N70, while increasing N rate significantly increased ($P<0.001$) weed N uptake.

Legume overwinter biomass was significantly higher ($P<0.001$) under CT than HINiT, followed by LINiT, while legume N uptake was significantly higher ($P<0.01$) under CT compared with HINiT and LINiT (Table 5.4). N fertilisation treatments significantly decreased ($P<0.001$) legume DM and N uptake when compared with N0.

There was no significant effects of undersowing on overwinter weed and legume DM and N uptake.


Table 5. 4. Weed and legume biomass and N uptake (Overwinter 2013)

	Total weed DM (t ha ⁻¹)	Total weed N uptake (kg N ha ⁻¹)	Legume DM (t ha ⁻¹)	Legume N uptake (kg N ha ⁻¹)
CT	0.242a	5.66a	0.4568c	13.948b
HINiT	0.403a	10.48a	0.1485b	4.925a
LINiT	0.905b	23.11b	0.0614a	1.999a
<i>SED (4 df)</i>	<i>0.070</i>	<i>1.853</i>	<i>0.030</i>	<i>1.554</i>
<i>P</i>	**	**	***	**
N0	0.377a	7.43a	0.6546b	20.647b
N70	0.463a	10.76b	0.1232a	3.714a
N140	0.575b	14.41c	0.0556a	1.878a
N210	0.650b	19.73d	0.0556a	1.626a
<i>SED (18 df)</i>	<i>0.046</i>	<i>1.203</i>	<i>0.059</i>	<i>1.22</i>
<i>P</i>	***	***	***	***

Values followed by same letter, do not differ significantly at $P<0.05$; **= $P<0.01$;
***= $P<0.001$

Due to significant interactions, CT and HINiT resulted in significantly higher weed N uptake with N210, compared specifically with N0, while LINiT increased weed N uptake with increasing N rate (Figure 5.2).


Figure 5. 2. Cultivation techniques and nitrogen fertilisation interaction effect on overwintering weed N uptake



Error bars representing average LSD ($P < 0.05$)

In addition, due to significant treatments interaction, under the Nus and WC regimes, with N70 and N140, LINiT resulted in higher weed N uptake compared with HINiT and CT (Figure 5.3). LINiT and HINiT resulted in higher weed N uptake than CT with N0 and BM, while with N140 and BM, the LINiT increased weed N uptake compared with HINiT and CT.

Figure 5. 3. Cultivation techniques, nitrogen fertilisation and undersowing interaction effect on overwintering weed N uptake



CT resulted in higher legume DM and N uptake compared with HINiT and LINiT under N0. Higher legume DM and N uptake were found with N70 under CT than HINiT and LINiT (Figure 5.4 & 5.5).

Figure 5. 4. Cultivation techniques and nitrogen fertilisation effect on overwinter legume biomass





Figure 5. 5. Cultivation techniques and nitrogen fertilisation effect on overwinter legume N uptake



5.3.2. Plant establishment, tiller numbers and total shoots

In the present study, overall plant establishment was 52% of the seed sown, which was lower compared with spring wheat in 2013 (77%). Between the cultivation techniques, there was significantly higher plant establishment ($P<0.05$) under CT (62.5%) and HINiT (56%) compared with LINiT (37.3%) (Table 5.5).

Tiller production per unit area was significantly higher ($P<0.01$) under CT and HINiT than LINiT (Table 5.5). However, HINiT and LINiT resulted in significant lower ($P<0.01$) total shoot number compared with CT.

There were no significant effects of N fertilisation, undersowing or any treatments interactions on plant establishment, tillers number and total shoots.

Table 5. 5. Effect of cultivation techniques on establishment, tiller number and total shoot of spring wheat

	Establishment (number m⁻²)	Tillers (number m⁻²)	Shoots (number m⁻²)
CT	300.00b	508.80b	469.20b
HINiT	269.00b	471.90b	406.80a
LINiT	179.00a	400.60a	382.60a
<i>SED (4 df)</i>	<i>30.06</i>	<i>15.66</i>	<i>9.47</i>
<i>P</i>	*	**	**

Values followed by same letter, do not differ significantly at $P<0.05$; *= $P<0.05$, and **= $P<0.01$

5.3.3. Mid-season wheat biomass and nitrogen uptake

At the May assessment, CT resulted in a significant higher ($P<0.001$) wheat DM and total wheat N uptake compared with HINiT, followed by LINiT (Table 5.6). N fertilisation did not have significant effects on wheat DM, although N210 resulted in significantly higher ($P<0.05$) total wheat N uptake compared with any other N rate (Table 5.6).

Furthermore, WC undersown resulted in a significantly higher ($P<0.05$) total wheat N uptake compared with Nus and BM undersown, while no significant differences were observed between undersowing treatments on wheat DM production.


Table 5. 6. Effect of cultivation techniques and nitrogen fertilisation on spring wheat biomass and N uptake (May 2014)

	Wheat DM (t ha ⁻¹) May	Total Wheat N uptake (kg N ha ⁻¹) May
CT	0.504c	21.39c
HINiT	0.429b	15.59b
LINiT	0.242a	9.68a
<i>SED (4 df)</i>	<i>0.015</i>	<i>0.53</i>
<i>P</i>	***	***
N0	0.382a	13.78a
N70	0.372a	14.69a
N140	0.358a	14.76a
N210	0.455a	18.97b
<i>SED (18 df)</i>	<i>0.038</i>	<i>1.66</i>
<i>P</i>	<i>ns</i>	*
BM	0.3817a	14.78a
Nus	0.3711a	14.80a
WC	0.4230a	17.08b
<i>SED (47 df)</i>	<i>0.02255</i>	<i>0.959</i>
<i>P</i>	<i>ns</i>	*

Values followed by same letter, do not differ significantly at $P<0.05$; *= $P<0.05$; ***= $P<0.001$; and ns= no significant

Due to significant interactions under WC, N210 resulted in a significantly higher total wheat N uptake compared with any other N rate treatment, at the May assessment (Figure 5.6). Under Nus, N application resulted in a significantly higher total wheat N uptake compared with unfertilised treatment.

Figure 5. 6. Cultivation techniques and undersowing interaction effect on wheat N uptake (May 2014)



At the July and August assessments, CT resulted in significantly higher wheat DM and total wheat N uptake compared with HINiT and LINiT (Table 5.7).

At the July assessment N140 and N210 significantly increased ($P < 0.01$) wheat DM compared with N0 and N70, while at the August assessment N210 resulted in a significantly higher ($P < 0.001$) wheat DM, compared specifically with N0. In both assessments times, total wheat N uptake significantly increased ($P < 0.001$) with increasing N fertilisation rate.

There were no significant effects of undersowing on wheat biomass and N uptake at the July and August assessments.


Table 5. 7. Effect of cultivation techniques and nitrogen fertilisation on spring wheat biomass and N uptake (July and August 2014)

	Wheat DM (t ha ⁻¹) July	Wheat N uptake (kg N ha ⁻¹) July	Wheat DM (t ha ⁻¹) August	Wheat N uptake (kg N ha ⁻¹) August
CT	6.724b	157.0b	11.412b	169.10b
HINiT	4.107a	76.3a	5.809a	80.90a
LINiT	4.287a	81.0a	6.724a	93.50a
<i>SED (4 df)</i>	0.384	7.89	0.417	8.86
<i>P</i>	**	***	***	***
N0	4.118a	65.20a	6.786a	86.80a
N70	4.854ab	92.20b	7.480ab	102.7b
N140	5.541b	120.60c	8.434bc	120.3c
N210	5.643b	141.00d	9.227c	148.2d
<i>SED (18 df)</i>	0.40	8.62	0.489	7.28
<i>P</i>	**	***	***	***

Values followed by same letter, do not differ significantly at $P < 0.05$; * = $P < 0.05$, ** = $P < 0.01$; *** = $P < 0.001$; and ns = no significant

Due to significant interactions, at the July assessment, CT resulted in higher wheat N content under Nus and WC than with BM undersown (Figure 5.7).

Figure 5. 7. Cultivation techniques and undersowing interaction effect on wheat N uptake (July 2014)



5.3.4. Mid-season weed and legume biomass and nitrogen uptake

May assessment

At this assessment time, weeds were separated by species and grouped into broadleaf and grass weeds. However, total weed DM, broadleaf and grass weeds, were not significantly affected by any treatment structure or interactions (Table 5.8). Nevertheless, HINiT resulted in a significantly higher ($P<0.05$) total weed N uptake compared with CT and LINiT. No significant effect of N fertilisation was observed on total weed N uptake.

Legume DM and N uptake were not significantly affected by cultivation techniques (Table 5.8). N applications, regardless of the N rate, significantly reduced ($P<0.05$) legume DM and N uptake compared with unfertilised treatment (Table 5.8).

Table 5. 8. Effect of cultivation techniques and nitrogen fertilisation on weed and legume biomass and N uptake (May 2014)

	Total weed DM (t ha⁻¹) May	Broadleaf weed DM (t ha⁻¹) May	Grass weed DM (t ha⁻¹) May	Total weed N uptake (kg N ha⁻¹) May	Legume DM (t ha⁻¹) May	Legume N uptake (kg N ha⁻¹) May
CT	0.0107a	0.0107a	0.00001a	0.350a	0.0000001a	0.00001a
HINiT	0.1832a	0.1832a	0.00001a	6.256b	0.00218a	0.0623a
LINiT	0.0573a	0.0530a	0.0043a	1.979a	0.00003a	0.0010a
<i>SED (4 df)</i>	<i>0.05231</i>	<i>0.0522</i>	<i>0.00347</i>	<i>1.489</i>	<i>0.00089</i>	<i>0.02534</i>
<i>P</i>	<i>ns</i>	<i>ns</i>	<i>ns</i>	*	<i>ns</i>	<i>ns</i>
N0	0.0754a	0.0752a	0.0002a	2.08a	0.002797b	0.07854b
N70	0.0978a	0.0922a	0.0055a	3.43a	0.000111a	0.00437a
N140	0.0804a	0.0804a	0.00001a	2.95a	0.000037a	0.00141a
N210	0.0814a	0.0813a	0.0001a	2.99a	0.0000001a	0.00001a
<i>SED (18 df)</i>	<i>0.03307</i>	<i>0.033</i>	<i>0.00387</i>	<i>1.006</i>	<i>0.00098</i>	<i>0.02754</i>
<i>P</i>	<i>ns</i>	<i>ns</i>	<i>ns</i>	<i>ns</i>	*	*

Values followed by same letter, do not differ significantly at $P<0.05$; *= $P<0.05$, and ns= no significant

Due to significant interactions ($P<0.01$), under N0, HINiT resulted in higher legume DM and N uptake compared with CT and LINiT (Figure 5.8 & 5.9). There were no significant effects of undersowing or any other treatments interactions on legume biomass in May.

Figure 5. 8. Cultivation techniques and nitrogen fertilisation interaction effect on legume biomass (May 2014)





Figure 5. 9. Cultivation techniques and nitrogen fertilisation interaction effect on legume N uptake (May 2014)



July assessment

CT resulted in a significantly lower ($P<0.01$) total weed DM compared with HINiT and LINiT, while total weed N uptake was significantly higher ($P<0.01$) under HINiT than LINiT, followed by CT (Table 5.9).

At this assessment time, weeds were also separated by species and grouped into broadleaf and grass weeds. Broadleaf weeds DM were significantly higher ($P<0.01$) under HINiT compared with LINiT, followed by CT, while no significant effects of cultivation techniques were observed on grass weeds DM (Table 5.9).

N fertilisation significantly increased ($P<0.001$) total weeds DM and broadleaf weeds DM compared with unfertilised treatment, while no significant effects were observed on grass weeds DM (Table 5.9). Total weed N uptake significantly increased ($P<0.001$) with increasing N rate application, although no significant differences were observed between N70 and N140.

There were no significant effects of cultivation techniques and N fertilisation on legume DM and N uptake.

Table 5. 9. Effect of cultivation techniques and nitrogen fertilisation on weed and legume biomass and N uptake (July 2014)


	Total weed DM (t ha ⁻¹) July	Broadleaf weed DM (t ha ⁻¹) July	Grass weed DM (t ha ⁻¹) July	Total weed N uptake (kg N ha ⁻¹) July	Legume DM (t ha ⁻¹) July	Legume N uptake (kg N ha ⁻¹) July
CT	0.196a	0.122a	0.073a	5.20a	0.0010a	0.005a
HINiT	1.909b	1.512c	0.397a	42.41c	0.0202a	0.277a
LINiT	1.309b	0.815b	0.494a	28.08b	0.0050a	0.102a
<i>SED (4 df)</i>	<i>0.21740</i>	<i>0.2458</i>	<i>0.1683</i>	<i>4.93</i>	<i>0.00705</i>	<i>0.1803</i>
<i>P</i>	**	**	<i>ns</i>	**	<i>ns</i>	<i>ns</i>
N0	0.412a	0.3379a	0.073a	6.25a	0.0134a	0.261a
N70	1.222b	0.9836b	0.239a	22.47b	0.0056a	0.110a
N140	1.332b	0.8981b	0.434a	30.08b	0.0152a	0.119a
N210	1.585b	1.0453b	0.540a	42.10c	0.0008a	0.022a
<i>SED (18 df)</i>	<i>0.2173</i>	<i>0.1565</i>	<i>0.2016</i>	<i>4.95</i>	<i>0.0063</i>	<i>0.0996</i>
<i>P</i>	***	***	<i>ns</i>	***	<i>ns</i>	<i>ns</i>

Values followed by same letter, do not differ significantly at $P<0.05$; ***= $P<0.01$;

***= $P<0.001$; and ns= no significant


Due to significant interactions, HINiT resulted in significant higher total weed DM under N application rates compared with N0 (Figure 5.10). LINiT resulted in significantly lower total weed DM under N0, compared specifically with N70 and N210 (Figure 5.10).

Figure 5. 10. Cultivation techniques and nitrogen fertilisation interaction effect on weed biomass (July 2014)




HINiT resulted in significant higher broadleaf weeds DM under N application rates compared with N0, due to significant interactions (Figure 5.11).

Figure 5. 11. Cultivation techniques and nitrogen fertilisation interaction effect on broadleaf weed biomass (July 2014)



Total weed N uptake was significantly higher under HINiT with N140 and N210, compared specifically with N0 (Figure 5.12). LINiT resulted in a significantly higher total weed N uptake under N210 than specifically with N0.

Figure 5. 12. Cultivation techniques and nitrogen fertilisation interaction effect on weed N uptake (July 2014)



August assessment

CT resulted in significantly lower ($P<0.01$) total weed DM and total weed N uptake than HINiT and LINiT (Table 5.10). Legume DM and N uptake were not significantly affected by cultivation techniques at this assessment time.

Unfertilised treatment resulted in a significantly lower ($P<0.001$) total weed DM compared with N application rates (Table 5.10). Total weed N uptake significantly increased ($P<0.001$) with N rate, although no significant differences were observed between N70 and N140. Legume DM and N uptake were significantly higher ($P<0.01$) under N0 than with any other N rate treatment (Table 5.10).

BM resulted in significantly higher ($P<0.05$) legume DM and N uptake when compared with WC while Nus was statistically similar (Table 5.10). No significant undersowing effect on weed DM and N uptake was observed.


Table 5. 10. Effect of cultivation techniques and nitrogen fertilisation on weeds and legumes biomass and N uptake (August 2014)

	Total weed DM (t ha ⁻¹) August	Total weed N uptake (kg N ha ⁻¹) August	Legume DM (t ha ⁻¹) August	Legume N uptake (kg N ha ⁻¹) August
CT	0.177a	3.90a	0.0063a	0.16a
HINiT	2.292b	34.98b	0.0483a	1.38a
LINiT	2.220b	34.18b	0.0184a	0.48a
<i>SED (4 df)</i>	<i>0.2569</i>	<i>4.00</i>	<i>0.0155</i>	<i>0.432</i>
<i>P</i>	**	**	<i>ns</i>	<i>ns</i>
N0	0.851a	12.12a	0.0733b	2.1258b
N70	1.970b	24.13b	0.0155a	0.3746a
N140	1.718b	27.19b	0.0080a	0.1949a
N210	1.713b	33.98c	0.0006a	0.0029a
<i>SED (18 df)</i>	<i>0.2075</i>	<i>2.65</i>	<i>0.0185</i>	<i>0.545</i>
<i>P</i>	***	***	**	**
BM	1.528a	23.50a	0.0409b	1.1909b
Nus	1.479a	23.80a	0.0226ab	0.5957ab
WC	1.683a	25.80a	0.0096a	0.2370a
<i>SED (48 df)</i>	<i>0.2233</i>	<i>2.90</i>	<i>0.0116</i>	<i>0.336</i>
<i>P</i>	<i>ns</i>	<i>ns</i>	*	*

Values followed by same letter, do not differ significantly at $P<0.05$; *= $P<0.05$, ** = $P<0.01$; *** = $P<0.001$; and ns = no significant

Furthermore, at the August assessment, due to significant interactions, HINiT and LINiT resulted in lower weed DM with N0, while CT did not show significant differences with any N treatments (Figure 5.13).


Figure 5. 13. Cultivation techniques and nitrogen fertilisation interaction effect on weed biomass (August 2014)



In addition, HINiT resulted in a lower total weed N uptake at N0, compared specifically with N140 and N210 (Figure 5.14). LINiT resulted in significantly lower total weed N uptake under N0 compared with any other N rate application, while CT did not show significant differences with any N rate treatments on weed N uptake (Figure 5.14).


Due to significant treatments interaction, HINiT resulted in higher legume biomass and N uptake than CT and LINiT with N0 (Figure 5.15 & 5.16).

Figure 5. 14. Cultivation techniques and nitrogen fertilisation interaction effect on weed N uptake (August 2014)




Error bars representing average LSD ($P < 0.05$)

Figure 5. 15. Cultivation techniques and nitrogen fertilisation interaction effect on legume biomass (August 2014)



Error bars representing average LSD ($P < 0.05$)

Figure 5. 16. Cultivation techniques and nitrogen fertilisation interaction effect on legume N uptake (August 2014)



5.3.5. Weed species composition

In the July assessment, weeds were separated by species, recording a total of 29 weed species, five were grass weed species and 24 broadleaf weed species. The list of all weed species recorded can be found in Appendix 2. The dominant weed species were *Sinapsis arvensis* L., *Avena sativa* L., *Stellaria media* L., and *Galium aparine* L., accounting 35.1%, 26.9%, 15.2% and 14.8% of the total weed biomass recorded.

Within the dominant weed species, *Sinapsis arvensis* and *Stellaria media*, were the only species significantly affected by management treatments (Table 5.11). Other species occurred too infrequently to permit treatments effects to be appropriately tested or were not significantly affected by any treatment.

Table 5. 11. Effect of cultivation techniques, nitrogen fertilisation and undersowing on *Sinapsis arvensis* L. and *Stellaria media* L. biomass (July 2014)


	<i>Stellaria media</i> L. DM (t ha ⁻¹)	<i>Sinapsis arvensis</i> L. DM (t ha ⁻¹)
CT	0.0081a	0.0607a
HINiT	0.3359b	0.9070b
LINiT	0.1557a	0.1874a
<i>SED</i> (4 <i>df</i>)	0.0579	0.1104
<i>P</i>	**	**
N0	0.0693a	0.1097a
N70	0.2701b	0.3671b
N140	0.1669ab	0.4331bc
N210	0.1601ab	0.6304c
<i>SED</i> (18 <i>df</i>)	0.0574	0.1103
<i>P</i>	*	***
BM	0.1575ab	0.269a
Nus	0.104a	0.408a
WC	0.2383b	0.477a
<i>SED</i> (48 <i>df</i>)	0.0523	0.0948
<i>P</i>	*	<i>ns</i>

Values followed by same letter, do not differ significantly at $P < 0.05$; * = $P < 0.05$, ** = $P < 0.01$; *** = $P < 0.001$; and *ns* = no significant

Stellaria media and *Sinapsis arvensis* DM were significantly higher ($P < 0.01$) under HINiT compared with CT and LINiT (Table 5.11). *Stellaria media* DM was significantly higher ($P < 0.05$) with 70 kg N ha⁻¹, specifically compared with N0, while WC significantly increased its DM ($P < 0.05$), compared particularly with Nus. *Sinapsis arvensis* DM was significantly higher ($P < 0.001$) with N210 compared with N70 and N0 ($P < 0.001$).


Due to significant interactions, HINiT increased *Sinapsis arvensis* DM when N was applied compared with unfertilised treatment, while no significant differences were observed between N rates under CT and LINiT (Figure 5.17).

Figure 5. 17. Cultivation techniques and nitrogen fertilisation interaction effect on *Sinapsis arvensis* L. biomass (July 2014)



In addition, HINiT resulted in significant higher *Sinapsis arvensis* DM with N70 and N210 under undersown WC or Nus, while no significant differences were observed with BM (Figure 5.18). HINiT increased *Sinapsis arvensis* DM compared with CT and LINiT under BM undersown and N140. Under WC and N140, HINiT increased *Sinapsis arvensis* DM compared with CT.

Figure 5. 18. Cultivation techniques, nitrogen fertilisation and undersowing interaction effect on *Sinapsis arvensis* L. biomass (July 2014)



Error bars representing average LSD ($P<0.05$)

5.3.6. Plant height and ears number

Wheat crop height was not significantly affected by cultivation techniques treatments (Table 5.12). However, N fertilisation treatments gave significantly ($P<0.001$) taller plants with increasing N rates, although no significant differences were observed between N140 and N210 (Table 5.12). There were no significant effects of undersowing treatment or any other treatment interaction on plant height.

Wheat ear number per unit area was significantly affected by cultivation treatments ($P<0.05$) resulting in higher number under CT, compared specifically with LINiT (Table 5.12). No significant effects were observed on ears number by N fertilisation or undersowing treatments.


Table 5. 12. Cultivation techniques and nitrogen fertilisation effect on spring wheat height and ears number

	Plant height (cm)	Ears (number m⁻²)
CT	88.26a	463.80b
HINiT	83.88a	367.90ab
LINiT	84.26a	318.60a
SED (4 df)	1.316	35.94
<i>P</i>	<i>ns</i>	*
N0	75.41a	370.00a
N70	84.65b	391.90a
N140	90.00c	376.90a
N210	91.81c	395.00a
SED (18 df)	1.324	26.96
<i>P</i>	***	<i>ns</i>

Values followed by same letter, do not differ significantly at $P<0.05$; *= $P<0.05$; ***= $P<0.001$; and ns= no significant

Due to significant interactions, CT produced taller plants than HINiT and LINiT under unfertilised conditions (Figure 5.19). No significant differences were observed between tillage treatments when N was applied.


Figure 5. 19. Cultivation techniques and nitrogen fertilisation interaction effect on spring wheat height



Error bars representing average LSD ($P<0.05$)

CT resulted in higher number of ears compared with HINiT and LINiT under BM undersown and Nus, due to significant interactions (Figure 5.20). Under undersown WC, a higher number of ears were recorded under CT, compared specifically with LINiT.

Figure 5. 20. Cultivation techniques and nitrogen fertilisation interaction effect on spring wheat ears number



5.3.7. Diseases scoring

The severity of two diseases was evaluated during the 2014 cropping season, *Zymoseptoria tritici* Desm. Quaedvlieg & Crous (leaf blotch of wheat) and *Puccinia triticina* Erikss. (leaf rust of wheat). Regardless of the management treatments, the mean severity of *Zymoseptoria tritici* and *Puccinia triticina* were 26.1% and 2.1% respectively. However, there were not statistically significant effects of any management treatment or treatments interactions on the severity of these diseases, even when data variability was reduced by transforming raw data.

5.3.8. Final biological harvest

Ear, straw and total wheat biomass

Ear, straw and total wheat biomass were significantly affected ($P<0.01$) by cultivation techniques, with CT resulting in higher biomass compared with LINiT and HINiT (Table 5.13). N140 and N210 significantly increased ear ($P<0.01$), straw ($P<0.001$) and total wheat DM ($P<0.01$), compared with N70 and N0 (Table 5.13). No significant effects were observed by undersowing treatments on ear, straw and total wheat DM.


Table 5. 13. Cultivation techniques and nitrogen fertilisation effect on ears, straw and total wheat biomass

	Ears DM (t ha⁻¹)	Straw DM (t ha⁻¹)	Total wheat DM (t ha⁻¹)
CT	6.888b	5.015b	11.821b
HINiT	3.359a	2.913a	6.272a
LINiT	3.991a	3.176a	7.167a
SED (4 df)	0.3986	0.2808	0.611
<i>P</i>	**	**	**
N0	4.101a	3.101a	7.203a
N70	4.184a	3.407a	7.591a
N140	5.316b	4.046b	9.254b
N210	5.382b	4.251b	9.633b
SED (18 df)	0.3849	0.2461	0.593
<i>P</i>	**	***	**

Values followed by same letter, do not differ significantly at $P<0.05$; **= $P<0.01$; and ***= $P<0.001$

Due to significant interactions, with BM undersown, CT resulted in a significant higher straw DM than LINiT, followed by HINiT. However, with WC and Nus, CT resulted in higher straw DM compared with non-inversion tillage treatments (Figure 5.21).

Figure 5. 21. Cultivation techniques and undersowing interaction effect on spring wheat straw biomass



TGW, grains per ear and grain yield

Thousand grain weight (TGW) was not significantly affected by cultivation techniques (Table 5.14). Increasing N rate significantly increased ($P < 0.01$) TGW, although no significant differences were observed between N70 and N140.

Numbers of grains per ear was significantly ($P < 0.001$) increased under CT than LINiT, followed by HINiT (Table 5.14). Additionally, N140 and N210 applications significantly increased ($P < 0.001$) grains number per ear compared with N0 and N70. Due to significant interactions, LINiT and HINiT significantly increased grain number per ear under N140 and N210 compared with N0 and N70, while CT did not differ between N rate applications (Figure 5.22).


Overall the final spring wheat grain yield mean obtained in the 2014 cropping season was 3.69 t ha⁻¹, which was lower than the 2013 final yield (5.65 t ha⁻¹). Among tillage treatments, CT resulted in a significantly higher ($P < 0.01$) grain yield compared with HINiT and LINiT (Table 5.14). N fertilisation significantly affected ($P < 0.01$) the final grain yield, with higher yield with 140 and 210 kg N ha⁻¹ compared with N70 and N0 (Table 5.14).

Table 5. 14. Cultivation techniques and nitrogen fertilisation effect on spring wheat TGW, grains per ear, final grain yield and harvest index

	TGW (g)	Grain per ears	Grain yield (t ha ⁻¹)	HI (%)
CT	32.37a	36.27c	5.37b	45.04b
HINiT	32.65a	21.44a	2.53a	39.96a
LINiT	33.69a	29.69b	3.20a	44.24b
<i>SED (4 df)</i>	0.764	0.941	0.266	0.695
<i>P</i>	ns	***	**	**
N0	34.60c	23.70a	3.10a	41.76a
N70	32.99b	25.53a	3.26a	42.90a
N140	32.83b	33.34b	4.18b	44.05a
N210	31.19a	33.96b	4.25b	43.61a
<i>SED (18 df)</i>	0.737	1.24	0.290	1.166
<i>P</i>	**	***	**	ns

Values followed by same letter, do not differ significantly at $P < 0.05$; ** = $P < 0.01$; *** = $P < 0.001$; and ns = no significant

Figure 5. 22. Cultivation techniques and nitrogen fertilisation interaction effect on spring wheat grains per ear




Error bars representing average LSD ($P < 0.05$)

Harvest index (HI) was significantly higher ($P < 0.01$) under CT and LINiT compared with HINiT (Table 5.14). However, N fertilisation did not significantly affect HI. Due to significant interactions, higher HI was observed under CT than HINiT and LINiT when BM was undersown. Under Nus, LINiT resulted in higher HI than CT, while with

WC, non-inversion tillage treatments had higher HI than CT (Figure 5.23). No significant effects of undersowing treatments were observed on TGW, grains per ear, final grain yield or HI.

Figure 5. 23. Cultivation techniques and undersowing interaction effect on spring wheat harvest index



Wheat nitrogen yield

CT resulted in significantly higher ($P < 0.001$) total wheat and grain N uptake compared with HINiT and LINiT (Table 5.15). Grain protein and N harvest index (NHI) were not significantly affected by cultivation techniques treatments.

Total wheat N uptake significantly increased ($P < 0.001$) with increasing N rate, although no significant differences were observed between N0 and N70 (Table 5.15). Total grain N uptake was significantly higher ($P < 0.001$) with N140 and N210 compared with N0 and N70. Grain protein significantly increased ($P < 0.001$) with N rate although no significant differences were observed between N0 and N70. NHI was not significantly affected by N fertilisation.

Table 5. 15. Cultivation techniques and nitrogen fertilisation effect on total wheat and grain N uptake, grain protein content, and N harvest index

	Total wheat N uptake (kg N ha⁻¹)	Total grain N uptake (kg N ha⁻¹)	Grain protein (%)	NHI (%)
CT	155.60b	120.60b	12.757a	77.05a
HINiT	71.90a	52.54a	11.986a	72.77a
LINiT	89.10a	67.25a	11.968a	74.51a
<i>SED (4 df)</i>	8.87	6.57	0.3173	1.858
<i>P</i>	***	***	<i>ns</i>	<i>ns</i>
N0	79.70a	61.38a	11.17a	75.41a
N70	88.20a	66.02a	11.51a	73.85a
N140	117.70b	91.96b	12.60b	76.81a
N210	136.70c	101.15b	13.66c	73.05a
<i>SED (18 df)</i>	7.00	5.93	0.3819	1.769
<i>P</i>	***	***	***	<i>ns</i>

Values followed by same letter, do not differ significantly at $P<0.05$; ***= $P<0.001$; and ns=no significant

Wheat nitrogen efficiency

N-use efficiency (NUE) was significantly higher ($P<0.05$) under CT and LINiT than HINiT (Table 18). N uptake efficiency (NUpE) was significantly higher ($P<0.001$) under CT than LINiT, followed by HINiT (Table 5.16), while no significant effects of cultivation techniques were observed on N utilisation efficiency (NUtE).

NUE and NUpE significantly decreased ($P<0.001$) when increasing N fertilisation rate, although no significant differences were observed between N70 and N140 (Table 5.16). NUtE was significantly lower ($P<0.001$) under N210 compared with any other N rate.

Table 5. 16. Cultivation techniques and nitrogen fertilisation effect on N- efficiency parameters

	NUE (kg kg⁻¹)	NU_pE (kg kg⁻¹)	NU_tE (kg kg⁻¹)
CT	22.85b	0.6358c	34.83a
HINiT	18.82a	0.5202a	35.39a
LINiT	21.54b	0.5739b	36.23a
<i>SED (4 df)</i>	<i>0.68</i>	<i>0.01257</i>	<i>1.053</i>
<i>P</i>	*	**	<i>ns</i>
N0	34.38c	0.8890c	38.69b
N70	19.83b	0.5301b	37.06b
N140	17.47b	0.4836b	35.24b
N210	12.60a	0.4037a	30.94a
<i>SED (18 df)</i>	<i>1.4</i>	<i>0.03188</i>	<i>1.677</i>
<i>P</i>	***	***	***

Values followed by same letter, do not differ significantly at $P<0.05$; * = $P<0.05$; ** = $P<0.01$; *** = $P<0.001$; and ns = no significant

Weeds and legumes biomass and nitrogen yield

At harvest time, CT resulted in a significantly lower ($P<0.05$) weed DM and total weed N uptake compared with the non-inversion tillage treatments (Table 5.17). Additionally, N fertilisation significantly increased ($P<0.01$) weed DM compared with unfertilised treatment. Total weed N uptake was significantly higher ($P<0.001$) under N210, compared specifically with N0. No significant effects of undersowing treatments were observed on total weed DM and N uptake.

Legume biomass production at harvest time was not significantly affected by cultivation techniques and N fertilisation treatments (Table 5.17). However, legume N uptake was significantly lower ($P<0.05$) with N fertilisation, compared with unfertilised treatment. BM resulted in significantly higher ($P<0.01$) legume DM and N uptake than WC and Nus (Table 5.17).

There were no significant treatments interaction effects on weed and legume DM and N uptake.

Table 5. 17. Cultivation techniques, nitrogen fertilisation and undersowing effects on weed and legume biomass and N uptake (Harvest 2014)

	Weed DM (t ha ⁻¹)	Weed N uptake (kg N ha ⁻¹)	Legume DM (t ha ⁻¹)	Legume N uptake (kg N ha ⁻¹)
CT	0.1403a	3.61a	0.049a	1.24a
HINiT	1.3848b	22.05b	0.119a	4.12a
LINiT	1.1888b	21.38b	0.112a	3.53a
<i>SED (4 df)</i>	0.2666	4.12	0.0519	2.09
<i>P</i>	*	**	<i>ns</i>	<i>ns</i>
N0	0.4972a	8.32a	0.241	8.125b
N70	1.0939b	15.81b	0.049	1.454a
N140	0.8732b	14.30ab	0.074	1.944a
N210	1.1542b	24.29c	0.009	0.335a
<i>SED (18 df)</i>	0.1639	2.987	0.0846	2.872
<i>P</i>	**	***	<i>ns</i>	*
BM	0.887a	15.47a	0.20528b	6.756b
Nus	0.920a	15.90a	0.04087a	1.149a
WC	0.907a	15.67a	0.03353a	0.988a
<i>SED (47 df)</i>	0.1284	2.197	0.0515	1.854
<i>P</i>	<i>ns</i>	<i>ns</i>	**	**



Values followed by same letter, do not differ significantly at $P < 0.05$; * = $P < 0.05$; ** = $P < 0.01$; *** = $P < 0.001$; and *ns* = no significant

5.3.9. Soil mineral nitrogen

Throughout the 2014 cropping season, cultivation technique effects on soil mineral N (SMN) were only statistically evident at the June, July and August assessments (Figure 5.24). In June, CT resulted in significant higher ($P < 0.001$) SMN content than LINiT, followed by HINiT, while in July and August, SMN content was significantly higher under CT than LINiT and HINiT.


N fertilisation significantly affected SMN from June till August, although differences between treatments diminished with time (Figure 5.25). In June, increasing N rate significantly increased ($P < 0.001$) SMN content, while in July, N0 and N70 resulted in significantly lower ($P < 0.001$) SMN, compared with N140 and N210. At the August assessment, higher ($P < 0.01$) SMN was observed with 210 kg N ha⁻¹ application compared with any other N rate.

No significant undersowing effect on SMN was observed at any assessment time.




Due to significant interactions, at the July assessment, under N210, CT resulted in higher SMN content than LINiT and HINiT, while with N140, CT resulted in similar SMN content than LINiT but higher than HINiT (Figure 5.26).

Figure 5. 26. Cultivation techniques and nitrogen fertilisation interaction effect on soil mineral nitrogen content (July 2014)




In August, under BM, CT resulted in similar SMN content than LINiT but higher than HINiT (Figure 5.27). Under Nus, higher SMN was observed with CT than LINiT, followed by HINiT. Under WC, CT resulted in higher SMN content than LINiT and HINiT.

Figure 5. 27. Cultivation techniques and undersowing interaction effect soil mineral nitrogen content (August 2014)



In addition also in August, with N140 and N210, CT resulted in higher SMN under Nus compared specifically with HINiT, while under WC under CT resulted in higher SMN than with HINiT and LINiT (Figure 5.28).


Figure 5. 28. Effect of cultivation techniques, nitrogen fertilisation and undersowing interaction on soil mineral nitrogen (kg N ha^{-1}) (August 2014)



Error bars representing average LSD ($P < 0.05$)

Figure 5.29 and 5.30 represent SMN content and total wheat N uptake under cultivation techniques and N fertilisation rates treatments at each month of assessment. Wheat N uptake, in May, was high under CT while SMN content was statistically similar between cultivation techniques treatments (Figure 5.29). In July and August, wheat N uptake and SMN content were significantly higher under CT compared with HINiT and LINiT.

Figure 5. 29. Soil mineral nitrogen (kg N ha⁻¹) and total wheat N uptake under three cultivation techniques at each month of assessment




SMN (LSD)	5.597 ^{ns}	12.107 ^{ns}	13.92 ^{ns}	5.3***	17.51*	9.67**
Wheat N uptake (LSD)	-	-	1.471***	-	21.91***	24.59***

LSD at $P < 0.05$ at each month of assessment; * = $P < 0.05$; ** = $P < 0.01$; *** = $P < 0.001$; and ns = no significant

N fertilisation in May increased wheat N uptake while SMN levels were not significantly affected (Figure 5.30). In July and August, wheat N uptake increased as SMN level increased by the N rates.

Figure 5. 30. Soil mineral nitrogen (kg N ha⁻¹) and total wheat N uptake under four N fertilisation rates at each month of assessment



LSD at $P < 0.05$ at each month of assessment; * = $P < 0.05$; *** = $P < 0.001$; and ns = no significant


5.3.10. Soil moisture (gravimetric) content

Soil moisture content was relatively stable across spring months when rainfall was higher than the long-term seasonal average (Figure 5.31). However, moisture content level decreased in July as a result of the warm and dry weather conditions experienced.

Throughout the assessment months, cultivation techniques only significantly affected soil moisture content in April, when CT resulted in lower moisture content than HINiT and LINiT (Figure 5.31). No significant effect of N fertilisation and undersowing treatments were observed at any assessment time.


At the June assessment, due to significant interactions, moisture content was significantly higher under N0 with LINiT than with HINiT and CT. LINiT resulted in higher moisture content with N70, specifically compared with CT. No significant differences were observed between cultivation techniques with N140 and N210 rates (Figure 5.32). At the August assessment, HINiT resulted in higher soil moisture content particularly than LINiT under Nus (Figure 5.33).

Figure 5. 31. Effect of cultivation techniques and nitrogen fertilisation on soil gravimetric moisture content (%) with mean precipitation during the 2014 experimental period in comparison to the 10-year average




Error bars representing LSD ($P < 0.05$) at each month of assessment

Figure 5. 32. Cultivation techniques and nitrogen fertilisation interaction effect on soil moisture content (%) (June 2014)



Error bars representing LSD ($P < 0.05$)

Figure 5. 33. Cultivation techniques and undersowing interaction effect on soil moisture content (%) (August 2014)



Error bars representing LSD ($P < 0.05$)


5.3.11. Soil pH

Soil pH was only measured in March 2014 in order to evaluate initial values for the cropping season. Overall soil pH was 7.0 and was not significantly affected by any treatment structure or interactions. No further measurements were carried due to no significant effect observed in 2013 cropping. The N fertilisation effect observed in 2013 was only temporal.

5.3.12. Soil penetration resistance


Soil penetration resistance after cultivation techniques operations (March 2014), during vegetative growth (June 2014), and before harvest (August 2014) were not significantly affected by tillage treatments. In addition, in all assessments time, soil penetration resistance was lower than 2MPa which is the upper limit for root restriction growth (Atwell, 1993). Nevertheless, penetration resistance showed a depth gradient effect. In the April and June assessments, penetration resistance was higher at 30cm depth compared with above soil layers (at 5cm, 10cm, and 15cm) (Figure 5.34 & 5.35).

Figure 5. 34. Effect of cultivation techniques and soil depth interaction on soil penetration resistance (kPa) (March 2014) with cultivation techniques soil disturbance depth




Error bar represents average LSD value at $P < 0.05$

Figure 5. 35. Effect of cultivation techniques and soil depth interaction on soil penetration resistance (kPa) (June 2014) with cultivation techniques soil disturbance depth



In August, soil penetration resistance was lower at 5cm than 10cm and 30cm while no significant differences were observed between 15cm and 30cm depth (Figure 5.36).

Figure 5. 36. Effect of cultivation techniques and soil depth interaction on soil penetration resistance (kPa) (August 2014) with cultivation techniques soil disturbance depth



5.4. Discussion

5.4.1. Overwinter growth assessment

In the 2013 cropping season, the undersown legume species had poor and low establishment, and slow growth. However after the 2013 harvest, the legume species undersown were allowed to grow during winter, resulting in higher legume DM under CT than non-inversion tillage systems, as also observed at the 2013 harvest time. Nevertheless, the overwinter legume DM was higher than the biomass recorded at harvest 2013, supporting the observation that the undersown legume species have slow growth but great biomass production and winter hardiness, as reported by others (Döring *et al.*, 2013; Jones, 1992; Smýkal *et al.*, 2015). Lower legume DM observed under LINiT was probably an effect of the increased weed biomass observed under this

treatment compared with CT and HINiT. This suggests that weed growth was more favourable under LINiT during winter time, perhaps due to higher weed seeds in the soil surface, germinating later in the season.

The greater weed DM under high N fertilisation rate treatment suggests that, regardless of the lower soil residual N level, weed N uptake competitiveness allows their growth, as also reported by Bergkvist (2003). However, the greater legume biomass and higher N uptake under unfertilised conditions corroborate that legume species can be more competitive under low N environments due to their capacity to fix N, and to produce great biomass which possibly reduces weed pressure. This supports Döring *et al.*, (2013) and Liebman & Dyck (1993) reporting that a weed suppression efficiency by a cover crop is mainly dependant on its biomass production. Increasing N supply may have reduced legumes competitiveness against weeds, resulting in higher weed N uptake under BM than with WC under N supply conditions. Döring *et al.*, (2013) and Squire (1997) suggested that BM produces lower biomass than WC which can negatively result in lower competitiveness against weeds.

5.4.2. Soil penetration resistance

According to Atwell (1993), soil penetration resistance higher than 2 MPa can disrupt root growth and adversely affect crop development. In the present study, penetration resistance at all assessment times and at any soil depth measured (up to 30 cm depth), seems to have no restrict root growth as it was lower than 2 MPa. Increasing penetration resistance was observed specifically when the first 15cm of soil was compared with the 30 cm soil depth layer, agreeing with other authors reporting a depth gradient (Grant & Lafond, 1993; Martinez *et al.*, 2008). In the present study, cultivation techniques did not significantly affect soil penetration resistance at any soil depth layer, in contrast with findings by other studies (Ardvisson *et al.*, 2013; Carter *et al.*, 1965; López-Bellido *et al.*, 1996).

5.4.3. Plant establishment, tillers and total shoot production

In the temperate climate of the UK, spring wheat has a relatively short growing season making necessary an early sowing in spring in order to the crop to produce an adequate crop canopy (Webb *et al.*, 1995). However, UK weather conditions are not always suitable for early sowing. High rainfall in April 2014 delayed spring wheat drilling operations, which probably negatively affected plant establishment. This resulted in fewer plants established, perhaps, compared with the 2013 season. Variations of tillage intensity influenced seedbed conditions affecting the crop establishment, growth and development (Strudley *et al.*, 2008). Increasing soil disturbance and reducing soil residues cover under CT and HINiT resulted in higher plant establishment, as also reported by Arvidsson *et al.* (2014). Less soil movement and the higher presence of wheat stubbles under LINiT produced a coarse seedbed. This may have affected seed-soil contact, reducing the crop establishment, as reported by others (Graham *et al.*, 1985; Känkänen *et al.*, 2011; Pietola & Tanni, 2003).

Tiller production is cultivar-dependant, but it can be modified by growing conditions and management operations (Peltonen-Sainio *et al.*, 2009). Cultivation techniques that affected plant establishment also influenced the number of tillers produced, with CT and HINiT having significantly higher tillers per unit area than LINiT. High crop residue retention in the soil surface under non-inversion tillage systems can affect soil conditions by reducing evaporation limiting soil temperature fluctuations, as reported by Morris *et al.*, (2011) and Reicosky *et al.* (1995). These conditions could perhaps have created cold soils under the non-inversion tillage systems and have adversely affected the onset of tillers, giving lower number of shoots compared with CT, as reported elsewhere by Rasmussen *et al.* (1997). In addition, increasing weed pressure under non-inversion tillage could have adversely affected shoot production in spite of the greater tillers number under HINiT.

Contrary to several authors (Cannell, 1985; Power & Alessi, 1978; Otteson *et al.*, 2008; Weisz *et al.*, 2001), in the present study, application of N fertiliser did not significantly affect crop establishment and early crop development.

5.4.4. Mid-season plant biomass and nitrogen uptake

5.4.4.1. Wheat

Cultivation techniques effects

Cultivation effects on initial crop establishment and early crop development seem to also have significantly affected the mid-season crop growth. In May, maximum tillage intensity under CT resulted in higher wheat DM, probably as a result of the high plant establishment and the increased tiller production compared specifically with LINiT. Although, HINiT treatment resulted in statistically similar establishment and tiller production than CT, wheat DM was lower when compared with ploughed plots. This was possibly as the soil cover produced a colder soil environment slowing down wheat biomass growth, as widely reported (Alakukku *et al.*, 2009; Bahrani *et al.*, 2007; Känkänen *et al.*, 2011). This was also evident as shoot number decreased under non-inversion tillage, probably accounting for the low wheat biomass production at the July and August assessments. Decreasing soil disturbance resulted in higher variability of seedbed conditions and greater weed pressure under non-inversion tillage, negatively affecting crop growth compared with CT across the mid-season growth assessments.

Intensifying soil disturbance by cultivation operations can often increase SMN content mainly by exposing organic matter to decomposition, as reported by Myrbeck *et al.* (2012) and Silgram & Shepherd (1999). Maximum soil disturbance and incorporation of plant residues into the soil under CT resulted in higher availability of N allowing greater N uptake by the crop, compared with non-inversion tillage across the cropping season. However, at the early assessments similar SMN levels between cultivation treatments suggests that higher wheat N uptake under CT than HINiT followed by LINiT, was likely to be related to plant establishment rather than influenced by the soil N availability. This agrees with Brennan *et al.* (2014) reporting that lower plant population results in less N uptake. Differences between non-inversion tillage treatments on wheat N uptake were less evident later in the season (July and August), suggesting that N availability reduced the initial differences as soil N levels were statistically similar.

Nitrogen fertilisation effect

For cereal plant growth there is an absolute requirement of N, although growth rate depends on N inputs and timing between N supplies and crop N demand (Justes *et al.*, 1994). However, wheat N and biomass accumulation depend on intra-regulation of crop physical processes (Gastal & Lemaire, 2002). During early growth assessment, N fertilisation significantly increased total wheat N uptake. It did not increase wheat DM, however, as no significant differences between N treatments were observed, as also found by Qiao *et al.* (2013) at an early crop growth stage. Conversely, at later growth assessments, the wheat DM production was significantly affected by N fertilisation treatments, although differences between N rates were more evident later in the season (August). Wheat DM in August significantly increased with 210 kg N ha⁻¹ applied, particularly compared with unfertilised plots, while N140 and N70 resulted in similar wheat DM. This was also observed by López-Bellido *et al.* (2005) and Pearman *et al.* (1977) who reported increases in crop biomass production with increasing N fertilisation supply. At this growth stage the crop was reaching its maximum growth under high N conditions (Justes *et al.*, 1994), but a similar N availability under N140 and N70 plots possibly triggered comparable wheat biomass production. In addition, increases of total wheat N uptake with increasing N fertilisation rate probably promoted differences in wheat biomass growth between high N rate treatments and unfertilised conditions, as also reported by others (Power & Alessi, 1978; Justes *et al.*, 1994; van Keulen & Seligman, 1987).

Interaction effect

At early crop growth the WC undersown significantly encouraged wheat N uptake compared with Nus and BM with the application of 210 kg N ha⁻¹. In July, WC and Nus resulted in higher crop N uptake under CT. The slow growth pattern of WC and its lower competitiveness under N-rich conditions (Moss *et al.*, 2004) possibly allowed more N for the crop and increased N uptake.

5.4.4.2. Legumes

Despite delayed wheat sowing, the environmental conditions in May 2014 were expected to be favourable for establishment and growth of the undersown legume species. Within a week after legume broadcast, however, the average air temperature and rainfall were 13.7°C and 1.7 mm. Such dry (1.7mm vs 97.30mm average in May 2014) and still warmer (13.7°C vs 12.8°C, average in May 2014) conditions were unfavourable for the legume species to germinate and establish, as also observed in the 2013 cropping season. This may have resulted in low legume DM during all growth assessments and perhaps triggered the lack of undersowing effect in a wide range of assessments.

Nitrogen fertilisation effect

In May and August higher wheat and weed biomass growth possibly reduced legume DM when N fertilisation was applied. This is the result of poor competitiveness of BM and WC under N-rich conditions, as reported by other authors (Döring *et al.*, 2013; Elgersma *et al.*, 2000; Gooding & Davies, 1997; Moss *et al.*, 2004; Soussana & Arregui, 1995).

Interaction effect

At the overwinter assessment, increasing tillage intensity resulted in higher legume DM as a result of lower weed competition. After herbicide applications, however, legumes that survived broad-spectrum herbicide application were incorporated into the soil under CT resulting in lower legume DM. In contrast, lower tillage intensity under non-inversion tillage increased legume biomass. This combined with less SMN content under HINiT than LINiT, favoured legume species growth, as observed in May and August. This agrees with Carr *et al.*, (2004), who reported higher total aboveground legume DM in soils with low N content compared with high N soil conditions.

5.4.4.3. Weeds

As for the 2013 cropping season, weed species observed during the 2014 season are commonly report in UK spring wheat (HGCA, 2010a) and their presence was again influenced by the agricultural management treatments and assessment date.

Herbicide application prior to cultivation operations appears to have provided a good head-start for the main crop, allowing its faster growth. It possibly contributed to reduced weed biomass production and mitigated any treatment effect in the early growth assessment (in May). This was more evident with grass weeds as the herbicide application seems to have particularly reduced their growth across the cropping season.

Cultivation techniques effect

In July and August, decreasing tillage intensity under non-inversion tillage led to higher total weed biomass compared with CT, as also reported by several authors (Cardina *et al.*, 1991; Clements *et al.* 1994a; Clements *et al.*, 1996a; Swanton *et al.*, 2000). In the present study, broadleaf weed species were the main contributors to total weed DM under non-inversion tillage. Weeds retained in soil under HINiT could have germinated after the herbicide application. Soil movement without soil inversion under HINiT, therefore, allowed more short-lived broadleaf weeds than CT and LINiT, as also observed in the 2013 cropping season. The high biomass of *Sinapsis arvensis* L. and *Stellaria media* L. under HINiT suggest their great contribution to total weed DM. These species accounted for 50.3% of the total weed biomass observed at the July assessment. This agrees with Ozpinar (2006) who reported the higher presence of *Sinapsis arvensis* under reduced tillage compared with CT. However, maximum reduction of tillage intensity under LINiT resulted in greater soil residues cover, creating shading and negatively affecting broadleaf weeds that survive herbicide application, as reported by other studies (Johnson *et al.*, 1993; Streit *et al.*, 2002; Teasdale *et al.*, 1991). Subsequent soil inversion after the foliar-contact herbicide application and greater crop competitiveness against weeds appear to have reduced broadleaf weeds biomass under CT, as observed by several authors (Demjanová *et al.*, 2009; Gruber *et al.*, 2012; Menalled *et al.*, 2001; Vijaya Bhaskar *et al.*, 2014b). Higher

weed N uptake under non-inversion tillage suggests that N uptake was a result of the greater weed DM under those treatments, rather than the soil N availability.

Nitrogen fertilisation effect

Total weed biomass was affected by N application, confirming that weed growth responds positively to higher N levels, as also reported by Blackshaw *et al.* (2005). Increasing broadleaf biomass was observed under N fertilised conditions compared with unfertilised plots, while grass weeds were not significantly affected. This confirms N supply differentially affects weed species growth, and can modify the competitive interactions between species (Iqbal & Wright, 1997). The dominant weed species, *Sinapsis arvensis* and *Stellaria media* contributed to the greater weeds biomass under N application rates, as these species seems to be highly more competitive under N-rich conditions (Iqbal & Wright, 1997; Moss *et al.*, 2004; Lal *et al.*, 2014). Weed N uptake, at the July and August assessments, was affected by N fertilisation treatments with N0 resulting in the lowest N weed uptake. This indicates that weed N uptake increased with increasing N supply as also observed on weed biomass, agreeing with Blackshaw *et al.* (2005) findings.

Undersowing effect

The increased competitive behaviour of *Stellaria media* under N-rich conditions (Moss *et al.*, 2004; Parchoma, 2002) possibly encouraged its biomass growth under WC as SMN was higher, particularly under N210 treatment.

Interaction effect

Cultivation techniques and N fertilisation interacted affecting weeds DM. N application under non-inversion tillage resulted in higher broadleaf and total weed biomass at the July and August assessments. HINiT increased *Sinapsis arvensis* biomass when N fertiliser was applied, regardless of the N rate. This support preference of this broadleaf weed species to N fertile conditions (Lundkvist & Verwijst, 2011). *Sinapsis arvensis* DM also positively responded to N70 and N210 applications under HINiT with WC undersown and Nus. However, when N140 was applied under HINiT, BM undersown resulted in higher *Sinapsis arvensis* DM. *Sinapsis arvensis* is more advantaged under

non-inversion tillage (Streit *et al.*, 2003), therefore, and its competitiveness was higher than the undersown legume species under N-rich conditions, particularly BM as this species has low biomass production and is disadvantageous to high N supply (Moss *et al.*, 2004).

5.4.5. Plant height and ears number

Cultivation techniques effect

As also observed in the 2013 cropping season, cultivation treatments did not significantly affect wheat plant height, indicating that weather conditions and N management were more relevant. Nevertheless, average plant height was relative shorter than in the 2013 cropping season, regardless of treatments. This is probably a result of later sowing, as the crop growth over a shorter period (Prasad *et al.*, 2008).

Tillage management affecting plant establishment, tiller production and shoot survival can also influence ear production (Mc-Master *et al.*, 1994). Soil conditions under CT were favourable for wheat growth promoting tillers and shoot production, and later greater number of ears per area (López-Bellido *et al.*, 1998; López-Bellido *et al.*, 2000). Similarly, Brennan *et al.* (2014) reported that high crop establishment under CT can contribute to high ear production. However, LINiT resulted in variable seedbed conditions and higher weed pressure compared with CT, inducing slower plant growth, affecting shoot survival and final ear number (Boomsma *et al.*, 2010). Higher tillage intensity and higher soil cover under HINiT could have resulted in similar ears number with LINiT and CT.

N fertilisation effect

Increases in plant height with increasing N supply is a well-known result of increasing N availability (Alijani *et al.*, 2012; Sourour *et al.*, 2014) and promoting the early vigorous wheat growth (Ayoub *et al.*, 1993; Lloveras *et al.*, 2001) by N fertilisation, particularly with high application rates, N140 and N210 in the present study.

Contrary to Abad *et al.* (2005) and Hay & Walker (1989) reporting increases in ear number by N fertilisation due to more tillering, the present study showed that N fertilisation did not significantly affect tillers production or ears number per area.

Interaction effect

Taller plants were observed in CT than LINiT and HINiT under unfertilised conditions. Suggesting that tillage treatment effects on nutrient availability promote plant growth as higher SMN level was recorded under CT, compared with non-inversion tillage treatments, as observed by Alijani *et al.* (2012).

Under BM undersown and Nus, CT resulted in higher ears number compared with non-inversion tillage treatments. While under WC, CT specifically resulted in higher ears number than LINiT. The higher wheat N uptake, observed at the July assessment, may have encouraged shoot survival and promoted more ears under those treatments interactions. Abad *et al.* (2005) also reported a positive effect on shoot survival and ears number with increasing wheat N uptake.

5.4.6. Ear, straw and total wheat biomass

Cultivation techniques effect

Under CT, higher ears number and the greater promotion of crop growth resulted in higher total wheat DM, compared with non-inversion tillage treatments. Greater variability of seedbed conditions and higher weed pressure observed under non-inversion tillage induced slower plant growth and ears number than CT resulting in poor total wheat DM, as observed by Boomsma *et al.* (2010) and Mehdi *et al.* (1999).

Nitrogen fertilisation effect

Increases in the early wheat biomass production by N fertilisation were translated to the final straw DM, with higher biomass with increasing N supply. The promoting N fertilisation effect on ear DM was a result of increasing number of grains per ear rather than an increase in the ears number per unit area, as previously reported by Pearman *et al.* (1977).

Interaction effect

Under BM, CT resulted in higher straw DM than LINiT, followed by HINiT. While CT with WC and Nus, resulted in higher straw DM compared with non-inversion tillage systems. These significant interaction effects on straw DM may have been triggered by the increase of earlier wheat N uptake previously discussed.

5.4.7. TGW, grains per ears, final grain yield and harvest index

Cultivation techniques effect

Differences in agricultural management, weather conditions, weed pressure and N availability influenced crop growth and final grain yield between cultivation techniques treatments. Higher plant establishment and plant growth followed by significantly higher ears number and grains per ear resulted in a significantly higher yield under CT than non-inversion tillage systems, regardless of the statistically similar TGW. In contrast, the higher variability of seedbed conditions combined with higher weed presence and lower N availability under non-inversion tillage systems negatively affected early crop growth and development. These negative constraints under non-inversion tillage resulted in less plant established and low plant growth followed by low ear number and grains per ear, finally causing a lower grain yield than CT. Several authors (Alvarez & Steinbach, 2009; Arvidsson *et al.*, 2013; Brennan *et al.*, 2014; Franchini *et al.*, 2012; López-Bellido *et al.*, 2000; Pietola, 2005; Vijaya Bhaskar *et al.*, 2013b) also reported higher yields under CT as with more favourable growing conditions compared with non-inversion tillage systems.

The higher number of grains per ear under CT followed by LINiT perhaps triggered similar harvest index (HI) under these treatments, as also reported by Hay (1995). However, it is evident that the higher total wheat DM under CT supported a higher grain yield compared with non-inversion tillage.

Nitrogen fertilisation effect

Thousand grain weight (TGW) is genetically determined, although its expression is affected by growing environment (De Vita *et al.*, 2007; Mogensen *et al.*, 1985). In the present study, increasing N fertilisation rate decreased TGW, although N70 and N140 resulted in statistically similar TGW. Lower TGW probably resulted from higher number of grains per ear and lower grain fill, perhaps, as reported by other authors (Alijani *et al.*, 2012; López-Bellido *et al.*, 2000; Pearman *et al.*, 1977; Rasmussen *et al.*, 1997).

Increasing N availability when 140 and 210 kg N ha⁻¹ were applied promoted higher number of grains resulting in significantly higher final grain yield. Several authors (Abad *et al.*, 2005; Alijani *et al.*, 2012; Halvorson *et al.*, 2001; Wang *et al.*, 2014) also reported a positive crop yield response with mineral N fertilisation.

Interaction effect

Non-inversion tillage resulted in more grains per ear when N140 and N210 were applied. These interactions suggest that the lower soil N availability under non-inversion tillage negatively affected grain production, making a higher N supply necessary to increase grains number per ear. This agrees with McConkey *et al.*, (2002) reporting that non-inversion tillage can require higher N supply to increase grain number compared with CT.

Cultivation techniques and undersowing treatments interaction effects on HI was strongly related to their effect on straw biomass. LINiT resulted in higher straw DM under BM specifically than Nus, reducing the HI under these treatments interactions.

5.4.8. Wheat N yield, grain protein and N efficiencies

Protein content in the present study was slightly lower compared with 2013 spring wheat. This might be due to interactions between environmental conditions (perhaps higher rainfall) and lower grain yields. These well-known observations support several authors (López-Bellido *et al.*, 1998 & 2001; Rao *et al.*, 1993; Stobard & Marshall,

1990) reporting that grain protein is highly determined by genotype, although still influenced by prevailing environmental conditions. This was evident with grain protein content not being significantly influenced by cultivation treatments, in spite of higher total wheat and grain N uptake under CT.

Cultivation techniques influences on SMN levels, final grain yield and N uptake seem to result from variations of N-use efficiency (NUE) and N-uptake efficiency (NUpE), as also reported by Håkansson (1994). Higher N availability, total wheat N-uptake and final crop yield under CT led to higher NUE and NUpE than with non-inversion tillage systems.

Nitrogen fertilisation effect

High N rate treatments significantly increased total wheat and grain N uptake compared with low N rate – the N70 and unfertilised conditions, indicating greater N availability at the time of crop high demand as reported by Campbell *et al.* (1977) and Halvorson *et al.* (2001). In the same way, higher grain protein content was also observed under N210 than N140, followed by N70 and N0, as reported by other authors (Garrido-Lestache *et al.*, 2004; Godfrey *et al.*, 2010; López-Bellido *et al.*, 2001). However, application of N fertiliser at 210 kg N ha⁻¹ would appear to be only justified for improving grain protein content, as this rate did not increase grain yield compared with 140 kg N ha⁻¹ application.

NUE and NUpE decreased with increasing N rate, probably as a result of increasing N supply increasing aboveground N relative to grain yield, as also reported by Huggins *et al.*, (2010). However, N utilisation efficiency (NUE) was significantly lower under N210 compared with any other N rate, suggesting lower efficiency in the use of the total wheat N uptake to produce the final yield. This was corroborated as N210 resulted in the highest total wheat N uptake, while grain yield was similar to that of N140.

5.4.9. Non-wheat biomass and N yield at harvest

5.4.9.1. Weeds

Weed biomass was reduced from the mid-season assessments to harvest time, probably as a result of natural decay of the weeds and warm and dry conditions during the summer. This was also observed during the 2013 cropping season.

Cultivation techniques effect

As previously observed during the mid-season growth assessments, seedbed conditions under non-inversion tillage treatments also resulted in high weed biomass at harvest time. The increase in tillage intensity and the soil inversion under CT resulted in less weed pressure, also favouring wheat growth and increasing wheat competitiveness against weeds, as also reported by Håkansson (2003) and Swanton *et al.* (2000). Lower N availability under non-inversion tillage suggests that the higher weed N uptake under these treatments was triggered by increases in weed DM, rather than by the soil N available.

Nitrogen fertilisation effect

Mineral N fertilisation affects soil fertility, influencing not only the main crop growth but also weeds growth (Jornsgård *et al.*, 1996; O'Donovan *et al.*, 1997). As also observed at the mid-season assessments, N application significantly increased weed pressure compared with unfertilised plots. Several authors (Ampog-Nyarko & De Datta, 1993; Blackshaw *et al.*, 2005; Moss *et al.*, 2004; Lal *et al.*, 2014) also reported a positive weed growth and N uptake response to N fertilisation, mainly due to differential competitiveness to N supply by the various weed species.

5.4.9.2. Legumes

Poor establishment of the undersown legume species resulted in low legume DM across the cropping season, regardless of management treatments. However, higher legume DM at harvest time compared with the mid-season assessments reinforces, as also observed in the 2013 season, the slow growth pattern but greater biomass production of

BM and WC (Döring *et al.*, 2013). Similarly, Amossé *et al.* (2013) reported that legume cover crops established under winter wheat can reach high levels of biomass at harvest time. Nevertheless, later legume growth was not significantly related to cultivation techniques and N fertilisation treatments, although unfertilised treatment resulted in higher legume N uptake. This emphasised the low response of legume species under fertile N conditions (Döring *et al.*, 2013). Higher legume biomass and N uptake under BM compared with WC and Nus agree with Döring *et al.*, (2013) reporting that BM aboveground biomass is significantly higher than WC, regardless of the slow establishment of BM (Wallace, 2001). However, Nus resulted in similar biomass than WC which perhaps is a result of natural regeneration of the legumes, in spite of previous herbicide application. Under Nus, legumes were separated from weeds even though they were considered legume weeds rather than the intended undersown legume.

5.4.10. Soil moisture content

Soil moisture content is highly affected by weather conditions and by soil management practices adopted (Fitter, 1991). In the present study, soil moisture (gravimetric) content seems to be fairly consistent across spring months. This is probably the result of increasing rainfall from March to May, which coincided with fast crop growth. Increasing crop growth increases demand for water, affecting moisture content level (Pietola & Tanni, 2003). Later crop growth combined with low rainfall and high temperatures during the summer months, however, resulted in a decline of moisture content in June which was highly evident in July. In contrast, increasing rainfall in August increased the soil moisture content.

Cultivation techniques effect

Cultivation techniques affected soil moisture content although this was only significantly evident in April, when CT resulted in lower moisture content compared with non-inversion treatments. Land preparations at the end of March and crop sowing in mid-April probably created differences between cultivation techniques. Maximum soil disturbance under CT breaks the water-related pores increasing water loss by evaporation (Reicosky *et al.*, 1999). In contrast, less soil disturbance and greater soil

residues cover under non-inversion tillage treatments reduce evaporation and run-off while also slowing soil drying (Baumhardt & Jones 2002; Beyaert *et al.*, 2002; Steiner, 1989). These attributes under non-inversion tillage treatments maintain or perhaps increase soil moisture content, as also reported by others (Fabrizzi *et al.*, 2005; Hatfield *et al.*, 2001; Pietola, 2005).

5.4.11. Soil mineral nitrogen

Soil mineral nitrogen (SMN) content in March 2014 was lower than the residual SMN observed after 2013 harvest. Despite the fact that no leaching assessment was performed, the low SMN content in March 2014 was thought to be the result of N losses by leaching and/or denitrification due to high rainfall occurring during winter months. Several authors (Abad *et al.*, 2005; Halvorson *et al.*, 2001; Lloveras *et al.*, 2001) have also reported decreases in residual SMN content as expected following high rainfall leaching soil N. Additionally, overwinter assessments showed that legumes and weed N uptake could have also contributed to reductions of SMN.

Cultivation techniques effect

Soil disturbance often increases SMN content as organic matter is exposed to decomposition, although such an effect is reportedly temporary (Myrbeck *et al.*, 2012; Silgram & Shepherd, 1999). In the present study, the cultivation techniques treatments only affected SMN content from June until August. At the June assessment, CT resulted in higher SMN than LINiT, followed by HINiT. Maximum soil disturbance with CT can greatly increase organic matter mineralisation increasing SMN to meet crop N demand, as widely reported (Gruber *et al.*, 2011; McConkey *et al.*, 2002; López-Bellido *et al.*, 2013; Soon *et al.*, 2001; Yagioka *et al.*, 2015). Less tillage intensity and soil cover presence can result in slow decomposition of crop residues and high N immobilisation or low rate of N releases. This can, therefore, result in lower SMN under non-inversion tillage than under CT, agreeing with several authors (Alvarez *et al.*, 1995; López-Bellido *et al.*, 2013; López-Bellido *et al.*, 1997). In addition, the reduction of tillage intensity, higher plant establishment and early plant growth under HINiT probably resulted in lower SMN level than CT and LINiT at the June assessment. Nevertheless,

later in the season (in July and August) CT continued to present higher SMN levels than non-inversion tillage in spite of higher plant biomass production (and N uptake). This may have been the result of increasing mineralisation and low N losses with high temperatures and low rainfall recorded in these months.

SMN availability and root distribution determine plant N uptake (Gastal & Lemaire, 2002). Greenwood *et al.* (1990) suggested that N uptake has often related either to crop demand or to soil N availability rather than both simultaneously. Throughout the present assessments, total wheat N uptake was higher under CT compared with non-inversion tillage treatments. This suggests that maximum soil disturbance increased SMN, and promoted better crop growth and productivity compared with reductions in tillage intensity, as also reported by Germon *et al.* (1994). Differences between cultivation techniques on wheat N uptake in the May assessment, despite SMN content being unaffected by tillage treatments, suggests that N uptake at this time was induced by differences in plant population levels affecting total wheat biomass.

Nitrogen fertilisation effect

Application of N fertiliser in late May seems to increase SMN levels even though this started to be evident in June rather than in May. This resulted from earlier soil assessment in May before the N applications. Similarly, several authors (Giacomini *et al.*, 2010; Glendining *et al.*, 1996; Liebig *et al.*, 2002; Zhao *et al.*, 2014) reported increases in SMN content with increasing N fertilisation supply. In the present study, the application of 210 kg N ha⁻¹ resulted in high SMN content from June till August. However, increases in wheat N uptake across growth assessments seem to have reduced differences between the low N rates as a result of increasing crop N demand as the crop approached maturity (Fuente *et al.*, 2003). This possibly depleted SMN levels when N supplies were low, agreeing with Zhao *et al.* (2014) reporting reductions of SMN content when crop uptake is higher than the N supplied.

Interaction effect

At the July assessment, cultivation techniques did not affect SMN content when there was a deficiency of N (N0 and N70). Differences were only evident at high N rates (N140 and N210) when SMN contents were higher under CT than under non-inversion

tillage, specifically HINiT. Similarly, López-Bellido *et al.* (2013) also reported higher SMN level under CT compared with non-inversion tillage when high N rates were supplied. At the August assessment, significantly higher legume N uptake under BM at harvest time perhaps reduced SMN. This was probably the effect of CT significantly interacting with BM and Nus, resulting in lower SMN content compared with the undersown WC.

5.5. Conclusions

Delayed sowing, variations of seedbed conditions induced by tillage, and greater weed prevalence, negatively affected spring wheat growth and reduced yield performance under non-inversion tillage treatments compared with CT. This agrees with other studies with similar findings (Arvidsson *et al.*, 2013; Brennan *et al.*, 2014; Franchini *et al.*, 2012; López-Bellido *et al.*, 1996). Key findings for the 2014 cultivation techniques effects are listed in Table 5.18.

Table 5. 18. Key outcomes for the cultivation techniques effects during 2014 cropping season

	CT	HINiT	LINiT
Tillage intensity	High	Intermediate	Low
Seedbed	Fine	Coarser	Much coarser
Condition	Level/uniform	Variable/Not uniform	Highly variable/Not uniform
Plant establishment	High	High	Low
Tiller production	High	High	Low
Plant height	Statistically not significant		
Ears number	High	Intermediate	Low
TGW	Statistically not significant		
Number of grains per ears	High	Low	Intermediate
Grain yield	High	Low	Low
SMN	High	Low	Low
Moisture content	Statistically not significant		

High rainfall patterns across the cropping season possibly reduced the level of residual SMN, allowing a positive effect of the N fertilisation application on final grain yield, up to 140 kg N ha⁻¹. Under this study weather conditions, it seems that 210 kg N ha⁻¹ rate is only required to increase grain protein content. Key outcomes for the N effects are listed in Table 5.19.

Table 5. 19. Key outcomes for the nitrogen fertilisation effects during 2014 cropping season

	N0	N70	N140	N210
Plant height	Low	Medium	High	High
Ears number	Statistically not significant			
TGW	High	Medium	Medium	Low
Number of grains per ears	Low	Low	High	High
Grain yield	Low	Low	High	High
SMN	Highly low	Low	Medium	High

Low precipitation at the time of undersowing caused failure of the undersown legume species as also observed in the 2013 cropping season.

CHAPTER SIX

Effect of weather on spring wheat performance following different cultivation regimes

6.1. Introduction

Shifts in weather patterns and increases in the frequency and magnitude of extreme weather events are a result of the global warming (Lobell *et al.*, 2012; Semenov & Shewry, 2011). According to the UK Meteorological Office report (Gosling *et al.*, 2011), the UK has been warming since 1960 with greater warming in the summer than winter. Gosling *et al.* (2011) report suggests that for the south of the UK an intensification of the frequency of water stress and drought is projected. The increase in air temperature and the incidence of drought associated with global warming are serious threats to crop production (IPCC, 2014a,b; Lobell *et al.*, 2013). In the UK, it has been estimated that a potential loss of around 10-20% of typical wheat yield could occur due to drought (Foulkes *et al.*, 2001). According to Spink *et al.* (2009), about 12% of the wheat crop in the UK is grown on land susceptible to yield limiting droughts in 2 out of 3 years. In the present study, the 2013 and 2014 cropping seasons experienced contrasting weather conditions, particularly rainfall patterns. The 2013 growing season, experienced higher rainfall conditions at beginning of the growing period in March, and much drier summer months compared with the 2014 season and the long-term seasonal average. The different weather conditions have highlighted the effect of soil management practices adopted reflected in subsequent differences in resultant wheat yields.

The aim of the present study, therefore, is to further investigate weather conditions, in particular rainfall, influences that impact on the effect of cultivation techniques including conventional tillage (CT), high intensity non-inversion tillage (HINiT) and low intensity non-inversion tillage (LINiT), on spring wheat production. Meta-analysis is the statistical combination and summarisation of results from multiples studies (Gurevitch & Hedges, 1999). In the current study a meta-analysis of trial results was conducted to compare and integrate the results of three experimental studies in order to identify general patterns.

6.2. Material and methods

The present study combined and contrasted results from a separate published study with 2013 and 2014 results previously discussed in Chapter 4 and Chapter 5. The published results were obtained from an adjacent organic experimental study (Vijaya Bhaskar *et al.*, 2013b) conducted from March to August 2012 at the Royal Agricultural University's Harnhill Manor Farm (NGR SP 075 006). The 2012 organic experiment followed a fully factorial treatment structure – *Spring wheat (block) x tillage systems (main plot) x +/- undersowing (subplot)*, with three replicates. All agricultural management treatments used in the organic experiment were present in each following year (2013 and 2014) except for herbicide application and N fertilisation management which were adopted for the 2013 and 2014 periods. For the 2013 and 2014 studies, only the results under an unfertilised N (N0) treatment were used. These studies followed then a fully factorial treatment structure.

Cultivation techniques in 2012 were named differently from the 2013 and 2014 studies. However, in order to maintain a uniform dataset, the terminology for the treatments is as previously used in the 2013 and 2014 studies, as tillage operations were the same. Treatment terminologies adopted in 2012 and their equivalents used in the present study are given in Table 6.1.

Table 6. 1. Cultivation techniques treatments terminology used by Vijaya Bhaskar *et al.* (2013b) and their equivalents in the present study

This item has been removed due to 3rd Party Copyright. The unabridged version of the thesis can be viewed in the Lanchester Library Coventry University.

The specification details for the cultivation techniques treatments were previously described in Chapter – 3, Material and Methods, § 3.2.2.1. Further details of the spring wheat cropping seasons are given in Table 6.2.

Table 6. 2. Details of the spring wheat cropping seasons

	2012	2013	2014
Cultivation technique treatments	CT, HINiT and LINiT		
Cultivation techniques performance date	09 March	20 March	24 March
Spring wheat	cv Paragon		
Seed rate	420 seed m ²	480 seed m ²	
Sowing date	14 March	10 April	18 April
Harvest date	22 August	27 August	31 August
Average grain yield (t ha ⁻¹)	2.86	5.79	3.07

Building the database

A database template was designed including the selected data from the 2012 organic study and the 2013 and 2014 data reported in the Chapters 4 and 5. In accordance with the objectives of the present study and the requirements of the meta-analysis, the database was formulated including the following categories: yield, cultivation techniques type, rainfall and standard deviations. The database is given in Appendix 3. The spring wheat growing season was from March to August, therefore, total season rainfall was only considered from those months.

Meta-analysis treatments and calculations

Meta-analysis allows experimental results reported to be combined for analysis towards evaluating overall treatment effects (Gurevitch & Hedges, 1999). To assess the overall effect and to determine the treatments constancy across the studies, mean differences were weighted. Initially, the 2013 and 2014 data were analysed separately, by analysis of variance. Comparisons between the mean of the HINiT treatment and the CT treatment, and their standard errors were calculated from each experiment, and used in a meta-analysis to obtain a consensus estimate combining information from all the experiments (see Whitehead, 2002, Section 4.2.). The same analysis was then done for the comparison between LINiT and CT. Seasonal rainfall was used as a covariant for the crop yield response to cultivation techniques managements. Seasonal rainfall was categorized into low (<300 mm), medium (300-500 mm) and high (>500 mm). The analysis performed produces a plot showing estimates of the comparisons from each

experiment together with the combined estimate, all with 95% confidence regions. The effect is significantly different from zero if its confidence region does not overlap zero. The analyses were performed using the META procedure of Genstat statistical software (15th Edition VSN International Ltd, Hemel Hempstead, UK).

6.3. Results and discussion

Cropping season (March-August) mean air temperatures for the studied years, 2012-2014, were below the 10- year average (2002-2012) temperature (13.9 °C) (Table 6.3). As air temperatures were quite similar between the cropping years studied, for the purpose of this study, only rainfall was evaluate as a covariant for the meta-analysis.

Table 6. 3. Monthly air temperature (°C) for the study period (2012-2014) and for the long-term records (2002-2012). Royal Agricultural University meteorological station, (NGR SP 42 004 011)

	2012	2013	2014	Average	Long term-average (2002 -2012)
March	8.34	3.09	7.18	6.20	7.48
April	6.81	7.31	9.71	7.94	10.28
May	11.84	9.97	12.08	11.30	13.05
June	13.60	13.60	15.25	14.15	16.34
July	15.71	18.94	18.03	17.56	18.27
August	16.55	17.14	14.74	16.15	18.43
Mean	12.14	11.68	12.83	12.22	13.98

Total seasonal rainfall for the 2013 cropping period (292.0 mm) was below the 10-year average seasonal rainfall (377.2 mm) (Table 6.4). Seasonal rainfall in 2012 (589.1 mm) and 2014 (400.5 mm) were above the 10-year average. Categories based on seasonal rainfall were used, instead of experimental year, to label the y-axis in Figure 6.1 and 6.2 below.


Table 6. 4. Monthly and cumulative rainfall (mm) for the study period (2012-2014) and the long-term records (2002-2012). Royal Agricultural University meteorological station, (NGR SP 42 004 011)

	2012	2013	2014	Average	Long term-average (2002 -2012)
March	24.9	76.8	39.5	47.07	51.39
April	126.3	31.5	65.9	74.57	45.37
May	50.9	76.6	97.3	74.93	68.15
June	175	42.5	49.7	89.07	64.60
July	99.8	31.3	56.6	62.57	82.57
August	112.2	33.3	91.5	79.00	65.15
Total	589.1	292.0	400.5	427.2	377.24

The long-term total rainfall for the last 60 years is presented in Figure 6.1. During this period, four extreme events of high rainfall (>1000 mm) occurred, with two events observed in the last 20 years. However, dry years with rainfall <600 mm were more common, recording three out of five in the last 20 years. Figure 6.1 shows the total rainfall divided into cropping season rainfall and off-season rainfall (October-February). From the seasonal rainfall, four events of rainfall >500 mm occurred in the last 60 years, although three out of four were recorded in the last 10 years. In addition, 11 cropping seasons experienced seasonal rainfall <300 mm, with six out of eleven happening in the last 20 years. These results show growing evidence that rainfall patterns are changing with drier and in some cases wetter seasons occurring more frequently.


The amount and distribution of rainfall are crucial for crop performance, including wheat (Mrabet, 2011). Moisture stress at critical growth stages can inhibit root growth, reduce tiller production, diminish wheat vegetative growth and number of grains per ear, and possibly cause poor grain filling (Gooding & Davies, 1997; Mrabet, 2011).

Figure 6. 1. Sixty years of long-term cropping season and off-season rainfall (mm). Royal Agricultural University meteorological station (NGR SP 42 004 011)



Analysis of the comparison between HINiT and CT showed strong evidence that the effects differ between seasonal rainfall (chi-square 93.6 on 2 df). Figure 6.2 shows that the overall spring wheat yield were 3.23 t ha⁻¹ and 0.56 t ha⁻¹ higher under CT compared with HINiT, when growing season rainfall were 300 – 500 mm and >500 mm, respectively. There was no evidence of any difference for rainfall <300 mm. Combined over all the studies, yield was higher by 1.34 t ha⁻¹ with CT compared with HINiT.


Figure 6. 2. Weighed mean differences in crop grain yield in HINiT compared with CT as affected by growing season rainfall (<300 mm; 300-500 mm; >500 mm)



Error bars represent 95% confidence intervals. Significant difference of the effect size is denoted by *.

The analysis of the comparison between LINiT and CT also showed strong evidence that the effect differ between seasonal rainfall (chi-square 33.1 on 2 df). CT resulted in 2.84 t ha⁻¹ and 1.41 t ha⁻¹ higher grain yield than LINiT when growing season rainfall were 300 – 500 mm and >500 mm (Figure 6.3). There was no evidence of any difference for rainfall <300 mm. Combined over all the studies, yields were higher by 1.80 t ha⁻¹ under CT compared with LINiT.

Figure 6. 3. Weighed mean differences in crop grain yield in LINiT compared with CT as affected by growing season rainfall (<300 mm; 300-500 mm; >500 mm)



Error bars represent 95% confidence intervals. Significant difference of the effect size is denoted by *.

The significant lower yield under non-inversion tillage under growing season rainfall >300 mm may be attributed to poor infiltration and drainage causing waterlogging problems, as reported by Anazodo *et al.* (1991), Rasmussen (1999) and Rusinamhodzi *et al.* (2011). This may have resulted in wetter and cooler soils under non-inversion tillage (Rasmussen *et al.*, 1993; Reicosky *et al.*, 1995; Riley *et al.*, 2005) reducing plant density and growth, and final grain yields compared with CT. Several authors also report reductions on plant establishment and initial crop growth under non-inversion tillage producing lower yields than CT, under conditions of excessive rainfall (Brennan *et al.*, 2014; Forristal & Murphy, 2009; López-Bellido & López-Bellido, 2001).

The low grain yield under non-inversion tillage in high rainfall conditions could also be related to low soil N availability. Tillage induced effects on soil properties (moisture content, temperature, structure and bulk density) can influence soil N status (Lipiec &

Stepniewski, 1995). Increasing tillage intensity can have several effects including reportedly intensifying soil organic matter decomposition and correspondingly increasing soil mineral N (SMN) content. On the other hand, increasing residue cover and reducing soil disturbance can lead to N immobilisation or slow N release under non-inversion tillage systems. This is supported by several authors (Chen *et al.*, 2007; Franzluebbers, 2004; López-Bellido *et al.*, 2013; McConkey *et al.*, 2002; Soon *et al.*, 2001) reporting higher SMN under CT compared with non-inversion tillage. In conclusion, low N mineralisation potential and high N leaching with high rainfall conditions can lead to low SMN level in soils under non-inversion. This could account for the low grain yield under LINiT and HINiT under high and medium rainfall conditions, compared with CT. Several authors (De Vita *et al.*, 2007; Hansen *et al.*, 2011; Wang *et al.*, 2012) reported, like in the present study, higher yield under CT with differences on grain yields between tillage systems mainly due to N availability.

The no significant difference between cultivation techniques on final grain yield when the growing seasonal rainfall was <300 mm is perhaps a result of the greater variability of rainfall during the growing season, with occurrence at times of dry periods. Presence of residues on the soil surface under non-inversion tillage could have reduced soil water evaporation, as demonstrated by Freebairn & Wockner (1983) and Stagnari *et al.* (2014). This effect can potentially maintain or increase soil moisture under non-inversion tillage systems reducing stress conditions for the crop, as reported by Cantero-Martínez *et al.* (2003), Kassam *et al.* (2009) and Šíp *et al.* (2013). Positive reduction of soil moisture loss, at time of water need by the crop, can potentially reduce yield differences between tillage treatments in scarcity of rainfall. This, like other studies (Hussain *et al.*, 1999; Lueschen *et al.*, 1991; Piggin *et al.*, 2015; Rusinamhodzi *et al.*, 2011), was particularly evident in the 2013 season when CT and LINiT resulted in similar grain yields, as discussed in Chapter 4.

6.4. Conclusions

Rainfall distribution and amount affected wheat production in the different soil management regimes explored. Meta-analysis of non-inversion tillage compared with

CT showed that grain yields are significantly lower under HINiT and LINiT when total seasonal rainfall is higher than 300 mm. Slow N mineralisation and immobilisation and possible N leaching result in low N availability under reduced tillage practices, while the presence of residues under these tillage systems could generate cool and wet soils negatively affecting crop establishment and growth. These characteristics resulted in the lower grain yield under non-inversion tillage compared with CT under conditions of excessive rainfall. Meta-analysis indicates a similar performance of the non-inversion tillage practices, however, with CT under dry rainfall patterns (<300mm). The potential of the soil under non-inversion tillage to maintain soil moisture can perhaps boost yields making them comparable to those under CT. This is particularly important as results for the 60-years rainfall data shows evidence of increasing frequency of dry seasons at this particular experimental site. In addition, climate change projections for southwest England weather suggest drier summers by up to 40% by 2080s and reduction in summer soil moisture around 30% by the 2050s (Jenkins *et al.*, 2008, 2009).

Integrating data across studies in a meta-analysis can provide an insight of the treatment effects with more precision compared with a single study (Liberati *et al.*, 2009). The present study can, therefore, provide a better understanding of the relative importance of the rainfall patterns and their interaction with crop yield under contrasting tillage practices. However, the number of data sets used is limited and further analysis including larger data set is needed, in order to confirm the present results.

CHAPTER SEVEN

Economic and energy-use evaluation of spring wheat production under different cultivation techniques, nitrogen fertilisation rates and undersowing

7.1.Introduction

In response to increasing global demand for food, energy-use in agriculture has been intensified to maximise yields; minimise labour intensive practices or both (Esengun *et al.*, 2007). The amount of arable land, mechanisation level and labour are amongst the most important factors influencing energy demand in agriculture (Alam *et al.*, 2005). The use of combined cultivation machines (for soil disturbance, levelling and seeding) has increased (Morris *et al.*, 2010) as the most effective way to save energy and reduce production costs (Hernández *et al.*, 1995). This is possible with reduced tillage operations and work rates potentially decreasing fuel consumption, as reported by Filipovic *et al.* (2006), Hobbs *et al.* (2008) and Koga *et al.* (2003). Non-inversion tillage adoption can potentially reduce the production costs and save operational time compared with conventional tillage (Clement *et al.*, 1995; Epplin *et al.*, 2005; Morris *et al.*, 2010). Saving time and cost makes the adoption of non-inversion tillage increasingly attractive to farmers, as reported by Harman *et al.* (1996) and Jones *et al.* (2006). However, irregular yield under reduced tillage systems is still reportedly a major concern (Kock *et al.*, 2009; Küstermann *et al.*, 2013; Rochecouste *et al.*, 2015). In contrast, and despite conventional tillage often producing high yields (Arvidsson *et al.*, 2013; Brennan *et al.*, 2014), it is also the greatest energy and labour consumer in arable crop production (Epplin *et al.*, 2005). The use of mineral N fertilisation can often result in higher yields, but it also requires high energy inputs and production costs (Dalgaard *et al.*, 2001; Hussain *et al.*, 2010; Refsgaard *et al.*, 1998; Rossner *et al.*, 2014; Sartori *et al.*, 2005). More efficient energy-use is one of the considerations towards more sustainable agriculture production. This could provide financial savings, reduction of fossil resources use and less air pollution (Uhlin 1998).

The aim of this study is to investigate energy input and outputs per hectare, and make cost and economic consideration of the adoption of contrasting cultivation techniques, application of N fertilisation and undersowing legumes on spring wheat production.

7.2. Material and methods

7.2.1. Energy considerations

The energy balance was analysed considering only the energy used in crop production, without considering environmental sources of energy (radiation, wind and water). The only energy output considered was the final marketable grain dry weight yield while wheat straw was left on the field and not considered as an economic output. The operations number and duration, seed rate, pesticide and fertiliser application, and human labour data was collected from field measurements and farm records (Royal Agricultural University's Harnhill Manor Farm records). Direct energy (operational energy) includes human energy and fuel consumption, while indirect energy includes machinery, fertiliser, herbicide and seeds. Using a process analysis (Fluck, 1992), total energy consumption was evaluated by summing direct and indirect energies. In the same way, renewable (human labour and seeds) and non-renewables energies (machinery, fuel, fertiliser and herbicides) were investigated. Specifications of the machinery used are included in Appendix 4. The amount of inputs utilised in all core experiments (human labour, machinery, fertilisers, herbicide, seeds and fuel) were specified in Figure 7.1. The amount of input per hectare was multiplied with the coefficient of energy equivalent obtained from the literature (cited in Table 7.1), in order to obtain the energy equivalents for this study. Figure 7.1 also shows the specific energy input by each agricultural management practice used in all core experiments. Energy-use efficiency, energy productivity, specific energy and net energy gain were calculated, as Demircan *et al.* (2006) and Sartori *et al.* (2005),

$$\text{Energy-use efficiency} = \frac{\text{Output energy (MJ ha}^{-1}\text{)}}{\text{Input energy (MJ ha}^{-1}\text{)}} \quad \text{Specific energy} =$$

$$\frac{\text{Input energy (MJ ha}^{-1}\text{)}}{\text{Grain yield (kg ha}^{-1}\text{)}}$$

$$\text{Energy productivity} = \frac{\text{Grain yield (kg ha}^{-1}\text{)}}{\text{Input energy (MJ ha}^{-1}\text{)}}$$

$$\text{Net energy gain} = \text{Grain yield (kg ha}^{-1}\text{)} - \text{Input energy (MJ ha}^{-1}\text{)}$$

Figure 7. 1. Amount and energy inputs assigned to various agricultural management practices for all core experiments across the cropping seasons




Table 7. 1. Energy equivalent of inputs and outputs

Particulars	Unit	Energy equivalent (MJ unit⁻¹)	References
Inputs			
Human labour	h ha ⁻¹	1.96	Ozkan <i>et al.</i> (2004); Yilmaz <i>et al.</i> (2005); Singh <i>et al.</i> (2002)
Machinery	h ha ⁻¹	62.70	Erdal <i>et al.</i> (2007); Ozkan <i>et al.</i> (2004)
Diesel fuel	l ha ⁻¹	56.31	Yilmaz <i>et al.</i> (2005); Erdal <i>et al.</i> (2007); Singh <i>et al.</i> (2002)
N fertilisers	kg ha ⁻¹	66.14	Alam <i>et al.</i> (2005); Esengun <i>et al.</i> (2007) Yilmaz <i>et al.</i> (2005); Pervanchon <i>et al.</i> (2002)
Herbicide	kg ha ⁻¹	238	Ozkan <i>et al.</i> (2007)
Seed (wheat)	kg	14.7	Ozkan <i>et al.</i> (2004)
Seed (legume)	kg	14.7	Kitani (1999)
Output			
Grain yield	kg	14.7	Ozkan <i>et al.</i> (2004); Pimentel (1980)

7.2.2. Economic analysis

Total production cost, expressed as £ ha⁻¹, was calculated by summing inputs cost (fertiliser, herbicide, spring wheat and legume (BM and WC) seeds), and contractor costs for land preparation and drilling (including fuel, labour and transportation of seeds) and combine harvesting (including carting and filling stage). Contractor costs were obtained from the Royal Agricultural University's Farm records. Fixed costs as labour, machinery power and overheads are included in the contractor costs, therefore, no fixed costs are detailed in the present study. Spring wheat grain price was based on the price quoted from the 2013 Farm Management Pocketbook (Nix, 2012) and adjusted by protein content, using grain yield and protein content raw data from each core experiment. Table 7.2 listed costs considered for all core experiments. Total production value, gross return, net return and benefit:cost ratio were calculated following Canakci *et al.* (2005) and Zangeneh *et al.* (2010).

$$\text{Total production value} = \text{wheat yield (t ha}^{-1}\text{)} \times \text{wheat price (£ t}^{-1}\text{)}$$

$$\text{Gross return} = \text{Total production value (£ ha}^{-1}\text{)} - \text{Total variable cost (£ ha}^{-1}\text{)}$$

$$\text{Benefit:cost ratio} = \frac{\text{Total production value } (\text{£ ha}^{-1})}{\text{Total production cost } (\text{£ ha}^{-1})}$$

Table 7. 2. Seeds cost, contractor costs and grain price considered for all core experiments

Parameter		Cost	Source
Spring wheat	197 kg ha ⁻¹ @ £400 t ⁻¹	£78.8 ha ⁻¹	Nix (2012)
White clover	7 kg ha ⁻¹ @ £9.30	£65.1 ha ⁻¹	Cotswolds Seeds Ltd.
Black medic	8 kg ha ⁻¹ @ £12.00	£96.0 ha ⁻¹	Cotswolds Seeds Ltd.
CT	Kverneland reversible plough + Power harrow combination seed drill	£50 ha ⁻¹ & £45 ha ⁻¹	Royal Agricultural University's Farm records
HINiT	2 passes of ST bar attached Simba X-press + Vaderstadt seed drill	£62 ha ⁻¹ & 31 ha ⁻¹	Royal Agricultural University's Farm records
LINiT	1 pass of ST bar attached Simba X-press + Eco-dyn integrated seed drill	£31 ha ⁻¹ & £34 ha ⁻¹	Royal Agricultural University's Farm records
Fertiliser sprayer	CASE IH SP3000	£12 ha ⁻¹	Royal Agricultural University's Farm records
N70	70 kg N ha ⁻¹ @ £0.74 kg N (£255 t ⁻¹ fertiliser)	£51.8 ha ⁻¹	Royal Agricultural University's Farm records
N140	140 kg N ha ⁻¹ @ £0.74 kg N (£255 t ⁻¹ fertiliser)	£103.6 ha ⁻¹	Royal Agricultural University's Farm records
N210	210 kg N ha ⁻¹ @ £0.74 kg N (£255 t ⁻¹ fertiliser)	£155.4 ha ⁻¹	Royal Agricultural University's Farm records
Herbicide sprayer	CASE IH SP3000	£12 ha ⁻¹	Royal Agricultural University's Farm records
Glyphosate (Roundup) herbicide	2 l ha ⁻¹ @ £6.3 l ⁻¹	£12.6 ha ⁻¹	Royal Agricultural University's Farm records
Combine harvesting		£80 ha ⁻¹	Royal Agricultural University's Farm records
Grain price	grain protein < 13%	£150 t ⁻¹	Nix (2012)
Grain price	grain protein > 13%	£164.4 t ⁻¹	Nix (2012)

Microsoft Excel – 2010 was used to perform general calculations. Energy budget and economic balance were analysed following a split-split plot analysis of variance (ANOVA) model in Genstat (15th Edition VSN International Ltd, Hemel Hempstead, UK). Results are reported as described in Chapter 3, Material and Methods, §3.5.

7.3. Results and discussion

7.3.1. Energy considerations

Input energy in agriculture can be classified as either direct or indirect (Mohtasebi *et al.*, 2008). Direct energy is the energy used directly by the operations, which are mainly human labour and fuel, while indirect energy include fertiliser, herbicide and crop seed inputs. The total energy inputs, including direct, indirect, and renewable and non-renewable energy inputs are summarised in Figure 7.3. Increasing tillage intensity increased the amount of direct and indirect energy used, hence total input energy, as also suggested by Knight (2004). This is due to higher work rate, machinery energy and mainly by higher fuel consumption under CT, compared with HINiT, and by LINiT (Figure 7.2). Non-renewable energy required by CT was higher, therefore, compared with non-inversion tillage. In contrast, renewable energy was relatively similar between tillage systems. Alhajj-Ali *et al.* (2013) also reported higher fuel consumption under CT compared with reduced tillage systems. Reducing tillage intensity and the use of a multi-tooled cultivation approach, therefore, can save field operations - including labour, diesel fuel and machinery energy. Several authors (Khaledian *et al.*, 2014; Rathke *et al.*, 2007; Ziaei *et al.*, 2015) agree that CT operations require higher energy compared with reduce tillage practices, although values differ due to soil type, field conditions and working depths.

Overall, the main source of energy input for spring wheat production was N mineral fertiliser (Figure 7.2). This agrees with several studies (Camargo *et al.*, 2013; Deike *et al.*, 2008; Piringer & Steinber, 2006; Safa *et al.*, 2011) reporting mineral fertilisation as the most important source of energy in conventional wheat production. Alhajj-Ali *et al.* (2013) also reported a linear relationship between increasing energy input and increasing N fertiliser rate. However, under unfertilised conditions diesel fuel was the

main energy source (Figure 7.2), as also reported under organic systems (Vijaya Bhaskar, 2014).

Figure 7. 2. Energy consumption for all core experiments (MJ ha⁻¹)





Figure 7. 3. Direct and indirect input energy, and renewable and non-renewable energy for all core experiments



Manufacturing one weight unit of herbicide active ingredient is energy intensive on principle (Green, 1987). However, in the present study, due to the low rate per hectare of herbicide application used, its contribution to the total energy consumption was small, as also reported by Clement *et al.* (1995). Nevertheless, unlike organic farming (e.g. Vijaya Bhaskar, 2014), the herbicide application added 476 MJ ha⁻¹ of the energy input increasing total energy consumption across all core experiments (Figure 7.2).

In 2013, LINiT had significantly higher output energy and energy gain than HINiT, and was statistically similar to CT (Table 7.3). Similarly, LINiT resulted in significantly higher energy-use efficiency and energy productivity than CT and HINiT. This suggests that in the case of LINiT, 0.546 kg of wheat yield was obtained per unit of energy used (MJ) (energy productivity). Hernandez *et al.* (1995) also reported greater energy productivity when reducing tillage intensity, compared with CT. The specific energy was significantly lower under LINiT than HINiT and statistically similar to CT. Differences among tillage systems resulted as LINiT produced significantly more yield with less total inputs than HINiT, whereas LINiT yielded statistically similar than CT but using less total inputs. This agrees with Zentner *et al.* (2004) reporting that energy efficiency and energy productivity can be increased either by increasing total energy output or by decreasing total energy input, and by both actions at the same time. In contrast, Borin *et al.* (1997) reported that decreasing tillage intensity increases energy efficiency due to lower output energy used.

Table 7. 3. Analysis of energy indices for spring wheat production in 2013

	Yield (kg ha ⁻¹)	Output energy (MJ ha ⁻¹)	Energy-use efficiency (MJ ha ⁻¹)	Energy productivity (kg MJ ⁻¹)	Specific energy (MJ kg ⁻¹)	Net gain (MJ ha ⁻¹)
CT	5576ab	81968ab	6.690a	0.4551a	2.659ab	68054ab
HINiT	5299a	77902a	6.605a	0.4493a	2.843b	63830a
LINiT	6075b	89309b	8.019b	0.5455b	2.413a	75822b
<i>SED (4df)</i>	206.8*	3040.0*	0.6953**	0.01704**	0.0934*	2846.2**
N0	5610a	82473a	12.077c	0.8215c	1.312a	75609a
N70	5498a	80815a	6.934b	0.4717b	2.267b	69129a
N140	5858a	86118a	5.285a	0.3595a	2.921b	69802a
N210	5635a	82833a	4.123a	0.2805a	4.054c	62401a
<i>SED (18df)</i>	435.2 ^{ns}	6397.9 ^{ns}	1.5434***	0.04997***	0.3397***	6544.2 ^{ns}

Values followed by same letter, do not differ significantly at $P < 0.05$. * = $P < 0.05$, ** = $P < 0.01$; *** = $P < 0.001$; and ns = no significant

In 2013, N fertilisation did not significantly affected final grain yield, and thus output energy. However, energy-use efficiency and energy productivity decreased with increasing N fertilisation rates. Unfertilised conditions resulted in 0.822 kg MJ⁻¹ of energy productivity compared to 0.281 kg MJ⁻¹ when 210 kg N ha⁻¹ was applied. This suggests that less total energy input under unfertilised conditions increased energy productivity. Accordingly, the specific energy increased with increasing N rate. Net gain was not, however, significantly affected by N fertilisation. These energy indices resulted as increasing N increased total inputs while, total outputs were similar between N rates, as also reported by Alhajj-Ali *et al.* (2013). In contrast to the 2013 results, Safa *et al.* (2011) reported that the consumption of indirect energy by N application is positively correlated with wheat yield.

Introducing a legume crop into the cropping systems can potentially reduce the demand for mineral fertiliser by the main crop (Lupwayi *et al.*, 2011; Schwenke *et al.*, 2011). The present study, however, revealed that undersown legume did not provide significant advantage to the cereal crop and resulted in similar energy-use efficiency than the no undersown treatment (Table 7.3). Nevertheless, undersown legumes significantly interacted with N fertilisation rates affecting energy-use (Figure 7.4 & 7.5). The additional energy inputs by BM and WC seeds combined to the extra energy added with high N rates resulted in lower energy-use efficiency and productivity under those treatments.

Figure 7. 4. Nitrogen fertilisation and undersowing treatment interaction effect on energy-use efficiency (2013)





Figure 7. 5. Nitrogen fertilisation and undersowing treatment interaction effect on energy productivity (2013)



In the 2014 season, CT resulted in a significantly higher yield, thus higher output energy than non-inversion tillage systems (Table 7.4). Energy-use efficiency, energy

productivity and net gain were higher under CT followed by LINiT, and HINiT. Despite the fact that direct and indirect input energies were lower under HINiT, output energy was significantly lower than CT, reducing its efficiency. This agrees with Borin *et al.* (1997) reporting that decreasing tillage intensity reduces output energy resulting in lower energy efficiency. Non-inversion tillage systems had less fuel consumption, machinery energy and have a higher work rate. Coarser seedbed conditions resulted in poorer plant establishment and plant growth, however, and greater weed competition (see Chapter 5) affecting the energy-use efficiency. This agrees with Küsterman *et al.* (2013) reporting that the benefit of reduced tillage systems over CT is mainly due to lower input energy as outputs are often lower.

Table 7. 4. Analysis of energy indices for spring wheat production in 2014

	Yield (kg ha⁻¹)	Output energy (MJ ha⁻¹)	Energy-use efficiency (MJ ha⁻¹)	Energy productivity (kg MJ⁻¹)	Specific energy (MJ kg⁻¹)	Net gain (MJ ha⁻¹)
CT	5371b	78951b	6.465c	0.4398c	2.60a	65037c
HINiT	2533a	37228a	2.883a	0.1962a	6.62a	23152a
LINiT	3196a	46981a	3.764b	0.2561b	4.43a	33492b
<i>SED (4df)</i>	265.8***	3907.0***	0.246***	0.01672***	1.193 ^{ns}	3700.5***
N0	3097a	45524a	6.548c	0.4455c	2.766a	38652a
N70	3264a	47983a	4.079b	0.2775b	4.413ab	36297a
N140	4182b	61478b	3.759b	0.2557b	4.709ab	45162a
N210	4256b	62562b	3.097a	0.2107a	6.316b	42131a
<i>SED (18df)</i>	290.1***	4264.6***	0.310***	0.0211***	0.953**	4241.3 ^{ns}

Values followed by same letter, do not differ significantly at $P < 0.05$. **= $P < 0.01$; ***= $P < 0.001$; and ns= no significant

The application of 140 and 210 kg N ha⁻¹ in the 2014 season resulted in significantly higher output energy in terms of grain yield, compared with unfertilised condition and low N rate (70 kg N ha⁻¹) (Table 7.4). However, increasing N rates increases total energy inputs as both direct and indirect energy inputs increases. This decreases the energy-use efficiency and energy productivity when raising N rates, although no significant differences were observed between N70 and N140. Sartori *et al.* (2005) also reported that reducing chemical inputs potentially increases energy-use efficiency. The specific energy was significantly higher when 210 kg N ha⁻¹ was applied compared

specifically with unfertilised conditions, as output energy increased with increasing total energy inputs.

In 2014 CT treatment combined with N fertilisation had a significantly lower energy-use efficiency and energy productivity, particularly when compared with unfertilised conditions (Figure 7.6 & 7.7). This resulted from those treatment combinations that produced high output energies but also required high direct and indirect energy inputs.

Figure 7. 6. Cultivation techniques and nitrogen fertilisation treatment interaction effect on energy productivity (2014)





Figure 7. 7. Cultivation techniques and nitrogen fertilisation treatment interaction effect on energy-use efficiency (2014)



7.3.2. Economic analysis

Irrespective of the year studied, no fixed costs were evaluated as machinery power and labour were included in the contractor cost for land preparation and drilling, and fertiliser and herbicide spraying operations. Total production cost, therefore, coincides with total variable cost. Overall, in the present study, production costs were higher comparing with organic farming (e.g. Vijaya Bhaskar, 2014) where mineral fertilisation and herbicide application would be proscribed.

Across experiments, total production cost was lower under LINiT, compared with HINiT, and by CT (Table 7.5 & 7.6), agreeing with Hernandez *et al.* (1995) and Knight (2004). Increasing N fertilisation rates increases production cost, while BM resulted in higher total production cost compared with WC, and compared with the no undersowing conditions, as BM seed price is higher.

In the 2013 spring wheat season, LINiT had a significantly higher gross return, compared with HINiT and CT (Table 7.5). Production value per unit of production cost (benefit:cost ratio) was also significantly higher with LINiT (2.59), than CT (2.15) and

HINiT (2.04). The lower contract cost for land preparation and drilling, and greater yield produced resulted in higher gross margin and also the benefit:cost ratio under LINiT, compared with HINiT and CT. Despite the statistically similar grain yields between CT and LINiT, high production cost under CT resulted in a lower gross margin and benefit:cost ratio. In the same context, low production cost under HINiT has not rewarded with a greater gross return.

Table 7. 5. Economic analysis for spring wheat production in 2013

	Yield (t ha⁻¹)	Total production value (£ ha⁻¹)	Total production cost (£ ha⁻¹)	Gross return (£ ha⁻¹)	Benefit:cost ratio
CT	5.58ab	863.4	409.8	453.6a	2.15a
HINiT	5.29a	814.6	407.8	406.7a	2.04a
LINiT	6.08b	950.7	379.8	570.9b	2.59b
<i>SED (4 df)</i>	0.207*	-	-	33.4*	0.09**
N0	5.61a	846.6	321.43	525.1a	2.64b
N70	5.50a	834.5	373.23	461.3a	2.29a
N140	5.86a	904.0	425.03	478.9a	2.16a
N210	5.63a	919.8	476.83	442.9a	1.95a
<i>SED (18 df)</i>	0.435 ^{ns}	-	-	67.2 ^{ns}	0.17**
BM	5.65a	882.7	441.43	441.3a	2.04a
Nus	5.60a	867.0	345.43	521.6a	2.56b
WC	5.70a	878.9	410.53	468.4a	2.17a
<i>SED (48 df)</i>	0.215 ^{ns}	-	-	33.8 ^{ns}	0.08***

Values followed by same letter, do not differ significantly at $P < 0.05$. * = $P < 0.05$, ** = $P < 0.01$; *** = $P < 0.001$; and ns = no significant

Between N rate treatments in 2013 the unfertilised conditions resulted in a significantly higher benefit:cost compared with any other N rate application. The N fertiliser increased production costs compared with unfertilised conditions. This, in addition to N fertilisation failing to increase 2013 grain yield (Chapter 4), resulted in a significantly lower benefit:cost ratio, compared with the unfertilised conditions (Table 7.5).

Higher production costs when using undersown legume species resulted in significantly lower benefit:cost than no undersowing, as the legume did not offer any advantage to grain yields in the drier weather conditions.

In 2014 the gross return was highly dependent on grain yield value, which was significantly higher under CT than non-inversion tillage treatments (Table 7.6). Despite high production costs relating to CT, increases in yield output resulted in a significantly high gross margin. Although production costs were quite similar between CT and HINiT, the more variable seedbed conditions and greater weed pressure under HINiT negatively affected yield. This reduced production value resulting in a negative gross margin under HINiT. Higher production value and lower production cost under LINiT resulted in a significantly higher benefit:cost ratio (1.28) than HINiT (0.95).

Table 7. 6. Economic analysis for spring wheat production in 2014

	Yield (t ha⁻¹)	Total production value (£ ha⁻¹)	Total production cost (£ ha⁻¹)	Gross return (£ ha⁻¹)	Benefit:cost ratio
CT	5.37b	844.7	409.8	434.9c	2.11c
HINiT	2.53a	388.2	407.8	-19.6a	0.95a
LINiT	3.20a	491.2	379.8	111.4b	1.28b
<i>SED</i>	0.266**	-	-	44.2**	0.114***
N0	3.10a	468.6	321.43	147.1a	1.48a
N70	3.26a	493.0	373.23	119.7a	1.32a
N140	4.18b	655.5	425.03	230.5a	1.56a
N210	4.25b	681.8	476.83	204.9a	1.44a
<i>SED</i>	0.290**	-	-	45.3 ^{ns}	0.109 ^{ns}
BM	3.75a	581.8	441.43	140.3a	1.31a
Nus	3.72a	580.8	345.43	235.4b	1.68b
WC	3.63a	561.5	410.53	151.0a	1.36a
<i>SED</i>	0.207 ^{ns}	-	-	33.1**	0.089***

Values followed by same letter, do not differ significantly at $P < 0.05$. ** = $P < 0.01$; *** = $P < 0.001$; and ns = no significant

N fertilisation, in the 2014 season, significantly increased production value but also total production costs, resulting in non-significant differences on the gross return and benefit:cost ratio between N rates (Table 7.6). This is more evident with the application of 210 kg N ha⁻¹ resulting in statistically higher protein content, compared with any other N rate (see Chapter 5). However, this N rate also increased production costs, resulting in no significant differences on the benefit:cost.

Increases in production costs by including undersown legumes resulted in a lower gross margin and benefit:cost ratio (Table 7.6).

7.4. Conclusion

The present study aimed to identify agricultural management operations where energy and cost savings could be realised. It appears that the energy and economic performances of the different agricultural management practices used in this case were dependent on specific characteristic affecting the final crop yield and grain protein content – such as weather conditions, agreeing with Gomiero *et al.* (2011).

It seems that regardless of high input energy and production costs, CT can potentially be efficient in the energy-use and economically viable. However, in considering low energy consumption and production costs, this study showed that LINiT is the most reliable alternative to CT systems. LINiT exhibited promising results in increasing productivity and economic returns when a resultant high crop yield is delivered.

The application of mineral N is energy consuming and costly. Results of the present study showed that N fertilisation is not always energy-efficient and economically viable, even when it can result in higher final yields. However, the yield and grain quality responses and grain value remain key, with higher crop performance providing greater justification.

For all experiments, undersowing BM and WC was economically less profitable than no undersowing. However, in terms of energy consumption, no differences between undersowing were found for either BM or WC.

In order to assess sustainability of the agricultural management practices used in the present study over the long-term, further investigation of wider economic and energy impacts (such as soil carbon sequestration and greenhouse gas emissions) could be addressed, together with consideration of the wheat straw output inclusion.

CHAPTER EIGHT


General discussion and conclusions

8.1.Introduction

The primary objective of the present study was to investigate the effects of selected cultivation techniques, N fertilisation and undersown legumes on spring wheat growth and development. To explore, in particular, the yield components that contribute to grain yield and quality, as well as weed pressure influences alongside changes in soil mineral N content. This was examined through field experiments on a clay soil using three cultivation techniques - from conventional tillage through high intensity non-inversion tillage to low intensity non-inversion tillage; four mineral N fertilisation rates of 0, 70, 140 and 210 kg N ha⁻¹ and two undersown legume species - black medic and white clover plus no undersowing treatment. Figure 8.1. shows links between all the thesis chapters, and how each study contributes and attempts to assess the most suitable agricultural management practices for increasing yield, and yet be the most efficient in energy-use and profitability.

Chapter 1 and 2 set the framework of the agricultural management practices adopted, and the effects on crop performance and weed prevalence reported in previous studies (Literature review). The methodologies and techniques utilised to establish and evaluate the two experimental trials are developed and described in Chapter 3. Data sets were collected across the two core experiments established in 2013 and 2014. As analysis combining core experiments over time did not show any progress compared to single data sets - core experiments were then analysed separately and presented in Chapter 4 and 5 (Core experiment 1 & 2). Based on results obtained from both core experiments, the effect of the weather conditions on spring wheat performance under contrasting cultivation regimes was evaluated by performing a meta-analysis (Chapter 6). Finally, Chapter 7 focuses on the energy-use and economic productivity of the management practices adopted. Chapter 8 deals with how the whole study is able to provide an improved understanding of the influences of the agricultural practices on crop development, soil mineral N content and weed infestation. This chapter also reaches key conclusions, and considers some implications and suggestions for further work.

Figure 8. 1. Chapters interlinking different agricultural management practices, weather conditions, and energy and economic considerations for spring wheat production



8.2. Cultivation techniques

8.2.1. Wheat performance

Key findings for the contrasting cultivation treatments adopted are summarised in Table 8.1. The use of the plough and power harrow under conventional tillage (CT) created a fine and uniform seedbed favouring plant germination and establishment in all of the seasons studied. This is mainly the result of the plough inverting the soil and incorporating plant residues. The power harrow breaks massive structure of clay soil leaving a fine and level seedbed, assumingly increasing soil-seed contact as previously reported (Atkinson *et al.*, 2007, 2009; Bell, 1996; Comia *et al.*, 1994). Seedbed conditions created by non-inversion tillage systems, such as high intensity non-inversion tillage (HINiT) and low intensity non-inversion tillage (LINiT) reduced crop emergence and final establishment when compared with CT, as also reported by Känkänen *et al.* (2011) and Pietola & Tanni (2003). This is likely to be due to large amount of plant residues on the soil surface and the increased presence of soil clods, leaving a much coarser and variable seedbed, as others have widely reported (Atkinson, 2008; Morris *et al.*, 2010; Känkänen *et al.*, 2011; Rieger *et al.*, 2008). HINiT and LINiT systems can also keep the soil surface wet and cold by reducing soil evaporation which negatively affects crop emergence and early growth, as Morris *et al.* (2010) and Reicosky *et al.* (1995) reported.

Table 8. 1. Key findings for contrasting cultivation techniques for all core experiments

	CT	HINiT	LINiT
Tillage intensity	High	Intermediate	Low
Seedbed	Fine	Coarser	Highly coarser
Seedbed evenness	Level / Uniform	Variable / Not uniform	Highly variable / Not uniform
Plant establishment	High	² High / ¹ Intermediate	Low
Tiller production	High	High	Low
Plant height		Statistically not significant	
Ears number	High	¹ High / ² Intermediate	Low
TGW		Statistically not significant	
Number of grains per ear	² High / ¹ Low	Low	¹ High / ² Intermediate
Grain yield	*Comparable / ² High	Low	¹ High / ² Low
SMN	² High / ¹ Low	¹ Intermediate / ² Low	¹ High / ² Low
Moisture content	¹ Low	¹ Intermediate	¹ High

¹Only observed in 2013 season (Chapter 4); ²Only observed in 2014 season (Chapter 5);

*Statistically comparable with LINiT and HINiT

One of the core objectives of the present study focussed on spring wheat performance and productivity. In all experiments, the performance of different cultivation techniques on spring wheat production was the result of complex interactions between seedbed conditions, moisture status, N availability, weed pressure and variable rainfall conditions (Table 8.1). This emphasises the difficulties of relating final grain yield to a particular yield limiting factor, as highlighted by Gooding & Davies (1997).

Contrary to the finding of several authors (e.g. Blake *et al.*, 2003; Ghaderi *et al.*, 2009), the present study did not always observe positive relationships between crop establishment and final grain yield. Weather conditions, especially rainfall, also exerted considerable influences on grain yield. The contrasting performance of the cultivation techniques treatments on wheat production in each core experiment have been attributed to various causes (Chapter 4, 5 & 6). The present study showed that under conditions of low rainfall, maintaining or increasing soil moisture can considerably positively influence final grain yield (Core experiment I). In the 2013 cropping season the ability to conserve soil moisture and the high resultant SMN content appear to compensate for

poor establishment and the initial slow crop growth under LINiT, finally making crop yield comparable to CT. This is particularly important as drought events in the UK are expected to increase (DEFRA, 2012; Spink *et al.*, 2009), and LINiT may, therefore, potentially reduce crop stress conditions. As in this study, several have reported the benefit of non-inversion tillage under low rainfall conditions (De Vita *et al.*, 2007; López-Bellido *et al.*, 2000; Martinez *et al.*, 2008; Stagnari *et al.*, 2014). Nevertheless, when water availability is not a limiting factor (as in Core experiment II), poor plant establishment, high variability of crop growth, low SMN content and high weed pressure under the non-inversion tillage adversely affected crop performance and resulted in lower crop yield compared with CT. These findings agree with other studies (Arvidsson *et al.*, 2013; Brennan *et al.*, 2014; Franchini *et al.*, 2012; McConkey *et al.*, 2002; Vijaya Bhaskar *et al.*, 2013b). The initial advantages of CT of better plant establishment and crop growth, mostly led to higher final grain yield than non-inversion tillage systems (Core experiment II).

Taking into account such factors as seedbed variability and complexity, and also weed pressure, the present study considers that conventional tillage (CT) has the greatest potential for ensuring a more reliable spring wheat yield performance in a given soil and location. However, considering climate uncertainty with dry seasons, Low Intensity Non-inversion Tillage (LINiT) also shows promising potential to be an optional practice to CT, in providing better soil moisture conditions in dry weather.

8.2.2. Weed pressure

The influences of cultivation techniques on weed growth have been documented by several authors (Froud-Williams *et al.*, 1981; Hakansson, 2003; Menalled *et al.*, 2001). Across the current core experiments, increasing weed pressure was generally observed under non-inversion tillage systems, although this was variable. Table 8.2. summarises the cultivation effects on weed pressure across core experiments. Overwinter assessment in 2013 showed that LINiT potentially allow weeds to grow, as more weed seeds stay in the soil surface after harvest practices, supporting the finding of other studies (Ball, 1992; Tuesca & Puricelli, 2007).

Table 8. 2. Trends in weed growth between cultivation techniques across core experiments

	CT	HINiT	LINiT
2013 overwinter weeds	Low	Low	High
Early total weed	¹ Low	¹ High	¹ Low
Mid-season total weed	Low	High	^{1#} Comparable / ² High
Broadleaf weed species	Low	High	¹ Low / ² Intermediate
Grass weed species	Statistically not significant		
Weed prevalence (from early growth assessments to harvest)	² Low	² High	² High

¹Only observed in 2013 season (Chapter 4); ²Only observed in 2014 season (Chapter 5);

[#]Statistically comparable with CT and HINiT

Even though the broad spectrum herbicide glyphosate controls many weed species (Norsworthy 2008), its application across the entire experimental site did not allow the separate testing of herbicide effect. However, in both cropping seasons, grass weed species were less prevalent than broadleaved weeds. Grass weeds were, therefore, presumably controlled by the pre-cultivation herbicide application, as also reported by Ewald & Aebischer (2000). Total weed biomass across seasons was dominated by broadleaf weed species and highly related to HINiT cultivation. This is contrary to several author reports relating broadleaf weed species prevalence more to CT (Froud-Williams *et al.*, 1983b; Tuesca & Puricelli, 2007; Tuesca *et al.*, 2001). Plant residue cover under HINiT may perhaps have protected weed seedlings from herbicide, as reported by Sadeghi *et al.* (1998), while follow-up soil movements created by HINiT allowed greater broadleaf weed germination, compared specifically with CT.

The present study, like Peigné *et al.* (2007) and Winkler & Chovancová (2014), revealed that the effectiveness of cultivation systems on weed control is also much influenced understandably by weather conditions. At an early growth assessment, under relatively warm and wet conditions (Core experiment II), cultivation techniques relevance in controlling early weed growth was reduced. This is an assumption, however, as soil conditions and weeds proportions were probably compensating, as suggested by Colbach *et al.* (2006). However, under relatively cold and wet conditions (Core experiment I), the non-inversion tillage, particularly HINiT, benefited weed growth. The lower plant residue cover under HINiT possibly allowed drier and warmer

soil conditions. While, with increasing soil disturbance without soil inversion encouraged weed germination, compared with LINiT, agreeing with Teasdale (1993). In contrast, maximum tillage intensity under CT reduced weed pressure, as reported elsewhere (Clements *et al.*, 1996a; Swanton *et al.*, 2000). Furthermore, weed incidence at harvest time varied across cropping seasons. Dry weather conditions (Core experiment I) seem to encourage the natural decay of weeds prior to harvest time, possibly reducing the initial tillage effects on weed growth, as reported elsewhere (Jørnsgård *et al.*, 1996; Mas & Verdú, 2003; Santín-Montoyá *et al.*, 2014). Nevertheless, under relatively wet and warm conditions prior to harvest time (Core experiment II), weed growth appears to have been encouraged. Those conditions revealed that CT is highly effective in controlling weeds, compared with non-inversion tillage systems, agreeing with others (Gruber *et al.*, 2012; Usman *et al.*, 2013) and even without herbicide applications as reported by (Vijaya Bhaskar *et al.*, 2014b).

The core experiments results (Chapter 4 & 5) showed an inverse relationship between high weed prevalence and crop performance, as reported previously by Clements *et al.* (1996a) and Stevenson *et al.* (1997). However, the present study was unable to relate that weeds alone were the only yield limiting factor. Gruber *et al.* (2012) reported that even though a high weed density was observed there was no evidence that weeds alone were restricting main crop yield. However, if the present study focuses on effects of cultivation techniques on weed pressure, then as others report (e.g. Clements *et al.*, 1996a; Froud-William *et al.*, 1983b; Gruber *et al.*, 2012; Hakansson, 2003; Swanton *et al.*, 2000; Tørresen & Skuterud, 2002; Yagioka *et al.*, 2015), CT controls weeds better.

8.2.3. Soil mineral nitrogen

Cultivation effects on SMN have been widely reported (Gruber *et al.*, 2011; McConkey *et al.*, 2002; Silgram & Shepherd, 1999; Soon *et al.*, 2001). In the present study, cultivation treatments effects were variable across and within the seasons. The contrasting effects were the result of complex interactions between tillage intensity, plant N uptake, residual soil mineral N and the prevailing weather conditions (Chapter 4 & 5). Maximum tillage intensity under CT resulted in higher SMN and provided greater

plant N uptake, compared with non-inversion tillage (Core experiment II). This is probably due to increasing tillage intensity increasing organic matter breakdown and enhancing soil mineral N availability, as reported elsewhere (López-Bellido *et al.*, 2013; Myrbeck *et al.*, 2012; Yagioka *et al.*, 2015). In contrast, greater plant residues and relative wet conditions under non-inversion tillage (Core experiment II) may have resulted in immobilisation and/or slow release of N, agreeing with Alvarez *et al.* (1995) and López-Bellido *et al.* (2013).

Under dry season conditions (Core experiment I), however, LINiT resulted in a higher SMN content compared with CT. The drier conditions perhaps limited the soil microbiota reducing N mineralisation, as reported by Jenkinson *et al.* (1987) and Rasmussen *et al.* (1998). Additionally, the lower plant populations under LiNiT probably left greater residual soil N, agreeing with others (e.g. Brennan *et al.*, 2014; Riley, 1998; Thomsen & Sørensen, 2006). These conditions, therefore, resulted in greater SMN content under LINiT in the drier season.

The present study revealed that maximum tillage intensity, under CT, increases N mineralisation and increased the soil mineral N content. However, under dry weather conditions this study suggests that LINiT can potentially allow high SMN content, mainly due to increases in residual N allowing more N availability to the crop.

8.2.4. Energy consumption and economic impact

The present study shows that high energy inputs and production cost can still be efficient and financially worthwhile if the final output of grain yield can compensate for the inputs applied (Chapter 7), agreeing with Borin *et al.* (1997) and Küsterman *et al.* (2013). This was observed under CT when high grain yield improved energy efficiency, productivity, and gross margin - compared with non-inversion tillage systems (2014 cropping season). In contrast, when higher energy output and production value was combined with low energy input and production costs, greater energy productivity and final gross return was obtained under LINiT, compared specifically with HINiT (2013 cropping season), as also reported by Alhajj-Ali *et al.* (2013) and Knight (2004).

In conclusion, CT can potentially be energy efficient and economically viable. However, this study shows that LINiT gave promising results as an alternative practice to CT, in terms of better energy-use and reliance on non-renewable sources even with low rate of herbicide application.

8.3. Nitrogen fertilisation

8.3.1. Wheat performance

Mineral N fertilisation influences on cereal growth and development, and final grain quantity and quality have been widely studied (Cossani *et al.*, 2009; Gooding & Davies, 1997; López-Bellido *et al.*, 1998; Ottesson *et al.*, 2008). The present study confirmed that N fertilisation increases crop height, particularly when comparing unfertilised conditions to the application of 210 kg N ha⁻¹, agreeing with Lloveras *et al.* (2001) and Sourour *et al.* (2014). However, these increases in crop growth were not always translated into greater yield. Several explanations and suggestions have been given for the mineral N fertilisation effects on spring wheat production in each core experiment (Chapter 4 & 5). Table 8.3 summarises key findings for N fertilisation treatments in all core experiments.

Mineral N fertilisation positively affected the production of grains per ear across core experiments, with increasing grain number per ear with up to 140 kg N ha⁻¹. This response with increasing N availability agrees with Alijani *et al.* (2012) and Ferrise *et al.* (2010). Several studies also report a positive correlation between grain number per ear and final grain yield (Halvorson *et al.*, 2001; Wang *et al.*, 2014), although this was not always observed in the present study. The N fertilisation effect on crop yield across core experiments was the result of interactions with prevailing weather conditions, particularly rainfall, and the residual soil N content, as also reported by Corbeels *et al.* (1998) and López-Bellido *et al.* (2012).

Table 8. 3. Key findings for different nitrogen fertilisation treatments for all core experiments

	N0	N70	N140	N210
Tiller production	¹ Low	^{1*} Comparable	¹⁺ Comparable	¹ High
Plant height	Low	Medium	^{1#} Comparable / High	High
Ears number	Statistically not significant			
TGW	High	Medium	Medium	Low
Number of grains per ear	Low	¹ High / ² Low	High	High
Grain yield	² Low	² Low	² High	² High
SMN	Highly low	Low	Medium	High

¹Only observed in 2013 season (Chapter 4); ²Only observed in 2014 season (Chapter 5);

*Statistically comparable with N0 and N140; +Statistically comparable with N70 and N210;

#Statistically comparable with N70 and N140

Increasing rainfall amount, particularly during winter months, can potentially increase soil N leaching and diminish soil N content (Core experiment II), as also reported by ADAS (2014), Halvorson *et al.* (2001) and Lloveras *et al.* (2001). Under these conditions of low SMN levels adding extra mineral N fertiliser, particularly up to 140 kg N ha⁻¹ increased crop grain yield (Core experiment II), as reported elsewhere (Abad *et al.*, 2005; Alijani *et al.*, 2012; Halvorson *et al.*, 2001; Wang *et al.*, 2014). Limited rainfall conditions and the occurrence of drought events across the cropping season (Core experiment I) showed that N fertilisation failed to encourage crop yield gain as higher soil mineral N was accumulated, which was perhaps enough to boost grain yield, as mentioned elsewhere (Abad *et al.*, 2005; Corbeels *et al.*, 1998; Miao *et al.*, 2015; López-Bellido *et al.*, 2000).

Despite the N fertilisation effect on grain yield, the present study revealed that grain protein content increased with up to 210 kg N ha⁻¹. Higher N rate, therefore, is required to increase grain protein rather than to increase grain yield, agreeing with Garrido-Lestache *et al.* (2004) and Godfrey *et al.* (2010). Additionally, in all core experiments, increasing the aboveground N relative to grain yield response reduced the efficiency in use of N, as Huggins *et al.* (2010) also reported.

One of the objectives of the present study was to evaluate spring wheat productivity under different mineral N fertilisation rates. The study confirms that spring wheat

response to N fertilisation may be absent if the accumulation of mineral N in the soil is substantial, as observed during a dry season on this particular soil type and location. However, under low soil N accumulation, the application of up to 140 kg N ha⁻¹ seems to support greater crop need by increasing grain yield.

8.3.2. Weed pressure

Across all core experiments, N fertilisation significantly affected weed growth which is consistent with other studies (Blackshaw *et al.*, 2005; Jørnsgård *et al.*, 1996; Moss *et al.*, 2004; Lal *et al.*, 2014; O’Donovan *et al.*, 1997). Table 8.4 shows trends in N rates effect on weed infestation across the experiments. Overwinter assessment in 2013 showed greater weed occurrence under high N conditions with increasing N availability, as also reported by Bergkvist (2003).

Table 8. 4. Trends in weed growth under different nitrogen fertilisation rates across core experiments

	N0	N70	N140	N210
2013 overwinter weeds	Low	Low	High	High
Early total weed	Statistically not significant			
Mid-season total weed	Low	High	High	High
Broadleaf weed species	Low	High	High	¹ Low / ² High
Grass weed species	Statistically not significant			
Weed prevalence (from early growth assessments to harvest)	² Low	² High	² High	² High

¹Only observed in 2013 season (Chapter 4); ²Only observed in 2014 season (Chapter 5).

In addition to being influenced by N, weed growth and prevalence are also affected by weather conditions influencing agricultural management effects on weed species, as reported by Peters *et al.* (2014). This was observed at early crop growth assessments when weather conditions seem to be more relevant than the N fertilisation effect. Dry weather conditions later in the season (Core experiment I) reduced weed biomass at harvest time, diminishing the N fertilisation effect. However, under wetter weather

conditions, N fertilisation increased weed prevalence compared with unfertilised conditions, as observed in the Core experiment II.

In terms of weed species, the present study showed that N fertilisation caused shifts in weed species. This was observed on the dominant species, *Stellaria media* L. and *Sinapsis arvensis* L., which were advantaged under N-rich conditions, agreeing with others (Maskell *et al.*, 2010; Stevens *et al.*, 2010; Storkey *et al.*, 2012).

If the focus is on N influences on weeds, mineral N fertilisation greatly increased weed growth. Nevertheless, this appears not to have affected final grain yield, as both the main crop and weeds increased their growth with N, as also reported by Jørnsgård *et al.* (1996) and O'Donovan *et al.* (1997).

8.3.3. Soil mineral nitrogen

In all core experiments, differences between N rates were initially marked, but decreased with time as a result of a proportional balance between N supply and crop N consumption, agreeing with Fuentes *et al.* (2003). The present study showed that application of mineral N increases SMN content, as widely reported elsewhere (Angás *et al.*, 2006; Giacomini *et al.*, 2010; Glendining *et al.*, 1996; Liebig *et al.*, 2002; Lu *et al.*, 2010; Zhao *et al.*, 2014).

8.3.4. Energy consumption and economic impact

The present study showed that the application of N fertiliser decreased energy efficiency, as the N application itself is also highly energy consuming, agreeing with Deike *et al.* (2008) and Safa *et al.* (2011). In addition, even when N fertilisation significantly increased the total production value, high production costs resulted in no greater gross margin compared with zero N. The present study, therefore, confirms that despite the final grain yield produced, N mineral fertilisation gives no great efficiency in energy-use and it does not always ensure higher economic returns.

8.4. Legume undersowing

The practice of undersowing legume species has been widely adopted (Fujita *et al.*, 1992; Shafi *et al.*, 2007; Thorsted *et al.*, 2006), due to their ability to fix atmospheric N (Bakht *et al.*, 2009; Kumar & Goh, 2002). The evaluation of overwinter growth in the present study showed that black medic (BM) and white clover (WC) undersown species have a slow growth pattern (Core experiment II), agreeing with Döring *et al.* (2014) and Moss *et al.* (2004).

The present study also used undersown BM and WC in order to evaluate their weed suppression effect and potential to encourage spring wheat production. Dry and warm conditions at the time of broadcasting the undersown legume species appeared to reduce legume establishment in all core experiments. Poor establishment and slow growth of the legume species, however, seem to diminish any undersowing effects on weed control and on the wheat performance (Chapter 4 & 5). If the focus is on cereal-legume bi-cropping, therefore, then this study alone cannot recommend inclusion of BM and WC for greater weed control and to encourage wheat production.

8.5. Interactions

Several authors (e.g. Alijani *et al.*, 2012; López-Bellido *et al.*, 2000) have reported interactions between cultivation techniques and N application on final grain yield. The present study shows, however, that grain yield was not greatly influenced by treatment interaction, despite of some small significant interactions between tillage practices and N on the mid-season wheat biomass. The present study revealed that cultivation techniques and N fertilisation were more important in defining final grain yield than their interactions, also agreeing with Feng *et al.* (2014).

8.6. Concluding remarks

- Conventional tillage can create a more uniform seedbed allowing better plant germination and crop establishment.

- Grain number per ear appears to be highly related to final grain yield of spring wheat under the contrasting cultivation techniques operations examined.
- Rainfall patterns can markedly affect spring wheat production under different cultivation techniques. The ability to save soil moisture under limited rainfall conditions can compensate for limited crop establishment and growth under low intensity non-inversion tillage, and result in similar grain yield to conventional tillage. Under high rainfall (> 300 mm) across the cropping season, non-inversion tillage can be detrimental, and cannot ensure high grain yield of spring wheat.
- Soil mineral N increases under conventional tillage. However, under drier conditions limiting N mineralisation, soil mineral N content is the result of the residual N left by the plants which is indirectly affected by cultivation techniques.
- Non-inversion tillage systems increase weed biomass, particularly broadleaf weed species under high intensity non-inversion tillage, even when herbicide is previously applied. Non-inversion tillage systems perhaps require higher herbicide rates to control weeds but in order to reduce reliance on non-renewable inputs as herbicides, these tillage systems probably need to be complemented with other management practices such as using cover crops and wider rotations.
- Regardless of the energy inputs and production cost, energy efficiency and economic return under contrasting cultivation techniques mostly depend on the final grain yield produced. Conventional tillage can be energy efficient and economically viable if high yield is ensured, despite considerable energy consumption and cost. Non-inversion tillage systems can save energy and production costs.
- Mineral N fertilisation effects on grain yield production are also indirectly affected by weather conditions affecting soil mineral N accumulation. In

conditions of high residual soil mineral N further crop response to N fertilisation can be limited.

- N fertilisation increases grain protein content separate to grain yield responses.
- Application of mineral N fertiliser boosts soil mineral N content, confirming what is already a well known relationship.
- Crop canopy and weed growth are highly advantaged with mineral N fertilisation, although increasing weed growth does not always seem to reduce final grain yield.
- Mineral N fertiliser application is no more efficient in energy-use, and can be no different in total economic bottom-line benefit to unfertilised environments, even when higher final grain yield is produced.
- Undersowing black medic and white clover in dry conditions can have no effect on spring wheat productivity or weed control.

8.7.Implications

- Transition of soil management agricultural practices towards reduced tillage systems is promoted by the European Union Common Agricultural Policy (CAP) and the United Nations Environment Programme (UNEP) to increase food security and profits while enhancing resource-use and sustaining productivity, and as being more resilient to climate and weather variability. The present study findings show that this may be possible for spring wheat on clay soil in dry years. Low Intensity Non-inversion Tillage (LINiT), even though it can result in low crop establishment, can allow the cereal crop to compensate without loss of grain yield. However, in conditions of high rainfall (>300 mm during cropping season), LINiT can disappoint due to low establishment and poor crop growth, and a high loss in yield caused by great variability of seedbed conditions, less soil N and greater weed pressures.

- In connection with the previous implication, the UK Climate Projections (2009) suggests that the UK is likely to undergo hotter and drier summers, and warmer and wetter winters, with increasingly occurrence of extreme weather events such as dry spells, heat waves, heavy rain and flooding (Jenkins *et al.*, 2009). The present study findings, therefore, indicate that reduced tillage systems such as Low Intensity Non-inversion Tillage (LINiT) can potentially be the best option to ensure yield production and adaptability to these climate change scenarios under drier season conditions. This is particularly important as the interest of farmers for adapting agricultural practices to climate change mitigation has been growing in the last decades (Gonzalez-Sanchez *et al.*, 2015; Olesen *et al.*, 2011; Peigné *et al.*, 2015; Rochecouste *et al.*, 2015).
- Several agri-environmental policies focus directly or indirectly on the reduction and greater efficiency of use of N fertilisation (United Nations Framework Convention on Climate Change; National Emissions Ceiling Directive; European Common Agricultural Policy; UK Nitrate Directive; UK Water Framework Directive). This study shows that mineral N fertilisation fails to improve final grain yield under dry seasons while, under conditions of more water availability, yield can be potentially increased. N application could perhaps be saved by foliar applications at a different timing when, and if, weather conditions are disadvantaged - with N fertilisation being more energy consuming and less economic compared to unfertilised practices.

8.8.Future work

- The variability of the responses to agricultural management practices used in the present study was the result of contrasting weather patterns across seasons. As this study was limited to only two cropping seasons, further study would be highly beneficial to a greater understanding of these cultivation techniques, N fertilisation and undersowing legumes effects on spring wheat productivity in the longer term. In addition, it would be useful to also evaluate the effects of the

agricultural management practices adopted under different soil types and weather conditions.

- Crop responses to water stress were mentioned throughout the thesis as a key factor affecting crop yield. How much of an influence crop root system growth and its water uptake within different seedbed conditions needs more detailed examination, and also these influences have on final grain yield also needs to be further considered.
- A further study of cultivation techniques and N fertilisation interaction effect on soil N leaching would also help to better explain N flow throughout the soil. In addition, the use of ^{15}N isotope labelling fertiliser method has been reported elsewhere as the most accurate method to evaluate both the soil N and N fertiliser relative contributions to plant N-uptake (López-Bellido *et al.*, 2012). It could also be beneficial, therefore, to undertake a further study using isotope labelling to clarify the N fertiliser contributions to effective plant uptake.
- This study could not evaluate potential interactions between cultivation techniques and herbicide application on weed control. Further study would be useful to determine how much of an effect these management practices have on weed infestation. Besides continuous changes in agricultural practices modifies weed community dynamics (Santín-Montanyá *et al.*, 2014). It would be useful to also evaluate changing farm management practices influences on weed species diversity for the longer term.
- During this study, it was not possible to more fully explore undersown legume effects on weed control and wheat performance. A further study designed specifically to evaluate broadcasting and drilling methods for establishing a greater range of undersowing species with spring wheat could be beneficial.
- In a wider future study, the measurement of soil carbon sequestration, losses of N and greenhouse gas emissions could provide further indices of sustainability of field management practices explored in the present study.

References

- Abad, A., Michelena, A. & Lloveras, J. (2005). Effect of nitrogen supply on wheat and soil nitrate. *Agronomy for Sustainable Development*. 25 (4), 439-446.
- Acharya, C.L. & Sharma, P.D. (1994). Tillage and mulch effects on soil physical environment, root growth, nutrient uptake and yield of maize and wheat on Alfisols in North-west India. *Soil and Tillage Research*. 32, 291-302.
- Agricultural Development Advisory Service. (2014). *Arable Crop Report, February 2014*. ADAS. Cambridge, UK.
- Ahn, P.M. & Hintze, B. (1990). No tillage, minimum tillage, and their influence on soil properties. In *Organic-matter Management and Tillage in Humid and Sub-humid Africa*. 341-349. IBSRAM Proceedings. 10. IBSRAM. Bangkok.
- Aikins, S.H.M. & Afuakwa, J.J. (2012). Effect of four different tillage practices on soil physical properties under cowpea. *Agriculture and Biology Journal of North America*. 3(1), 17-24.
- Akbarnia A., Alimardani R. & Baharloeyan, Sh. (2010). Performance comparison of three tillage systems in wheat farms. *Australian Journal of Crop Science*. 4(8):586-589
- Alakukku, L., Ristolainen, A. & Salo, T. (2009). Grain yield and nutrient balance of spring cereals in different tillage systems. in *18th Triennial Conference Sustainable Agriculture*, ISTRO. June 15–19, 2009. Izmir, Turkey.
- Alam, M.S., Alam, M.R. & Islam, K.K. (2005). Energy Flow in Agriculture: Bangladesh. *American Journal of Environmental Sciences*. 1 (3), 213–220.
- Alam, S.M. (1990). Effect of wheat straw extract on the germination and straw growth of wheat (cv. Pavon). *Wheat Information Service*. 71, 16-18.
- Alexandratos, N. & Bruinsma, J. (2012). World agriculture towards 2030/2050: the 2012 revision. Rome, FAO.
- Alhajj-Ali, S., Tedone, L. & De Mastro, G. (2013). A comparison of the energy consumption of rainfed durum wheat under different management scenarios in southern Italy. *Energy*. 61, 308-318.
- Alijani, K., Bahrani, M.J. & Kazemeini, S.A. (2012). Short-term responses of soil and wheat yield to tillage, corn residues management and nitrogen fertilization. *Soil and Tillage Research*. 124, 78-82.

- Al-Khasi, M.M., Yin, X. & Licht, M.A. (2005). Soil carbon and nitrogen changes as affected by tillage systems and crop biomass in a corn-soybean rotation. *Applied Soil Ecology*. 30, 174-191.
- Allmaras, R.R., Rickman, R.W., Ekin, L.G. & Kimball, B.A. (1977). Chiselling influences on soil hydraulic properties. *Soil Science Society of America Journal*. 41, 796-803.
- Alvarez, R., Diaz, R., Barbero, N., Santanatoglia, O.J. & Blotta, L. (1995). Soil organic carbon, microbial biomass, and CO₂-C production from three tillage system. *Soil and Tillage Research*. 33, 17-28.
- Alvarez, R. & Steinbach, H.S. (2009). A review of the effects of tillage systems on some soil physical properties, water content, nitrate availability and crops yield in the Argentine Pampas. *Soil and Tillage Research*. 104, 1–15.
- Amossé, C., Jeuffroy, M-H. & David C. (2013). Relay intercropping of legumes cover crops in organic winter wheat: Effects on performance and resource availability. *Field Crop Research*. 145, 78-87.
- Ampong-Nyarko, K. & De Datta, S.K. (1993). Effects of nitrogen application on growth, nitrogen use efficiency and rice-weed interaction. *Weed Research*. 33, 269-276.
- Anazodo, U.G.N., Onwualu, A.P. & Watts, K.C. (1991). Evaluation of alternative tillage systems in the absence of herbicides for maize production in a savannah loamy sand. *Journal of Agricultural Engineering Research*. 49, 259-272.
- Andersen, M.K., Hauggaard-Nielsen, H., Ambus P. & Jensen, E.S. (2004). Biomass production, symbiotic nitrogen fixation and inorganic N use in dual and tri-component annual intercrops. *Plant and Soil*. 266, 73-287.
- Andersen, M.K., Hauggaard-Nielsen, H., Weiner, J. & Jensen, E.S. (2007). Competitive dynamics in two-and three-component intercrops. *Journal of Applied Ecology*. 44, 545-551.
- Andreasen, C. & Stryhn, H. (2012). Increasing weed flora in Danish beet, pea and winter barley fields. *Crop Protection*. 36, 11-17.
- Angas, P., Lampurlanás, J. & Cantero-Martínez, C. (2006). Tillage and N fertilization effects on N dynamics and Barley yield under semiarid Mediterranean conditions. *Soil and Tillage Research*. 87, 59-71.
- Anil, L., J. Park, R.H. Phipps, & F.A. Miller. (1998). Temperate intercropping of cereals for forage: A review of the potential for growth and utilization with particular reference to the UK. *Grass Forage Science*. 53 (4), 301–317.

- Anken, T., Weisskopf, P., Zihlmann, U., Forrer, H., Jansa, J. & Perhacov, K. (2004). Long-term tillage system effects under moist cool conditions in Switzerland. *Soil and Tillage Research*. 78, 171-183.
- Arshad, M.A., Gill, K.S. & Coy, G.R. (2003). Wheat yield and weed population as influenced by three tillage systems on a clay soil in temperate continental climate. *Soil and Tillage Research*. 28 (3-4), 227-238.
- Arvidsson, J. (2010). Energy use efficiency in different tillage systems for winter wheat on clay and silt loam in Sweden. *European Journal of Agronomy*. 33, 250-256.
- Arvidsson, J., Westlin, A. & Sörensson, F. (2013). Working depth in non-inversion tillage-Effects on soil physical properties and crop yield in Swedish field experiments. *Soil and Tillage Research*. 126, 259-266.
- Arvidsson, J., Etana, A. & Rydberg, T. (2014). Crop yield in Swedish experiments with shallow tillage and no-tillage 1983-2012. *European Journal of Agronomy*. 52, 307-315.
- ASAE. (1994). ASAE Standards engineering practices data (soil cone penetrometer, S313.2), St. Joseph. American Society of Agriculture Engineers (ASAE).
- Atkinson, B.S. (2008). Identification of optimum seedbed preparation and establishment using soil structural visualisation. PhD thesis. University of Nottingham.
- Atkinson, B.S., Sparkes, D.L. & Mooney, S.J. (2009). The impact of soil structure on the establishment of winter wheat (*Triticum aestivum*). *European Journal of Agronomy*. 30, 243-257.
- Atkinson, B.S., Sparkes, D.L. & Mooney, S.J. (2007). Using selected soil physical properties of seedbeds to predict establishment. *Soil and Tillage Research*. 97 (2), 218-228.
- Atwell, B.J. (1993). Response of roots to mechanical impedance. *Environmental and Experimental Botany*. 33, 27-40.
- Ayoub, M., Guertin, S., Lussier, S. & Smith, D.L. (1993). Timing and level of nitrogen fertility effects on spring wheat yield in Eastern Canada. *Crop Science*. 34 (3), 748-756.
- Aziz, I., Mahmood, T., Raut, Y., Lewis, W., Islam, R. & Weil, R.R. (2009). Active organic matter as a simple measure of field soil quality. In *ASA International Meetings*, Pittsburg, PA. USA.
- Badaruddin, M. & Meyer, D.W. (1990). Green-manure legume effects on soil nitrogen, grain yield and nitrogen nutrition of wheat. *Crop Science*. 30, 819-825.

- Bahrani, M.J., Raufat, M.H. & Ghadiri, H. (2007). Influence of wheat residues management on irrigated corn grain production in a reduced tillage system. *Soil and Tillage Research*. 94, 305-309.
- Bailey, K.L. & Duczek, L.J. (1996). Managing cereal diseases under reduced tillage. *Canadian Journal of Plant Pathology*. 18, 159-167.
- Bakht, J., Shafi, M., Jan, M.T. & Shah, Z. (2009). Influence of crop residue management, cropping system and N fertilizer on soil N and C dynamics and sustainable wheat (*Triticum aestivum* L.) production. *Soil and Tillage Research*. 104, 233-240.
- Ball, B.C., Dobbie, K.E., Parker, J.P. & Smith, K.A. 1997. The influence of gas transport and porosity on methane oxidation in soils. *Journal of Geophysical Research*, 102, 23301–23308.
- Ball, D.A. (1992). Weed seed bank response to tillage, herbicide and crop rotation sequence. *Weed Science*. 40, 654-659.
- Ball, D.A. & Miller, S.D. (1993). Influence of cropping history and reduced tillage practices on weed flora in corn (*Zea mays* L.). *Agronomy Journal*. 85 (4), 817-821.
- Banik, P., Midya, A., Sarkar, B.K. & Ghose, S.S. (2006). Wheat and chickpea intercropping systems in an additive series experiments: Advantages and weed smothering. *European Journal of Agronomy*. 24, 325-332.
- Barraclough, P.B., Howarth, J.R., Jones, J., López-Bellido, R., Parmar, S., Sheperd, C.E. & Hawkesford, M.J. (2010). Nitrogen efficiency of wheat: Genotypic and nevironmental variation and prospects for improvement. *European Journal of Agronomy*. 33, 1-11.
- Barraclough, P.B., Kuhlmann, H. & Weir A.H. (1989). The effects of prolonged drought and nitrogen-fertilizer on root and shoot growth and water uptake by winter-wheat. *Journal of Agronomy and Crop Science*. 163 (5), 352-360.
- Barraclough, P.B., López-Bellido, R. & Hawkesford, M.J. (2014). Genotypic variation in the uptake, partitioning and remobilisation of nitrogen during grain-filling in wheat. *Field Crops Research*. 156, 242-248.
- Basch, G. (2005). Conservation agriculture experiences in the European Union. In *Proceedings of Congreso Internacional sobre Agricultura de Conservación*. 9-11 November 2005. Cordoba, Spain.
- Bassett, L.M., Allan, R.E. & Rubenthaler, G.L. (1989). Genotype x environment interactions on soft white winter wheat quality. *Agronomy Journal*. 81, 955–960.

- Baumhardt, R.L. & Jones, O.R. (2002). Residue management and tillage effects on soil-water storage and grain yield of dryland wheat and sorghum for a clay loam in Texas. *Soil and Tillage Research*. 68, 71-82
- Bélanger, G., & F. Gastal. (2000). Nitrogen utilization by forage grasses. *Canadian Journal of Plant Science*. 80, 11–20.
- Bell, B. (1996). Farm machinery. 4th Edition. Farming Press Books, Ipswich.
- Bengough, A.G., Bransby, M.F., Hans, J., McKenna, S.J., Roberts, T.J. & Valentine, T.A. (2006). Root responses to soil physical conditions; growth dynamics from field to cell. *Journal of Experimental Botany*. 57, 437-447.
- Bergkvist, G. (2003). Effect of white clover and nitrogen availability on the grain yield of winter wheat in a three-season intercropping system. *Acta Agriculturae Scandinavica, Section B – Soil & Plant Science*. 53 (3), 97–109.
- Berry, P.M., Sylvester-Bradley, R., Philipps, L., Hatch, D.J., Cuttle, S.P., Rayns, F.W. & Gosling, P. (2002). Is the productivity of organic farms restricted by the supply of available nitrogen? *Soil Use and Management*. 18, 248-255.
- Bescansa, P., Imaz, M.J., Virto, I., Enrique, A. & Hoogmoed, W.B. (2006). Soil water retention as affected by tillage and residue management in semiarid Spain. *Soil and Tillage Research*. 87, 19-27.
- Beyaert, R.P., Schott, J.W. & White, P.H. (2002). Tillage effects on corn production in a coarse-textured soil in Southern Ontario. *Agronomy Journal*. 94, 767-774.
- Bhattachary, D. & Hunter, A. (2012). Global food security programme – Exploring public views. Global Food Security (GFS) Programme, TNS- BMRB.
- Blackshaw, R.E., Brandt, R.N., Janzen, H.H., Entz, T., Grant, C.A. & Derksen, D.A. (2003). Differential response of weed species to added nitrogen. *Weed Science*. 51 (4), 532-539.
- Blackshaw, R.E., Molnar, L.J. & Janzen, H.H. (2004). Nitrogen fertilizer timing and application method affect weed growth and competition with spring wheat. *Weed Science*. 52 (4), 614-622.
- Blackshaw, R.E., Molnar, L.J. & Larney, F.J. (2005). Fertilizer, manure and compost effect on weed growth and competition with winter wheat in western Canada. *Crop Protection*. 24 (11), 971-980.
- Blackshaw, R.E., Moyer, J.R., Doram, R.C. & Boswell, A.L. (2001). Yellow sweetclover, green manure, and its residues effectively suppress weeds during fallow. *Weed Science*. 49, 406-413.

- Blackman, J.A., Payne, P.I. (1987). Grain quality in wheat breeding. In: Lupton, F.G.H. (ed.). *Grain quality in wheat breeding - It's scientific basis*.
- Blake, L., Goulding, K.W.T., Mott, C.J.B. & Johnston, A.E. (1999). Changes in soil chemistry accompanying acidification over more than 100 years under woodland and grass at Rothamsted Experimental Station, UK. *European Journal of Soil Science*. 50, 401–412.
- Blake, J.J., Spink, J.H. & Dyer, C. (2003). Factors affecting cereal establishment and its prediction. *HGCA Research Review*. 51. Home-Grown Cereal Authority (HGCA).
- Blaustein, R.J. (2008). The Green Revolution arrives in Africa. *BioScience*. 58 (1), 8-14.
- Blevins, D.D. (1989). An overview of nitrogen metabolism in higher plants. In: Poulton, J.E., Romeo, J.T. & Conn, E.C. (eds.). *Plant Nitrogen Metabolism*. Plenum, New York, pp 1-41.
- Blevins, R.L. & Frye, W.W. (1993). Conservation tillage: an ecological approach to soil management. *Advances in Agronomy*. 51, 33–78.
- Blevins, R.L., Thomas, G.W. & Cornelius, P.L. (1977). Influence of no-tillage and nitrogen fertilization on certain soil properties after 5 years of continuous corn. *Agronomy Journal*. 69, 383-386.
- Blevins, R.L., Thomas, G.W., Smith M.S., Frye W.W. & Cornelius P.L. (1983). Changes in soil properties after 10 years continuous non-tilled and conventionally tilled corn. *Soil and Tillage Research*. 3, 135-146.
- Blumenthal, C.S., Batey, I.L., Bekes, F., Wrigley, C.W. & Barlow, E.W.R. (1991). Seasonal changes in wheat grain quality associated with high temperatures during grain filling. *Australian Journal of Agricultural Research*. 42, 21–30.
- Bock, B.R. (1984). Efficient use of nitrogen in cropping systems. in Hauck, R.D. (ed.) *Nitrogen in crop production*. ASA, CSSA, and SSSA, Madison, WI.
- Bockus, W.W. & Shroyer, J.P. (1998). The impact of reduced tillage on soil borne plant pathogens. *Annual Review of Phytopathology*. 36, 485-500.
- Bond, W., Davies, G. & Turner, R. (2007). The biology and non-chemical control of Common chickweed (*Stellaria media* L.). The Organic Organisation, UK. [Online] Available from <http://www.gardenorganic.org.uk/sites/www.gardenorganic.org.uk/files/organic-weeds/stellaria%20media.pdf> Accessed (06-04-2013)

- Bonfil, D.J., Mufradi, I., Klitman, S. & Asido, S. (1999). Wheat grain yield and soil profile water distribution in a no-till arid environment. *Agronomy Journal*. 91, 368–373.
- Boomsma, C.R., Santini, J.B., West T.D., Brewer, J.C. McIntyre, L.M. & Vyn, T.J. (2010). Maize grain yield response to plant height variability resulting from crop rotation and tillage system in a long-term experiment. *Soil and Tillage Research*. 106, 227-240.
- Borghi, B. (2000). Nitrogen as determinant of wheat growth and yield. In: Satorre, E.H. & Slafer, G.A. (eds). *Wheat ecology and physiology of yield determination*. Food Products Press, New York. 67-84.
- Borghi, B., Corbellini, M., Minoia, C., Palombo, M., Di Fonzo, N. & Perenzin, M. (1997). Effects of Mediterranean climate on wheat bread-making quality. *European Journal of Agronomy*. 6, 145–154
- Borin, M., Merini, C., Sartori, L. (1997). Effect of tillage systems on energy and carbon balance in north-eastern Italy. *Soil and Tillage Research*. 40 (3), 209-226.
- Borlaugh, N.E. (1971). Mankind and civilization at another crossroad. McDougall Memorial Lecture. 1971. In *Conference Food and Agriculture Organization of the United Nations*. FAO.
- Børresen, T. & Njos, A. (1990). The effects of three tillage systems combined with different compaction and mulching treatments on soil temperature and soil thermal properties. *Norwegian Journal of Agricultural Sciences*. 4, 363–371.
- Bouyoucos, G.J. (1962). Hydrometer method improved for making particle size analysis of soils. *Agronomy Journal*. 54, 464-465.
- Bradford, J.M. (1986). Penetrability. In Klute, A. (ed.). *Methods of soil analysis, Part I. Physical and mineralogical methods – Agronomy Monograph*. 9. 2nd Edition. ASA-SSSA, Madison, USA. 463-478.
- Brady, N.C. & Weil, R.R. (1999). The nature and properties of soils. 12th edition. Prentice Hall Inc., Jersey.
- Braim, M.A., Chaney, K. & Hodgson, D.R. (1992). Effects of simplified cultivation on the growth and yield of spring barley on a sandy loam soil. 2. Soil physical properties and root growth; root:shoot relationship, inflow rates of nitrogen; water use. *Soil and Tillage Research*. 22, 173-187.
- Brassington, T. (1986). *Ploughs and plough attachments – their operation and use*. ADAS Mechanisation Occasional Note. 335.

- Braunack, M.V. & Dexter, A.R. (1989). Soil aggregation in the seedbed – a review. 1. Properties of aggregates and beds of aggregates. *Soil and Tillage Research*. 14, 259-279.
- Breland, T.A. (1996). Green manuring with clover rye grass catch crops undersown in small grains: crop development and yields. *Soil and Plant Science*. 46, 30-40.
- Brennan, E.B. & Smith, R.E. (2005). Winter cover crop growth and weed suppression on the central cost of California. *Weed Technology*. 19, 1017-1024.
- Brennan, J., Hackett, R., McCabe, T., Grant, J., Fortune, R.A & Forristal, P.D. (2014). The effect of tillage system and residue management on grain yield and nitrogen use efficiency in winter wheat in a cool Atlantic climate. *European Journal of Agronomy*. 54, 61-69.
- Bristow, K.L. & Horton, R. (1996). Modelling the impact of partial surface mulch on soil heat and water flow. *Theoretical and Applied Climatology*. 54, 85-98.
- Brooks, R.J., Semenov, M.A., Jamieson, P.D. (2001). Simplifying Sirius: sensitivity analysis and development of a meta-model for wheat yield prediction. *European Journal of Agronomy*. 14, 43-60.
- Brown, L.R. (2012). *Full planet, empty plates. The new geopolitics of food scarcity*. Earth Policy Institute. New York, USA.
- Buhler, D.D. (1995). Influence of tillage systems on weed population dynamics and management in corn and soybean in the central USA. *Crop Science*. 35, 1247-1258.
- Buhler, D.D., Stoltenberg, D.E., Becker, R.L. & Gunsolus, J.L. (1994). Perennial weed population after 14 years of variable tillage and cropping practices. *Weed Science*. 42, 205-209.
- Burger, M. & Jackson, L.E. (2003) Microbial immobilisation of ammonium and nitrate in relation to ammonification and nitrification rates in organic and conventional cropping systems. *Soil Biology and Biochemistry*, 35, 29-36.
- Cabrera-Bosquet, L., Albrizio, R., Araus, J.L. & Nogués, S. (2009). Photosynthetic capacity of field-grown durum wheat under different N availabilities: A comparative study from leaf to canopy. *Environmental and Experimental Botany*. 67 (1), 145-152.
- Calado, J.M.G., Basch, G. & Carvalho, M. (2010). Weed management in no-till winter wheat (*Triticum aestivum* L.). *Crop Protection*. 29 (1), 1-6.
- Camargo, G.G.T., Ryan, M.R. & Richard T.L. (2013). Energy use and greenhouse gas emissions from crop production using the farm energy analysis tool. *BioScience*. 63 (4), 263-273.

- Campbell, C.A., Davidson, H.R. & Warder, F.G. (1977). Effects of fertilizer N and soil moisture on yield, yield components, protein content and N accumulation in the aboveground parts of spring wheat. *Canadian Journal of Soil Science*. 57, 311-327.
- Campbell, D.J. & Henshall, J.K. (1991). Bulk density. In Smith, K.A. & Mullins, C.E. (eds.). *Soil analysis. Physical methods*. Marcel Dekker, New York, USA. 329-366.
- Campbell, D.J. & O'Sullivan, M.F. (1991). The cone Penetrometer in relation to trafficability, compaction and tillage. In Smith, K.A. & Mullins, C.E. (eds.). *Soil analysis. Physical methods*. Marcel Dekker, New York, USA. 329-366.
- Canakci, M., Topakci, M., Akinci, I. & Ozmerzi, A. (2005). Energy use pattern of some field crops and vegetable production: case study for Antalya region, Turkey. *Energy Conversion and Management*. 46 (4), 655-666.
- Cannell, R.Q. (1985). Reduced tillage in North-west Europe – a review. *Soil and Tillage Research*. 5, 129-177.
- Cannell, R.Q. & Hawes, J.D. (1994). Trends in tillage practices in relation to sustainable crop production with special reference to temperate climates. *Soil and Tillage Research*. 30, 245-282.
- Cantero-Martínez, C., Angas, P. & Lampurlanes, J. (2003). Growth, yield and water productivity of barley (*Hordeum vulgare* L.) affected by tillage and N fertilization in Mediterranean semiarid, rainfed conditions of Spain. *Field Crops Research*. 84, 341–357.
- Cardina, J., Herms, C.P. & Doohan, D.J. (2002). Crop rotation and tillage system effects on weed seedbanks. *Weed Science*. 50, 448–460.
- Cardina, J., Regnier, E. & Harrison, K. (1991). Long-term tillage effects on seedbanks in 3 Ohio soils. *Weed Science*. 39, 186-194.
- Carof, M., De ourdonnet, S., Saulas, P., Le Floch, D. & Roger-Estrade, J. (2007). Undersowing wheat with different living mulches in a no-till system II. Competition for light and nitrogen. *Agronomy for Sustainable Development*. 27, 357-365.
- Carr, P.M., Horsley, R.D. & Martin, G.B. (2006). Impact of tillage and crop rotation on grain yield of spring wheat I. Tillage effect. *Crop Management*. 5 (1).
- Carr, P.M., Horsley, R.D. & Poland W.W. (2003). Tillage and seeding rate effects on wheat cultivars: I. Grain production. *Crop Science*. 43, 202-209.
- Carr, P.M., Horsley, R.D. & Poland, W.W. (2004). Barley, oat and cereal-pea mixtures as dryland forages in the Northern Great Plains. *Agronomy Journal*. 96, 677–684.

- Carter, L.M., Stockton, J.R., Tavernetti, J.R. & Colwick, R.F. (1965). Precision tillage for cotton production. *Transactions of the American Society of Agricultural Engineers*. (Trans ASAE). 2, 177-179.
- Carter, M.R. (1991). Evaluation of shallow tillage for spring cereals on a fine sandy loam. 1. Growth and yield components, N accumulation and tillage economics. *Soil and Tillage Research*. 21, 23-35.
- Carter, M.R. (1988). Temporal variability of soil macroporosity in a fine sandy loam under mouldboard ploughing and direct drilling. *Soil and Tillage Research*. 12 (1), 37-51.
- Carter, M.R. & Ball, B.C. (1993). Soil porosity. In Carter, M.R. (ed.). *Soil sampling methods of soil analysis*. Lewis Publishers. 581-588.
- Carter, A., Jordan, V. & Stride, C. (2003). *A guide to managing crop establishment*. Soil management Initiative, Chester.
- Casa, R. & Castrignanò, A. (2008). Analysis of spatial relationships between soil and crop variables in a durum wheat field using a multivariate geostatistical approach. *European Journal of Agronomy*. 28 (3), 331-342.
- Cassel, D.K. (1982). Tillage effects on soil bulk density and mechanical impedance. In *Predicting tillage effects on soil physical properties and processes*. ASA-SSSA, Madison, WI, USA: 45-67.
- Cassel, D.K., Raczkowski, C.W. & Denton, H.P. (1995). Tillage effects on corn production and soil physical properties. *Soil Science Society of America Journal*. 59, 1436-1443.
- Celik, I., Barut, Z.B., Ortas, I., Gok, M., Demirbas, A., Tulun, Y. & Akpinar, C. (2011). Impacts of different tillage practices on some soil microbiological properties and crop yield under semi-arid Mediterranean conditions. *International Journal of Plant Production*. 5 (3), 237-254.
- Chamen, W.C.T. & Parkin, A.B. (1995). The impact of tillage practices on the soil and its linked environment, with particular reference to arable cropping in the United Kingdom. *ADAS-SRI Joint Contract Report No. CR/678/95/0209*.
- Chan, K.Y., Heenana, D.P. & Oayes A. (2002). Soil carbon fraction and relationship to quality under different tillage and stubble management. *Soil Tillage Research*. 63 (3-4), 133-139.

Charles, A.H. (1958). The effect of undersowing on the cereal cover crop. *Field Crop Abstracts*. 11, 233-239.

Chen, Ch., Wang, E. & Yu, Q. (2010). Modelling the effects of climate variability and water management on crop water productivity and water balance in the North China Plain. *Agricultural Water Management*. 97 (8), 1175-1184.

Chen, S.Y., Zhang, X.Y., Pei, D., Sun, H.Y. & Chen, S.L. (2007). Effects of straw mulching on soil temperature, evaporation and yield of winter wheat: field experiments on the North China Plain. *Annals of Applied Biology*. 150, 261-268.

Christensen, J.H. & Christensen, O.B. (2007). A summary of PRUDENCE model projections of changes in European climate by the end of this century. *Climate Change*. 81, 7-30.

Christensen, N.W. & Meints, V.W. (1982). Evaluating N fertilizer sources and timing for winter wheat. *Agronomy Journal*. 74, 840-844.

Christian, D.G. (1994). Experience with direct drilling and reduced cultivation in England. In Tebrugge, F. (ed.). *EC Workshop: Experience with the Applicability of No-tillage Crop Production in the West European Countries*. Giessen, Langons Wissenschaftlicher Fachverlag, June 27-28. 25-31.

Clapham, A.R., Tutin, T.G., Moore, D.M. (1987). *Flora of the British Isles*. 3rd edition. Cambridge, UK: Cambridge University Press.

Clark, C.J., Poulsen, J.R., Levey, D.J. & Osenberg, C.W. (2007). Are plant populations seed limited? A critique and meta-analysis of seed addition experiments. *American Naturalist*. 170, 128-142.

Clark, J.M., Campbell, C.A., Cutforth, H.W., DePauw, R.M. & Wilkleman, G.E. (1990). Nitrogen and phosphorus uptake, translocation, and utilization efficiency of wheat in relation to environment and cultivar yield and protein levels. *Canadian Journal of Plant Science*. 70, 965-977.

Clark, M.S., Horwath, W.R., Shennan, C., Scow, K.M., Lantni, W.T. & Ferris, H. (1999). Nitrogen, weeds and water as yield-limiting factors in conventional, low-input and organic tomato systems. *Agriculture, Ecosystems and Environment*. 73, 257-270.

Clements, C.R., & Williams, T.E. (1967). Leys and organic matter II. The accumulation of nitrogen in soils under different leys. *Journal of Agricultural Science*. 69, 133-138.

- Clements, D.R., Benoit, D.L., Murphy, S.D. & Swanton, C.J. (1996a). Tillage effects on weed seed return and seed bank composition. *Weed Science*. 44, 314-322
- Clements, D.R., Weise, S.F., Brown, R., Stonehouse, D.P., Hume, D.J. & Swanton, C.J. (1995). Energy analysis of tillage and herbicide inputs in alternative weed management systems. *Agriculture, Ecosystems and Environment*. 52 (2-3), 119–128.
- Clements, D.R., Weise, S.F. & Swanton, C.J. (1994a). Integrated weed management and weed-species diversity. *Weed Science*. 42, 205-209
- Clements, D.R., Weise, S.F. & Swanton, C.J. (1996b). Integrated weed management and weed species diversity. *Phytoprotection*. 75 (1), 1-18.
- Clements, R.O. & Donaldson, G. (1997). Clover and cereal: low input bi-cropping. *Farming and Conservation*. 3 (4), 12-14.
- Clements, R.O., Kendall, D.A., Asteraki, E.J. & George, S. (1994b). Progress in development of a cereal-clover bi-cropping system. *British Grassland Society 4th research conference*, pp. 77-78.
- Colbach, N., Busset, H., Yamada, O., Dürr, C. & Caneill, J. (2006). ALOMYSYS: Modelling black-grass (*Alopecurus myosuroides* Huds.) germination and emergence, in interaction with seed characteristics, tillage and soil climate: II. Evaluation. *European Journal of Agronomy*. 24 (2), 113-128.
- Comia, R.A., Stenberg, M., Nelsom, P., Rydberg, T. & Hakansson, I. (1994). Soil and crop response to different tillage systems. *Soil and Tillage Research*. 29, 335-355.
- Conant, R.T., Easter, M., Paustian, K., Swan, A. & Williams, S. (2007). Impacts of periodic tillage on soil C stocks: a synthesis. *Soil and Tillage Research*. 95, 1–10.
- Conway, G. (1998). *The Doubly Revolution. Food for all in the Twenty-First century*. Cornell University Press. New York, USA.
- Conway, K.E. (1996). An overview of the influence of sustainable agricultural systems on plant diseases. *Crop Protection*. 15, 223-228.
- Conway, G. & Waage, J. (2010). *Science and innovation for development*. UK Collaborative on Development Science (UKCDS). London, UK.
- Cook, S.K., Swain, A.J., Clarke, J., Moss, S., Hughes, Z., Orson, J., Powell, L., Creasey, A., Norman, K. & Alford, J. (2006). Improving crop profitability by using minimum cultivation and exploiting grass weed ecology. *Home Grown Cereals Authority* (HGCA). Project Report No. 381. London.

- Corbeels, M., Hofman, G. & Van Cleemput, O. (1998). Analysis of water use by wheat grown on a cracking clay soil in a semi-arid Mediterranean environment: weather and nitrogen effects. *Agriculture and Water Management*. 38, 147-167.
- Cossani, C.M., Slafer, G.A. & Savin, R. (2009). Yield and biomass in wheat and barley under a range of conditions in a Mediterranean site. *Field Crops Research*. 112 (2-3), 205-213.
- Cox, M.C., Qualset, C.O. & Rains, D.W. (1985). Genetic variation for nitrogen assimilation and translocation in wheat. 2. Nitrogen assimilation in relation to grain yield and protein. *Crop Science*, 25, 435-440.
- Cox, D.J. & Shelton, D.R. (1992). Genotype-by-tillage interactions in hard red winter wheat quality evaluation. *Agronomy Journal*. 84, 627–630.
- Craig, K. & Logan, R. (eds.) (2013). *Farm management handbook 2013/14*. 34th Edition. Scottish Agricultural College (SAC) Consulting.
- Crawley, M.J. (1997). *Plant ecology*. Blackwell Science, Cambridge.
- Cussans, F.E. (1968). The growth and development of *Agropyron repens* (L.) Beauv., in competition with cereal, field beans and oilseed rape. In *Proceedings of the 9th British Weed Control Conference*. Brighton. 131-136.
- Daepf, M., Nösberger, J. & Lüscher, A. (2001). Nitrogen fertilization and developmental stage alter the response of *Lolium perenne* to elevated CO₂. *New Phytologist*. 150, 347-358.
- Dakora, F.D. (1996). Using indigenous knowledge to increase agricultural productivity in Africa. In Normann H, Snyman I, Cohen M (eds). *Indigenous Knowledge and its Uses in Southern Africa*. Human Science Research Council (HSRC) Press, Pretoria, South Africa.
- Dalgaard, T., Halberg, N. & Porter, J.R. (2001). A model for fossil energy use in Danish agriculture used to compare organic and conventional farming. *Agriculture, Ecosystem and Environment*. 87, 51–65
- Das, T.K. & Yaduraju, N.T. (1999). Effect of weed competition on growth, nutrient uptake and yield of wheat as affected by irrigation and fertilizers. *The Journal of Agricultural Science*. 133, 4551
- Davies, B. & Finney, B. (2002). Reduced cultivation for cereals: Research, development and advisory needs under changing economic circumstances. *Home-Grown Cereals Authority* (HGCA), London, UK.

- Davis, A.S. & Liebman, M. (2001). Nitrogen source influences wild mustard growth and competitive effect on sweet corn. *Weed Science*. 49, 558-566.
- Day, G.E., Paulsen, G.M. & Sears, R.G. (1985). Nitrogen relations in winter wheat cultivars differing in grain protein percentage and stature. *Journal of Plant Nutrition*. 8, 555-566
- Deike, S., Pallutt, B. & Christen, O. (2008). Investigations on the energy efficiency of organic and integrated farming with specific emphasis on pesticide use intensity. *European Journal of Agronomy*. 28, 461-470.
- Delogu, G., Cattivelli, L., Pecchioni, N., Defalcis, D., Maggiore, T. & Stanca, A.M. (1998). Uptake and agronomic efficiency of nitrogen in winter barley and winter wheat. *European Journal of Agronomy*. 9, 11-20.
- Demircan, V., Ekinçi, K., Keener, H.M., Akbolat, D. & Ekinçi, C. (2006). Energy and economic analysis of sweet cherry production in Turkey: A case study from Isparta province. *Energy Conversion Management*. 47, 1761–1769.
- Demjanová, E., Macák, M., Đalović, I., Majerník, F., Týr, Š. & Smatana, J. (2009). Effects of tillage systems and crop rotation on weed density, weed species composition and weed biomass in maize. *Agronomy Research*. 7 (2), 785-792.
- DEFRA. (2013). *Agriculture in the UK – Tables & Charts*. Department for Environment Food and Rural Affairs. [Online] Available from: <http://archive.defra.gov.uk/evidence/statistics/foodfarm/general/auk/latest/excel/> Accessed (21-11-13)
- DEFRA. (2012). *Agriculture in the UK 2012. National Statistics*. Department for Environment Food and Rural Affairs. [Online] Available from: <https://www.gov.uk/government/statistical-data-sets/agriculture-in-the-united-kingdom> Accessed (05-12-12)
- DEFRA. (2014a). *Organic Statistics 2013*. United Kingdom. National Statistics. Department for Environment Food and Rural Affairs. [Online] Available from: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/317366/organics-statsnotice-05jun14.pdf Accessed (17-05-14)
- DEFRA. (2010). *Fertiliser Manual*. 8th Edition (RB 209). Department for Environment Food and Rural Affairs.
- DEFRA. (2012). *UK Climate Change Risk Assessment: Government Report*. Department for Environment Food and Rural Affairs. [Online] Available from <https://www.gov.uk/government/publications/uk-climate-change-risk-assessment-government-report> Accessed (05-12-12)

DEFRA. (2011). The Natural Choice: securing the value of nature. Department for Environment Food and Rural Affairs. [Online] Available from <http://www.gov.uk/government/publications/the-natural-choice-securing-the-value-of-nature> Accessed (05-12-12)

DEFRA. (2014b). The Natural Environment White Paper Implementation Update. Department for Environment Food and Rural Affairs. [Online] Available from: <http://www.gov.uk/government/publications/natural-environment-white-paper-implementation-updates> Accessed (15-03-14)

Derksen, D.A., Thomas, A.G., Lafond, G.P., Loeppky, H.A. & Swanton, C.J. (1995). Impact of post-emergence herbicides on weed community diversity within conservation-tillage systems. *Weed Research*. 35, 311-320.

De Vita, P., Di Paolo, E., Fecondo, D., Di Fonzo, N. & Pisante, M. (2007). No-tillage and conventional tillage effects on durum wheat yield, grain quality and soil moisture content in southern Italy. *Soil and Tillage Research*. 92, 69-78.

Dimmock, J.P.R.E. & Gooding, M.J. (2002). The influence of foliar diseases, and their control by fungicides, on the protein concentration in wheat grain: A review. *Journal of Agricultural Science*. 138, 349-366.

Ding, X., Zhang, B., Zhang, X., Yang, X. & Zhang, X. (2011). Effects of tillage and crop rotation on soil microbial residues in a rainfed agroecosystem of northeast China. *Soil and Tillage Research*. 114, 43-49.

Di Tomaso, J. (1995). Approaches for improving crop competitiveness through the manipulation of fertilization strategies. *Weed Science*. 43, 491-491.

Donald, C.M. & Hamblin, J. (1976). The biological yield and harvest index of cereals as agronomic and plant breeding criteria. *Advance Agronomy*. 28, 361-405.

Döring, T.F., Baddeley, J.A., Brown, R., Collins, R., Crowlet, O., Cuttle, S., Howlett, S.A., Jones, H.E., McCalman, H., Measures, M., Pearce, B.D., Pearce, H., Roderick, S., Stobart, R., Strokey, J., Tilston, E., Topp, K., Watson, C., Winkler, L.R. & Wolfe, M.S. (2013). *Using legume-based mixture to enhance the nitrogen use efficiency and economic viability of cropping systems*. Final Report (LK09106/HGCA3447). HGCA Project Report No 513. Home-Grown Cereal Authority. Kenilworth, UK.

Duda, G.P., Guerra, J.M.G., Monteiro, M.T., De-Polli, H. & Teixaira, M.G. (2003). Perennial herbaceous legume as live soil mulches and their effects on C, N and P of the microbial biomass. *Scientific Agricola*. 60, 139-147.

Duncan, S.R. & Schapaugh, W.T. (1997). Relay-intercropped soybean in different water regimes, planting patterns and winter wheat cultivars. *Journal of Production Agriculture*. 10, 123-129.

Edwards, C. (2006). *Review of HGCA-funded Weed Research (including LINK projects) 1994–2005*. Home-Grown Cereals Authority (HGCA), London. Research Review No. 57.

Elgersma, A., Schleepers, H. & Nassiri, M. (2000). Interactions between perennial ryegrass (*Lolium perenne* L.) and white clover (*Trifolium repens* L.) under contrasting nitrogen availability: productivity, seasonal patterns of species composition, N₂ fixation, N transfer and N recovery. *Plant and Soil*. 221 (2), 281-299.

Elliot, L.F., McCalla, T.M. & Waiss, A.J. (1976). Phytotoxicity associated with residue management. In Oschwald, W.R. (ed.). *Crop Residue Management Systems*. Houston, Texas. American Society of Agronomy. 28 November – 3 December. pp, 131-146.

El Titi (ed.). (2003). *Soil Tillage in Agroecosystems*. CRC Press, Boca Raton.

Ennaifar, S., Lucas, P., Meynard, J.M. & Makowski, D. (2005). Effects of summer fallow management on take-all of winter wheat caused by *Gaeumannomyces graminis* var. *tritici*. *European Journal of Plant Pathology*. 112 (2); 167-181.

Epplin, F.M., Stock, C.J., Kletke, D.D. & Peeper, T.F. (2005). Economic of size for conventional tillage and no-till wheat production. In *Southern Agricultural Economics Association*, Annual Meeting Feb 5-9, Little Rock, AR.

Erdal, G., Esengun, K., Erdal, H. & Gunduz, O. (2007). Energy use and economical analysis of sugar beet production in Tokat province of Turkey. *Energy*. 32, 35–41.

Esengun, K., Erdal, G., Gunduz, O. & Erdal, H. (2007). An economic analysis and energy use in stake-tomato production in Tokat province of Turkey. *Renewable Energy*. 32: 1873–1881.

EUROSTAT. 2013. *Agriculture, forestry and fishery statistics*. Eurostat pockbooks. 2013 Edition. [Online] Available from http://epp.eurostat.ec.europa.eu/cache/ITY_OFFPUB/KS-FK-13-001/EN/KS-FK-13-001-EN.PDF Accessed (04-02-14)

EUROSTAT. 2014. [Online] Available from http://epp.eurostat.ec.europa.eu/portal/page/portal/agriculture/agricultural_production/database Accessed (04-02-14)

Ewald, J.A. & Aebischer, N.J. (2000). Trends in pesticide use and efficacy during 26 years of changing agriculture in southern England. *Environmental Monitoring and Assessment*. 64, 493-529.

- Fabrizzi, K.P., Garcia, F.O., Costa, J.L. & Picone, L.I. (2005). Soil water dynamics, physical properties and corn and wheat response to minimum and no-tillage systems in the southern Pampas of Argentina. *Soil and Tillage Research*. 81, 57-69.
- Fageria, N.K., Baligar, V.C. & Li, Y.C. (2008). The role of nutrient efficient plants in improving crop yields in the twenty-first century. *Journal of Plant Nutrition*. 31, 1121-1157.
- Faithfull, N.T. (2002). *Methods in agricultural chemical analysis*. CABI Publishing, Wallingford, UK.
- FAO. (2013a). *FAO statistical yearbook*. World food and agriculture. Food and Agriculture Organization of the United Nations (FAO). Rome, Italy.
- FAO. (2013b). *FAOSTAT*. Food and Agriculture Organization of the United Nations (FAO). [Online] Available from: <http://faostat3.fao.org/faostat-gateway/go/to/home/E> Accessed (19-03-14)
- FAO. (2009). *How to Feed the World in 2050*. High Level Expert Forum. Food and Agriculture Organization of the United Nations (FAO). Rome, Italy.
- FAO. (2003). *World Agriculture: Towards 2015/2030. A FAO Perspective*. Food and Agriculture Organization of the United Nations (FAO). Rome, Italy.
- Farrand, E.A. (1972). Potential milling and baking value of home grown wheat. *Journal of the National Institute of Agricultural Botany*. 12, 464-470.
- Feng, Y., Ning, T., Li, Z., Han, B., Han, H., Li, Y., Sun, T. & Zhnag, X. (2014). Effects of tillage practices and rate of nitrogen fertilization on crop yield and soil carbon and nitrogen. *Plant and Soil Environment*. 60 (3), 100-104.
- Ferguson, W.S. (1967). Growth and development of wheat. In: Nielsen, K.F. (ed). *Canadian Centenary Wheat Symposium*. 197-206
- Ferrante, A., Savin, R. & Slafer, G.A. (2010). Floret development of durum wheat in response to nitrogen availability. *Journal of Experimental Botany*. 61, 4351-4359.
- Ferrise, R., Triossi, A., Stratonovitch P., Bindi, M. & Martre, P. (2010). Sowing date and nitrogen fertilisation effects on dry matter and nitrogen dynamics for durum wheat: an experimental and simulation study. *Field Crop Research*. 117, 245-257.
- Filipovic, D., Kosutic, S., Gospodaric, Z., Zimmer, R. & Banaj, D. (2006). The Possibilities of fuel savings and the reduction of CO₂ emissions in the soil tillage in Croatia. *Agriculture, Ecosystems and Environment*. 115. 290-294.

Finch, H.J.S., Samuel, A.M. & Lane, G.P.F. (eds). (2014). *Lockhart & Wiseman's Crop husbandry including grassland*. 9th edition. Woodhead Publishing series in Food Science, Technology and Nutrition. 277.

Finney, J.R. & Knight, B.A.G. (1973). The effect of soil physical conditions produced by various cultivation systems on the root development of winter wheat. *Journal of Agricultural Science*. 80, 435-442

Fitter, A.H. (1991). The ecological significance of root system architecture, an economic approach. In: Atkinson, D. (Ed.) *Plant Root Growth: An Ecological Perspective*. Oxford, Blackwell Scientific Publications. 229–243.

Fleming, A. & Vanclay, F. (2010) Farmer responses to climate change and sustainable agriculture. A review. *Agronomy for Sustainable Development*. 30, 11–19.

Fluck, R. C. (1992) Energy Analysis for Agriculture. In Fluck, R.C. (ed.). *Energy in Farm Production*. Energy in World Agriculture, 6. Elsevier. 45-52

Fois, S., Motzo, R. & Giunta, F. (2009). The effect of nitrogenous fertiliser application on leaf traits in durum wheat in relation to grain yield and development. *Field Crops Research*. 110 (1), 69-75.

Forristal, D. & Murphy, K. (2009). Can we reduce costs and increase profits with Min Till? In *Proceedings of the National Tillage Conference*. Teasgasc, Oak Park, Carlow, Ireland. pp, 48-66

Foulkes, M.J., Scott, R.K. & Sylvester-Bradley, R. (2001). A comparison of the ability of wheat varieties to withstand drought in UK conditions: resource capture. *Journal of Agricultural Science*. 137, 1-16.

Fox, R.L. & Bandel, V.A. (1986). Nitrogen utilization with no tillage. In Sprague, M.A. & Triple, G.B. (eds.). *No-tillage and Surface-Tillage Agriculture – The tillage Revolution*. Wiley Interscience, New York, USA.

Fowler, D.B., Brydon, J. (1989). No-till winter wheat production on the Canadian prairies: Timing of nitrogen fertilization. *Agronomy Journal*. 81, 817-825.

Franchini, J.C., Debiassi, H., Balnibot-Junior, A.A., Tonon, B.C., Bouças-Farias, J.R., de Oliveira, M.C.N. & Torres, E. (2012). Evolution of crop yields in different tillage and cropping systems over two decades in southern Brazil. *Field Crop Research*. 137, 178-185.

Francis, C.A. (1989). Biological efficiencies in multiple cropping systems. *Advance in Agronomy*. 42, 1-42.

Francis, C.A. (ed.). (1986). *Multiple cropping systems*. Macmillan, New York, New York, USA.

Franzluebbers, A.J. (2004). Tillage and residue management effects on soil organic matter. In Magdoff, F. & Weil, R.R (eds.). *Soil organic matter in Sustainable agriculture*. CRS Press, New York. 2004. 227-268.

Franzluebbers, A.J., Hons, F.M. & Zuberer, D.A. (1995). Soil organic carbon, microbial biomass, and mineralizable carbon and nitrogen in sorghum. *Soil Science Society of America Journal*. 59, 460-466.

Freebairn, D.M. & Wockner, G.H. (1983). Soil erosion control research provides management answers. *Queensland Agricultural Journal*. 109, 277-234.

Freibauer, A., Rounsevell, M.D.A., Smith, P. & Verhagen, J. (2004). Carbon sequestration in the agricultural soils of Europe. *Geoderma*. 122, 1–23.

Froud-Williams, R.J., Chancellor, R.J. & Drennan, D.S.H. (1983a). Influence of cultivation regime upon buried weed seeds in arable cropping systems. *Journal of Applied Ecology*. 20, 199–208.

Froud-Williams, R.J., Drennan, D.S.H & Chancellor, R.J. (1983b). Influence of cultivation regime on weed floras of arable cropping systems. *Journal of Applied Ecology*. 20, 187-197.

Froud-Williams R.J., Chancellor, R.J., Drennan, D.S.H. (1981). Potential changes in weed floras associated with reduced-cultivation systems for cereal production in temperate regions. *Weed Research*. 21, 99-109.

Froud-Williams, R.J., Chancellor, R.J. & Drennan, D.S.H. (1984). The effects of seed burial and soil disturbance on emergence and survival of arable weeds in relation to minimal disturbance. *Journal of Applied Ecology*. 21 (2), 629–641.

Froud-Williams, R.J., Drennan, D.S.H. & Chancellor, R.J. (1983b). Influence of cultivation regime on weed floras of arable cropping systems. *Journal of Applied Ecology*. 20, 187–197.

Fuentes, J.P., Flury, M., Huggins, D.R. & Bezdicek, D.F. (2003). Soil water and nitrogen dynamics in dryland cropping systems of Washington State, USA. *Soil and Tillage Research*. 71, 33-47.

Fujita, K., Ofosu, K.G. & Ogata, S. (1992). Biological nitrogen fixation in mixed legume-cereal cropping system. *Plant and Soil*. 144, 155-175

Fukai, S. & Trenbath, B.R. (1993). Processes determining intercrop productivity and yields of component crops. *Field Crop Research*. 34, 247-271.

G20-led Wheat Initiative. [Online] Available from <http://urgi.versailles.inra.fr/Species/Wheat/Projects2/Wheat-Initiative> Accessed (05-11-14).

Gajri, P.R., Arora, V.K. & Prihar, S.S. (2002). *Tillage for sustainable cropping*. Food Product Press, New York.

Gao, X., Lukow, O.M. & Grant, C.A. (2012). Grain concentrations of protein, iron and zinc and bread making quality in spring wheat as affected by seeding date and nitrogen fertilizer management. *Journal of Geochemical Exploration*. 121, 36-44

Garrido-Lestache, E., López-Bellido, R.J. & López-Bellido L. (2004). Effect of N rate, timing and splitting and N type on bread-making quality in hard red spring wheat under rainfed Mediterranean conditions. *Field Crops Research*. 85, 213-236.

Gastal, F. & Lemaire, G. (2002). N uptake and distribution in crops: an agronomical and ecophysiological perspective. *Journal of Experimental Botany*. 53 (370). Inorganic Nitrogen Assimilation Special Issue. 789-799.

Gerba, L., Getachew, B. & Walelign, W. (2013). Nitrogen fertilization effects on grain quality of durum wheat (*Triticum turgidum* L. var. durum) varieties in central Ethiopia. *Agricultural Science*. 4 (3), 123-130.

Germon, J.C., Taureau, J.C. & Thomas, J.M. (1994). Effects des méthodes simplifiées de travail du sol sur les transformations de l'azote et leurs conséquences sur le lessivage des nitrates. In INRA (ed.). *Simplification du travail du sol*. Paris, France.

Ghaderi, M.G., Zeinalikhanghan, H., Hosseinzadeh, A.H., Taleei, A.R. & Naghavi, M.R. (2009). Evaluation of relationships between grain yield, yield components, and the other characteristics associated with grain yield in bread wheat using multivariate statistical analysis. *Iran Journal of Crop Research*. 7 (2), 573-582.

Ghaley, B.B., Hauggaard-Nielsen, H., Høgh-Jensen, H. & Jensen, E.S. (2005). Intercropping of wheat and pea as influenced by nitrogen fertilization. *Nutrient Cycling in Agroecosystems*. 73. 201-212.

Ghuman, B.S. & Sur, H.S. (2001). Tillage and residue management effects on soil properties and yields of rainfed maize and wheat in a subhumid subtropical climate. *Soil and Tillage Research*. 58, 1-10.

Giacomini, S.J., Machet, J.M., Boizard, H. & Recous, S. (2010). Dynamics and recovery of fertilizer ¹⁵N in soil and winter wheat crop under minimum versus conventional tillage. *Soil and Tillage Research*. 108, 51-58.

Giller, K.E. (2001). *Nitrogen fixation in tropical cropping systems*. CAB International, Wallingford, UK.

Giller, K.E. & Wilson, K.J. (1991). *Nitrogen fixation in tropical cropping systems*. CAB International, Wallingford, UK.

Glendining, M.J., Powlson, D.S., Poulton, P.R., Bradbury, N.J., Palazzo, D. & Li, X. (1996). The effects of long-term applications of inorganic nitrogen fertilizer on soil nitrogen in the Broadbalk Wheat Experiment. *Journal of Agricultural Science*. 127, 347-363.

Godfray, H.C.J. (2014). The Challenge of feeding 9-10 billion people equitably and sustainably. *Journal of Agricultural Science*. 152, S2-S8.

Godfrey, D., Hawkesford, M.J., Powers, S.J., Millar, S. & Shewry, P.R. (2010). Effects of crop nutrition on wheat grain composition and end use quality. *Journal of Agriculture and Food Chemistry*. 58, 3012-3021.

Gomiero, T., Pimentel, D. & Paoletti, M.G. (2011). Environmental impact of different agricultural management practices: Conventional vs. Organic agriculture. *Critical Reviews in Plant Sciences*. 30, 95-124.

Gonzalez-Sanchez, E.J., Veroz-Gonzalez, O., Blanco-Roldan, G.L., Marquez-Garcia, F. & Carbonell-Bjollo, R. (2015). A renewed view of conservation agriculture and its evolution over the last decade in Spain. *Soil and Tillage Research*. 146, 204-212.

Gooding, M.J. & Davies, W.P. (1997). *Wheat production and utilization: Systems, quality and the environment*. CAB International.

Gooding, M.J., Kasyanova, E., Ruske, R., Hauggaard-Nielsen, H., Jensen, E.S., Dahlmann, C., Von Fragstein, P., Dibet, A., Corre-Hellou, G., Crozat, Y., Pristeri, A., Romeo, M., Monti, M. & Launay, M. (2007). Intercropping with pulses to concentrate nitrogen and sulphur in wheat. *Journal of Agricultural Science*. 145, 469-479.

GO-Science. (2011). *The future of food and farming: Challenges and choices for global sustainability*. Final Project Report. The Government Office for Science (GO-Science). Foresight. London, UK.

Gosling, S., Dunn, R., Carrol, F., Christidis, N., Fullwood, J., Gusmao, D., Golding, N., Good, L., Hall, T., Kendon, L., Kennedy, J., Lewis, K., McCarthy, R., McSweeney, C., Morice, C., Parker, D., Perry, M., Stott, P., Willett, K., Allen, M., Arnell, N., Bernie,

- Da., Betts, R., Bowerman, N., Brak, B., Caesar, J., Challinor, A., Dankers, R., Hewer, F., Huntingford, C., Jenkins, A., Klingaman, N., Lowe, J., Lloyd-Hughes, B., Miller, J., Nicholls, R., Noguer, M., Otto, F., van der Linden, P., Warren, R. (2011). *Climate: observations, projections and impacts*. UK Met Office (UKMO).
- Graham, J.P., Ellis, F.B., Christian, D.G. & Cannell, R.Q. (1985). Effect of straw residues on the establishment, growth and yield of autumn-sown cereals. *Journal of Agriculture Engineering Research*. 33 (1), 39-49.
- Grant, C.A. & Lafond, G.P. (1993). The effects of tillage systems and crop sequences on soil bulk density and penetration resistance on a clay soil in southern Saskatchewan. *Canadian Journal of Soil Science*. 73, 223-232.
- Graybosch, R.A., Peterson, C.J., Shelton, D.R. & Baenziger, P.S. (1996). Genotypic and environmental modification of wheat flour protein composition in relation to end-use quality. *Crop Science*. 36, 296–300.
- Green, M.B. (1987). Energy in pesticide manufacture, distribution and use. In Helsel, Z.R. (ed.). *Energy in Plant Nutrition and Pest Control*. Elsevier, Amsterdam. 165–177.
- Greenwood, D.J. (1982). Modelling of crop response to nitrogen fertiliser. *Philosophical Transactions of the Royal Society, London, Series B*, 296, 351- 362.
- Greenwood, D.J., Lemaire, G., Gosse, G., Cruz, P., Draycott, A. & Neeteson, J.J. (1990). Decline in percentage N of C₃ and C₄ crops with increasing plant mass. *Annals of Botany*. 66, 425-436.
- Gregory, P.J. (1994). Root growth and activity. In Boote, K.J., Bennett, J.M., Sinclair, T.R. & Paulsen, G.M. (eds.). *Physiology and determination of crop yield*. ASA-CSSA-SSSA, Madison, USA. 65-93.
- Griffith, D.R., Mannering, J.V. & Moldenhauer, W.C. (1977). Conservation tillage in the eastern corn belt. *Journal of Soil and Water Conservation*. 32 (1), 20-28.
- Gruber, S., Möhring, J., Claupein, W. (2011). On the way towards conservation tillage-soil moisture and mineral nitrogen in a long-term field experiment in Germany. *Soil and Tillage Research*. 115-116, 80-87.
- Gruber, S., Pekrun, C., Möhring, J., Claupein, W. (2012). Long-term yield and weed response to conservation and stubble tillage in SW Germany. *Soil and Tillage Research*. 121, 49-56.
- Grundy, A.C. (2003) Predicting weed emergence: a review of approaches and future challenges. *Weed Research*. 43, 1-11.

- Grundy, A.C., Froud-Williams, R.J. & Boatman, N.D. (1993). The use of cultivar, crop see rate and nitrogen level for the suppression of weeds in winter wheat. *Brighton Crop Protection Conference – Weeds*. 997-1002.
- Guérif, J., Richard, G., Dürr, C., Machet, J.M., Recous, S. & Roger-Estrade, J. (2001). A review of tillage effects on crop residue management, seedbed conditions and seedling establishment. *Soil and Tillage Research*. 61, 13-32.
- Guo, J.H., Liu, X.J., Zhang, Y., Shen, J.L., Han, W.X., Zhang, W.F., Christie, P., Gouling, K.W.T., Vitousek, P.M. & Zhang, F.S. (2010). Significant acidification in major Chinese croplands. *Science*. 327, 1008–1010.
- Gurevitch, J. & Hedges, L.V. (1999). Statistical issues in ecological meta-analyses. *Ecology*. 80, 1142-1149.
- Gürsoy, S., Sessiz, A. & Malhi, S.S. (2010). Short-term effects of tillage and residue management following cotton on grain yield and quality of wheat. *Field Crop Research*. 119 (2-3), 260-268.
- Guy, S.O. & Lauver, M.A. (2007). No-till and conventional-till effects on spring wheat in the Palouse. *Crop management*. 6 (1). doi:10.1094/CM-2007-0508-01-RS
- Haas, G., Wetterich, F., and Kopke, U. 2001. Comparing intensive, extensified and organic grassland farming in southern Germany by process life cycle assessment. *Agriculture, Ecosystem and Environment*. 83, 43–53.
- Håkansson, I. (1994). Soil tillage for crop production and for protection of soil and environmental quality: A Scandinavian point of view. *Soil and Tillage Research*. 30, 109-124.
- Halvorson, A.D., Wienhold, B.J. & Black, A.L. (2001). Tillage and nitrogen fertilization influence grain and soil nitrogen in an annual cropping system. *Agronomy Journal*. 93, 836–841.
- Hamdi, Y.A. (1995). Potential and assessment of BNF (Biological Nitrogen Fixation) and its direct contribution in selected cropping systems and ecological conditions. In Dudal, R. & Roy, R.N. (eds.). *Integrated plant nutrition systems. Report of an Expert Consultation, Rome, Italy 13-15 December 1993*. FAO Fertilizer and Plant Nutrition Bulletin No 12. Rome, Italy.
- Hamilton, R. (2009). Agriculture’s sustainable future: Breeding better crops. In *Scientific America*. [Online] Available from: <http://www.scientificamerican.com/article/agricultures-sustainable-future/> Accessed (23-11-13)

- Hansen, E.M., Munkholm, L.J. & Olesen, J.E. (2011). N-utilization in non-inversion tillage systems. *Soil and Tillage Research*. 113, 55-60.
- Harman, W.L., Daves, R.M., Morrison, J.E. & Potter, K.N. (1996). Economics of wide-bed farming systems for Vertisol (clay) soils: chisel tillage vs. no-tillage. *Journal of American Farm Management Rural Appraise*. 19, 129-134.
- Harper, S.H.T. (1989). Management of crop residues. In: Bäumer, K. & Ehler, W. (eds.). EC Workshop: *Agriculture: Energy saving by reduced soil tillage*, Gottingen, Luxembourg: Commission of the European Communities, 10-11 June 1987, 127-132.
- Hartl, W. (1989). Influence of undersown clovers on weeds and on the yield of winter wheat in organic farming. *Agriculture, Ecosystems and Environment*. 27, 389-396.
- Hartmann, S., Rößl, G., Forster, M. (2009). Gelbklees – *Medicago lupulina* L. in: für Pflanzenbau und Pflanzenzüchtung, Bayerische Landesanstalt für Landwirtschaft. [Online]. Available from <http://www.flbayernde/ipz/gruenland/09663/gelbkleepdfInstitut> Accessed (25-11-13)
- Hartwig, N.L. & Ammon, H.U. (2002). Cover crops and living mulches. *Weed Science*. 50, 688-699.
- Hatfield, J.L. & Prueger, J.H. (2004). Nitrogen over-use, under-use, and efficiency. In *Proceedings of the 4th International Crop Science Congress*. Brisbane, Australia, 26 September – 1 October.
- Hatfield, J.L., Sauer, T.J. & Prueger, J.H. (2001). Managing soils to achieve greater water use efficiency: A review. *Agronomy Journal*. 93, 271-280.
- Hauggaard-Nielsen, H., Ambus, P. & Jensen, E. S. (2001). Interspecific competition, N use and interference with weeds in pea-barley intercropping. *Field Crops Research*. 70, 101-109.
- Hauggaard-Nielsen, H., Andersen, M.K., Jornsgaard, B. & Jensen, E. S. (2006). Density and relative frequency effects on competitive interactions and resource use in pea-barley intercrops. *Field Crops Research*. 95, 256–267.
- Hauggaard-Nielsen, H., Jornsgaard, B., Kinane, J. & Jensen, E.S. (2008). Grain legume-cereal intercropping: The practical application of diversity, competition and facilitation in arable and organic cropping systems. *Renewable Agriculture and Food Systems*. 23 (1), 3-12.
- Hauggaard-Nielsen, H., Jornsgaard, B. & Steen, J.E. (2003). Legume-cereal intercropping system as a weed management tool. In: *Proceedings of the 4th European Weed Research Society Workshop: Crop weed competition interaction*. Universita Tusca, Viterbro, Italy, 10–12th April.

- Hawkesford, M.J., Araus, J.L., Park, R., Calderini, D., Miralles, D., Shen, T., Zhang, J. & Parry, M.A.J. (2013). Prospect of doubling global wheat yields. *Food and Energy Security*. 2 (1), 34-48.
- Hay, R.K.M. (1995). Harvest index- a review of its use in plant breeding and crop physiology. *Annals of Applied Biologists*. 126, 197-216.
- Hay, R.K.M. & Walker, A.J. (1989). *An introduction to the physiology of crop yield*. Longman Scientific and Technical, Essex, UK.
- Haymes, R. & Lee, H.C. (1994). Agronomic aspects of wheat-bean intercropping in a low input system. In *Proceedings of 3rd ESA Congress on Sustainable Agriculture*. Abavo-Pavoda. 706-707.
- Hayward, C.F. (1990). The effect of sowing date on grain quality of winter wheat. *Aspect of Applied Biology*. Cereal Quality II. 25, 163-170.
- Heitholt, J.J., Croy, L.I., Maness, N.O. & Nguyen, H.T. (1990) Nitrogen partitioning in genotypes of winter wheat differing in grain N concentration. *Field Crops Research*. 23, 133-144.
- Hemmat, A. & Taki, O. (2001). Grain yield of irrigated winter wheat as affected by stubble-tillage management and seeding rates in central Iran. *Soil and Tillage Research*. 63 (1-2), 57-64.
- Hernanz, J.L, Giron, V.S. & Cerisola, C. (1995). Long-term energy use and economic evaluation of three tillage systems for cereal and legume production in central Spain. *Soil and Tillage Research*. 35, 183–198.
- HGCA. (2012). *Desk study to evaluate contributory causes of the current 'yield plateau' in wheat and oilseed rape*. Project Report No. 502. Home-Grown Cereals Authority (HGCA). London, UK.
- HGCA. (2010). *Managing weeds in arable rotation – a guide*. Home-Grown Cereals Authority (HGCA). London, UK.
- HGCA. (2014). *Recommended List Spring Wheat (for spring sowing)*. Home-Grown Cereals Authority (HGCA). London, UK.
- HGCA. (2006). *Take-all in winter wheat- management guidelines*. Home-Grown Cereals Authority (HGCA), London, UK.
- HGCA. (2008a). *The encyclopaedia of cereal diseases*. Home-Grown Cereals Authority (HGCA), London, UK.

HGCA. (2008b). *The wheat growth guide*. Spring 2008. 2nd edition. Home-Grown Cereals Authority (HGCA). London, UK.

Hickman, M.V. (2007). Long-term tillage and crop rotation effects on soil chemical and mineral properties. *Journal of Plant Nutrition*. 25 (7), 1457-1470.

Hiltbrunner, J., Bloch, L., Richner, W. & Streit, B. (2002). Development of weed populations with no-tillage of winter wheat into living mulches. *Journal of Plant Disease and Protection*. XVIII, 581-587.

Hiltbrunner, J., Liedgens, M., Bloch, L., Stamp, P. & Streit, B. (2007). Legume cover crop as living mulches for winter wheat: components of biomass and the control of weeds. *European Journal of Agronomy*. 26, 21-29.

Hill, R.L. (1990). Long-term conventional and no-tillage effects on selected soil physical properties. *Soil Science Society of America Journal*. 54, 161-166

Hirel, B., Le Gouis, J., Ney, B. & Gallais, A. (2007). The challenge of improving nitrogen use efficiency in crop plants: Towards a more central role for genetic variability and quantitative genetics within integrated approaches. *Journal of Experimental Botany*. 58, 2369-2387.

Hobbs, P.R., Sayre, K. & Gupta, R. (2008). The role of conservation agriculture in sustainable agriculture. *Philosophical Transactions of the Royal Society B*. 363, 543-555.

Holland, J.M. (2004). The environmental consequences of adopting conservation tillage in Europe: reviewing the evidence. *Agriculture, Ecosystems and Environment*. 103, 1-25.

Horwith, B. (1985). A role for intercropping in modern agriculture. *BioScience*. 35, 286-291.

Houx, IIIJ. H., Wiebold, W.J., Fritschi, F.B. (2011). Long-term tillage and crop rotation determines the mineral nutrient distributions of some elements in a Vertic Epiaqualf. *Soil and Tillage Research*. 112, 27-35.

Hruskova, M. & Famera, O. (2003) Prediction of wheat and flour Zeleny sedimentation value using NIR technique. *Czech Journal of Food Science*. 21, 91-96.

Huggins, D.R., & Pan, W.L. (1993). Nitrogen efficiency component analysis: An evaluation of cropping system differences in productivity. *Agronomy Journal*. 85, 898-905.

Huggins, D.R. & Pan, W.L. (2003). Key indicators for assessing nitrogen use efficiency in cereal-based agroecosystems. *Journal of Crop Production*. 8 (1-2), 157-185.

- Huggins, D., Pan, W. & Smith, J. (2010). Yield, protein and nitrogen use efficiency of spring wheat: Evaluating field-scale performance. *CSANR Research Report 2010-2001*. Precision N field performance.
- Hunter, A.S. & Stanford G. (1973). Protein content of winter wheat in relation to rate and time of nitrogen fertilizer application. *Agronomy Journal*. 65, 772-774.
- Hulme, M., Jenkins, G.J., Lu, X., Turnpenny, J.R., Mitchell, T.D., Jones, R.G., Lowe, J., Murphy, J.M., Hassell, D., Boorman, P., McDonald, R. & Hill, S. (2002). *Climate Change Scenarios for the United Kingdom: The UKCIP02 Scientific Report*. Tyndall Centre for Climate Change Research, School of environmental Sciences, University of East Anglia, Norwich, UK. 120.
- Hussain, I., Olson, K.R. & Ebelhar, S.A. (1999). Impacts of tillage and no-till on production of maize and soybean on an eroded Illinois silt loam soil. *Soil and Tillage Research*. 52 (1-2), 37-49.
- Hussain, Z., Khan, M.A. & Irfan, M. (2010). Water energy and economic analysis of wheat production under raised bed and conventional irrigation systems: A case study from semi-arid area of Pakistan. *Soil and Tillage Research*. 109, 61-67.
- IPCC. (2007). *Climate Change 2007: The Physical Science Basis*. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change (IPCC). Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor & H.L. Miller (eds.). Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 996.
- IPCC. (2014a). *Climate change 2014: Mitigation of Climate change*. Working group III contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (IPCC). WMO, UNEP, Geneva.
- IPCC (2014b). Summary for policymakers. in: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects*. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (IPCC). Field, C.B., V.R. Barros, D.J. Dokken, K.J. Mach, M.D. Mastrandrea, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S. MacCracken, P.R. Mastrandrea, & L.L. White (eds.). Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. 1-32.
- Iqbal, J. & Wright, D. (1997). Effects of nitrogen supply on competition between wheat and three annual weed species. *Weed Research*. 37, 391-400.
- James, W.C. (1971). An illustrated series of assessment keys for plant diseases, their preparation and usage. *Canadian Plant Disease Survey*. 51 (2), 39-65.

Jenkins, G. J., Murphy, J. M., Sexton, D. M. H., Lowe, J. A., Jones, P. & Kilsby, C. G. (2009). *UK Climate Projections: Briefing report*. Met Office Hadley Centre, Exeter, UK.

Jenkins, G.J., Perry, M.C. & Prior, M.J. (2008). *The climate of the United Kingdom and recent trends*. Met Office Hadley Centre, Exeter, UK

Jenkinson, D. (2010). *Climate change*. [Online]. Available from: http://www.rothamsted.bbsrc.ac.uk/aen/reviews/Climate_Change_Draft_B03.pdf
Accessed (08-08-14)

Jenkinson, D.S., Hart, P.B.S., Rayner, J.H. & Parry, L.C. (1987). Modelling the turnover of organic matters in long-term experiments at Rothamsted. *INTECOL Bulletin*. 15. 1-8.

Jenkyn, J.F., Gutteridge, R.J. & Bateman, G.L. (2004). Effects of cultivation method and crop debris on wheat stem-base diseases and take-all. In *HGCA Conference: Managing soil and roots for profitable productions*.

Jensen, E.S. (1996) Grain yield and symbiotic N₂ fixation and interspecific competition for inorganic N in pea–barley intercrops. *Plant and Soil*. 182, 25–38.

Jensen, E.S. (1991). Nitrogen accumulation and residual effects of nitrogen catch crops. *Acta Agriculturae Scandinavica*. 41, 333-344.

Johnson, G.A., Defelice, M.S. & Helsel, Z.R. (1993). Cover crop management and weed control in corn (*Zea mays*). *Weed Technology*. 7, 425–430.

Johnson, V.A. & Mattern, P.J. (1987). Wheat, rye and triticale. In Olson, R.A., Frey, K.J. (eds.). *Nutritional quality of cereals grains: Genetics and agronomic improvements*. 38. American Society of Agronomy, Inc., Madison, WI, USA. 133-182.

Johnson, V.A., Mattern, P.J., Peterson, C.T. & Kurh, S.L. (1985). Improvement of wheat protein by traditional breeding and genetic techniques. *Cereal Chemistry*. 62, 350-355.

Jones, C.A., Basch, G., Baylis, A.D., Bazzoni, D., Bigs, J., Bradbury, R.B., Chaney, K., Deeks, L.K., Field, R., Gomez, J.A., Jones, R.J.A., Jordan, V., Lane, M.C.G., Leake, A., Livermore, M., Owens, P.N., Ritz, K., Sturny, W.G. & Thomas, F. (2006). *Conservation agriculture in Europe: an approach to sustainable crop production by protecting soil and water?* The Soil and Water Project (SOWAP). Jealott's Hill, Bracknell, UK.

Jones, L. (1992). Preliminary trials using a white clover (*Trifolium repens* L.) understorey to supply the nitrogen requirements of a cereal crop. *Grass and Forage Science*. 47, 366-374.

Jones, L. & Clements, R.O. (1993). Development of low input system for growing wheat (*Triticum vulgare*) in a permanent understorey of white clover (*Trifolium repens*). *Annals of Applied Biology*. 123, 109-119.

Jones, O.R. (1981). Land forming effects on dryland sorghum production in the Southern Great Plain. *Soil Science Society of America Journal*. 45, 606-611.

Jørnsgård, B., Rasmussen, K., Hill, J. & Christiansen, J.L. (1996). Influence of nitrogen on competition between cereals and their natural weed populations. *Weed Research*. 36: 461–470

Justes, E., Mary, B., Meynard, J.M., Machet, J.M. & Thelier-Huche, L. (1994). Determination of a critical nitrogen dilution curve for winter wheat crops. *Annals of Botany*. 74, 397.

Känkänen, H., Alakukku, L., Salo, Y., Pitkänen, T. (2011). Growth and yield of spring cereals during transition to zero tillage on clay soils. *European Journal of Agronomy*. 34, 35–45.

Känkänen, H., Eriksson, C., Räkköläinen, M. & Vuorinen, M. (2001). Effect of annually repeated undersowing on cereal grain yields. *Agricultural and Food Science in Finland*. 10. 197-208.

Kamiji, Y., Pang, J., Milroy, S.P. & Palta, J.A. (2014). Shoot biomass in wheat is the driver for nitrogen uptake under low nitrogen supply, but not under high nitrogen supply. *Field Crops Research*. 165, 92-98.

Kapusta, G., Krausz, R.F. & Matthews, J.L. (1996). Maize yield is equal in conventional, reduced, and no tillage after 20 years. *Agronomy Journal*. 88, 812- 817.

Karlen, D.L., Kovar, J.L., Cambardella, C.A. & Colvin, T.S. (2013). Thirty-year tillage effects on crop yield and soil fertility indicators. *Soil and Tillage Research*. 130, 24-41.

Kaspar, T.C. & Bland, W.L. (1992). Soil-temperature and root growth. *Soil Science*. 154, 290-299.

Kassam, A., Friedrich, T., Shaxson, F. & Pretty, J. (2009). The spread of conservation agriculture: justification, sustainability and uptake. *International Journal of Agricultural Sustainability*. 7, 292-320.

Keep, M. (2009). *Agriculture: historical statistics*. House of Commons Library. UK.

Khaledian, M., Mailjol, J.C. & Ruelle, P. (2014). Diesel oil consumption, work duration, and crop production of corn and durum wheat under conventional and no-tillage in southeastern France. *Archives of Agronomy and Soil Science*. 60 (8), 1067-1076.

- Kitani, O. (1999). *CIGR Handbook of Agricultural Engineering*. Energy and biomass engineering. Vol-V. American Society of Agricultural Engineers (ASAE) Publications. ST Joseph, MI.
- Knight, S.M. (2004). Plough, minimal till or direct drill? – Establishment method and production efficiency. In *Managing soil and roots for profitable production*. HGCA Conference. 2004.
- Koch, H.J., Dieckmann, J., Büchse, A. & Märländer, B. (2009). Yield decrease in sugar beet caused by reduced tillage and direct drilling. *European Journal of Agronomy*. 30, 101–109.
- Koga, N., Tsuruta, H., Tsuji, H. & Nakano, H. (2003). Fuel consumption-derived CO₂ emissions under conventional and reduced tillage cropping systems in northern Japan. *Agriculture, Ecosystems and Environment*. 99, 213-219.
- Kosmolak, F.G. & Crowle, W.L. (1980). An effect of nitrogen fertilization on the agronomics traits and dough mixing strength of five Canadian hard red spring wheat cultivars. *Canadian Journal of Plant Science*. 60, 1071-1076.
- Kramer, T. (1979). Environmental and genetic variation for protein content in winter wheat (*Triticum aestivum*). *Euphytica*. 28, 209-218.
- Kumar, K. & Goh, K.M. (2002). Management practices of antecedent leguminous and non-leguminous crop residues in relation to winter wheat yields, nitrogen uptake, soil nitrogen mineralization and simple nitrogen balance. *European Journal of Agronomy*. 16 (4), 295-308.
- Kumar, V., Saharawat, Y.S., Gathala, M.K., Jat A.S., Singh, S.K., Chaudhary, N. & Jat, M.L. (2013). Effect of different tillage and seeding methods on energy use efficiency and productivity of wheat in the Indo-Gangetic Plains. *Field Crop Research*. 142, 1-8.
- Küstermann, B., Munch, J.C. & Hülsbergen, K.J. (2013). Effects of soil tillage and fertilization on resource efficiency and greenhouse gas emissions in a long-term field experiment in Southern Germany. *European Journal of Agronomy*. 49, 61-73.
- Ladha, J.K., Pathak, H., Krupnik, T.J., Six, J. & Van Kessel, C. (2005). Efficiency of fertiliser nitrogen in cereal production: Retrospects and prospects. *Advances in Agronomy*. 87, 85–156.
- Lafond, G.P., Loepky, H. & Derksen, D.A. (1992). The effects of tillage systems and crop rotations on soil water conservation, seedling establishment and crop yield. *Canadian Journal of Plant Science*. 72, 103-115.

- Lafond, G.P., May, W.E., Stevenson, F.C. & Derken, D.A. (2006). Effects of tillage systems and rotations on crop production for a thin Black Chernozem in the Canadian Praires. *Soil and Tillage Research*. 89, 232-245.
- Lal, B., Gautam, P., Raja, R., Nayak, A.K., Shahid, M., Tripathi, R., Bhattacharyya, P., Mohanty, S., Puri, C., Kumar, A. & Panda, B.B. (2014). Weed community composition after 43 years of long-term fertilization in tropical rice-rice system. *Agriculture, Ecosystems and Environment*. 197, 301-308.
- Lal, R. (1979). Influence of six years of no-tillage and conventional ploughing on fertiliser response of maize on an Alfisol the tropics. *Soil Science Society of America Journal*. 43, 399-403.
- Lal, R., Reicosky, D.C. & Hanson, J.D. (2007). Evolution of the plough over 10,000 years and the rationale for no-till farming. *Soil and Tillage Research*. 93, 1-12.
- Lampurlanés, J., Angás, P. & Cantero-Martínez, C. (2001). Root growth, soil water content and yield of barley under different tillage systems on two soils in semiarids conditions. *Field Crop Research*. 69 (1), 27-40.
- Larson, W.E. & Osborne, G.J. (1982). Tillage accomplishments and potential. In Unger, P.W., Van Doren Jr, D.M., Whisler, F.D. & Skidmore, E.L. (eds.). *Predicting tillage effect on soil tillage effects on soil physical properties and processes*. America Society of Agronomy Special Publication. 44.
- Lauk, R. & Lauk, E. (2008). Pea-oat intercrops are superior to pea-wheat and pea-barley intercrops. *Acta Agriculturae Scandinavica Section B, Plant and Soil Science*. 58, 139-144.
- Lawlor, D.W. (1995). The effects of water deficit on photosynthesis. In Smirnoff, N. (ed.). *Environment and Plant metabolism*. Oxford: BioScience publishers. 129-160
- Lawlor, D.W., Boyle, F.A. Keys, A.J. Kendall, A.C. & Young, A.T. (1988). Nitrate nutrition and temperature effects on wheat: a synthesis of plant growth and nitrogen uptake in relation to metabolic and physiological processes. *Journal of Experimental Botany*. 39 (3), 329-343.
- Le Gouis, J., Beghin, D., Heumez, E. & Pluchard, P. (2000) Genetic differences for nitrogen uptake and nitrogen utilisation differences in winter wheat. *European Journal of Agronomy*. 12, 163-173.
- Lemaire, G. & Chartier M. (1992). Relationships between growth dynamics and nitrogen uptake for individual sorghum plants growing at different plant densities. In *Proceedings 2nd Congress of European Society for Agronomy*, Warwick, UK. 98-99

- Lemaire, G., Onillon, B., Gosse, G., Chartier, M. & Allirand, J.M. (1991). Nitrogen distribution within a Lucerne canopy during regrowth: Relation with light distribution. *Annals of Botany*. 68, 483-488.
- Liao, M.T., Fillery, I.R.P. & Palta, J.A. (2004). Early vigorous growth is a major factor influencing nitrogen uptake in wheat. *Functional Plant Biology*. 31, 121–129.
- Liberati, A., Altman, D.G. Tetzlaff, J., Mulrow, C., Gøtzsche, P.C., Ioannidis, J.P.A., Clarke, M., Devereaux, P.J., Kleijnen, J. & Moher, D. (2009). The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *Journal of Clinical Epidemiology*. 62 (10), e1–e34.
- Liebig, M.A., Varvel, G.E., Doran, J.W. & Wienhold, B.J. (2002). Crop sequence and nitrogen fertilization effects on soil properties in the Western corn Belt. *Soil Science of American Journal*. 66, 596-601.
- Liebman, M.Z. (1986). Ecological suppression of weeds in intercropping systems: Experiments with barley, pea, and mustard. PhD dissertation, University of California, Berkeley, CA.
- Liebman, M., & Dyck, E. (1993). Crop-rotation and intercropping strategies for weed management. *Ecological Applications*. 3, 92–122.
- Lindstrom, M.J., Koehler, F.E. & Papendick, R.I. (1974). Tillage effects on fallow water storage in the eastern Washington dryland region. *Agronomy Journal*. 66, 312-316.
- Lipiec, J. & Stepniewski, W. (1995). Effects of soil compaction and tillage systems on uptake and losses of nutrients. *Soil and Tillage Research*. 35, 37–52.
- Lithourgidis, A.S., Dordas, C.A., Damalas, C.A. & Vlachostergios, D.N. (2011). Annual intercrops: An alternative pathway for sustainable agriculture. *Australian Journal of Crop Science*. 5 (4), 396-410.
- Liu, B., Tu, C., Hu, S., Gumpertz, M., & Ristaino, J. B. (2007). Effect of organic, sustainable, and conventional management strategies in grower fields on soil physical, chemical, and biological factors and the incidence of Southern blight. *Applied Soil Ecology*. 37, 202–214.
- Liu, L., Hu, C., Olesen, J., Ju, Z., Yang, P. & Zhang, Y. (2013). Warming and nitrogen fertilization effect on winter wheat yields in northern China varied between four years. *Field Crops Research*. 151, 56-64.

- Lobell, D.B., Hammer, G.L., McLean, G., Messina, C., Roberts, M.J. & Schlenker, W. (2013). The critical role of extreme heat for maize production in the United States. *Nature Climate Change*. 3, 497-501.
- Lobell, D.B., Sibley, A. & Ortiz-Monasterio, J.I. (2012). Extreme heat effects on wheat senescence in India. *Nature Climate Change*. 2, 186-189.
- Lloveras, J., Lopez, A., Ferran, J., Espachs, S. & Solsona, J. (2001). Bread-making wheat and soil nitrate as affected by nitrogen fertilisation in irrigated Mediterranean conditions. *Agronomy Journal*. 93 (6), 1183-1190.
- Loffler, C.M., Rauch, T.L. & Busch, R. H. (1985). Grain and plant protein relationships in hard red spring wheat. *Crop Science*. 25, 521-524.
- Long, S.P., Zhu, X.G., Naidu, S.L. & Ort, D.R. (2006). Can improvement in photosynthesis increase crop yields? *Plant Cell Environment*. 29, 315–330.
- López, M.V., Arrúe, J.L. & Sánchez-Girón, V. (1996). A comparison between seasonal changes in soil water storage and penetration resistance under conventional and conservation tillage systems in Aragon. *Soil and Tillage Research*. 37 (4), 251-271
- López-Bellido, L., Fuentes, M., Castillo, J.E. & López-Garrido, F.J. (1998). Effects of tillage, crop rotation and nitrogen fertilization on wheat-grain quality grown under rainfed Mediterranean conditions. *Field Crop Research*. 57(3). 265-276.
- López-Bellido, L., Fuentes, M., Castillo, J.E., López-Garrido F.J. & Fernández, E.J. (1996). Long-term tillage, crop rotation, and nitrogen fertilizer effects on wheat yield under Mediterranean conditions. *Agronomy Journal*. 88, 783-791.
- López-Bellido, R.J. & López-Bellido, L. (2001). Efficiency of nitrogen in wheat under Mediterranean conditions: effect of tillage, crop rotation and N fertilization. *Field Crop Research*. 71, 31-46.
- López-Bellido, R.J., López-Bellido, L., Benítez-Vega, J. & López-Bellido, F.J. (2007a). Tillage systems, preceding crop, and nitrogen fertilizer in wheat crop: I. Soil water content. *Agronomy Journal*. 99, 59-67.
- López-Bellido, R.J., López-Bellido, L., Benítez-Vega, J. & López-Bellido, F.J. (2007b). Tillage systems, preceding crop, and nitrogen fertilizer in wheat crop: II. Water utilization. *Agronomy Journal*. 99, 66-72.
- López-Bellido, L., López-Bellido, R.J., Castillo, J.E. & López-Bellido, F.J. (2000). Effects of tillage, crop rotation, and nitrogen fertilization on wheat under rainfed Mediterranean conditions. *Agronomy Journal*. 92, 1054-1063.

- López-Bellido, L., López-Bellido, R.J., Castillo, J.E., López-Bellido, F.J. (2001). Effect of long-term tillage, crop rotation and nitrogen fertilization on bread-making quality of hard red spring wheat. *Field Crop Research*. 72, 197-210
- López-Bellido, L., López-Bellido, R.J & Redondo, E. (2005). Nitrogen efficiency in wheat under rainfed Mediterranean conditions as affected by Split nitrogen application. *Field Crop Research*. 94, 86-97.
- López-Bellido, L., López-Garrido, F.J., Fuentes, M., Castillo, J.E. & Fernández, E.J. (1997). Influence of tillage, crop rotation and fertilization on soil organic matter and nitrogen under rain-fed Mediterranean conditions. *Soil and Tillage Research*. 43, 277-293.
- López-Bellido, L., Muñoz-Romero, V., Benítez-Vega, J., Fernández-García, P., Redondo, R. & López-Bellido, R.J. (2012). Wheat response to nitrogen splitting applied to a Vertisol in different tillage systems and cropping rotations under typical Mediterranean climatic conditions. *European Journal of Agronomy*. 43, 24-32.
- López-Bellido, L., Muñoz-Romero, V. & López-Bellido, R.J. (2013). Nitrate accumulation in the soil profile: Long-term effect of tillage, rotation and N rate in Mediterranean Vertisol. *Soil and Tillage Research*. 130, 18-23.
- López-Fando C. & Pardo M.T. (2009). Changes in soil chemical characteristics with different tillage practices in a semi-arid environment. *Soil and Tillage Research*. 104, 278–284.
- Lovett, J.V. & Jessop, R.S. (1982). Effects of residue of crop plants on germination and early growth of wheat. *Australian Journal of Agricultural Research*. 33, 909-916.
- Lu, C., Ma, J., Chen, X., Zhang, X., Shi, Y. & Huang, B. (2010). Effect of nitrogen fertilizer and maize straw incorporation on $\text{NH}_4^{+15}\text{N}$ and NO_3^{-15} accumulation in black soil of Northeast China among three consecutive cropping cycles. *Journal of Soil Science and Plant Nutrition*. 10 (4), 444-453.
- Lueschen, W.E., Evans, S.D., Ford, J.H., Hoverstad, T.R., Kanne, B.K., Orf, J.H., Stienstra, W.C., Warnes, D.D. & Hicks, D.R. (1991). Soybean production as affected by tillage in a corn and soybean management system: I. Cultivar response. *Journal of Production Agriculture*. 4, 571-579
- Lundkvist, A. & Verwijst, T. (2011). Weed Biology and Weed Management in Organic Farming, Research in Organic Farming. Nokkoul, R. (ed.). [Online], Available at <http://www.intechopen.com/books/research-in-organic-farming/weed-biology-and-weed-management-inorganic-farming> Accessed (09-05-13)

- Lungu, O.I.M. & Dynoodt, R.F.P. (2008). Acidification from long-term use of urea and its effect on selected soil properties. *African Journal of Food, Agriculture, Nutrition and Development*. 8 (1), 63-76.
- Lupwayi, N.Z., Kennedy, A.C. & Chirwa, R.M. (2011). Grain legume impacts on soil biological processes in sub-Saharan Africa. *African Journal of Plant Science*. 5 (1), 1-7.
- Lutman, P.J.W., Moss, S.R., Cook, S. & Welham, S.J. (2013). A review of the effects of crop agronomy on the management of *Alopecurus myosuroides*. *Weed Research*. 53 (5), 299-313.
- Machado, S. (2009). Does intercropping have a role in modern agriculture? *Journal of Soil and Water Conservation*. 64 (2), 55A-57A
- Machado, S., Petrie, S., Rhinhart, K. & Qu, A. (2007). Long-term continuous cropping in the Pacific Northwest: Tillage and fertilizer effects on winter wheat, spring wheat, and spring barley production. *Soil and Tillage Research*. 94, 473-481.
- MAFF. (1968). *Century of Agricultural Statistics. Great Britain 1866-1966*. HMSO. Ministry of Agriculture Fisheries and Food (MAFF). London, UK.
- MAFF. (1976). *Manual of plant growth stages and disease assessment keys*. Fisheries and food. (Looseleaf binder). Ministry of Agriculture Fisheries and Food (MAFF). London, UK.
- MAFF. (1988). *Agricultural land classification of England and Wales*. Ministry of Agriculture Fisheries and Food (MAFF). pp, 52.
- Magdoff, F., Lanyon, I. & Liebhardt, B. (1997). Nutrient cycling, transformation and flows: Implications for a more sustainable agriculture. *Advance Agronomy*. 60, 1-73.
- Mahler, R.L., Koehler, F.E. & Lutcher, L.K. (1994). Nitrogen source, timing of application and placement: Effects on winter wheat production. *Agronomy Journal*. 86, 637-642.
- Mahn, E.G. (1984). Structural changes of weed communities and populations. *Vegetatio*. 58, 79-85.
- Małecka I. & Blecharczyk, A. (2008). Effect of tillage systems, mulches and nitrogen fertilization on spring barley (*Hordeum vulgare*). *Agronomy Research*. 6 (2), 517-529.
- Malhi, S.S. Johnston, A.M., Schoenau, J.J., Wang, Z.H. & Vera, CL. (2007). Seasonal biomass accumulation and nutrient uptake of pea and lentil on a black chernozem soil in Saskatchewan. *Journal of Plant Nutrition*. 30, 721-737.

- Malhi, S.S. & Kutcher, H.R. (2007). Small grains stubble burning and tillage effects on soil organic C and N, and aggregation in north-eastern Saskatchewan. *Soil and Tillage Research*. 94, 353–361.
- Malhi, S.S., McAndrew, D.W. & Carter, M.R. (1992). Effect of tillage and N fertilization of a Solonchic soil on barley production and some soil properties. *Soil and Tillage Research*. 22, 95-107.
- Marks, H.F. (1989). *A hundred years of British food and farming. A statistical Survey*. Britton, D.K. (ed.). London, UK.
- Marshall, E.J.P., Brown, V.K., Boatman, N.D., Lutman, P.J., Squire, G.R. & Ward, L.K. (2003). The role of weeds in supporting biological diversity within crop fields. *Weed Research*. 43, 77-89.
- Marshall, E.P.J. & Nowakowski, J.B. (1996). Interactions between cutting and a graminicide on a newly-sown grass and wild flower field margin strip. *Aspect of Applied Biology*. 44, 307-312.
- Marshall, T.J. & Holmes, J.W. (1988). *Soil physics*. 2nd edition. Cambridge University Press, Cambridge.
- Martin-Rueda, I., Muñoz-Guerra, L.M., Yunta, F., Esteban, E., Tenorio, J.L. & Lucena J.J. (2007). Tillage and crop rotation effects on barley yield and soil nutrients on a Calcicortidic Haplixeralf. *Soil and Tillage Research*. 92, 1-9.
- Martínez, E., Fuentes, J.P., Silva, P., Valle, S. & Acevedo, E. (2008). Soil physical properties and wheat root growth as affected by no-tillage and conventional tillage systems in a Mediterranean environment of Chile. *Soil and Tillage Research*. 99, 232-244.
- Mas, M.T. & Verdú, A.M.C. (2003). Tillage system effects on weed communities in a 4-year crop rotation under Mediterranean dryland conditions. *Soil and Tillage Research*. 74, 15-24.
- Maskell, L.C., Smart, S. M., Bullock, J.M., Thompson, K. & Stevens, C.J. (2010). Nitrogen deposition causes widespread loss of species richness in British habitats. *Global Change Biology*. 16, 671–679.
- Mavunganidze, Z., Madakadze, I.C., Nyamangara, J. & Mafongoya, P. (2014). The impact of tillage system and herbicides on weed density, diversity and yield of cotton (*Gossipium hirsutum* L.) and maize (*Zea mays* L.) under the smallholder sector. *Crop Protection*. 58, 25-32.

- McConkey, B.G., Campbell, C.A., Zentner, R.P., Dyck, F.B. & Selles, F. (1996). Long-term tillage effects on spring wheat production on three soil textures in the Brown soil zone. *Canadian Journal of Plant Science*. 74 (4), 747-756.
- McConkey, B.G., Curtin, D., Campbell, C.A., Brandt, S.A. & Selles, F. (2002). Crop and soil nitrogen status of tilled and no-tillage systems in semiarid regions of Saskatchewan. *Canadian Journal of Soil Science*. 82, 489-498.
- Mc-Master, G.S., Wilhelmm G.S., Bartling, P.N.S. (1994). Irrigation and culm contribution to yield and yield components of winter wheat. *Agronomy Journal*. 86, 1123-1127.
- McNeal, F.H., McGuire, C.F. & Klindworth, D.L. (1982). Agronomic and quality characteristics of spring wheat lines selected for protein content and protein yield. *Euphytica*. 31, 377-381.
- McVal, K.A., Budde, J.A., Fabrizzi, K., Mikha, M.M., Rice, C.W., Schlegel, A.J., Peterson, D.E., Sweeney, D.W. & Thompson, C. (2006). Management effects on soil physical properties in long-term studies in Kanas. *Soil Science Society of America Journal*. 70, 434-438.
- Mehdi, B.B., Madramooto, C.A. & Mehuys, G.R. (1999). Yield and nitrogen content of corn under different tillage practices. *Agronomy Journal*. 91, 631-636.
- Menalled, F.D., Gross, K.L. & Hammond, M. (2001). Weed above ground and seed bank community responses to agricultural management systems. *Ecological Applications*. 11, 1586-1601.
- Mercier, S. & Hyberg, B. (1995). Grain quality revised. *American Agricultural Economic Asssociation. Choises*. 1, 35-38.
- Mester, T.C. & Buhler, D.D. (1991). Effects of soil temperature, seed depth, and cyanazine on giant foxtail (*Setaria faberi*) and velvetleaf (*Abutilon theoprastris*) seedling development. *Weed Science*. 39, 204-209.
- Miao, Y.F., Wang, Z.H. & Li, S.X. (2015). Relation of nitrate N accumulation in dryland soil with wheat response to N fertilizer. *Field Crops Research*. 170, 119-130.
- Mitchell, J.P., Jackson, L. & Miyao, G. (2004). *Minimum tillage vegetable crop production in California*. Oakland: University of California Agriculture and Natural Resources Publication. 8132. UC ANR CS. [Online] Available from <http://anrcatalog.ucdavis.edu/VegetableCropProductioninCalifornia/8132.aspx>. Accessed (26-08-13)
- Moakes, S. & Lampkin, N. (2011). *Organic Farm Incomes in England and Wales 2009/2010*. IBERS. Aberystwyth University, UK.

- Mogensen, V.O., Jensen, H.E. & Rab, A. (1985). Grain yield, yield components drought sensitivity, and water use efficiency of spring wheat subjected to water stress at various growth stages. *Irrigation Science*. 6, 131-140.
- Mohtasebi, S.S., Lar M.B., Safa, M. & Chaichi, M.R. (2008). Comparison of direct and indirect energy coefficients for seeding and fertilizing in irrigated wheat production. *World Applied Science*. 3 (3), 353-358.
- Moll, R.H., Kamprath, E.J. & Jackson, W.A. (1982). Analysis and interpretation of factors which contribute to efficiency of nitrogen utilization. *Agronomy Journal*. 74 (3), 562-564.
- Morris, N.L., Miller, P.C.H, Orson, J.H. & Froud-Williams, R.J. (2010). The adoption of non-inversion tillage systems in the United Kingdom and the agronomic impact on soil, crops and the environment- A review. *Soil and Tillage Research*. 108,1-15.
- Moss, S.R. (2013). *Black-grass (Alopecurus myosuroides)*. Rothamsted technical publication. [Online]. Available from: <http://www.rothamsted.ac.uk/sites/default/files/attachments/2014-01-09/2%20Black-grass%20leaflet%20revised%20easy%20read%20format%2028May13.pdf> Accessed (02-12-14)
- Moss, S.R., Storkey, J., Cussans, J.W., Perryman, S.A.M. & Hewitt, M.V. (2004). The Broadbalk long-term experiment at Rothamsted: what has it told us about weeds? *Weed Science*. 52, 864–873.
- Mrabet, R. (2011). Effects of residue management and cropping systems on wheat yield stability in a semiarid Mediterranean clay soil. *American Journal of Plant Science*. 2, 202-216.
- Munkholm, L.J., Schjønning, P., Rasmussen, K.J. & Tanderup, K. (2003). Spatial and temporal effects of direct drilling on soil structure in the seedling environment. *Soil and Tillage Research*. 71, 163-173.
- Murillo, J.M., Moreno, F., Pelegrín, F. & Fernández, J.E. (1998). Response of sunflower to traditional and conservation tillage under rainfed conditions in southern Spain. *Soil and Tillage Research*. 49, 233-241.
- Myrbeck, A., Stenberg, M., Arvidsson, J. & Rydberg, T. (2012). Effects of autumn tillage of clay soil on mineral N content, spring cereal yield and soil structure over time. *European Journal of Agronomy*. 37 (1), 96-104.
- NABIM. (2014). *Wheat Guide*. National Association of British and Irish Millers (NABIM). London, UK.

- Neugschwandtner, R.W., Liebhard, P., Kaul, H.P. & Wagentrist, H. (2014). Soil chemical properties as affected by tillage and crop rotation in a long-term field experiment. *Plant Soil Environment*. 60 (2), 57-62.
- Nix, J. (2012). *Farm Management Pocketbook*. 43th edition. Agro Business Consultants.
- Nix, J. (2013). *Farm Management Pocketbook*. 44th edition. Agro Business Consultants.
- Norsworthy, J.K. (2008). Effect of tillage intensity and herbicide programs on changes in weed species density and composition in the southeastern coastal plains of the United States. *Crop Protection*. 27, 151-160.
- Novoa, B. & Loomis, R.S. (1981). Nitrogen and plant production. *Plant and Soil*. 58, 177–204.
- O'Donovan, J.T., Mandrew, D.W. & Thomas, A.G. (1997). Tillage and nitrogen influence weed population dynamics in barley. *Weed Technology*. 11, 502–509
- Ochoa-Hueso, R. & Manrique, E. (2010). Nitrogen fertilization and water supply affect germination and plant establishment of the soil seed bank present in a semi-arid Mediterranean scrubland. *Plant Ecology*. 210 (2), 263-273.
- Ofori, F. & Stern, W.R. (1987). Cereal-legume intercropping systems. *Advances in Agronomy*. 41, 41-90.
- Olesen, J.E., Trnka, M., Kersebaum, K.C., Skjelvåg, A.O., Seguin, B., Peltonen-Sainio, P., Rossi, F., Kozyra, J. & Micalle, F. (2011). Impact and adaptation of European crop production systems to climate change. *European Journal of Agronomy*. 34, 96-112.
- Olsen, S.R., Cole, C.V., Watanabe, F.S. & Dean, L.A. (1954). Estimation of available phosphorus in soils by extraction with sodium bicarbonate. in, Banderis, A.D., Barter, D.H. & Anderson, K. (eds.) *U.S. Department of Agriculture Circular No. 939*. Agricultural and Advisor.
- Olson, R.A., Frank, K.D., Deibert, E.J., Dreier, A.F., Sander, D.H. & Johnson, V.A. (1976). Impact of residual mineral N in soil on grain protein yields of winter wheat and corn. *Agronomy Journal*. 68, 769-772.
- Oorts, K., Laurent, F., Mary, B., Thiébeau, P., Labreuche, J. & Nicolardot, B. (2007). Experimental and simulated soil mineral N dynamics for long-term tillage systems in northern France. *Soil and Tillage Research*. 94 (2), 441-456.

- Ordoñez-Fernández, R., González- Fernández, P., Giráldez-Cervera, J.V. & Perea-Torres, F. (2007). Soil properties and crop yields after 21 years of direct drilling trials in southern Spain. *Soil and Tillage Research*. 94, 47-54.
- Ortiz-Monasterio, J.I., Sayre, K.D., Rajaram, S. & McMahon, M. (1997) Genetic progress in wheat yield and nitrogen use efficiency under four nitrogen rates. *Crop Science*. 37, 898-904.
- Osborne, T.B. (1907). The proteins of the wheat kernel. Carnegie Institute of Washington. Washington DC, USA. pp. 103.
- Otteson, B.N., Mergoum, M., Ransom, J.K. & Schatz, B. (2008). Tiller contribution to spring wheat yield under varying seeding and nitrogen management. *Agronomy Journal*. 100 (2), 406-413.
- Ozkan, B., Akcaoz, H. & Fert, C. (2004). Energy input–output analysis in Turkish agriculture. *Renewable Energy*. 29, 39–51.
- Ozkan, B., Fert, C. & Karadeniz, C.F. (2007). Energy and cost analysis for greenhouse and open-field grape production. *Energy*. 32, 1500-1504.
- Ozpinar, S. (2006). Effects of tillage systems on weed population and economics for winter wheat production under the Mediterranean dryland conditions. *Soil and Tillage Research*. 87, 1-8.
- Ozpinar, S. & Çay, A. (2005). Effect of minimum and conventional tillage systems on soil properties and yield of winter wheat (*Triticum aestivum* L.) in clay-loam in the Canakkale Region. *Turkish Journal of Agriculture*. 29, 9-18.
- Palta, J.A. & Fillery, I.R.P. (1993). Post-anthesis remobilisation and losses of nitrogen in wheat in relation to applied nitrogen. *Plant and Soil*. 155, 179-181.
- Papadakis, J. S. (1941). Small grains and winter legumes grown mixed for grain production. *Journal of the American Society of Agronomy*. 33, 504-511.
- Parchoma, G. (2002). *A Guide to Weeds in British Columbia*. Ministry of Agriculture, Food and Fisheries. British Columbia.
- Parry, M.A.J., & Hawkesford, M.J. (2012). An integrated approach to crop genetic improvement. *Journal of Integrated Plant Biology*. 54 (4). 250-259.
- Parry, M.A.J., & Hawkesford, M.J. (2010). Food security: increasing yield and improving resource use efficiency. *Proceedings of the Nutrition Society*. 69, 1–9.
- Pearman, I., Thomas, S.M. & Thorne, G.N. (1977). Effect of nitrogen fertilizer on growth and yield of spring wheat. *Annals of Botany*. 41 (1), 93-108.

- Pearman, I., Thomas, S.M. & Thorne, G.N. (1978). Effect of nitrogen fertilizer on growth and yield of semi-dwarf and tall varieties of winter wheat. *Journal of Agricultural Science*. 91, 31-45.
- Peigné, J., Ball, B.C., Roger-Estrade, J. & David, C. (2007). Is conservational tillage suitable for organic farming? A review. *Soil Use and Management*. 23 (2), 129-144.
- Peigné, J., Casagrande, M., Payet, V., David, C., Sans, F.X., Blanco-Moreno, J.M., Cooper, J., Gascoyne, K., Antichi, D., Bàrberi, P., Bigongiali, F., Surböck, A., Kranzler, A., Beckman, A., Willekens, K., Luik, A., Matt, D., Grosse, M., Heß J., Clerc, M., Dierauer, H., & Mäder, P. (2015). How organic farmers practice conservation agriculture in Europe. *Renewable Agriculture and Food Systems*. [Online] Available from <http://dx.doi.org/10.1017/S1742170514000477> Accessed (02-02-15)
- Pekrun, C., Kaul, H.P. & Claupein, W. (2003). Soil tillage for sustainable nutrient management. In El Titi, A. (ed.). *Soil Tillage in Agroecosystems*. CRC Press, Boca Raton, 38-113.
- Pelegriñ, F., Moreno, F., Martin-Aranda, J. & Camps, M. (1990). The influence of tillage methods on soil physical properties and water balance for a typical crop rotation in SW Spain. *Soil & Tillage Research*. 16, 345-358.
- Peltonen-Sainio, P., Jauhiainen, L., Rajala, A. & Muurinen, S. (2009). Tiller traits of spring cereals under tiller-depressing long day conditions. *Field Crops Research*. 113, 82-89.
- Penny, A., Widdowson, F.N. & Jenkyn, J.E. (1978). Spring top-dressings of 'Nitro-Chalk' and late sprays of liquid N-fertilizer and a broad spectrum fungicide for consecutive crops of winter wheat at Saxmundham, Suffolk. *Journal of Agricultural Science*. Cambridge. 90, 509-516.
- Pervanchon, F., Bockstaller, C. & Girarding, P. (2002). Assessment of energy use in arable farming systems by means of an agro-ecological indicator: the energy indicator. *Agricultural Systems*. 72, 149-172.
- Peters, K., Breitsameter, L. & Gerowitt, B. (2014). Impact of climate change on weeds in agriculture: a review. *Agronomy for Sustainable Development*. 34, 707-721.
- Peters, N.C.B., Froud-Williams, R.J. & Orson, J.H. (1993). The rise of barren brome *Bromus sterilis* in UK cereal crops. In *Proceedings of the British Crop Protection Conference on Weeds*. 2. Brighton, UK, 22–25 November 1993. 773–779.
- Pierce, F.J. & Rice, C.W. (1988). Crop rotation and its impact of efficiency of water and nitrogen use. In Hargrove (ed.). *Cropping strategies for efficient use of water and nitrogen*. ASA, Special publication. No 15, Wisconsin. 101-113.

- Pietola, L.M. (2005). Root growth dynamics of spring cereals with discontinuation of mouldboard ploughing. *Soil and Tillage Research*. 80, 130-114.
- Pietola, L. & Tanni, R. (2003). Response of seedbed physical properties, soil N and cereal growth to peat application during transition to conservation tillage. *Soil and Tillage Research*. 74, 65–79.
- Piggin, C., Haddad, A., Khalil, Y., Loss, S. & Mustafa, P. (2015). Effects of tillage and time of sowing on bread wheat, chickpea, barley and lentil grown in rotation in rainfed systems in Syria. *Field Crops Research*. 173, 57-67.
- Pimentel, D. (1980). *Handbook of energy utilization in agriculture*. CRC press. Boca Raton, FL, USA.
- Pimentel, D., Hepperly, P., Hanson, J., Douds, D. & Seidel, R. (2005). Environmental, energetic, and economic comparisons of organic and conventional farming systems. *BioScience*. 55, 573–582.
- Piringer, G. & Steinberg, L.J. (2006). Reevaluation of energy use in wheat production in the United States. *Journal of Industrial Ecology*. 10 (1-2), 149-167.
- Pollard, F., Elliot, J.G., Ellis, F.B. & Barnes, B.T. (1981). Comparison of direct drilling, reduced cultivation and ploughing on the growth of cereals. *Journal of Agricultural Science*. 97 (3), 677-684.
- Power, J.F. & Alessi, J. (1978). Tiller development and yield of standard and semidwarf spring wheat varieties as affected by nitrogen fertilizer. *Journal of Agricultural Science*. 90, 97-108.
- Powlson, D.S., Hart, P.B.S., Poulton, P.R., Johnston, A.E. & Jenkinson, D.S. (1992). Influence of soil type, crop management and weather on the recovery of ¹⁵N-labelled fertilizer applied to winter wheat in spring. *Journal of Agricultural Science*. 118, 83–100.
- Prasad, P.V.V., Pisipati, S.R., Ristic, Z., Bukovnik, U. & Fritz, A.K. (2008). Impact of night time temperature on physiology and growth of spring wheat. *Crop Science*. 48, 2372-2380.
- Pridham, J.C. & Entz, M.H. (2008). Intercropping spring wheat with cereal grains, legumes and oilseeds fails to improve productivity under organic management. *Agronomy Journal*. 100 (5), 1436-1442.
- Pushman, F.M. & Bingham, J. (1975). Components of test weight of ten varieties of winter wheat grown at two rates of nitrogen fertiliser application. *Journal of Agricultural Science*. 85, 559-563.

- Qian, B., De Jong, R. & Gameda, S. (2009). Multivariate analysis water related agroclimatic factors affecting spring wheat yields on the Canadian prairies. *European Journal of Agronomy*. 30, 140-150.
- Qiao, J., Yang, L., Yan, T., Xue, F. & Zhao, D. (2013). Ride dry matter and nitrogen accumulation, soil mineral N around root and N leaching, with increasing application rates of fertilizer. *European Journal of Agronomy*. 49, 93-103.
- Quaedvlieg, W., Kema, G.H.J., Groenewald, J.Z., Verkley, G.J.M., Seifbarghi, S., Razavi, M., Mirzadi Gohari, A., Mehrabi, R., Crous, P.W. (2011). *Zymoseptoria* gen. nov.: a new genus to accommodate *Septoria*-like species occurring on graminicolous host. *Persoonia*. 26, 57-69.
- Radford, B.J., Gibson, G., Nielsen, R.G.H., Butler, D.G., Smith, G.D. & Orange, D.N. (1992). Fallowing practices, soil water storage, plant-available soil nitrogen accumulation and wheat performance in south west Queensland. *Soil and Tillage Research*. 22, 73-93.
- Rao, A.C.S., Smith, J.L., Jandhyala, V.K., Papendick, R.L. & Parr, J.F. (1993). Cultivar and climatic effects on the protein content of soft white winter wheat. *Agronomy Journal*. 85, 1023-1028.
- Rasmussen, P.E. (1999). Impact of ploughless soil tillage on yield and soil quality. A Scandinavian review. *Soil and Tillage Research*. 53, 3-14.
- Rasmussen, P.E. (1993). Surface residue and nitrogen fertilization effects on no-till wheat. In Barrow, N.J. (ed.). *Plant Nutrition-from Genetic Engineering to Field Practice*. Kluwer Academic Publishers. 555-558.
- Rasmussen, P.E., Albercht, S.L. & Smith, R.W. (1998). Soil C and N changes under tillage and cropping systems in semi-arid Pacific Northwest agriculture. *Soil and Tillage Research*. 47, 197-205.
- Rasmussen, P.E. & Collins, H.P. (1991). Long-term impacts of tillage, fertilizer and crop residue on soil organic matter in temperate semi-arid regions. *Advance Agronomy*. 45, 93-134.
- Rasmussen, P.E., Rickman, R.W. & Klepper, B.L. (1997). Residue and fertility effects on yield of no-till wheat. *Agronomy Journal*. 89, 563-567.
- Rathke, G.W., Wienhold, B.J., Wilhelm, W.W. & Diepenbrock, W. (2007). Tillage and rotation effect on corn-soybean energy balances in Eastern Nebraska. *Soil and Tillage Research*. 97 (1), 60-70.
- Raun, W.R., & Johnson, G.V. (1999). Improving nitrogen use efficiency for cereal production. *Agronomy Journal*. 91, 357-363.

- Raun, W.R., Solie, J.B. & Stone, M.L. (2010). Independence of yield potential and crop nitrogen response. *Precision Agriculture*. 12,508–518.
- Refsgaard, K., Halberg, N. & Kristensen, E.S. (1998). Energy utilization in crop and dairy production in organic and conventional livestock production systems. *Agricultural Systems*. 57 (4), 599-630.
- Reicosky, D.C., Evans, S.D., Cambardella, C.A., Allmaras, R.R., Wilts, A.R. & Huggins, D.R. (2002). Continuous corn with moldboard tillage: Residue and fertility effects on soil carbon. *Journal of Soil and Water Conservation*. 57 (5), 277-284.
- Reicosky, D.C., Kemper, W.D., Langdale, G.W., Douglas-Jr, C.L. & Rasmussen, P.E. (1995). Soil organic matter changes resulting from tillage and biomass production. *Journal of Soil and Water Conservation*. 50, 253–261.
- Reicosky, D.C., Reeves, D.W., Prior, S.A., Runion, G.B., Rogers, H.H. & Raper, R.L. (1999). Effects of residue management and controlled traffic on carbon dioxide and water loss. *Soil and Tillage Research*. 52, 153-165.
- Reynolds, M., Bonnett, D., Chapman, S.C., Furbank, R.T., Manes, Y., Mather, D.E. & Parry, M.A.J. (2011). Raising yield potential of wheat: (I) Overview of a consortium approach and breeding strategies. *Journal of Experimental Botany*. 62, 439–452.
- Reynolds, M.P. & Bourlaug, N.E. (2006a). Applying innovations and new technologies for international collaborative wheat improvements. Centenary Review. *Journal of Agricultural Science*. 144, 95-110.
- Reynolds, M.P. & Bourlaug, N.E. (2006b). Impacts of breeding on international collaborative wheat improvement. Centenary Review. *Journal of Agricultural Science*. 144, 3-17
- Reynolds, M., Foulkes, J., Furbank, R., Griffiths, S., King, J., Murchie, E., Parry, M. & Slafer, G. (2012). Achieving yield gains in wheat. *Plant, Cell & Environment*. 35, 1799–1823.
- Reynolds, M., Foulkes, M.J., Slafer, G.A., Berry, P., Perry, M.A.J., Snape J.W. & Angus, W.J. (2009). Raising yield potential in wheat. *Journal of Experimental Botany*. 60 (7), 1899-1918.
- Reynolds, M.P., Rajaram, S. & Sayre, K.D. (1999). Physiological and genetic changes of irrigated wheat in the post-Green Revolution period and approaches for meeting projected global demand. *Crop Science*. 39, 1611–1621.
- Reynolds, M.P., Sayre, K.D. & Vivar, H.E. (1994). Intercropping wheat and barley with N-fixing legume species: a method for improving ground cover, N-use efficiency and productivity in low input systems. *Journal of Agricultural Science*. 123, 175-183.

- Rhoton, F.E. (2000). Influence of time on soil response to no-till practices. *Soil Science Society of America Journal*. 77, 700-709.
- Rice, M.C. (1984). *Allelopathy*. 2nd edition. Orlando, FL, Academic Press.
- Rieger, S., Richner, W., Streit, B., Frossard, E. & Liedgens, M. (2008). Growth, yield, and yield components of winter wheat and the effects of tillage intensity, preceding crops and N fertilisation. *European Journal of Agronomy*. 28, 405-411.
- Riley, H.C.F. (1998). Soil mineral-N and N-fertilizer requirements of spring cereals in two long-term tillage trials on loam soil in southeast Norway. *Soil and Tillage Research*. 48, 265-274.
- Riley, H.C.F., Bleken, M.A., Abrahamsen, S., Bergjord, A.K. & Bakken, A.K. (2005). Effects of alternative tillage systems on soil quality and yield of spring cereals on silty clay loam and sandy loam soils in the cool, wet climate of central Norway. *Soil and Tillage Research*. 80, 79–93.
- Riley, H., Børresen, T., Ekeberg, E. & Rydberg, T. (1994). Trends in reduced tillage research and practice in Scandinavia. In Carter, M.R. (ed.). *Conservation Tillage in Temperate Agroecosystems*. Lewis Publishers, pp. 23-45.
- Robinson, F.E., Cudney D.W. & Lehman, W.F. (1979). Nitrogen fertilizer timing, irrigation, protein and yellow berry in durum wheat. *Agronomy Journal*. 61, 755-759.
- Rocheouste, J-F., Dargusch, P., Cameron, D. & Smith, C. (2015). An analysis of the socio-economic factors influencing the adoption of conservation agriculture as a climate change mitigation activity in Australian dryland grain production. *Agricultural Systems*. 135. 20-30.
- Rosenfeld, A., Rayns, F. & Wilkinson, I. (2011). *Sort out your soil: A practical guide to green manure*. Milner, I. (ed.). Cotswold Grass Seeds Direct. [Online] Available from <https://www.cotswoldseeds.com/files/cotswoldseeds/Sort%20Out%20Your%20Soil.pdf> Accessed (23-10-13)
- Ross, S.M., King, J.R., Izaurrealde, R.C. & O'Donovan, J.T. (2001). Weed suppression by seven clover species. *Agronomy Journal*. 93, 820-827.
- Rossner, H., Ritz, C. & Astover, A. (2014). Optimisation of fertiliser rates in crop production against energy use indicators. *European Journal of Agronomy*. 55, 72-76.
- Rozas, S., Echeverria, H.E. & Studdert, G.A. (1999). No-tillage maize nitrogen uptake and yield: Effect of urease inhibitor and application time. *Agronomy Journal*. 91, 950-955.

- Ryan, M.H., Derick, J.W. & Dann, P.R. (2004). Grain mineral concentration and yield of wheat grown under organic and conventional management. *Journal of the Science, Food and Agriculture*. 84, 207-216.
- Rusinamhodzi, L., Corbeels, M., Wijk, M.T., Rufino, M.C., Nyamangara, J. & Giller, K.E. (2011). A meta-analysis of long-term effects of conservation agriculture on maize grain yield under rain-fed conditions. *Agronomy for Sustainable Development*. 31 (4), 657-673.
- Sadeghi, A.M., Isensee, A.R. & Shelton, D.R. (1998). Effect of tillage age on herbicide dissipation: a side-by side comparison using microplots. *Soil Science*. 163, 883-890.
- Safa, M., Samarasinghe, S. & Mohssen, M. (2011). A field study energy consumption in wheat production in Canterbury New Zealand. *Energy Conversion and Management*. 52, 2526-2532.
- Sainju, U.M., Lenssen, A.W., Caesar-TonThat, T., Jabro, J.D., Lartey, R.T., Evans, R.G. & Allen, B.T. (2012). Tillage, crop rotation, and cultural practice effects on dryland soil carbon fractions. *Open Journal of Soil Science*. 2, 242-255.
- Samuel, A. (2005). Cropping in the UK. In Soffe, R. (ed.). *The countryside notebook*. Oxford, UK.
- Sánchez-Girón V., Serrano A., Hernanz J.L., Navarrete L. (2004). Economic assessment of three long-term tillage systems for rainfed cereal and legume production in semiarid central Spain. *Soil and Tillage Research*. 78, 35-44.
- Santín-Montoyá, M.I., Zambrana, E., Fernández-Getino, A.P. & Tenorio, J.L. (2014). Dry pea (*Pisum sativum* L.) yielding and weed infestation response, under different tillage conditions. *Crop Protection*. 65, 122-128.
- Sarandon, S.J. & Caldiz, D.O. (1990). Effects of varying nitrogen supply at different growth stages on nitrogen uptake and nitrogen partitioning efficiency in two wheat cultivars. *Nutrient Cycling in Agroecosystems*. 22, 21-27.
- Sarandon, S.J. & Gianibelli, M.C. (1990). Effect of foliar urea spraying and nitrogen application at sowing upon dry matter and nitrogen distribution in wheat (*Triticum aestivum* L.). *Agronomie*. 10, 183-189.
- Sartori, L., Basso, B., Bertocco, M. & Oliviero, G. (2005). Energy Use and Economic Evaluation of a Three Year Crop Rotation for Conservation and Organic Farming in NE Italy. *Biosystems Engineering*. 91 (2), 245–256.
- Sayili, M., Akca, H. & Önen, H. (2006). Economic analysis of herbicide usage in wheat fields. *Journal of Plant Diseases and Protection*. XX, 755-760.

- Schomberg, H.H., Ford, P.B., Hargrove, W.L. (1994). Influence of crop residues on nutrient cycling and soil chemical properties. In Unger, P.W. (ed.). *Managing Agricultural Residues*. CRC Press Inc, Boca Raton, FL, USA. 1-6.
- Schroder, J.L., Zhang, H.L., Girma, K., Raun, W.R., Penna, C.J. & Payton, M.E. (2011). Soil acidification from long-term use of nitrogen fertilizers on winter wheat. *Soil Science Society of America Journal*. 75, 957–964.
- Schulten, H-R. & Schnitzer, M. (1998). The chemistry of soil organic nitrogen: A review. *Biology and Fertility of Soils*. 26, 1-15.
- Schwenke, G., Herridge, D., McMullen, G., Leedham, K. & Haigh, B. (2011). Nitrous oxide emissions from dryland cropping in Northwest NSW. in Rural Climate Change Solutions Symposium. University of New England, Armidale, NSW, Australia.
- Semenov, M.A., Jamieson, P.D. & Martre, P. (2007). Deconvoluting nitrogen use efficiency in wheat: A simulation study. *European Journal of Agronomy*. 26, 283-294.
- Semenov, M.A. & Shewry, P.R., (2011). Modelling predicts that heat stress, not drought, will increase vulnerability of wheat in Europe. *Scientific Reports*. 1, 66.
- Semenov, M.A., Stratonovitch, P., Alghabari, F. & Gooding, M.J. (2014). Adapting wheat in Europe for climate change. *Journal of Cereal Science*. 59, 245-256.
- Sexsmith, J.J. & Russell, G.C. (1963). Effect of nitrogen and phosphorus fertilization on wild oats and spring wheat. *Canadian Journal of Plant Science*. 43, 64-69.
- Shafi, M., Bakht, J., Jan, M.T. & Shah, Z. (2007). Soil C and N dynamics and maize (*Zea may* L.) yield as affected by cropping systems and residue management in North-western Pakistan. *Soil and Tillage Research*. 94, 520- 529.
- Sharma, P., Abrol, V. & Sharma, R.K. (2011). Impact of tillage and mulch management on economics, energy requirement and crop performance in maize– wheat rotation in rainfed subhumid inceptisols, India. *European Journal of Agronomy*. 34, 46–51
- Sheibani, S. & Ghadiri, H. (2012). Integration effects of split nitrogen fertilization and herbicide application on weed management and wheat yield. *Journal of Agriculture, Science and Technology*. 14, 77-86.
- Shinners, K.J., Nelson, W.S. & Wang, R. (1994). Effects of residue-free band width on soil temperature and water content. *Transactions of the American Society of Agricultural Engineers* (Trans ASAE). 37 (1), 39–49.
- Shrestha, A., Knezevic, Z., Roy, R.C., Ball-Coelho, B. & Swanton, C.J. (2002). Effect of tillage, cover crop and crop rotation on the composition of weed flora in sandy soil. *Weed Research*. 42, 76-87.

- Sieling, K. Schröder, H. Finck, M. & Hanus, H. (1998). Yield, N uptake, and apparent N-use efficiency of winter wheat and winter barley grown in different cropping systems. *Journal of Agricultural Science*. 131, 375-387.
- Siemens, M.C., Wilkins, D.E. & Correa, R.F. (2004). Development and evaluation of a residue management wheel of hoe-type no-till drills. *Transactions of the American Society of Agricultural Engineers* (Trans ASAE). 47 (2), 397-404.
- Silgram, M. & Shepherd, M. (1999). The effects of cultivation on soil nitrogen mineralisation. *Advances in Agronomy*. 65, 267-311.
- Simpson, R.J., Lambers, H. & Dalling, M.J. (1983). Nitrogen redistribution during grain growth in wheat (*Triticum aestivum* L.). IV. Development of a quantitative model of the translocation of nitrogen to the grain. *Plant Physiology*. 71, 7-14.
- Singh, B., Chanasyk, D.S. & McGill, W.B. (1998). Soil and water regimen under barley with long-term tillage-residue systems. *Soil and Tillage Research*. 45. 59-74.
- Singh, H., Mishra, D. & Nahar, N.M. (2002). Energy use pattern in production agriculture of a typical village in Arid Zone India—Part I. *Energy Conversion Management*. 43 (16), 2275–2286.
- Šíp, V., Vavera, R., Chrpová, J., Kusá, H. & Růžek, P. (2013). Winter wheat yield and quality related to tillage practice, input level and environmental conditions. *Soil and Tillage Research*. 132, 77-85.
- Six, J., Conant, R.T., Paul, E.A. & Paustian, K. (2002). Stabilization mechanisms of soil organic matter: Implications for C-saturation of soil. *Plant and Soil*. 241, 155-176.
- Six, J., Elliot, E.T. & Paustian, K. (2000). Soil macroaggregate turnover and microaggregate formation a mechanics for C sequestration under no-tillage agriculture. *Soil Biology & Biochemistry*. 32, 2099-2103.
- Slafer, G.A., Satorre, E.H. & Andrade, F.H. (1994). Increases in grain yield in bread wheat and associated physiological changes. In *Genetic improvement of field crops*. Slafer, G.A. (ed.). 1-68.
- SMI. (2003). A guide to managing crop establishment and improved soil management for agronomic and environmental gain. *Soil Management Initiative* (SMI). UK.
- Smith, G. & Gooding, M. (1999). Models of wheat grain quality considering climate, cultivar and nitrogen effects. *Agricultural and forest meteorology*. 94 (3), 159-170.
- Smith, P., Martino, D., Cai, Z., Gwary, D., Janzen, H.H., Kumar, P., McCarl, B., Ogle, S., O'Mara, F., Rice, C., Scholes, R.J., Sirotenko, O., Howden, M., McAllister, T., Pan, G., Romanenkov, V., Schneider, U., Towprayoon, S., Wattenbach, M. & Smith J.U.

(2007). Greenhouse gas mitigation in agriculture. *Philosophical Transactions of the Royal Society B*. 363. Singh, B., Chanasyk, D.S. & McGill, W.B. (1998). Soil water regime under barley with long-term tillage-residue systems. *Soil and Tillage Research*. 45, 59-74

Smith, S.P., Davies, W.P., Bulman, C. & Evans, E.J. (1990). Response of grain protein in winter wheat to nitrogen. In *Aspect of Applied Biology. Cereal Quality II*. Association of Applied Biologists, Warwick. 25, 255-260.

Smykal, P., Coyne, C.J., Ambrose, M.J., Maxted, N., Schaefer, H., Blair, M.W., Berger, J., Greene, S.L., Nelson, M.N., Besharat, N., Vymyslický, T., Toker, C., Saxena, R.K., Roorkiwal, M., Pandey, M.K., Hu, J., Li, Y.H., Wang, L.X., Guo, Y., Qiu, L.J., Redden, R.J. & Varshney, R.K. (2015). Legume crops phylogeny and genetic diversity for science and breeding. *Critical Reviews in Plant Sciences*. 34 (1-3), 43-104.

Soane, B.D., Ball, B.C., Arvidsson, J., Basch, G., Moreno, F. & Roger-Estrade, J. (2012). No-till in northern, western and south-western Europe. A review of problems and opportunities for crop production and the environment. *Soil Tillage and Research*. 118, 66–87.

Soffe, R.J. (ed.) (2003). *The agricultural notebook*. 20th edition. Blackwell Science.

Soil Association. (2011). *The Lazy Man of Europe*. [Online] Available from <http://www.soilassociation.org/LinkClick.aspx?fileticket=0DLNEXzSIJk%3d&tabid=1315> Accessed (10-09-13)

Soon, Y.K., Clayton, G.W. & Rice, W.A. (2001). Tillage and previous crop effects on dynamics of nitrogen in a wheat–soil system. *Agronomy Journal*. 93, 842–849.

Soon, Y.K. & Lupwayi, N.Z. (2012). Straw management in a cold-arid region: Impact on soil quality and crop productivity. *Field Crops Research*. 139, 39-46.

Sourour, A., Mohsen, R., Afef, O., Ali, B.M., Arbi, M., Mounir, R. & Mongi, B.Y. (2014). Effect of nitrogen fertilization on durum wheat (*Triticum turgidum* var. *durum*) in semi-arid environment. *Canadian Journal of Plant Breeding*. 2 (2), 66-75.

Sousa, W.P. (1984). The role of disturbance in natural communities. *Annual Review of Ecology and Systematics*. 15, 353-91.

Soussana, J.F. & Arregui, M.C. (1995). Impact de l'association sur le niveau de nutrition azotée et la croissance du ray-grass anglais et du trèfle blanc. *Agronomie*. 15, 81–96.

Sowers, K.E., Pan, W.L., Miller, B.C. & Smith, J.L. (1994). Nitrogen use efficiency of split nitrogen applications in soft white winter wheat. *Agronomy Journal*. 86, 942-948.

- Spiegel, H., Dersch, G., Hosch, J. & Baumgarten, A. (2007). Tillage effects on soil organic carbon and nutrients availability in a long-term field experiment in Austria. *Die Bodenkultur*. 58 (1), 47-58.
- Spink, J., Street, P., Sylvester-Bradley, R. & Berry, P. (2009). The potential to increase productivity of wheat and oilseed rape in the UK. in *Report to the Government Chief Scientific Adviser, Prof. John Beddington*. 1-136.
- Squire, J.M. (1997). *Annual medics: Use as a cover crop in corn; weed control; herbicide tolerance*. Ph.D dissertation. Michigan State University. East Lansing, USA.
- Stagnari, F., Galieni, A., Specca, S., Cafiero, G. & Pisante, M. (2014). Effects of straw mulch on growth and yield of durum wheat during transition to Conservation Agriculture in Mediterranean environment. *Field Crops Research*. 167, 51-63.
- Staley, T.E. (1999). Soil microbial biomass alterations during the maize silage growing season relative to tillage method. *Soil Science Society of America Journal*. 63 (6), 1845-1847.
- Steiner, K.G. (1984). *Intercropping in tropical smallholder agriculture with special reference to West Africa*. 2nd edition. Deutsche Gesellschaft für Technische Zusammenarbeit (GTZ), Eschborn, Germany
- Steiner, J.L. (1989). Tillage and surface residue effects on evaporation from soils. *Soil Science Society of America Journal*. 53, 911–916.
- Stevens, C.J., Duprè, C., Dorland, E., Gaudnik, C., Gowing, D.J.G., Bleeker, A., Diekmann, M., Alard, D., Bobbink, R., Fowler, D., Corcket, E., Mountford, J.O. Vandvik, V., Aarrestad, P.A., Muller, S & Dise, N.B. (2010). Nitrogen deposition threatens species richness of grasslands across Europe. *Environmental Pollution*. 158, 2940-2945.
- Stevenson, F.C., Légère, A., Simard, R.R., Angers, D.A., Pangeau, D. & Lafond, J. (1997). Weed species diversity in spring barley varies with crop rotation and tillage, but not with nutrient source. *Weed Science*. 45, 798–806.
- Stodbard, F.L. & Marshall, D.R. (1990). Variability in grain protein in Australia hexaploid wheats. *Australian Journal of Agricultural Research*. 41, 277-288
- Stokes, D.T., Scott, R.K., Tilston, C.H., Cowie, G. & Sylvester-Bradley, R. (1992). The effect of soil disturbance on nitrate mineralisation. *Aspects of Applied Biology*. 30, 279-282.
- Storkey, J. (2006). A functional group approach to the management of UK arable weeds to support biological diversity. *Weed Research*. 46, 513–522.

- Storkey, J., Meyer, S., Still, K.S. & Leuschner, C. (2012). The impact of agricultural intensification and land-use change on the European arable flora. in *Proceedings of the Royal Society B*. 279, 1421-1429.
- Streit, B., Rieger, S.B., Stamp, P. & Richner, W. (2002). The effect of tillage intensity and time of herbicide application on weed communities and populations in maize in central Europe. *Agriculture, Ecosystems and Environment*. 92, 211-224.
- Streit, B., Rieger, S.B., Stamp, P. & Richner, W. (2003). Weed populations in winter wheat as affected by crop sequence, intensity of tillage and time of herbicide application in a cool humid climate. *Weed Research*. 43, 20-32.
- Strudley, M.W., Green, T.R. & Ascough, J.C. (2008). Tillage effects on soil hydraulic properties in space and time. *Soil and Tillage Research*. 99, 4-48.
- Sumner, D.R., Doupnik, B.Jr. & Boosalis, M.G. (1981). Effect of reduced tillage and multiple cropping on plant diseases. *Annual Review of Phytopathology*. 19, 167-187.
- Sutton, J.C. & Vyn, T.J. (1990). Crop sequence and tillage practices in relation to diseases of winter wheat in Ontario. *Canadian Journal of Plant Pathology*. 12, 358-368.
- Swaminathan, M.S. (2007). Can science and technology feed the world in 2025?. *Field Crops Research*. 104, 3-9.
- Swaminathan, M.S. (2004). Ever-Green Revolution and sustainable food security. in Eaglesham, A., Wilderman, A.G. & Hardy, R.W.F. (eds.). *Agricultural Biotechnology: Finding common international goals*. National Agricultural Biotechnology Council. U.S. 63-75.
- Swanton, C.J., Shrestha, A., Knezevic, S.Z., Roy, R.C. & Ball-Coelho, B.R. (2000). Influence of soil tillage type on vertical weed seedbank distribution in sandy soil. *Canadian Journal of Plant Science*. 80, 455-457.
- Sweeney, A.E., Renner, K.A., Laboski, C. & Davis, A. (2008). Effect of fertilizer nitrogen on weed emergence and growth. *Weed Science*. 56, 714-721.
- Tang, L., Cheng, C., Wan, K., Li, R., Wang, D., Tao, Y., Pan, J., Xie, J. & Chen, F. (2014). Impact of fertilizing pattern on the biodiversity of a weed community and wheat growth. *PLoS ONE*. 9 (1), e84370.
- Teasdale, J.R. (1996). Contribution of cover crops to weed management in sustainable agriculture systems. *Journal of Production Agriculture*. 9, 475-479.
- Teasdale, J.R. (1993). Interaction of light, soil moisture, and temperature with weed suppression by hairy vetch residue. *Weed Science*. 41, 46-51.

- Teasdale, J.R., Beste, C.E. & Potts, W.E. (1991). Response to weeds to tillage and cover crop residue. *Weed Science*. 39, 195-199.
- Teasdale, J.R. & Mohler, C.L. (1993). Light transmittance, soil temperature, and soil moisture under residue of hairy vetch and rye. *Agronomy Journal*. 85, 673-680.
- Tebrügge, F. & Düring, R.A. (1999). Reducing tillage intensity- A review of results from a long-term study in Germany. *Soil and Tillage Research*. 53 (1), 15-28.
- Terman, G.L., Raming, R.E., Dreier, A.F. & Olson, R.A. (1996). Yield potential relationships in wheat grain, as affected by nitrogen and water. *Agronomy Journal*. 61. 755-759.
- Theunissen, J. (1997). Intercropping in field vegetables as an approach to sustainable horticulture. *Outlook on Agriculture*. 26, 95–99.
- Thiessen-Martens, J.R., Entz, M.H. & Hoepfner, J.W. (2005). Legume cover crops with winter cereals in southern Manitoba: Fertilizer replacement values for oat. *Canadian Journal of Plant Science*. 85, 645-648.
- Thiessen-Martens, J.R., Hoepfner, J.W. & Entz, M.H. (2001). Legume cover crops with winter cereals in southern Manitoba: Establishment, productivity and microclimate effect. *Agronomy Journal*. 93, 1086-1096.
- Thomas, G.A., Dalal, R.C. & Standley, J. (2007). No-till effects on organic matter, pH, cation exchange capacity and nutrient distribution in a Luvisol in the semi-arid subtropics. *Soil and Tillage Research*. 94, 295–304.
- Thompson, K., Grime, J.P. & Mason, G. (1977). Seed germination in response to diurnal fluctuations of temperature. *Nature*. 267, 147-149.
- Thomsen, I.K. & Sørensen, P. (2006). Tillage-induced N mineralization and N uptake in winter wheat on a coarse sandy loam. *Soil and Tillage Research*. 89, 58-69.
- Thorsted, M.D., Olesen, J.E. & Weiner, J. (2006). Width of clover strip and wheat rows influence grain yield in winter wheat/white clover intercropping. *Field Crops Research*. 95, 280-290.
- Thurston, J. M. (1959). A comparative study of the growth of wild oats (*Avena fatua* L. and *Avena ludoviciana* Dur.) and of cultivated cereals with varied nitrogen supply. *Annals of Applied Biology*. 47, 716-739.
- Tørresen, K.S. & Skuterud R. (2002). Plant protection in spring cereal production with reduced tillage. IV: Changes in the weed flora and weed seedbank. *Crop Protection*. 21 (3), 179-193.

- Triplett, G.B. & Dick, W.A. (2008). No-tillage crop production: a revolution in agriculture! *Agronomy Journal*. 100, S153-S165.
- Troccoli, A., Borrelli, G.M., De Vita, P., Fares, C. & Di Fonzo, N. (2000). Durum wheat quality: A multidisciplinary concept. *Journal of Cereal Science*. 32, 99-113.
- Tuesca, D. & Puricelli, E. (2007). Effect of tillage systems and herbicide treatments on weed abundance and diversity in a glyphosate resistant crop rotation. *Crop Protection*. 26, 1765-1770.
- Tuesca, D., Puricelli, E. & Papa J.C. (2001). A long-term study of the weed flora shifts in different tillage systems. *Weed Research*. 41, 369-382.
- Turley, D.B., Phillips, M.C., Johnson, P., Jones, A.E. & Chambers, B.J. (2003). Long-term straw management on yields of sequential wheat (*Triticum aestivum* L.) crops in clay and silty clay loam soils in England. *Soil and Tillage Research*. 71, 59-69
- Uhlen, H. (1998). Why energy productivity is increasing: an analysis of Swedish agriculture. *Agricultural Systems*. 56 (4), 443-465.
- Unger, P.W. & Cassel, D.K. (1991). Tillage implement disturbance effects on soil properties related to soil and water conservation – a literature review. *Soil and Tillage Research*. 19, 363–382.
- United Nations, Department of Economic and Social Affairs, Population Division (2013). *World Population Prospects: The 2012 Revision, Volume I: Comprehensive Tables*. ST/ESA/SER.A/336. [Online] Available online at http://esa.un.org/unpd/wpp/Documentation/pdf/WPP2012_Volume-I_ComprehensiveTables.pdf Accessed (08-01-14).
- United Nations, Department of Economic and Social Affairs, Population Division (2014). *The World Population Situation in 2014. A concise report*. ST/ESA/SER.A/354. New York.
- Unkovich, M., Herridge, D., Peoples, M., Cadish, G., Boddey, R., Giller, K., Alves, B. & Chalk, P. (2008). *Measuring plant-associated nitrogen fixation in agricultural systems*. Australian Centre for International Agricultural Research (ACIAR)
- Usman, K., Khan, N., Khan, M.U., ur Rehman, A. & Ghulam, S. (2013). Impact of tillage and herbicide on weed density, yield and quality of cotton in wheat based cropping systems. *Journal of Integrative Agriculture*. 12 (9), 1568-1579.
- Valkama, E., Salo, T., Esala, M. & Turtola, E. (2013). Nitrogen balances and yields of spring cereals as affected by nitrogen fertilization in northern conditions: A meta-analysis. *Agriculture, Ecosystems and Environment*. 164, 1-13.

- Van Bemmelen, J.W. (1890). Die Zusammenetzung der Ackererde nach Anleitung der in dem vorigen Abhandlungen mitgeleiteten Analysen von gewöhnlichen und vulkanischen Thonboden. Die landwirthschaftlichen Versuchs-Stationen. *Band. 37*, 347.
- Van de Geijn, S., Vos, J.C., Groenwold, J., Goudriaan, J. & Leffelaar, P.A. (1994). The Wageningen Rhizolab – a facility to study soil–root–atmosphere interactions in crops. I. Description of main functions. *Plant and Soil*. 161, 275-287.
- Vandermeer, J. (1989). *The Ecology of Intercropping*. Cambridge University Press, New York, NY.
- Van der Mensbrugghe, D., Osorio-Rodarte, I., Burns, A. & Baffes, J. (2011). Macroeconomic Environment and Commodity Markets: A Longer Term Outlook. In FAO (Food and Agriculture Organization). *Looking ahead in world food and agriculture: perspective to 2050, an Expert Meeting and high level expert forum around the question 'How to feed the world in 2050'*. FAO. Rome, Italy. 191-230.
- van Keulen, H. & Seligman, N.G. (1987). Simulation of water use, nitrogen nutrition and growth of a spring wheat crop. *Simulation Monographs*. Pudoc, Wageningen, 310.
- Vicent, E.M. & Roberts, E.H. (1977). Interaction of light, nitrate and alternative temperatures in promoting the germination of dormant seeds of common weed species. *Seed Science and Technology*. 5, 659-670.
- Vijaya Bhaskar, A.V. (2014). *Cultivation regimes and legumes cover crops for organic wheat (Triticum aestivum) production*. PhD dissertation. Coventry University. UK
- Vijaya Bhaskar, A.V., Davies, W.P., Cannon, N.D. & Conway, J.S. (2013a). Characterisation of soil quality following conventional and non-inversion tillage for organic farming. *Aspect of Applied Biology. Rethinking Agricultural Systems in UK*. 121, 181-185.
- Vijaya Bhaskar, A.V., Davies, W.P., Cannon, N.D. & Conway, J.S. (2013b). Organic wheat performance following conventional and non-inversion tillage systems. *Biological Agriculture and Horticulture*. 29 (4), 236-243.
- Vijaya Bhaskar, A.V., Cannon N. D. & Davies, W. P. (2013c). Investigation of sustainable legume species for under sowing in organic spring wheat. *Aspect of Applied Biology. Rethinking Agricultural Systems in UK*. 121, 199-202.
- Vijaya Bhaskar, A.V., Davies, W.P., Cannon, N.D. & Conway, J.S. (2013d). Tillage and undersowing effect on organic wheat yield components and yield. *Aspect of Applied Biology. Rethinking Agricultural Systems in UK*. 121, 173-180.

- Vijaya Bhaskar, A.V., Davies, W.P., Cannon, N.D. & Conway, J.S. (2014a). Soil cultivation influences on organic spring wheat performance in contrasting weather. *Aspects of Applied Biology*. 125, 224-232.
- Vijaya Bhaskar, A.V, Rial Lovera, K, Davies, W.P. & Cannon, N.D. (2014b). Weed prevalence under organic and low input cultivation systems. *Aspect of Applied Biology*. Agronomic decision making in an uncertain climate. 125, 129-134.
- Vilich-Meller, V. (1992). Mixed cropping of cereals to suppress plant diseases and omit pesticide applications. *Biological Agriculture and Horticulture*. 8, 299–308.
- Vyn, T.J. & Raimbault, B.A. (1993). Long-term effect of five tillage systems on corn response and soil structure. *Agronomy Journal*. 85, 1074-1079.
- Wade, A., Spink, J.H. & Orson, J. (2006). The effect of cultivation method on optimum plant population in winter wheat. *HGCA Project Report No. 405*. Home-Grown Cereals Authority, London, UK.
- Walker, S., Stiger, C.J., Ofori, E. & Kyei-Baffour, N. (2011). Intercropping and its implications for soil management. In Hatfield, J.L. & Sauer, T.J. (eds.). *Soil management: Building a stable base for agriculture*, ASA-SSSA, Madison, USA.
- Wall, J.S. (1979). The role of wheat proteins in determining baking quality. In Laidman, D.L. & Wyn Jones, R.G. (eds.). *Recent advances in the biochemistry of cereals*. London, New York Academic. 275-311.
- Wallace, J. (2001). Black medic: a promising green manure for dryland areas. *Eco-Farm & Garden*. 4, 18-19.
- Wallace, S.U., Bacanamwo, M., Palmer, J.H. & Hull, S.A. (1996). Yield and yield components of relay-intercropped wheat and soybean. *Field Crops Research*. 46, 161-168.
- Wang, G.L., Ye, Y.L., Chen, X.P. & Cui, Z.L. (2014). Determining the optimal nitrogen rate for summer maize in China by integrating agronomic, economic and environmental aspects. *Biogeosciences*. 11, 3031-3041.
- Wang, H., Guo, Z., Shi, Y., Zhang, Y., Yu, Z. (2015). Impact of tillage practices on nitrate accumulation and translocation in wheat and soil nitrate-nitrogen leaching in drylands. *Soil and Tillage Research*. 153. 20-27.
- Wang, X., Wu, H., Dai, K., Zhang, D., Feng, Z., Zhao, Q., Wu, Z., Jin, K., Cai, D., Oenema, O. & Hoogmoed, W.B. (2012). Tillage and crop residue effects on rainfed wheat and maize production in northern China. *Field Crop Research*. 132, 106-116.

Watkins, J.E. & Boosalis, M.G. (1994). Plant disease incidence as influence by conservation tillage systems. In Unger, P.W. (ed.) *Managing Agricultural Residues*. Lewis Publisher Inc, Boca Raton, FL, USA. 261-283.

Webb, J., Sylvester-Bradley, R. & Wafford, J.D. (1995). Influence of sowing date on the uptake of and responses to soil and fertilizer nitrogen by the spring wheat cultivar Tonic. *Journal of Agricultural Science*. 125, 25-37.

Weisz, R., Crozier, C.R. & Heinige, R.W. (2001). Optimizing nitrogen application timing in no-till soft red winter wheat. *Agronomy Journal*. 93, 435-442.

Wheat Yield consortium. [Online] Available from <http://www.cimmyt.org/en/newsletter/38-2009/461-wheatwarriors-the-struggle-to-break-the-yield-barrier> Accessed (10-01-14)

White, P.F. (1990). The influence of alternative tillage systems on the distribution of nutrients and organic carbon in some common Western Australia wheat belt soil. *Australia Journal of Soil Research*. 28, 95-116.

Whitehead, A. (2002). *Meta-analysis of Controlled Clinical Trials*. Wiley, Chichester.

Wibberley, E.J. (1984). Spring wheat: strategic potential in cropping systems including a special study of Cotswold crops. *National Agricultural Centre. Arable Unit*.

Wicks, G.A., Smika, D.E. & Hergert, G. (1988). Long-term effect of no-tillage in a winter wheat (*Triticum aestivum*)-sorghum (*Sorghum bicolor*)-fallow rotation. *Weed Science*. 36, 384-393.

Wieser, H. & Seilmeier, W. (1998). The influence of nitrogen fertilisation on quantities and proportions of different protein types in wheat flour. *Journal of Science Food and Agriculture*. 76, 49-55.

Wild, A. (1988). *Russell's soils conditions & plant growth*. 11th Ed. Longmand Scientific/John Wiley, Harlow, UK.

Wilkins, D.E., Klepper, B. & Rickman, R.W. (1989). Measuring wheat seedling response to tillage and seeding systems. *Transactions of the American Society of Agricultural Engineers*. (Trans ASAE). 32. 795-800

Winkler, J. & Chovancová, S. (2014). The effect of precipitation on weed infestation of spring barley under different tillage conditions. *International Journal of Biological, Veterinary, Agricultural and Food Engineering*. 8 (8), 805-812.

Wojciechowski, W. & Sowiński, J. (2005). Changes in the number of weed seeds in soil under different tillage systems of winter wheat. *Journal of Plant Protection Research*. 45 (2), 83-92.

- Woźniaka, A. & Sorokab, M. (2014). Effects of a 3-year reduced tillage on the yield and quality of grain and weed infestation of spring triticale (*Triticosecale* Wittmack). *International Journal of Plant Production*. 8 (2), 231-242.
- Xu, Z.Z., Yu, Z.W., Wang, D. & Zhang, Y.L. (2005). Nitrogen accumulation and translocation for winter wheat under different irrigation regimes. *Journal of Agronomy and Crop Science*. 191, 439-449.
- Yagioka, A., Komatsuzaki, M., Kaneko, N. & Ueno, H. (2015). Effect of no-tillage with weed cover mulching versus conventional tillage on global warming potential and nitrate leaching. *Agriculture, Ecosystems and Environment*. 200, 42-53.
- Yilmaz, I., Akcaoz, H. & Ozkan, B. (2005). An analysis of energy use and input costs for cotton production in Turkey. *Renewable Energy*. 30, 145–155.
- Young, D.L., Kwon, T.J. & Young, F.L. (1994). Profit and risk for integrated conservation farming systems in the Palouse. *Journal of Soil Water Conservation*. 49, 601-606.
- Zadoks, J.C., Chang, T.T. & Konzak, C.F. (1974). A decimal code for the growth stages of cereals. *Weed Research*. 14, 415-421.
- Zangeneh, M., Omid, M. & Akram, A. (2010). A comparative study on energy use and cost analysis of potato production under different farming technologies in Hamadan province of Iran. *Energy*. 35, 2927-2933.
- Zentner, R.P., Lafond, G.P., Derksen, D.A., Nagy, C.N., Wall, D.D. & May, W.E. (2004). Effect of tillage method and crop rotation on non-renewable energy use efficiency for a thin Black Chernozem in the Canadian Prairies. *Soil and Tillage Research*. 77, 125-136.
- Zhang, F. & Li, L. (2003). Using competitive and facilitative interactions in intercropping systems enhances crop productivity and nutrient-use efficiency. *Plant and Soil*. 248, 305-312.
- Zhang, H., Turner, N.C. & Poole, M.L. (2012). Increasing the harvest index of wheat in the high rainfall zones of southern Australia. *Field Crops Research*. 129, 111-123.
- Zhao, S., Qiu, S., Cao, C., Zheng, C., Zhou, W. & He, P. (2014). Response of soil properties, microbial community and crop yields to various rates of nitrogen fertilization in a wheat-maize cropping system in north-central China. *Agriculture, Ecosystems and Environment*. 194, 29-37.

Zheng, C., Jiang, Y., Chen, C., Sun, Y., Feng, J., Deng, A., Song, Z. & Zhang, W. (2014). The impacts of conservation agriculture on crop yield in China depend on specific practices, crops and cropping regions. *The Crop Journal*. 2 (1), 289-296.

Zhu, J. & Khan, K. (2001). Effects of genotype and environment on glutenin polymers and breadmaking quality. *Cereal Chemistry*. 78, 125–130.

Ziadi, N., Bélanger, G., Claessens, A., Lefebvre, L., Cambouris, A.N., Tremblay, N., Nolin, M.C. & Parent, L-E. (2010). Determination of a critical nitrogen dilution curve for spring wheat. *Agronomy Journal*. 102 (1), 241-250.

Ziaei, S.M., Mazloumzadeh, S.M. & Jabbari, M. (2015). A comparison of energy use and productivity of wheat and barley (case study). *Journal of the Saudi Society of Agricultural Sciences*. 14 (1), 19-25.

Appendices

Appendix 1. Illustrated keys used for disease assessment

Appendix 2. Weed species biomass across core experiments

Appendix 3. Data set for meta-analysis

Appendix 4. Energy balance

Appendix 5. Publications prepared during this investigation

Appendix 1. Illustrated keys used for disease assessment

Appendix 1.1. Illustrated key for leaf rust of wheat.

This item has been removed due to 3rd Party Copyright. The unabridged version of the thesis can be viewed in the Lanchester Library Coventry University.

Appendix 1.2. Illustrated key for septoria leaf blotch of wheat.

This item has been removed due to 3rd Party Copyright. The unabridged version of the thesis can be viewed in the Lanchester Library Coventry University.

Appendix 1.3. Illustrated key for take-all disease of wheat
This item has been removed due to 3rd Party Copyright. The unabridged version of the thesis can be viewed in the Lanchester Library Coventry University.

Appendix 2. Weed species biomass across core experiments

Appendix 2.1. Grass weed species for the 2013 core experiment

Grass weed species	Mean DM (kg ha ⁻¹)
1. <i>Avena fatua</i>	170.0 ± 47.20
2. <i>Lolium perenne</i>	116.0 ± 19.40
3. <i>Alopecurus myosu.</i>	19.90 ± 3.01
4. <i>Avena sativa</i>	0.412 ± 0.294
5. <i>Hordeum vulgare</i>	0.329 ± 0.329

Appendix 2.2. Broadleaf weed species for the 2013 core experiment

Broadleaf weed species	Mean DM (kg ha ⁻¹)	Broadleaf weed species	Mean DM (kg ha ⁻¹)
1. <i>Stellaria media</i>	425.0 ± 44.7	18. <i>Cirsium arvense</i>	2.44 ± 1.30
2. <i>Fallopia convolvulus</i>	173.0 ± 16.20	19. <i>Veronica persica</i>	1.39 ± 0.42
3. <i>Sinapsis arvensis</i>	138.0 ± 28.40	20. <i>Veronica hederifolia</i>	0.931 ± 0.33
4. <i>Polygonum aviculare</i>	117.0 ± 15.60	21. <i>Urtica urens</i>	0.745 ± 0.498
5. <i>Galium aparine</i>	82.10 ± 13.20	22. <i>Angallis arvensis</i>	0.616 ± 0.215
6. <i>Aethusa cynapium</i>	70.90 ± 6.49	23. <i>Rumex obtusifolius</i>	0.597 ± 0.26
7. <i>Sonchus oleraceus</i>	37.50 ± 9.55	24. <i>Sonchus arvensis</i>	0.597 ± 0.374
8. <i>Persicaria maculosa</i>	33.60 ± 8.29	25. <i>Lamium amplexicaule</i>	0.403 ± 0.182
9. <i>Geranium dissectum</i>	13.30 ± 3.18	26. <i>Fumaris officinalis</i>	0.366 ± 0.258
10. <i>Lapsana communis</i>	6.53 ± 2.07	27. <i>Brassica napus olifera</i>	0.333 ± 0.333
11. <i>Sinapsis alba</i>	4.38 ± 2.22	28. <i>Scandix pecten-veneris</i>	0.301 ± 0.201
12. <i>Atriplex patula</i>	3.32 ± 1.30	29. <i>Convolvulus arvensis</i>	0.292 ± 0.292
13. <i>Galeopsis tetrahip</i>	3.32 ± 3.32	30. <i>Rumex spp</i>	0.241 ± 0.241
14. <i>Viola tricolor</i>	3.16 ± 1.17	31. <i>Sonchus arvensis</i>	0.093 ± 0.065
15. <i>Senecio vulgare</i>	2.98 ± 1.25	32. <i>Cirsum vulgare</i>	0.051 ± 0.042
16. <i>Capsella bursa-pastoris</i>	2.98 ± 2.12	33. <i>Rumex crispus</i>	0.037 ± 0.037
17. <i>Chenopodium spp</i>	2.84 ± 1.96	34. <i>Vicia sativa</i>	0.005 ± 0.005

Appendix 2.3. Grass weed species for the 2014 core experiment

Grass weed species	Mean DM (kg ha⁻¹)
1. <i>Lolium perenne</i>	22 ± 10.7
2. <i>Avena sativa</i>	295 ± 62.0
3. <i>Alopecurus myosuroides</i>	3.8 ± 1.31
4. <i>Hordeum vulgare</i>	0.204 ± 0.204
5. <i>Poa annua</i>	0.0556 ± 0.0556

Appendix 2.4. Broadleaf weed species for the 2014 core experiment

Broadleaf weed species	Mean DM (kg ha⁻¹)	Broadleaf weed species	Mean DM (kg ha⁻¹)
1. <i>Aethusa cynapium</i>	3.75 ± 0.643	13. <i>Polygonum aviculare</i>	1.23 ± 0.462
2. <i>Persicaria maculosa</i>	30 ± 12.70	14. <i>Cirsium vulgare</i>	0.171 ± 1.39
3. <i>Galium aparine</i>	162 ± 31.70	15. <i>Rumex crispus</i>	0.606 ± 0.31
4. <i>Stellaria media</i>	167 ± 25.80	16. <i>Veronica agrestis</i>	4.94 ± 1.35
5. <i>Sinapsis arvensis</i>	385 ± 60.90	17. <i>Sonchus arvensis</i>	2.78 ± 0.943
6. <i>Veronica persica</i>	6.4 ± 6.20	18. <i>Brassica nigra</i>	0.102 ± 0.102
7. <i>Geranium dissectum</i>	18.1 ± 4.88	19. <i>Viola arvensis</i>	0.37 ± 0.48
8. <i>Lapsana communis</i>	11.7 ± 3.22	20. <i>Myosotis arvensis</i>	0.0648 ± 0.0437
9. <i>Sonchus arvensis</i>	4.75 ± 2.01	21. <i>Lapsana communis</i>	2.60 ± 1.65
10. <i>Fallopia convolvulus</i>	4.15 ± 0.91	22. <i>Atriplex patula</i>	3.36 ± 3.07
11. <i>Convolvulus arvensis</i>	3.66 ± 2.01	23. <i>Senecio vulgare</i>	1.49 ± 0.627
12. <i>Phalaris paradoxa</i>	0.648 ± 0.468	24. <i>Cirsium arvense</i>	2.37 ± 1.14

Appendix 3. Data set for meta-analysis

Appendix 3.1. Organic study used

Reference	Crop	Tillage treatments	Seasonal rainfall (mm)
Vijaya Bhaskar <i>et al.</i> (2013b)	Spring wheat (2012)	CT, HINiT and LINiT	589.1

Appendix 3.2. Data set with spring wheat yield (t ha⁻¹) under contrasting cultivation techniques across experiments

	2012	2013	2014
CT	3.52	5.09	5.12
HINiT	2.96	5.1	1.89
LINiT	2.11	6.2	2.28
Mean	2.86	5.46	3.10
SED	0.152	1.132	0.233

Appendix 3.3. HINiT weighed yield against CT

Seasonal rainfall type	Tillage	Weighed yield (t ha ⁻¹)	SED
<300 mm	HINiT	0.009	1.132
300-500 mm	HINiT	-3.23	0.233
>500	HINiT	-0.56	0.152

Appendix 3.4. LINiT weighed yield against CT

Seasonal rainfall type	Tillage	Weighed yield (t ha ⁻¹)	SED
<300 mm	LINiT	1.11	1.132
300-500 mm	LINiT	-2.84	0.233
>500	LINiT	-1.41	0.152

Appendix 4. Energy balance

Appendix 4.1. Amount of inputs in all core experiments

Inputs	Quantity per unit	Inputs	Quantity per unit
Human labour (h ha⁻¹)		Diesel fuel (l ha⁻¹)	
CT	1.97	CT	38
HINiT	2	HINiT	34
LINiT	1.58	LINiT	24
BM	1.5	BM	0
WC	1.5	WC	0
Nus	0	Nus	0
N0 application	0	N0 application	0
N70 application	0.8	N70 application	2.4
N140 application	0.8	N140 application	2.4
N210 application	0.8	N210 application	2.4
Herbicide spraying	0.4	Herbicide spraying	1.2
Harvesting	1.05	Harvesting	23
Machinery (h ha⁻¹)		Fertiliser (kg ha⁻¹)	
CT	1.97	N0 application	0
HINiT	2	N70 application	70
LINiT	1.58	N140 application	140
BM	0	N210 application	210
WC	0	Herbicide (kg ha⁻¹)	2
Nus	0	Seeds (kg ha⁻¹)	
N0 application	0	Wheat	200
N70 application	0.8	BM	8
N140 application	0.8	WC	7
N210 application	0.8		
Herbicide sprayer	0.4		
Harvesting	1.05		

Appendix 4.2. Specification of all machinery used in all core experiments

	Tractor type	Implement type	Implement width (m)	Working depth (cm)	Speed of work (km h⁻¹)	Work rate (ha h⁻¹)	Time (h ha⁻¹)	Fuel (L ha⁻¹)
CT	MF 5465 (120 HP, 4wd) (5080 kg)	5 furrow Kverneland plough + press	1.8	20	7	1	1	23
		Power harrow seed drill	3	8	8	1.7	0.97	15
HINiT	TM 155 (154 HP, 4wd) (5642 kg)	2 passes of ST bar attached Simba X-press	3	25 & 12	10	1.2	1.4	28
		Vaderstadt with seed drill	4	8	10	2.8	0.6	6
LINiT	TM 155 (154 HP, 4wd) (5642 kg)	1 passes of ST bar attached Simba X-press	3	25 & 12	10	1.2	0.7	14
		Eco-dyn seed drill	3	26	9	1.9	0.88	10
Fertiliser sprayer		2 passes CASE IH SP3000 (150HP)	6				0.8	2.4
Herbicide sprayer		CASE IH SP3000 (150HP)	6				0.4	1.2
Harvest combine							1.05	23

Appendix 5. Publications prepared during this investigation

Vijaya Bhaskar, A.V., Rial Lovera, K., Davies, W.P. & Cannon, N.D. (2014). Weed prevalence under organic and low input cultivation systems. *Aspect of Applied Biology*. Agronomic decision making in an uncertain climate. 125, 129-134.

This item has been removed due to 3rd Party Copyright. The unabridged version of the thesis can be viewed in the Lanchester Library Coventry University.

This item has been removed due to 3rd Party Copyright. The unabridged version of the thesis can be viewed in the Lanchester Library Coventry University.

This item has been removed due to 3rd Party Copyright. The unabridged version of the thesis can be viewed in the Lanchester Library Coventry University.

This item has been removed due to 3rd Party Copyright. The unabridged version of the thesis can be viewed in the Lanchester Library Coventry University.

This item has been removed due to 3rd Party Copyright. The unabridged version of the thesis can be viewed in the Lanchester Library Coventry University.

This item has been removed due to 3rd Party Copyright. The unabridged version of the thesis can be viewed in the Lanchester Library Coventry University.