

1 **An evaluation of persistent meteorological drought using a**
2 **homogeneous Island of Ireland precipitation network**

3

4

5 Wilby¹, R.L., Noone², S., Murphy², C., Matthews³, T., Harrigan², S. and Broderick², C.

6

7 ¹ Department of Geography, Loughborough University, LE11 3TU, UK

8 ² Department of Geography, National University of Ireland Maynooth, Ireland

9 ³ School of Natural Sciences & Psychology, Liverpool John Moores University, L3 3AF, UK

10

11

12

13 Main text word count: 4035

14

15 2 September 2015

16

17 Re-submitted to: *International Journal of Climatology*

18

19 Corresponding author: Robert Wilby (email: r.l.wilby@lboro.ac.uk)

20

21 **Abstract**

22 This paper investigates the spatial and temporal properties of persistent meteorological
23 droughts using the homogeneous Island of Ireland Precipitation (IIP) network. Relative to a
24 1961-1990 baseline period it is shown that the longest observed run of below average
25 precipitation since the 1850s lasted up to 5 years (10 half-year seasons) at sites in southeast
26 and east Ireland, or 3 years across the network as a whole. Dry- and wet-spell length
27 distributions were represented by a first-order Markov model which yields realistic runs of
28 below average rainfall for individual sites and IIP series. This model shows that there is
29 relatively high likelihood ($p=0.125$) of a 5 year dry-spell at Dublin, and that near unbroken
30 dry runs of 10 years or more are conceivable. We suggest that the IIP network and attendant
31 rainfall deficit modelling provide credible data for stress testing water supply and drought
32 plans under extreme conditions.

33

34 *Key words:*

35 Drought duration; Markov model; homogeneous rainfall series; water planning; Ireland.

36

37

38 **1. Introduction**

39 Drought is hardly synonymous with perceptions about the climate of Ireland. Nonetheless,
40 the *Freeman's Journal*¹ provides numerous reports of potable water shortages in Dublin
41 during severe dry spells over the period 1763-1924 and Barrington (1888) gives a rich
42 account of impacts of the 1887 drought on Irish agriculture. Other notable events such as the
43 pan-European drought of 1976 caused heat- and moisture-stress-related problems for
44 Ireland's agricultural sector (Stead, 2014). Likewise, some future climate scenarios foresee
45 loss of production for crops such as potatoes linked to rising temperatures and summer aridity
46 (Holden *et al.*, 2003); reduced grass growth and heat stress on livestock which could impact
47 meat and dairy exports (Hunt *et al.*, 2014); and decreased river flow in summer (Steele-
48 Dunne *et al.*, 2008; Bastola *et al.*, 2011).

49 Given these vulnerabilities, surprisingly little has been published on the drought climatology
50 of Ireland. O'Laoghog (1979) provides a summary of rainfall anomalies alongside impacts on
51 agriculture and public water supply of the 1974-1976 drought. Brogan and Cunnane (2005)
52 contend that 1976 may have witnessed the lowest recorded river flows since the 1930s. They
53 also cite droughts in 1934, 1949, 1955, 1959, 1975, 1989, 1990, 1991 and 1995.

54 MacCarthaigh (1996) compared 1995 with droughts back to 1975 whilst Dooge (1985)
55 provides a synopsis of droughts in Irish history beginning with accounts in the Annals of
56 Ulster and of Clonmacnoise (for the period AD 759 to 1408). Symons (1887) documents five
57 droughts in the 1850s, two in the 1860s and three in the 1870s. Garcia-Suarez and Butler
58 (2006) find periods with persistently negative annual Palmer Drought Severity Index at
59 Armagh in the 1880s, 1890s, 1930s, 1970s and 1990s. Mandal (2011) estimated low flows for
60 125 Irish rivers using catchment properties. Aside from these sources, there is little
61 quantitative information on which to base rigorous assessments of long-term drought risk and
62 water planning for Ireland.

63 This *Short Communication* addresses this knowledge gap by using the homogeneous Island
64 of Ireland Precipitation (IIP) network of Noone *et al.* (2015) to evaluate the *occurrence* and
65 *persistence* of meteorological droughts since 1850. Here, a straightforward definition of
66 drought is used for half-year periods or longer that have below average precipitation, at both
67 site and regional scales. We accept that the term 'drought' is ambiguous and that runs of
68 seasonal rainfall deficiency do not necessarily translate into periods of agricultural,

¹ *Freeman's Journal* available through Irish Newspaper Archives www.irishnewsarchive.com/

69 hydrological or environmental drought (Wilhite and Glantz, 1985). Nonetheless, our
70 interrogation uses seasonal rainfall and persistence metrics not applied in the homogenization
71 process to quality-assure the integrity of series within the IIP network. We first reprise the
72 methods used to homogenise the IIP series and list other homogeneous rainfall products for
73 comparison. We then describe and apply statistical techniques for simulating occurrence and
74 persistence of below average rainfall across the IIP network. This leads into an account and
75 interpretation of the key findings before concluding with a few suggestions for further
76 research.

77

78 **2. Data**

79 Our analysis draws on several data sets. The homogeneous IIP network contains monthly
80 totals for 25 stations (Figure 1) covering the common years 1850 to 2010 (Noone *et al.*,
81 2015). Precipitation data underpinning the IIP network were drawn from four sources: long-
82 term series held by the Climatic Research Unit (UK) and Centre for Environmental Data
83 Archival (UK) updated to 2010 (16 stations); the record of Armagh Observatory (UK) (1
84 station); plus digital and paper records of varying completeness held by Met Éireann (IE) (8
85 stations). Raw data exist for all 25 stations from 1908; 23 from the 1890s; 19 from the 1880s;
86 and 8 in the 1850s. The longest continuous record is for Belfast which begins in 1812.

87 In preparation of the IIP network, for each station, detailed information about correction
88 factors, nearest neighbours, observer practices, meteorological site and gauge condition was
89 transcribed to a master file of meta-data to help interpret break points detected during
90 homogenisation. The HOMegenisation softwarE in R (HOMER) package (Mestre *et al.*,
91 2013) was used to detect and correct inhomogeneity in the monthly series and to infill/extend
92 records to the period 1850-2010 (see: Noone *et al.*, 2015). HOMER compares differences
93 between candidate and reference sites within a network to identify likely break points that can
94 then be ratified against meta-data. Given the low density of long-term stations available for
95 IIP, at least 12 correlated reference stations for each candidate series were identified for
96 pairwise comparison and break detection. Annual correction factors were applied to
97 confirmed break points using an ANOVA model (Caussinus and Mestre, 2004; Mestre *et al.*,
98 2013; Venema *et al.* 2012). Missing data were infilled using the same method such that all
99 records start in 1850 with correction factors based on the adjustment amplitude applied until

100 the first detected change point of the series (Noone *et al.*, 2015). Finally, the regional IIP
101 series was constructed as the un-weighted monthly mean of the 25 series.

102 Monthly precipitation totals for England and Wales (EWP), Scotland (SP) and Northern
103 Ireland (NIP) were obtained from the Met Office Hadley Centre. The EWP series begins in
104 1766, whereas SP and NIP start in 1931. All series are based on long-running meteorological
105 stations weighted to provide spatially and temporally homogeneous, area-averaged
106 precipitation totals (Alexander and Jones, 2001). These precipitation series were used
107 alongside the meta-data and archival material described above to quality check the
108 provenance of the IIP series and major dry-spells detected therein.

109

110 **3. Methods**

111 Following Wilby *et al.* (2015) the homogeneous IIP series were processed in four ways. First,
112 mean monthly precipitation totals were derived for a baseline period 1961-1990 with
113 averages consolidated into mean winter (October to March) and summer (April to September)
114 half years. Seasonal anomalies were then calculated for the 1850-2010 series relative to 1961-
115 1990 half-year means, recognising that spell lengths are sensitive to choice of baseline period
116 (Sen, 1980). As will be shown later, 1961-1990 was a relatively dry period. Therefore, any
117 negative anomalies referenced to this baseline are indeed noteworthy.

118 The number of stations with below average precipitation was counted for each half year to
119 establish the spatial coherence of dry-spells, accepting that this is a crude metric because of
120 the sparse and uneven distribution of sites (Figure 1). When more than two thirds of stations
121 in the IIP network report a dry season the event is regarded as widespread and unlikely to be
122 due to a local anomaly or suspect data. Dates of spells lasting three or more half-year seasons
123 were then cross-referenced to EWP and SP to establish coherence at the scale of the British-
124 Irish Isles.

125 Second, conditional dry-to-dry (Pdd) and wet-to-wet (Pww) first-order Markov model
126 transition probabilities were determined from series of seasonal anomalies. This involved
127 counting the frequency with which a below average season is followed by another dry season.
128 Pdd is the proportion of transitions that are dry-to-dry out of all transitions (i.e. dry-to-dry
129 plus dry-to-wet). Similarly, Pww was derived from the proportion of wet-to-wet transitions.

130 Unbroken dry- and wet-season runs were used to construct frequency distributions of spell
131 lengths and to identify the most persistent dry-spells in each record. Pdd and Pww were also
132 estimated for 30-year moving blocks to establish whether there has been any long-term
133 change in dry- and/or wet-spell persistence throughout the IIP, EWP, NIP and SP series.

134 Third, as in Wilby (2007), Sharma and Panu (2012; 2014a;b) and Wilby *et al.* (2015), Pdd
135 and Pww transition probabilities based on 1850-2010 observations were used to
136 stochastically simulate series of seasons with above or below average rainfall. The process
137 begins by seeding with a uniform random number $r[0,1]$ to determine whether there is a
138 change from the initial state (assumed to be below average rainfall). If $r \leq Pdd$ the dry-spell
139 continues; if $r > Pdd$ then the new state is wet and Pww is applied at the next time step. In
140 this way, a single 10,000 season Markov model simulation was performed to generate a
141 distribution of synthetic spell lengths. The two-sample, nonparametric Kolmogorov-Smirnov
142 (KS) test was applied to determine whether the largest discrepancy (Dstat) between observed
143 and simulated cumulative distributions of spell-length was significantly ($p < 0.05$) greater than
144 expected by chance.

145 Finally, 1000 boot-strap, Markov model simulations were performed to generate 100- and
146 160-year (i.e. 200 and 320 season) sequences for each site and region. Maximum dry- and
147 wet-spell lengths were retained from all 1000 realisations to construct distributions of
148 synthetic 100- and 160-year spells for comparison with observations. The 160-year event was
149 generated for equivalence with observed record lengths; the 100-year event enables
150 comparison with Wilby *et al.* (2015). Both sets of distributions were used to estimate
151 likelihoods of a 10-season spell with below average precipitation at each site. This provides
152 an upper bound (yet plausible) dry-spell that is much longer than the single season (1995)
153 design drought applied in, for example, the Dublin City Council (2010) Water Plan.

154

155 **4. Results**

156 Figure 2 shows seasonal anomalies as percentages of the 1961-1990 mean for IIP and EWP
157 since 1850. The IIP series is significantly correlated with both EWP ($r = +0.75$) and SP ($r =$
158 $+0.57$) (not shown). Seasonal anomalies of IIP vary between -40% (winter 1879/80) and
159 +49% (summer 1924). Most (84%) seasons lie within $\pm 20\%$ of the 1961-1990 average
160 precipitation. Overall, the driest 30-year period in IIP was 1884-1913 with 2% less

161 precipitation than 1961-1990. Hence, our chosen standard period was close to the very driest
162 continuous run in the IIP series and any negative anomalies would certainly have been
163 indicative of dry seasons. However, as noted at the outset, meteorological droughts do not
164 necessarily coincide with significant agricultural, water resource or environmental stress.

165 Nine dry-spells lasting longer than three seasons (and simultaneously occurring at more than
166 two thirds of stations) were identified (Table 1). Persistent events stand out in 1853-1856,
167 1886-1888 (followed shortly by 1892-1894) and 1970-1973 (tailed by 1974-1976). Dry spells
168 in the 1850s and late nineteenth century have been reported previously for Ireland
169 (Barrington, 1888; Symons, 1887; Tabony, 1980) and for England and Wales (Barker *et al.*,
170 2004; Burt and Howden, 2011; Burt and Horton, 2007; Jones *et al.*, 2006; Marsh *et al.*, 2007;
171 Wilby and Quinn, 2013). Likewise, dry-spells in the 1970s achieved notoriety for their
172 drought orders, rota cuts and standpipes across parts of south Wales, central and southern
173 England, water rationing in the Channel Islands, and even nightly shut-offs in Belfast (Rodda
174 and Marsh, 2011). Another noteworthy feature is the relatively quiet period 1908-1950 for
175 widespread, multi-year droughts (Table 1). Our criteria (i.e. two thirds of stations reporting
176 anomalies lasting at least three seasons) exclude well-known intense, but short-lived droughts
177 of 1921, 1933/34 and 1941-1943 that are evident in both IIP and EWP (Marsh *et al.*, 2007).

178 Overall, the longest unbroken runs of dry half-years in IIP were 14 seasons at Waterford
179 (1912-1919/1920); 12 at Cork Airport (1905-1911); 11 at both Markree Castle (1882-
180 1887/1888) and Mullingar (1904-1909) (Figure 3). All four dry-spells occurred prior to the
181 era of digital records (1941) but overlap with the coherent rainfall deficits (Table 1) of 1886-
182 1888 (at Markree Castle) and 1905-1907 (at Cork Airport and Mullingar). While no breaks in
183 this period were detected by Noone *et al.* (2015) the exceptionally long run of seasons with
184 below average precipitation at Waterford may be explained in part by documented
185 movements and sheltering of the rain gauge at Gortmore (used to bridge the record by
186 Tabony (1980)). Meta-data further signal that the Markree Castle gauge was 'leaking' and the
187 provenance of a 10 season dry spell at Belfast is questionable due to a large change of
188 correction factor applied by Tabony (1980) for bridging stations in the 1850s.

189 The most persistent dry-spell recorded anywhere in the IIP network since record digitisation
190 (1941) lasted 9 seasons at Cappoquin (1969-1973). This period partially overlaps with the
191 longest dry runs in the central and southeast part of the network covering Athboy, Birr Castle,
192 Drumsna, Enniscorthy, Foulkesmills, Portlaw and Roches Point (Table 2). The single season

193 drought in 1995 is noteworthy for the large precipitation anomaly (-35%) averaged across all
194 stations in the IIP network. The most recent 20 years have witnessed only a few single and
195 two season dry-spells (in 1996-97, 2001-02 and 2003-04) consistent with wetter and stormier
196 conditions (Sutton and Dong, 2012; Matthews *et al.*, 2014; 2015).

197 The relative quiescence of droughts since the 1990s is reflected by the moving average Pdd
198 and Pww indices (Figure 4). In particular, Pww shows non-stationary behaviour towards
199 more persistent wet-spells that is also evidenced by the EWP and SP series. The 30-year
200 mean Pww for IIP peaked in 2009 whereas Pdd is now lower than at any time since the
201 period 1938-1967, consistent with trends in SP. [Note that IIP and NIP are not independent
202 series because the former contains some records used to construct the latter. With this in
203 mind, the divergence in persistence behaviours over the last decade could reflect the
204 influence of the comparatively high density of stations within IIP along the east and southeast
205 seaboard].

206 The KS test indicates that simulated and observed spell distributions are statistically ($p < 0.05$)
207 indistinguishable across all regional (Figure 5) and station (Figure 6) series. The model tends
208 to overstate the frequency of single-season dry-spells and underestimate the occurrence of
209 two-season events. Overall, the geometric distribution yielded by the Markov model provides
210 good representations of the observed dry-spell length distribution. The closest match for dry-
211 spells is for Shannon airport (KS = 0.017) and greatest discrepancy for Ardara (KS = 0.111).
212 Note, however, that the KS results are for the whole distribution whereas the fit to tails is
213 more relevant for estimating low frequency events. Validation data are limited for this part of
214 the distribution so we are restricted to assessing the ability of the model to generate the
215 maximum observed dry-spell length at each site.

216 Table 2 shows the extent to which bootstrap Markov model simulations replicate the most
217 extreme runs of dry-spells in the 160-year series. The model overestimates the duration of the
218 160-year dry-spell by less than one season at 7 sites and by more than one season at 2 sites.
219 The largest discrepancy is for Portlaw where the model simulates a 7.6 season dry-spell
220 compared with 5 season run in observations. Meta data suggest that some precipitation totals
221 are too high at this site due to incorrect conversion between inches and mm. Conversely, the
222 maximum observed dry-spell is underestimated by less than one season at 5 sites and more
223 than one season at 13 sites. The largest difference is at Waterford with 14 observed and 7.5

224 simulated seasons. As noted above, this mismatch may be explained by the likely under-catch
225 and site changes affecting the Waterford record.

226 The 160- and 100-year simulations also provide likelihoods for a 10-season dry-spell for each
227 site and region (Table 2). This outcome is over three times more likely across EWP ($p=0.044$)
228 than for IIP ($p=0.012$). To date, the maximum observed dry-spell for Dublin is 8 seasons
229 (1903-1907), however, the Markov model suggests a relatively high likelihood ($p=0.125$) of a
230 10-season run of rainfall deficiencies in a 160-year record. A slightly higher likelihood is
231 estimated for Markree Castle ($p=0.127$) but this could be due to fitting the model to a record
232 with possible rainfall under-catch in the early part of the series. On the other hand, a 10-
233 season dry-run is least likely at Armagh ($p=0.008$), Birr Castle ($p=0.005$), Foulkesmills
234 ($p=0.008$) and Roches Point ($p=0.009$).

235 Comparison of probability distributions for simulated maximum dry- and wet-spell lengths
236 reveals three distinct patterns (Figure 7). There are sites with greater dry-spell persistence
237 than wet-spell persistence (Dublin, Enniscorthy, Markree Castle, Mullingar, Phoenix Park);
238 sites where wet- and dry-spell lengths have similar likelihoods (Ardara, Athboy, Belfast,
239 Cork, Derry, Drumsna, Malin Head, Portlaw, Rathdrum, Strokestown, UC Galway,
240 Waterford); and sites where a given wet-spell length is more likely than the same length dry-
241 spell (Armagh, Birr, Cappoquinn, Foulkesmills, Killarney, Roches Point, Shannon, Valentia).
242 Across all sites and 100-year simulations, the longest dry-spell was generated for Dublin and
243 persisted 24 seasons (not shown in Figure 7). This might appear implausible but Dublin
244 observations contain near unbroken runs exceeding 20 seasons in 1850-1868 (26/36), 1928-
245 1946 (23/36) and 1961-1978 (23/35).

246

247 **5. Discussion**

248 Using 1961-1990 as the reference period (and excluding Waterford and Markree Castle for
249 reasons noted above) we found that the longest observed run of below average precipitation
250 persisted 12 seasons at Cork Airport (1905-1911). Noone *et al.* (2015) note that this record
251 was originally constructed by Tabony (1980) using a composite of stations with data prior to
252 1962 based on a lower elevation gauge at University College Cork. This station change is
253 thought to explain lower early seasonal totals and a detected break point in 1958. The break

254 was adjusted by Noone *et al.* (2015) but the same correction factor was applied across all
255 months which could affect dry run persistence for this station.

256 The next longest run lasted 11 seasons at Mullingar (1904-1909) but, again, Noone *et al.*
257 (2015) report break points in 1937 and 1950 that could be due to a station change in the latter
258 case. Correction of the 1950 break resulted in a large downward adjustment, again potentially
259 affecting dry run persistence. The 10 season dry-spell at Belfast (1853-1858) has already
260 been queried, so the longest run now becomes 9 seasons at Ardara (1927-1932), Cappoquinn
261 (1969-1973), Phoenix Park (1903-1908) and Strokestown (1919-1912) (Table 2). The Ardara
262 record is based on a composite of stations with a small amplitude break point in 1983. While
263 Strokestown has been bridged from 1961, the years 1908-1961 represent a stable period in
264 the record (Noone *et al.*, 2015). No breaks were detected for Cappoquinn and there are no
265 issues of note from metadata. There are documented station moves early in the record at
266 Phoenix Park but these pre-date the identified dry run and a station inspection in 1903 noted a
267 very clear/open site. Therefore, having accounted for break points, station/instrument changes
268 and reported measurement errors the most credible, conservative upper bound continuous
269 dry-spell length for the IIP network is 9 seasons.

270 Our sub-annual analysis interrogated data that were homogenized at annual scales and thus
271 represents a stringent test of the IIP network. Anecdotal accounts, proxy sources and data
272 from neighbouring regions, all provide a basis for quality assuring our catalogue of
273 widespread multi-year rainfall deficits (Table 1). We find issues with two stations (Waterford
274 and Markree Castle) that were not picked up in the annual homogenisation of Noone *et al.*
275 (2015). While the confounding issues identified by metadata may have negligible effect at
276 annual resolution they can evidently become important when examining long duration
277 rainfall deficits. Additionally, suspicion is raised at Cork, Mullingar and Belfast that high
278 persistence of negative rainfall anomalies may be an artefact of using a single correction
279 factor equally across several months. Both issues arise despite application of best-practice
280 methods for homogenisation and emphasise the need for cautious use of homogenous series,
281 particularly when examining sequences of sub-annual extremes. [Note that snowfall is only a
282 small component of total precipitation across Ireland and thus any underestimation normally
283 associated with snowy climates is a minor concern]. Our analysis shows how metadata are
284 critical for increasing confidence in the authenticity of long-term precipitation indices.

285 There is strong independent evidence of persistent, regional droughts in the 1850s and 1880s
286 but bridging and homogenization techniques increase dependency between records as the
287 network density decreases further back in time. This is particularly the case for the 1850s
288 where only eight stations were active; by the 1880s this increases to 19. Thus, greater drought
289 coherence would be expected at the beginning of the IIP series than at the end due to the
290 smaller number of active stations. Hence, when evaluating the realism of Markov model
291 simulations there is ambiguity about whether inability to replicate dry-spells (>10 seasons) at
292 some sites is due to model deficiency, uncertainty in homogenized data, or both.

293 There is plenty of scope for developing more elaborate Markov model simulations for
294 Ireland. For example, seasonal Pdd and Pww parameters could be conditioned by the phase of
295 the North Atlantic Oscillation, Atlantic Multidecadal Oscillation, or El Niño Southern
296 Oscillation to replicate low-frequency variations (evident in Figure 4) and hence more
297 realistic clustering of dry-spells at decadal time-scales (e.g. Wilby et al., 2002). The
298 distribution of seasonal precipitation anomalies could be simulated using gamma or normal
299 functions. There is also scope for multi-site simulation of meteorological drought occurrence
300 and severity across the network as a whole and/or within homogeneous precipitation regions.
301 Such tools could be used to simulate groundwater recharge, river flow and reservoir levels for
302 vulnerable water supply zones, as well as for assessing potential environmental stress.

303 An important finding of our analysis is that recent decades have been relatively benign in
304 terms of widespread, multi-year sequences of below average rainfall in Ireland. This reflects
305 a return to generally stormier and wetter summers since the 1990s (Matthews et al., 2015).
306 Nonetheless, there is no room for complacency about drought risk given rising water
307 demands. Routinely updating the Pdd and Pww indices offers a simple way of tracking the
308 long-term propensity for seasonal rainfall deficits in Ireland.

309

310 **6. Conclusions**

311 We have investigated the spatial and temporal properties of long-lasting negative rainfall
312 anomalies across the Island of Ireland at site and regional scales with half-year granularity.
313 Our aim was to create the first coherent picture of multi-season rainfall deficit occurrence and
314 persistence across the region and, in the process, subject the IIP network to stringent
315 appraisal. Our preliminary analysis has highlighted the immense value of carefully

316 cataloguing station meta-data – an essential resource for interpreting break-points and
317 exceptional runs of below/above average precipitation. We acknowledge that interpretations
318 of spatial patterns are hindered by the sparse and uneven distribution of sites, as well as by
319 the range of issues picked up by meta-data, so we were restricted to describing three types of
320 spell-length regime. Further work is needed to determine whether these regimes form
321 coherent clusters in space.

322 Overall, we find that the Island of Ireland is surprisingly prone to runs of seasonal rainfall
323 deficiency and that major dry spells in the 1850s, 1880s and 1970s were far more persistent
324 than any episodes experienced in the last 40 years. These events could provide useful
325 analogues for stress testing the robustness of water supply and drought plans; a practice that
326 is finding favour elsewhere (e.g. Spraggs *et al.*, 2015). As Irish Water embarks on a period of
327 major investment in water infrastructure, stress testing designs against episodes with negative
328 rainfall anomalies lasting up to 9 seasons offers an altogether different risk assessment than
329 ability to cope with single season deficiencies. We also show that there is relatively high
330 likelihood ($p=0.125$) of a continuous 5 year (10 season) dry-spell at Dublin, a region in which
331 population growth and aging infrastructure has resulted in a water system operating at the
332 edge of its capacity.

333 In practice, water resource system vulnerability depends on a host of factors including the
334 type(s) of resource (i.e. groundwater, river intake, reservoir, or combination of sources);
335 amount of raw and treated water storage; connectivity of the system linking points of supply
336 to demand; water quality and treatment constraints. Such issues would clearly modulate any
337 assessment of droughts based on the analysis of meteorological data alone. Homogenised
338 rainfall series would need to be fed into more elaborate rainfall-runoff models and then, in
339 turn, simulated inflows input to water system models. Markov modelling, as demonstrated for
340 IIP, offers a way of generating severe drought sequences for evaluating water supply system
341 performance under combinations of long duration and intense rainfall deficits.

342 We have only begun to speculate about the underlying physical drivers of dry-spells lasting 5
343 or even 10 years. This is an area of active research, not least because of the potential to apply
344 such insights to long range drought forecasting (Folland *et al.*, 2015; Kingston *et al.*, 2015).
345 Assembling homogeneous meteorological records from paper and digital records (with
346 accompanying meta-data) is a laborious but critical part of this process. Creation of the IIP
347 series (Noone *et al.*, 2015), reference networks for river flow (Murphy *et al.*, 2013) and

348 attendant analytical tools (Wilby *et al.*, 2015) is bringing together ingredients needed for a
349 deeper understanding of multi-decadal hydroclimatic variability and change at a sentinel
350 location of Europe.

351

352 **Acknowledgements**

353 SN and SH are funded by the Irish Research Council. CM, TM and CB acknowledge funding
354 provided by the Irish Environmental Protection Agency under project 2014-CCRP-MS.16 .
355 The authors thank the anonymous referees for their diligent and constructive remarks.

356

357 **References**

358 Alexander, L.V. and Jones, P.D. 2001. Updated precipitation series for the UK and discussion
359 of recent extremes. *Atmospheric Science Letters*, doi:10.1006/asle.2001.0025.

360 Barker, P.A., Wilby, R.L. and Borrows, J. 2004. A 200-year precipitation index for the central
361 English Lake District. *Hydrological Sciences Journal*, **49**, 769-785.

362 Barrington, R. M. 1888. The drought of 1887, and some of its effects on Irish agriculture.
363 *Journal of the Statistical and Social Inquiry Society of Ireland*, Vol. IX Part LXVII, 223-247.

364 Bastola, S., Murphy, C. and Sweeney, J. 2011. The role of hydrological modelling
365 uncertainties in climate change impact assessments of Irish river catchments. *Advances in*
366 *Water Resources*, **34**, 562-576.

367 Brogan, L. and Cunnane, C. 2005. Low flows and low flow distributions for Ireland.
368 Understanding and Managing Hydrological Extremes. Irish National Committees of the IHP
369 and ICID National Hydrology Seminar 2005, 15th November, pp85-92.

370 Burt, T.P. and Howden, N.J.K. 2011. A homogeneous daily rainfall record for the Radcliffe
371 Observatory, Oxford, from the 1820s. *Water Resources Research*, **47**, W09701.

372 Burt, T.P., and Horton, B.P. 2007. Inter-decadal variability in daily rainfall at Durham (UK)
373 since the 1850s. *International Journal of Climatology*, **27**, 945-956.

374 Caussinus, H. and Mestre, O. 2004. Detection and correction of artificial shifts in climate
375 series. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, **53**, 405-425.

376 Dooge, J.C.I. 1985. Droughts in Irish history. In de Buitléar, É., (Ed.) *Irish Rivers*, Country
377 House Press, Dublin, pp26-28.

378 Dublin City Council, 2010. *The Plan: Water Supply Project – Dublin Region*. Report by RPS
379 and Veolia Water UK. Dublin City Council, Dublin, 125pp.

380 Folland, C.K., Hannaford, J., Bloomfield, J.P., Kendon, M., Svensson, C., Marchant, B.P.,
381 Prior, J. and Wallace, E. 2015. Multi-annual droughts in the English Lowlands: a review of
382 their characteristics and climate drivers in the winter half year. *Hydrology and Earth System
383 Sciences Discussion*, **11**, 12933-12985.

384 Garcia-Suarez, A.M. and Butler, C.J. 2006. Soil temperatures at Armagh Observatory,
385 Northern Ireland, from 1904 to 2002. *International Journal of Climatology*, **26**, 1075-1089.

386 Holden, N.M., Brereton, A.J., Fealy, T. and Sweeney, J. 2003. Possible change in Irish
387 climate and its impact on barley and potato yields. *Agricultural and Forest Meteorology*, **116**,
388 181-196.

389 Hunt, A.S.P., Wilby, R.L., Dale, N., Sura, K. and Watkiss, P. 2014. Embodied water imports
390 to the UK under climate change. *Climate Research*, **59**, 89-101.

391 Jones, P.D., Lister, D.H., Wilby, R.L. and Kostopoulou, E. 2006. Extended river flow
392 reconstructions for England and Wales, 1865-2002. *International Journal of Climatology*, **26**,
393 219-231.

394 Kingston, D.G., Stagge, J.H., Tallaksen, L.M. and Hannah, D.M. 2015. European-scale
395 drought: Understanding connections between atmospheric circulation and meteorological
396 drought indices. *Journal of Climate*, **28**, 505-516.

397 MacCarthaigh, M., 1996. *An assessment of the 1995 drought including a comparison with
398 other drought years*. Environmental Protection Agency, Dublin, 70pp.

399 Mandal, U.K. 2011. *Studies in low and flood flow estimation for Irish river catchments*.
400 Unpublished PhD thesis. National University of Ireland, Galway.

401 Marsh, T., Cole, G. and Wilby, R.L. 2007. Major droughts in England and Wales, 1800-2006.
402 *Weather*, **62**, 87-93.

403 Matthews, T., Murphy, C., Wilby, R.L. and Harrigan, S. 2014. Stormiest winter on record for
404 Ireland and UK. *Nature Climate Change*, **4**, 738-740.

405 Matthews, T., Murphy, C., Wilby, R.L. and Harrigan, S. 2015. A cyclone climatology of the
406 British-Irish Isles 1871-2012. *International Journal of Climatology*, doi:10.1002/joc.4425.

407 Mestre, O., Domonkos, P., Picard, F., Auer, I., Robin, S., Lebarbier, E., Böhm, R., Aguilar,
408 E., Guikarro, J., Vertachnik, G., Klan-car, M., Dubuisson, B., Stepanek, P. 2013. HOMER: A
409 Homogenization Software - Methods and Applications. *Idojaras*, **117**, 47-67.

410 Murphy, C., Harrigan, S., Hall, J. and Wilby, R.L. 2013. Assessing climate driven trends in
411 mean- and high- river flows from a network of reference stations in Ireland. *Hydrological
412 Sciences Journal*, **58**, 755-772.

413 Noone, S., Murphy, C., Coll, J., Matthews, T., Mullan, D., Wilby, R.L. and Walsh, S. 2015.
414 Homogenisation and analysis of an expanded monthly rainfall network for the Island of
415 Ireland (1850-2010). *International Journal of Climatology*, submitted.

416 O’Laoghog, S.S. 1979. *The dry period October 1974 to August 1976*. Meteorological Service,
417 Internal Memorandum 88/79, Dublin.

418 Rodda, J.C. and Marsh, T.J. 2011. *The 1975-76 Drought - a contemporary and retrospective
419 review*. Centre for Ecology & Hydrology, Wallingford.

420 Sen, Z. 1980. Statistical analysis of hydrologic critical droughts. *ASCE Journal of the
421 Hydraulics Division*, **106**, 99-115.

422 Sharma, T.C. and Panu, U.S. 2012. Prediction of hydrological drought durations based on
423 Markov chains: case of the Canadian prairies. *Hydrological Sciences Journal*, **57**, 705-722.

424 Sharma, T.C. and Panu, U.S. 2014a. Modeling of hydrological drought durations and
425 magnitudes: Experiences on Canadian streamflows. *Journal of Hydrology: Regional Studies*,
426 **1**, 92-106.

427 Sharma, T.C. and Panu, U.S. 2014b. A simplified model for predicting drought magnitudes: a
428 case of streamflow droughts in Canadian Prairies. *Water Resource Management*, **28**, 1597-
429 1611.

430 Spraggs, G., Peaver, L., Jones, P. and Ede, P. 2015. Re-construction of historic drought in the
431 Anglian Region (UK) over the period 1798-2010 and the implications for water resources and
432 drought management. *Journal of Hydrology*, **526**, 231-252.

433 Stead, D.R. 2014. Irish agriculture and agricultural policy during the hot, dry summer of
434 1976. *Agricultural History Review*, **62**, 337-359.

435 Steele-Dunne, S., Lynch, P., McGrath, R., Semmler, T., Wang, S., Hanafin, J. and Nolan, P.
436 2008. The impacts of climate change on hydrology in Ireland. *Journal of Hydrology*, **356**, 28-
437 45.

438 Sutton, R.T. and Dong, B. 2012. Atlantic Ocean influence on a shift in European climate in
439 the 1990s. *Nature Geoscience*, **5**, 788-792.

440 Symons GJ. 1887. *British Rainfall*. Edward Stanford, London.

441 Tabony, R.C. 1980. *A Set of Homogeneous European Rainfall Series*, Meteorological 13
442 Branch Memorandum No. 104, Meteorological Office, Bracknell.

443 Venema, V., Mestre, O., Aguilar, E., Auer, I. *et al.* 2012. Benchmarking homogenization
444 algorithms for monthly data. *Climate of the Past*, **8**, 89-115.

445 Wilby, R.L. 2007. Experimental seasonal rainfall forecasts for the River Medway, UK. *British*
446 *Hydrological Society National Meeting on Drought Forecasting*, London.

447 Wilby, R.L. and Quinn, N.W. 2013. Reconstructing multi-decadal variations in fluvial flood
448 risk using atmospheric circulation patterns. *Journal of Hydrology*, **487**, 109-121.

449 Wilby, R.L., Conway, D. and Jones, P.D. 2002. Prospects for downscaling seasonal
450 precipitation variability using conditioned weather generator parameters. *Hydrological*
451 *Processes*, **16**, 1215-1234.

452 Wilby, R.L., Prudhomme, C., Parry, S. and Muchan, K.G.L. 2015. Persistence of
453 hydrometeorological droughts in the United Kingdom: A regional analysis of multi-season
454 rainfall and river flow anomalies. *Journal of Extreme Events (Special Issue)*, in press.

455 Wilhite, D.A. and Glantz, M.H. 1985. Understanding the drought phenomenon: The role of
456 definitions. *Water International*, **10**, 111-120.

457

458

459 **Tables**

460

461 **Table 1** Periods with more than 2/3 of all IIP stations reporting below average seasonal
462 rainfall for at least three continuous seasons compared with dry-spell lengths in IIP, EWP and
463 SP. Note that NIP was not included because of the risk of double-counting with IIP.

Period	Number of seasons		
	IIP	EWP	SP
1853-1856	6	5	-
1858-1860	3	1	-
1886-1888	3	2	-
1892-1894	3	3	-
1905-1907	3	3	-
1951-1953	3	2	2
1962-1964	3	3	3
1970-1973	5	4	5
1974-1976	4	3	3

464

465 **Table 2** Observed maximum dry-spell duration (seasons) compared with simulated mean
 466 160- and 100-year events at each site as well as for IIP, NIP, EWP and SP. Likelihoods of a
 467 simulated 10-season dry-spell are also given.

Record	Observed Maximum event		Simulated 160-year event		Simulated 100-year event	
	Period(s)	Length (seasons)	Length (seasons)	Likelihood (10 season)	Length (seasons)	Likelihood (10 season)
Ardara	1927-32	9	6.7	0.028	6.1	0.017
Armagh	1892-95	7	5.5	0.008	5.1	0.002
Athboy	1969-73	8	7.0	0.044	6.4	0.027
Belfast	1853-58	10	7.4	0.058	6.6	0.029
Birr Castle	1970-73	5	5.2	0.005	4.7	0.001
Cappoquin	1969-73	9	6.4	0.021	5.8	0.015
Cork Airport	1905-11	12	7.8	0.086	7.1	0.045
Derry	1885-88	6	6.3	0.022	5.7	0.014
Drumsna	1966-70	7	6.6	0.024	6.0	0.018
Dublin Airport	1903-07	8	8.7	0.125	7.8	0.069
Enniscorthy	1969-73	8	7.8	0.068	7.0	0.057
Foulkesmills	1969-73	8	5.6	0.008	5.2	0.001
Killarney	1853-56, 1939-42	6	5.8	0.010	5.2	0.005
Malin Head	1950-54	7	6.9	0.033	6.3	0.022
Markree Castle	1882-88	11	8.6	0.127	7.9	0.089
Mullingar	1904-09	11	8.0	0.111	7.3	0.065
Phoenix Park	1903-08	9	7.9	0.070	7.2	0.053
Portlaw	1855-58, 1888-91, 1904-07, 1948-50, 1969-71, 2003-06	5	7.6	0.067	7.0	0.046
Rathdrum	1853-56	6	6.3	0.017	5.8	0.015
Roches Point	1941-43, 1961-63, 1969-71, 1974-76, 1990-92	4	5.7	0.009	5.2	0.008
Shannon Airport	1904-07	6	6.8	0.032	6.2	0.018
Strokestown	1919-22	9	6.2	0.027	5.8	0.008
UC Galway	1887-91	7	7.2	0.048	6.6	0.033
Valentia	1908-11, 1970-73	6	6.6	0.023	6.0	0.021
Waterford	1912-20	14	7.5	0.061	6.9	0.048
IIP	1969-73	8	6.0	0.012	5.6	0.009
NIP	1970-73	7	6.0	0.014	5.5	0.009
EWP	1900-03, 1904-07, 1941-44	8	7.0	0.044	6.5	0.034
SP	1970-73	6	5.6	0.008	5.0	0.004

468

469




471

472 **Figure 1** Map of station locations

473


474



475

476 **Figure 2** Above (blue) and below (red) long-term average precipitation totals in winter
 477 (October to March) and summer (April to September) half years (seasons) across a) the Island
 478 of Ireland and b) England and Wales for the years 1850-2010. All deviations are percentage
 479 anomalies with respect to the 1961-1990 mean.


480



481

482 **Figure 3** Long-run dry-spells at a) Waterford (1912-1919/1920), b) Cork Airport (1905-
 483 1911), c) Markree Castle (1882-1887/1888) and d) Mullingar (1904-1909)

484




485

486 **Figure 4** a) Dry-to-dry (Pdd) and b) wet-to-wet (Pww) season persistence for the Island of
487 Ireland, Northern Ireland, England and Wales, and Scotland. All series are based on 30-year
488 moving windows with anomalies referenced to the 1961-1990 mean. Adapted from Wilby et
489 al. (2015).

490

491

492



493

494 **Figure 5** Observed and modelled likelihood of dry-spells of duration 1 to 12 seasons in the
495 Island of Ireland (1850-2010), Northern Ireland (1931-2014), England and Wales (1766-
496 2014) and Scotland (1931-2014). Adapted from Wilby et al. (2015).




Figure 6 As in Fig.5 but 25 sites across the Island of Ireland.




Figure 7 Probability distributions of maximum simulated 100-year dry- (red lines) and wet- (blue lines) spells for Island of Ireland sites.