

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Dynamin function is important for CC-chemokine receptor induced cell migration

Richard O Jacques^a, Shirley C Mills^a, Paula Cazonatto Zerwes^a, Feyisope O
Fagade^a, John E Green^a, Scott Downham^a, Darren W Sexton^{b,c} and Anja Mueller^{a*}

a) School of Pharmacy, University of East Anglia, Norwich Research Park, Norwich
NR4 7TJ, UK, Telephone: 0044(0)1603-593138, Fax: 0044(0)1603-592003, email:
anja.mueller@uea.ac.uk

b) Norwich Medical School, University of East Anglia, Norwich Research Park,
Norwich NR4 7TJ, UK

c) current address: School of Pharmacy and Biomolecular Science, Liverpool John
Moores University, Byrom Street, Liverpool, L3 3HF

*Correspondence to this author

Abstract

The HIV viral entry co-receptors CCR5 and CXCR4 function physiologically as typical chemokine receptors. Activation leads to cytosolic signal transduction that results in a variety of cellular responses such as cytoskeletal rearrangement and chemotaxis (CTX). Our aim was to investigate the signalling pathways involved in CC- and CXC-receptor mediated cell migration. Inhibition of dynamin I and II GTPase with dynasore completely inhibited CCL3 stimulated chemotaxis in THP-1 cells, whereas the dynasore analogue dyngo-4a, which is a more potent inhibitor, showed reduced ability to inhibit CC-chemokine induced CTX. In contrast, dynasore was not able to block cell migration via CXCR4. The same activation/inhibition pattern was verified in activated-T-lymphocytes for different CC- and CXC-chemokines. Cell migration induced by CC- and CXC-receptors is not relying on active internalisation processes driven by dynamin since the blockade of internalisation does not affect migration, but they might rely on dynamin interaction with the cytoskeleton. We identify here a functional difference in how CC- and CXC-receptor migration is controlled, suggesting that specific signalling networks are being employed for different receptor classes and potentially specific therapeutic targets to prevent receptor migration can be identified.

Keywords

Chemokine receptor, chemotaxis, dynamin, signalling, internalisation

Introduction

Cellular migration can be activated by chemokine receptors, which are part of the G protein coupled receptors family (GPCRs) [1]. In different disease settings and different cancer types it has been shown that chemokine receptors play a crucial role in promoting cell migration and even cancer growth [2, 3]. Several chemokines (CCL5, CCL8) act as agonists for CCR1, CCR3 and CCR5, whereas a few chemokines, like CCL2, only activate CCR2 and CCR4, but not CCR5 [4, 5]. It has been shown that chemokine receptor activation leads to activation of heterotrimeric G proteins and phosphorylation of the receptor via G protein coupled receptor kinases (GRKs), which in turn leads to binding of β -arrestins to the receptor and is followed by receptor internalisation [4]. Activation is also followed by actin polymerisation but the signalling networks become which activated to allow this to happen have yet to be fully defined.

In recent years it has become clear that GPCRs do not signal solely via G proteins [6]. The so called receptosome of these receptors, which includes β -arrestins and other associating proteins makes signalling of GPCRs comparatively complex. β -arrestins can associate directly with a range of proteins including ERK1/2, cofilin, filamin and Jnk3 and therefore activate a variety of cellular responses without the involvement of G proteins [7]. Ligand biased signalling is important for chemokine receptors and it allows different ligands to activate different signalling cascades by encouraging specific ligand-receptor conformations. Receptors adopting such ligand dependent conformations then display specificity or bias towards certain signalling pathways dependent on which ligand is binding to the receptor [8] and which

1
2
3 receptor class is involved. There is also a marked difference in the regulation of
4
5 CXC-receptors and CC-receptors expression on the cell surfaces. Whereas CCR5,
6
7 CCR2 and CCR4 internalise via clathrin-coated-pits and caveolae, CXCR3 and
8
9 CXCR4 only use clathrin-coated pits [9-13]. Similarly, CCR5 recycles back to the cell
10
11 surface, whereas CXCR3 and CXCR4 are targeted to the late endosomes and
12
13 lysosomes [10, 12, 13].
14

15
16 Traditionally it is thought that $\beta\gamma$ -subunits of the G proteins induce migration via
17
18 activation of PI3K [14], however we have recently shown that this seems not to be
19
20 the case for CCL3 induced migration in THP-1 cells [15]. For CXCR4 it has been
21
22 shown that migration under certain circumstances is dependent on β -arrestins as
23
24 well as filamin-A, a protein, which can bind actin and interacts with β -arrestins [16,
25
26 17]. This raises the possibility that β -arrestin and actin interacting proteins are
27
28 activated downstream of different types of chemokine receptors. One of these actin
29
30 interacting proteins is dynamin. Several groups have shown that dynamin, an
31
32 enzyme which has traditionally been linked to internalisation of receptors via clathrin-
33
34 coated pits, is important for the integral structure of actin polymers [18-20].
35
36 Dynamins are large multi-domain proteins (~100 kDa) that constitute an N-terminal
37
38 GTPase domain, a middle domain, a PH domain, a GTPase effector domain (GED),
39
40 and a C-terminal proline-rich domain (PRD), which interacts with proteins that
41
42 contain SH3 domains [21] and there are several types of the protein: dynamin I is
43
44 primarily found in neurones where it is involved in synaptic vesicle endocytosis [22,
45
46 23] and it has been linked with several neurological processes such as long-term
47
48 memory formation [24]. Dynamin II is ubiquitously expressed and is found in all cell
49
50 types, dynamin III is primarily found in the testis. Dynamin II interacts with numerous
51
52 GPCRs as well as non-GPCR receptors, including the chemokine receptor CCR5 [25]
53
54
55
56
57
58
59
60

1
2
3 and various cytokine receptors [26] and is, therefore, an interesting target protein to
4
5 investigate chemokine receptor triggered migration.
6

7
8 Here we analysed different small molecule inhibitors for their effects on chemokine
9
10 receptor-induced migration and release of intracellular calcium. We investigated
11
12 whether dynamin plays a role for both CC-receptor- as well as CXC-receptor-
13
14 induced migration or whether distinct signalling pathways are activated by different
15
16 subsets of receptors.
17

18 19 20 **Methods**

21 *Cells and materials*

22
23
24
25 Culture conditions for THP-1 cells have previously been described [9]. Jurkat cells
26
27 were obtained from ATCC and grown in RPMI containing 10% FCS and 2 mM L-
28
29 glutamine. Blood was sampled from healthy normal subjects according to a protocol
30
31 approved by a local ethics committee (reference number 2008042). Peripheral blood
32
33 mononuclear cells (PBMCs) were subsequently isolated as previously described by
34
35 Sabroe et al. [27]. Lymphocytes were separated from monocytes by allowing the
36
37 latter to adhere to a tissue culture flask for 2 hours at 37 °C and 5% CO₂ and were
38
39 activated by culture in the presence of IL-2 (200 mg/mL) and concanavalin A (30
40
41 mg/mL) for at least 10 days. The chemokine used for CCR5/CCR1 activation was
42
43 human CCL3 (D26A) and has been described before [9, 28]. CXCL11 and CXCL12
44
45 were from Peprotech (UK). Dynamin inhibitors dynasore, dyngo-4a, MiTMAB,
46
47 OcTMAB, dynole-34-2, dynole-31-2 (negative control), iminodyn-22 and iminodyn-17
48
49 (negative control), pyrimidin-7 were purchased from Abcam (for an overview of
50
51 dynamin inhibitors see Table 1). Clathrin-mediated endocytosis inhibitor pitstop 2
52
53
54
55
56
57
58
59
60

1
2
3 and the corresponding negative control were from Abcam. All other chemicals were
4
5 from Fisher Scientific.
6
7

8 9 *Chemotaxis Assays*

10
11 Cells were harvested and washed twice with pre-warmed, sterile PBS, then
12
13 resuspended in serum-free RPMI 1640 which contained 0.1% BSA. The
14
15 concentration of cells was adjusted to 6.25×10^7 cells/mL. Chemoattractants were
16
17 loaded in a final volume of 31 μ L at indicated concentrations in the lower
18
19 compartment and 20 μ L of resuspended cells were loaded onto the upper
20
21 compartment of a microchemotaxis chamber (Receptor Technologies, Adderbury,
22
23 UK). The two compartments were separated by a polyvinylpyrrolidone-free
24
25 polycarbonate filter with 5 μ m pores. For inhibitor treatment, cells were incubated for
26
27 30 minutes with the inhibitor or with vehicle control before loading onto the upper
28
29 compartment of the chamber. Chambers were incubated at 37°C and 5% CO₂ for 4
30
31 hours before cells were counted. Data was analysed as previously described [15].
32
33
34
35
36
37
38
39

40 *Analysis of data*

41
42 Data were analysed using GraphPad Prism 5 (GraphPad Software). Statistical
43
44 analyses were performed using a one-way ANOVA with a Bonferroni multiple
45
46 comparison test as post-hoc test with a p value <0.05 deemed significant. In all
47
48 figures, data represent the mean \pm S.E.M. of at least three independent experiments.
49
50
51

52 **Results**

53
54
55
56
57
58
59
60

1
2
3 Chemokine receptors are expressed on different cell types, and THP-1 cells express
4 naturally CCR1, CCR2 and CCR5 as well as CXCR4 and migrate towards stimuli
5 with CCL2, CCL3, CCL8, CCL23 and CXCL12 whereas Jurkat cells express CXCR4
6 and migrate towards stimuli with CXCL12, but not towards CCL3. Activated T-cells
7 have been shown to express functional CXCR3 and CCR5 and migrate towards
8 CXCL11 and CCL3 [29]. We therefore used these different cells to investigate the
9 effect of dynamin inhibitors on cell migration with the view to differentiate between
10 CC-receptor and CXC-receptor family behaviour. In our hand Jurkat cells do not
11 migrate towards CCL3 and hence we used THP-1 cells for both CCL3 and CXCL12.
12 Dynasore blocks migration towards CCL3 in THP-1 cells in a dose dependant
13 manner (Figure 1a). At a concentration where dynasore clearly blocks CCL3 induced
14 migration (40 μ M), it does not affect CXCL12 induced migration in THP-1 (Figure 1b).
15 To rule out any ambiguities, we used the higher concentration of dynasore (80 μ M) in
16 activated T-cells. Dynasore does not block migration of activated T-cells towards
17 CXCL11, whereas, there is a clear trend of inhibition towards CCL3 induced
18 migration (Figure 1 c,d), showing a distinct difference in the activation pattern of
19 CXC- and CC-receptors. Confirming the differences between CC- and CXC-
20 receptors are results with CXCL12 in Jurkat cells, where dynasore has no effect at
21 all on migration, even at 80 μ M (Figure 1 e). At the concentration used, none of the
22 dynamin inhibitors showed any cytotoxic effects in the experimental set up, as shown
23 by MTS assays (data not shown).
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50
51 The dynasore analogue dyngo-4a, which is more potent than dynasore (dyngo-4a
52 IC_{50} 16 \pm 1.2 μ M versus dynasore 79.3 \pm 1.3 μ M [30]) blocks migration towards CCL3
53 in THP-1 cells to a lesser degree than dynasore. Remarkably dyngo-4a blocks
54
55
56
57
58
59
60

1
2
3 migration towards CXCL11 in activated T-cells and CXCL12 in THP-1 and Jurkat
4 cells (Figure 2). Dyngo-4a shows selectivity towards dynamin I versus dynamin II
5 whereas dynasore is non-selective and therefore these results might reflect a
6 different usage of the dynamin isoforms by different receptors (see Table 1). We
7 further investigated which domains of the dynamin proteins are essential for cell
8 migration and whether they are equally important for different receptor families. In
9 the first instance, we used iminodyn-22 and dynole-34-2 which are both non-
10 selective dynamin I and II inhibitors, and their negative controls which are iminodyn-
11 17 and dynole-31-2. Neither iminodyn-22 nor dynole-34-2 block migration in THP-1
12 cells towards CCL3 or in Jurkat cells towards CXCL12 (Figure 3 a-d). However there
13 is a distinct difference between CCL3- and CXCL12-induced chemotaxis for the non-
14 selective MiTMAB and OcTMAB inhibitors which bind to the dynamin PH domain [30],
15 and completely block any migration in Jurkat cells towards CXCL12 (Figure 3 f), but
16 have no significant effect on CCL3-induced migration in THP-1 cells (Figure 3 e,g)
17 but still effect CXCL12 migration in THP-1, even though with less of an effect than in
18 Jurkat cells (Figure 3 h). Again this data shows a clear difference in the reliance of
19 CC- and CXC-receptors on dynamin usage. We also used pyrimidyn-7, which
20 competitively inhibits both GTP and phospholipid binding and is the only inhibitor
21 available up to now which targets two distinct domains of dynamin. There is no effect
22 on CCL3 induced migration in THP-1 cells (Figure 4 a), but CXCL12 induced
23 migration is significantly blocked in Jurkat cells (Figure 3 b) as well as THP-1 cells
24 (data not shown).

25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

Dynamin is classically known as being of importance for clathrin-coated pit triggered internalisation of receptors, even though recently its importance for actin-dynamics have become more apparent [19]. We previously showed that CCR5 can use

1
2
3 clathrin-coated pits for internalisation [9, 31] and indeed dynamin inhibition via
4
5 dynasore completely abrogates internalisation on CHO.CCR5 as well as THP-1 cells
6
7 as analysed via immunofluorescence (data not shown). To investigate whether it is
8
9 the prohibition of internalisation which prevents cell migration, we used another
10
11 clathrin-coated pit endocytosis inhibitor, pitstop 2 and its negative control analogue in
12
13 THP-1 cells for CCL3 activation as well as in Jurkat cells for CXCL12 activation.
14
15 Pitstop 2 does not block cell migration (Figure 4 c,d). The concentration of pitstop 2
16
17 used for migration assays actually inhibits internalisation of CCR5 receptor in THP-1
18
19 cells (data not shown). An increase of the concentration of pitstop 2 used in THP-1
20
21 cells, actually increased the number of migrating cells by a small, but significant
22
23 amount.
24
25
26
27
28
29

30 Discussion

31
32
33
34 In this study we investigated the role of dynamin in the signalling events that occur
35
36 after the activation of CC- and CXC-receptors. Dynamin involvement in cell migration
37
38 is related to its role as a focal adhesion regulator and it has been shown that
39
40 inhibition of dynamin 2 inhibits focal adhesion disassembly and impairs cell migration.
41
42 We, therefore, used different dynamin inhibitors, which either have a higher potency
43
44 for dynamin I over II (dyngo-4a) or are non-selective dynamin I and II inhibitors
45
46 (dynasore, dynole-34-2, MiTMAB, OcTMAB, iminodyn-22, pyrimidyn-7) [30, 32, 33].
47
48 Dynasore has been shown previously to block endocytosis via clathrin-coated pits
49
50 [34-36] and indeed it blocks CCL3-induced endocytosis of CCR5 in CHO.CCR5 cells.
51
52 Dynasore blocks CCL3 induced migration in THP-1 cells and activated T-
53
54 lymphocytes, but it has no effect on either CXCL12-induced migration of THP-1 or
55
56
57
58
59
60

1
2
3 Jurkat cells or CXCL11-induced migration of activated T-lymphocytes. These results
4
5 point towards a significant difference between CC- and CXC-receptor activated
6
7 signalling networks. Dyngo-4a, a close analogue of dynasore, which is more potent
8
9 than dynasore and has a higher potency for dynamin I over dynamin II [30], is less
10
11 effective in blocking CCL3-induced migration in THP-1 cells, however it blocks
12
13 CXCL11- and CXCL12-induced migration in activated T-lymphocytes and Jurkat
14
15 cells, respectively. Similarly, dynasore can significantly reduce CCL2 induced
16
17 migration in THP-1 cells, whereas dyngo-4a shows a trend to inhibit migration, but
18
19 does not reach significance. The functional differences between dynasore and
20
21 dyngo-4a have not been fully analysed yet, but with the knowledge available today,
22
23 our data points towards either a different usage of dynamin isoforms by chemokine
24
25 receptor subtypes or to the usage of a varying set of dynamin interacting protein by
26
27 different receptor subfamilies. This difference between CC- and CXC-receptors was
28
29 further highlighted by the use of dynole-34-2, MiTMAB, OcTMAB, iminodyn-22 and
30
31 pyrimidyn-7. None of those blocked CCL3-induced migration in THP-1 cells, but
32
33 pyrimidyn-7, MiTMAB and OcTMAB block CXCL12-induced migration in THP-1 and
34
35 Jurkat cells. Unlike MiTMAB and OcTMAB, which block dynamin recruitment to the
36
37 membranes, dynole-34-2, dyngo-4a and dynasore block dynamin function after its
38
39 recruitment [30]. MiTMAB, OcTMAB also bind to the PH-domain of the dynamin
40
41 molecule, unlike the other inhibitors, which bind to the G domain. An obvious reason
42
43 for the prevention of migration after the use of dynamin inhibitors is the potential
44
45 importance of internalisation for receptor activation and ultimately signal transduction.
46
47 We, therefore, employed a different clathrin-coated pits endocytosis inhibitor, pitstop
48
49 2 and its negative control compound to analyse whether internalisation is a pre-
50
51 requisite for migration. In both THP-1 cells and Jurkat cells pitstop 2 did not prevent
52
53
54
55
56
57
58
59
60

1
2
3 CCL3- and CXCL12-induced migration, respectively, which is evidence that receptor
4
5 internalisation is not necessary to activate cell migration as had been described
6
7 already for the CCR2b receptor [37].
8

9
10 In our study we detect distinct differences between the CC- and CXC-receptors.
11
12 Traditionally it has been shown that CXCR4 activation leads to chemotaxis in a β -
13
14 arrestin 2, ERK 1/2, G β γ dependent manner and is PI3K dependent [38, 39].
15
16 Therefore the implication of the dynamin PH domain in cell migration for CXCL12 is
17
18 in line with the already published signalling networks, whereas the PH domain is not
19
20 necessary for CCL3-induced migration, since this migration is independent of PI3K
21
22 activation [15]. Overall our study showed that there are distinct differences in
23
24 signalling networks used by CC-receptors compared to CXC-receptors which will
25
26 yield novel therapeutic targets to prevent cell migration triggered by specific
27
28 receptors.
29
30
31
32
33

34 **Acknowledgments:**

35
36 We thank the UEA for providing a studentship for ROJ and we acknowledge the
37
38 support for SCM from the Novartis studentship fund.
39
40
41
42

43 **References**

- 44
45
46
47 [1] M. Thelen, Dancing to the tune of chemokines, *Nat Immunol*, 2 (2001) 129-134.
48
49 [2] B. Homey, A. Muller, A. Zlotnik, Chemokines: agents for the immunotherapy of
50
51 cancer?, *Nat Rev Immunol*, 2 (2002) 175-184.
52
53
54 [3] A. Zlotnik, Chemokines and cancer, Ernst Schering Res Found Workshop, (2004)
55
56 53-58.
57
58
59
60

- 1
2
3 [4] F. Bachelierie, A. Ben-Baruch, A.M. Burkhardt, C. Combadiere, J.M. Farber, G.J.
4 Graham, R. Horuk, A.H. Sparre-Ulrich, M. Locati, A.D. Luster, A. Mantovani, K.
5 Matsushima, P.M. Murphy, R. Nibbs, H. Nomiya, C.A. Power, A.E. Proudfoot,
6 M.M. Rosenkilde, A. Rot, S. Sozzani, M. Thelen, O. Yoshie, A. Zlotnik, International
7 Union of Pharmacology. LXXXIX. Update on the Extended Family of Chemokine
8 Receptors and Introducing a New Nomenclature for Atypical Chemokine Receptors,
9 Pharmacological reviews, 66 (2014) 1-79.
10
11 [5] A. Zlotnik, O. Yoshie, The chemokine superfamily revisited, Immunity, 36 (2012)
12 705-716.
13
14 [6] A.C. Magalhaes, H. Dunn, S.S. Ferguson, Regulation of GPCR activity, trafficking
15 and localization by GPCR-interacting proteins, Br J Pharmacol, 165 (2012) 1717-
16 1736.
17
18 [7] S. Rajagopal, K. Rajagopal, R.J. Lefkowitz, Teaching old receptors new tricks:
19 biasing seven-transmembrane receptors, Nature reviews. Drug discovery, 9 (2010)
20 373-386.
21
22 [8] S. Rajagopal, D.L. Bassoni, J.J. Campbell, N.P. Gerard, C. Gerard, T.S.
23 Wehrman, Biased agonism as a mechanism for differential signaling by chemokine
24 receptors, J Biol Chem, 288 (2013) 35039-35048.
25
26 [9] A. Mueller, E. Kelly, P.G. Strange, Pathways for internalization and recycling of
27 the chemokine receptor CCR5, Blood, 99 (2002) 785-791.
28
29 [10] A. Meiser, A. Mueller, E.L. Wise, E.M. McDonagh, S.J. Petit, N. Saran, P.C.
30 Clark, T.J. Williams, J.E. Pease, The Chemokine Receptor CXCR3 Is Degraded
31 following Internalization and Is Replenished at the Cell Surface by De Novo
32 Synthesis of Receptor, J Immunol, 180 (2008) 6713-6724.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 [11] M. Mariani, R. Lang, E. Binda, P. Panina-Bordignon, D. D'Ambrosio, Dominance
4 of CCL22 over CCL17 in induction of chemokine receptor CCR4 desensitization and
5 internalization on human Th2 cells, *Eur J Immunol*, 34 (2004) 231-240.
6
7
8
9 [12] E.M. Borroni, A. Mantovani, M. Locati, R. Bonecchi, Chemokine receptors
10 intracellular trafficking, *Pharmacology & therapeutics*, 127 (2010) 1-8.
11
12 [13] Y. Zhang, A. Foudi, J.F. Geay, M. Berthebaud, D. Buet, P. Jarrier, A. Jalil, W.
13 Vainchenker, F. Louache, Intracellular localization and constitutive endocytosis of
14 CXCR4 in human CD34+ hematopoietic progenitor cells, *Stem Cells*, 22 (2004)
15 1015-1029.
16
17
18 [14] G. Germena, E. Hirsch, PI3Ks and small GTPases in neutrophil migration: two
19 sides of the same coin, *Mol Immunol*, 55 (2013) 83-86.
20
21
22 [15] J.S. Kerr, R.O. Jacques, C. Moyano Cardaba, T. Tse, D. Sexton, A. Mueller,
23 Differential regulation of chemotaxis: role of Gbetagamma in chemokine receptor-
24 induced cell migration, *Cell Signal*, 25 (2013) 729-735.
25
26
27 [16] B. Lagane, K.Y. Chow, K. Balabanian, A. Levoye, J. Harriague, T. Planchenault,
28 F. Baleux, N. Gunera-Saad, F. Arenzana-Seisdedos, F. Bachelierie, CXCR4
29 dimerization and beta-arrestin-mediated signaling account for the enhanced
30 chemotaxis to CXCL12 in WHIM syndrome, *Blood*, 112 (2008) 34-44.
31
32
33 [17] M.G. Scott, V. Pierotti, H. Storez, E. Lindberg, A. Thuret, O. Muntaner, C.
34 Labbe-Jullie, J.A. Pitcher, S. Marullo, Cooperative regulation of extracellular signal-
35 regulated kinase activation and cell shape change by filamin A and beta-arrestins,
36 *Molecular and cellular biology*, 26 (2006) 3432-3445.
37
38
39 [18] S.M. Ferguson, A. Raimondi, S. Paradise, H. Shen, K. Mesaki, A. Ferguson, O.
40 Destaing, G. Ko, J. Takasaki, O. Cremona, O.T. E, P. De Camilli, Coordinated
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 actions of actin and BAR proteins upstream of dynamin at endocytic clathrin-coated
4 pits, *Developmental cell*, 17 (2009) 811-822.

5
6
7 [19] J. Chua, R. Rikhy, J. Lippincott-Schwartz, Dynamin 2 orchestrates the global
8 actomyosin cytoskeleton for epithelial maintenance and apical constriction, *Proc Natl*
9 *Acad Sci U S A*, 106 (2009) 20770-20775.

10
11
12 [20] G. Gallo, More than one ring to bind them all: recent insights into the structure of
13 the axon, *Developmental neurobiology*, 73 (2013) 799-805.

14
15
16 [21] H.S. Shpetner, R.B. Vallee, Dynamin is a GTPase stimulated to high levels of
17 activity by microtubules, *Nature*, 355 (1992) 733-735.

18
19
20 [22] P. De Camilli, K. Takei, P.S. McPherson, The function of dynamin in endocytosis,
21 *Current opinion in neurobiology*, 5 (1995) 559-565.

22
23
24 [23] K. Tomizawa, S. Sunada, Y.F. Lu, Y. Oda, M. Kinuta, T. Ohshima, T. Saito, F.Y.
25 Wei, M. Matsushita, S.T. Li, K. Tsutsui, S. Hisanaga, K. Mikoshiba, K. Takei, H.
26 Matsui, Cophosphorylation of amphiphysin I and dynamin I by Cdk5 regulates
27 clathrin-mediated endocytosis of synaptic vesicles, *J Cell Biol*, 163 (2003) 813-824.

28
29
30 [24] T. Casoli, G. Di Stefano, P. Fattoretti, B. Giorgetti, M. Balialetti, F. Lattanzio, G.
31 Aicardi, D. Platano, Dynamin binding protein gene expression and memory
32 performance in aged rats, *Neurobiology of aging*, 33 (2012) 618 e615-619.

33
34
35 [25] K. Tadagaki, D. Tudor, F. Gbahou, P. Tschische, M. Waldhoer, M. Bomsel, R.
36 Jockers, M. Kamal, Human cytomegalovirus-encoded UL33 and UL78 heteromerize
37 with host CCR5 and CXCR4 impairing their HIV coreceptor activity, *Blood*, 119 (2012)
38 4908-4918.

39
40
41 [26] K. Kawada, G. Upadhyay, S. Ferandon, S. Janarthanan, M. Hall, J.P. Vilardaga,
42 V. Yajnik, Cell migration is regulated by platelet-derived growth factor receptor
43 endocytosis, *Molecular and cellular biology*, 29 (2009) 4508-4518.

1
2
3 [27] I. Sabroe, A. Hartnell, L.A. Jopling, S. Bel, P.D. Ponath, J.E. Pease, P.D. Collins,
4
5 T.J. Williams, Differential regulation of eosinophil chemokine signaling via CCR3 and
6
7 non-CCR3 pathways, *J Immunol*, 162 (1999) 2946-2955.
8

9
10 [28] A. Mueller, N.G. Mahmoud, P.G. Strange, Diverse signalling by different
11
12 chemokines through the chemokine receptor CCR5, *Biochem Pharmacol*, 72 (2006)
13
14 739-748.
15

16 [29] M. Loetscher, B. Gerber, P. Loetscher, S.A. Jones, L. Piali, I. Clark-Lewis, M.
17
18 Baggiolini, B. Moser, Chemokine receptor specific for IP10 and mig: structure,
19
20 function, and expression in activated T-lymphocytes, *J Exp Med*, 184 (1996) 963-969.
21

22 [30] C.B. Harper, M.R. Popoff, A. McCluskey, P.J. Robinson, F.A. Meunier, Targeting
23
24 membrane trafficking in infection prophylaxis: dynamin inhibitors, *Trends in cell*
25
26 *biology*, 23 (2013) 90-101.
27

28 [31] A. Mueller, P.G. Strange, Mechanisms of internalization and recycling of the
29
30 chemokine receptor, CCR5, *Eur J Biochem*, 271 (2004) 243-252.
31

32 [32] A.B. McGeachie, L.R. Odell, A. Quan, J.A. Daniel, N. Chau, T.A. Hill, N.N.
33
34 Gorgani, D.J. Keating, M.A. Cousin, E.M. van Dam, A. Mariana, A. Whiting, S.
35
36 Perera, A. Novelle, K.A. Young, F.M. Deane, J. Gilbert, J.A. Sakoff, M. Chircop, A.
37
38 McCluskey, P.J. Robinson, Pyrimidyn compounds: dual-action small molecule
39
40 pyrimidine-based dynamin inhibitors, *ACS chemical biology*, 8 (2013) 1507-1518.
41

42 [33] T.A. Hill, A. Mariana, C.P. Gordon, L.R. Odell, M.J. Robertson, A.B. McGeachie,
43
44 N. Chau, J.A. Daniel, N.N. Gorgani, P.J. Robinson, A. McCluskey, Iminochromene
45
46 inhibitors of dynamins I and II GTPase activity and endocytosis, *J Med Chem*, 53
47
48 (2010) 4094-4102.
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 [34] S. Lee, K.Y. Jung, J. Park, J.H. Cho, Y.C. Kim, S. Chang, Synthesis of potent
4 chemical inhibitors of dynamin GTPase, *Bioorg Med Chem Lett*, 20 (2010) 4858-
5 4864.
6
7

8
9
10 [35] C.B. Harper, S. Martin, T.H. Nguyen, S.J. Daniels, N.A. Lavidis, M.R. Popoff, G.
11 Hadzic, A. Mariana, N. Chau, A. McCluskey, P.J. Robinson, F.A. Meunier, Dynamin
12 inhibition blocks botulinum neurotoxin type A endocytosis in neurons and delays
13 botulism, *J Biol Chem*, 286 (2011) 35966-35976.
14
15
16

17
18 [36] A. McCluskey, J.A. Daniel, G. Hadzic, N. Chau, E.L. Clayton, A. Mariana, A.
19 Whiting, N.N. Gorgani, J. Lloyd, A. Quan, L. Moshkanbaryans, S. Krishnan, S.
20 Perera, M. Chircop, L. von Kleist, A.B. McGeachie, M.T. Howes, R.G. Parton, M.
21 Campbell, J.A. Sakoff, X. Wang, J.Y. Sun, M.J. Robertson, F.M. Deane, T.H.
22 Nguyen, F.A. Meunier, M.A. Cousin, P.J. Robinson, Building a better dynasore: the
23 dyngo compounds potently inhibit dynamin and endocytosis, *Traffic*, 14 (2013) 1272-
24 1289.
25
26
27
28
29
30
31
32
33

34 [37] H. Arai, C.L. Tsou, I.F. Charo, Chemotaxis in a lymphocyte cell line transfected
35 with C-C chemokine receptor 2B: evidence that directed migration is mediated by
36 betagamma dimers released by activation of Galphai-coupled receptors, *Proc Natl*
37 *Acad Sci U S A*, 94 (1997) 14495-14499.
38
39
40
41
42

43 [38] L.J. Drury, J.J. Ziarek, S. Gravel, C.T. Veldkamp, T. Takekoshi, S.T. Hwang, N.
44 Heveker, B.F. Volkman, M.B. Dwinell, Monomeric and dimeric CXCL12 inhibit
45 metastasis through distinct CXCR4 interactions and signaling pathways, *Proc Natl*
46 *Acad Sci U S A*, 108 (2011) 17655-17660.
47
48
49
50

51 [39] C. Delgado-Martin, C. Escribano, J.L. Pablos, L. Riol-Blanco, J.L. Rodriguez-
52 Fernandez, Chemokine CXCL12 uses CXCR4 and a signaling core formed by
53 bifunctional Akt, extracellular signal-regulated kinase (ERK)1/2, and mammalian
54
55
56
57
58
59
60

1
2
3 target of rapamycin complex 1 (mTORC1) proteins to control chemotaxis and
4
5 survival simultaneously in mature dendritic cells, J Biol Chem, 286 (2011) 37222-
6
7 37236.
8
9

10 11 12 13 14 15 **Figures and legends**

16
17
18
19 **Figure 1:** Cell migration towards CCL3 but not towards CXCL11 or CXCL12 is
20 blocked by dynasore. a) THP-1 cells were treated with 16, 40 or 80 μ M of dynasore.
21
22 b) THP-1 cells were treated with 40 μ M of dynasore. Migration was induced with 1
23
24 nM CXCL12. c) Activated T-lymphocytes were treated with 80 μ M of dynasore and
25
26 migration was induced with 20 nM CCL3. d) Activated T-lymphocytes were treated
27
28 with 80 μ M of dynasore and migration was induced with 1 nM CXCL11. e) Jurkat
29
30 cells were treated with 80 μ M of dynasore and migration was induced with 1 nM
31
32 CXCL12. Base level of migration was determined in the absence of chemokines.
33
34 Statistical analysis were performed using a one-way ANOVA with a Bonferroni
35
36 Multiple comparison test as post-test ** showing a p value <0.01 and *** <0.001 .
37
38 Data represent the mean \pm S.E.M. of at least three independent experiments.
39
40
41
42
43
44
45

46
47 **Figure 2:** Cell migration towards CCL3, CXCL11 and CXCL12 is blocked by dyngo-
48
49 4a. a) THP-1 cells were treated with 80 μ M of dyngo-4a. Migration was induced with
50
51 1 nM CCL3. b) Activated T-lymphocytes were treated with 80 μ M of dyngo-4a and
52
53 migration was induced with 20 nM CXCL11. c) THP-1 cells were treated with 80 μ M
54
55 of dyngo-4a and migration was induced with 1 nM CXCL12. d) Jurkat cells were
56
57 treated with 80 μ M of dyngo-4a and migration was induced with 1 nM CXCL12.
58
59
60

1
2
3 Statistical analysis were performed using a one-way ANOVA with a Bonferroni
4
5 Multiple comparison test as post-test with * showing a p value <0.05, ** showing a p
6
7 value <0.01 and *** showing a p value < 0.001. Data represent the mean \pm S.E.M. of
8
9 at least three independent experiments.
10

11
12
13
14 **Figure 3:** Effect of different dynamin inhibitors on migration towards CCL3 and
15
16 CXCL12. a) THP-1 cells were treated with 1 μ M of iminodn-17 and 1 μ M of
17
18 iminodn-22. Migration was induced with 1 nM CCL3. b) Jurkat cells were treated
19
20 with 1 μ M of iminodn-17 and 1 μ M of iminodn-22. Migration was induced with 1 nM
21
22 CXCL12. c) THP-1 cells were treated with 15 μ M dynole-31-2 and 15 μ M dynole-34-
23
24 2 and migration was induced with 1 nM CCL3 d) Jurkat cells were treated with 15 μ M
25
26 dynole-31-2 and 15 μ M dynole-34-2 and migration was induced with 1 nM CXCL12.
27
28 e) THP-1 cells were treated with 10 μ M of MiTMAB and migration was induced with 1
29
30 nM CCL3. f) Jurkat cells were treated with 10 μ M of MiTMAB and 5 μ M of OcTMAB.
31
32 Migration was induced with 1 nM CXCL12. g) THP-1 cells were treated with 5 μ M of
33
34 OcTMAB. Migration was induced with 1 nM CCL3. h) THP-1 cells were treated with
35
36 10 μ M of MiTMAB and 5 μ M of OcTMAB. Migration was induced with 1 nM CXCL12.
37
38 Statistical analysis were performed using a one-way ANOVA with a Bonferroni
39
40 Multiple comparison test as post-test with *** showing a p value < 0.001. Data
41
42 represent the mean \pm S.E.M. of at least three independent experiments.
43
44
45
46
47
48

49
50 **Figure 4:** Effect of endocytosis inhibitors on migration towards CCL3 and CXCL12. a)
51
52 THP-1 cells were treated with 10 μ M of pyrimidyn-7. Migration was induced with 1
53
54 nM CCL3. b) THP-1 cells were treated with 10 μ M of pyrimidyn-7. Migration was
55
56 induced with 1 nM CXCL12 and migrated cells were counted after 4 hours. c) THP-1
57
58
59
60

1
2
3 cells were treated with 1.25 μM of pitstop 2 and pitstop 2 negative control compound
4
5 and 30 μM of pitstop 2 and pitstop 2 negative control compound, respectively.
6
7 Migration was induced with 1 nM CCL3. d) Jurkat cells were treated with 30 μM of
8
9 pitstop 2 and pitstop 2 negative control compound, respectively and migration was
10
11 induced with 1 nM CXCL12. Statistical analysis were performed using a one-way
12
13 ANOVA with a Bonferroni Multiple comparison test as post-test with ** showing a p
14
15 value < 0.01 and *** showing a p value < 0.001. Data represent the mean \pm S.E.M.
16
17 of at least three independent experiments.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1

190x254mm (96 x 96 DPI)




Figure 2

190x254mm (96 x 96 DPI)




Figure 3

190x254mm (96 x 96 DPI)




Figure 4

190x254mm (96 x 96 DPI)

Table 1: Overview of dynamin inhibitors

Inhibitor	Mode of action	IC _{50(SVE)} ^a (μM)	Specificity Dynamin I versus II (μM)	Refs
Dyngo-4a TM	G domain: allosteric site	16±1.2	Dynamin I selective 0.38±0.05 vs 2.6±0.12	[30, 35]
Dynasore	Unknown: non-competitive inhibition	79.3±1.3	Non-selective	[35]
Dynole-34-2 TM	G domain: uncompetitive with GTP	105	Non-selective	[36]
Iminodyn 22 TM	G domain: uncompetitive with GTP	99.5±1.7	Non-selective	[33]
Pyrimidin-7 TM	Competitively inhibits both GTP and phospholipid binding	Not reported	Non-selective 1.1vs1.8	[32]
MiTMAB TM	PH domain: competitive with lipid and non-competitive with GTP	105	Non-selective	[36]
OcTMAB TM	PH domain: competitive with lipid and non-competitive with GTP	Not reported	Non-selective	[36]

^a FM4-64 uptake in synaptosomes