

The Identification of Nuclear Receptors Associated with Hepatic Steatosis to Develop and Extend Adverse Outcome Pathways

C.L. Mellor, F.P. Steinmetz and M.T.D. Cronin*

School of Pharmacy and Biomolecular Sciences, Liverpool John Moores University, Byrom Street, Liverpool L3 3AF, England

*Corresponding author: Tel. +44 151 231 2402; e-mail address: M.T.Cronin@ljmu.ac.uk
(Mark Cronin)

Keywords

Adverse Outcome Pathway (AOP), hepatic toxicity, mechanisms of toxicity, nuclear receptor, steatosis

Abstract

The development of Adverse Outcome Pathways (AOPs) is becoming a key component of 21st century toxicology. AOPs provide a conceptual framework that links the molecular initiating event to an adverse outcome through organised toxicological knowledge, bridging the gap from chemistry to toxicological effect. As nuclear receptors (NRs) play essential roles for many physiological processes within the body, they are used regularly as drug targets for therapies to treat many diseases including diabetes, cancer and neurodegenerative diseases. Due to the heightened development of NR ligands there is increased need for the identification of related AOPs to facilitate their risk assessment. Many NR ligands have been linked specifically to steatosis. This paper reviews and summarises the role of NR and their importance with links between NR examined to identify plausible putative AOPs. The following NRs are shown to induce hepatic steatosis upon ligand binding: aryl hydrocarbon receptor, constitutive androstane receptor, oestrogen receptor, glucocorticoid receptor, farnesoid X receptor, liver X receptor, peroxisome proliferator-activated receptor, pregnane X receptor, and the retinoic acid receptor. A preliminary, putative AOP was formed for NR binding linked to hepatic steatosis as the adverse outcome.

Table of Contents

Abstract

Introduction

AOP Development

Hepatic Steatosis

Nuclear Receptors

Androgen Receptor (AR)

Aryl Hydrocarbon Receptor (AHR)

Constitutive Androstane Receptor (CAR)

oEstrogen Receptor (ER)

Farnesoid X receptor (FXR)

Glucocorticoid Receptor (GR)

Liver X Receptor (LXR)

Peroxisome Proliferator-Activated Receptor (PPAR)

Progesterone Receptor (PR)

Pregnane X Receptor (PXR)

Retinoic Acid Receptor (RAR)

Retinoid X Receptor (RXR)

Thyroid Hormone Receptor (THR)

Vitamin D Receptor (VDR)

AOP Formation

Acknowledgement

Declaration of Interest

References

Introduction

AOP Development

The impact of 21st century life has led to increased demand for safer and more sustainable chemical products, whilst at the same time reducing the amount of animal testing (Patlewicz et al., 2015). Animal testing has been integral to toxicology studies for over 50 years; however, the need for more modern alternative toxicological approaches is now a key focus for toxicologists and regulatory scientists alike (NRC, 2007). For instance, the results gained from animal testing require many presumptions and extrapolations in order to predict human health effects and this process is still controversial due to both scientific and ethical reasons (NRC, 2007). Current chemical safety assessments require faster testing with the use of fewer animals, at the same time the number of chemicals being tested are rapidly increasing worldwide, making the cost of chemical safety assessment substantial (Patlewicz et al., 2015). The constraints of the modern chemical safety assessments, along with the increased cost of testing, have induced a new mind-set among toxicologist aiming to replace the more traditional extensive phenotypic animal testing with a more mechanistic based approach relying on the use of toxicokinetics, computational models and *in vitro* testing (Patlewicz et al., 2015; Vinken, 2013). The momentum of this change has led to big strides in the development of *in vitro* assays for use in high throughput screening and high content screening (seen in projects such as the United States Environmental Protection Agency's (US EPA's) ToxCast programme) (Cohen et al., 2010; NRC, 2007) and the development of computational approaches. Computational models are integral to the development of integrated alternative methods to identify organ level toxicity and ultimately leading towards the replacement of animal testing (Cronin and Livingstone, 2004). These models traditionally include the use of (quantitative) structure-activity relationships ((Q)SARs). QSARs are mathematical models that predict biological activity of chemicals from structure or physicochemical properties, whereas a SAR is a

qualitative association between a specific molecular substructure and biological activity (Cronin and Livingstone, 2004).

(Q)SARs are currently used successfully for the prediction of single dose acute toxicity and specific endpoints such as mutagenicity, however, the use of QSARs for the prediction of repeat dose systemic toxicity, carcinogenicity and reproductive toxicity poses a real challenge for computational scientists (Alder et al., 2010; Gocht et al., 2015; Hartung et al., 2011). This has led to the development of integrated testing strategies which use a combination of computational models (such as structural alerts, read across, QSARs and modelling) along with *in vitro* testing in order to develop alternative methods. The integration of alternative methods with pathways of toxicity has, in part, also catalysed the formation of the Adverse Outcome Pathway (AOP) framework concept (Ankley et al., 2010; Groh et al., 2015; Vinken, 2013, 2015). An AOP describes the causal linkage between a molecular initiating event (MIE) and an adverse outcome at individual or population levels (Patlewicz et al., 2015). The data-richness of an AOP is critical for driving its practical application therefore the process of data mining to find mechanistic connections is receiving considerable interest (Patlewicz et al., 2015). Repeat dose systemic organ toxicity is currently an area of considerable interest for AOP development, with the liver being prominently researched due to its integral links to drug/chemical metabolism and effects. The structure of an AOP consists of a molecular initiating event (MIE) which is then linked via key events and ends in a particular adverse effect (as summarised in Figure 1) (Ankley et al, 2010; AOP wiki, 2015). As seen in Figure 1, the key events recorded within an AOP usually consist of the transition from the initiating event, to cellular, tissue and organ-level responses and may also be extrapolated to organism (or even population / ecosystem for environmental effects) (Vinken, 2013, 2015).

FIGURE 1 HERE

A number of free-to-use tools are available for the scientific community to contribute towards the development of AOPs. These have recently been formalised into the Adverse Outcome Pathway Knowledge Base (AOP-KB) which was released in September 2014 (AOP-KB, 2015). The AOP-KB is an online space dedicated to gathering all the work undertaken worldwide on the development of AOPs into one easily accessible web-site. The AOP-KB is an Organisation for Economic Co-operation and Development (OECD) initiative involving collaborations with the European Commission's Joint Research Centre (JRC), US EPA and the US Army Engineer Research and Development Center (ERDC). It gathers four individually developed platforms in one place; the AOP wiki, AOP explorer, intermediate effects database and effectopedia, as well as allowing for third party applications and plug-ins. The AOP-KB allows a contributor to build an AOP within the AOP wiki by entering information about MIE and key events associated with a particular AOP. As scientists understand that mechanistic pathways are diverse and not homologous, the AOP-KB allows the links between different MIEs, key events and AOPs forming a branching connection which can be visualised via the AOP explorer platform. The AOP-KB allows users to gain insight into currently known AOPs, add information, comment on particular AOPs and review work that has already been carried out by other stakeholders/ users. The AOP wiki has been successful at engaging scientists to add details for mechanistic pathways they have data/ knowledge on allowing the progression of AOP development.

Due to their involvement in many essential processes within the body, the search for novel ligands for Nuclear Receptors (NR) has recently been intensified in order to elucidate possible preventative / therapeutic treatments for a wide range of diseases including diabetes, cancer, cardiovascular diseases, atherosclerosis, neurodegenerative diseases and obesity (Love, 2006). However, the induction of some NR ligands has been linked to the development of drug

induced liver injury (DILI) such as liver steatosis, due to the bio-activation of drugs (or metabolites) and / or the induction of hepatotoxic pathways (Love, 2006). The mechanisms behind these hepatotoxic pathways and the chemical structures of the ligands that induce them must first be understood before the definition of the characteristics for binding (which would be essential for the development of structural alerts) can take place. NRs are linked to onset of hepatic toxicity, especially hepatic steatosis.

Hepatic Steatosis

Hepatic steatosis is induced via the excessive accumulation of fats (triglycerides) within the hepatic parenchymal cells (hepatocytes) of the liver (Reddy and Roa, 2006; Zafrani, 2004; Nguyen et al., 2008). Hepatic steatosis usually occurs as the first stage of fatty liver disease, if the cause persists steatosis typically progresses to steatohepatitis (inflammation of the liver cells), cirrhosis (scarring of the liver) and liver cancer (Reddy and Roa, 2006). The onset of hepatic steatosis is associated with many different causes including alcoholism, diabetes, obesity and mitochondrial dysfunction (Reddy and Roa, 2006). Morphologically, hepatic steatosis presents as the accumulation of both large (macrovesicular) and small (microvesicular) intra-cytoplasmic fat droplets within hepatocytes which cause cytoplasm displacement. The macrovesicular fat droplets are found in cases of alcoholic, diabetic or obese patients and also present in cases of malnutrition such as immune deficiency syndrome. Macrovesicular steatosis gives rise to a large single vacuole of fat which fills the cytoplasm of hepatocytes and leads to the displacement of the nucleus (Zafrani, 2004). Microvesicular steatosis is present mostly in steatosis induced via β -oxidation of fatty acids (either mitochondrial or peroxisomal) and presents as the formation of many smaller fat droplets leaving the nucleus at the centre of the hepatocytes (Zafrani, 2004). The diagnosis of steatosis is made when lipid content in the liver exceeds 5–10% of the liver's total weight (Zafrani, 2004).

This critical review assesses NR induced hepatic steatosis considering the availability of information from extant AOPs. The purpose is to demonstrate, through relevant AOPs, the linkage of the receptor mediated molecular initiating event to adverse outcome. There are currently two AOPs listed on the AOP wiki that have hepatic steatosis as their adverse outcome (AOP wiki, April 2015). The first is aryl hydrocarbon receptor (AHR) leading to hepatic steatosis, for which there is strong evidence but it is currently under construction. The second is liver X receptor (LXR) activation to liver steatosis, which is under construction and the weight of evidence is unspecified (AOP wiki, April 2015).

NRs linked previously to the onset of liver injury are summarised in Table 1. The aim of this study was to utilise the information captured in Table 1, gathering the current knowledge of the mechanistic pathways of NRs, in order to investigate which are associated with the adverse outcome of hepatic steatosis.

TABLE 1 HERE

Nuclear Receptors

Those NRs associated with liver injury (Table 1) were studied with the objective of compiling knowledge to support the development and extension of an AOP with NR activation being the MIE and hepatic steatosis being the adverse outcome and ultimately allowing for the creation of (chemistry-based) structural alerts (although this is not the purpose of this investigation). The following summarises the pertinent information found within the literature for each NR:

Aryl Hydrocarbon Receptor (AHR)

The AHR is a ligand-activated transcription factor that is involved in the regulation of the biological response to aromatic hydrocarbons (Nebert et al., 2004). Research has shown the AHR regulates xenobiotic metabolising enzymes, for example cytochrome P450. AHR is located within the cytosol when in its inactive form bound to its co-chaperones. Upon ligand

activation, the co-chaperones dissociate and AHR translocates to the nucleus. Once at the nucleus AHR undergoes dimerisation to AHR nuclear translocator (ARNT) which elicits a change to gene transcription (Elferink et al., 1990). Activation of the AHR is known to induce many toxic responses such as teratogenicity, immunotoxicity, tumour promotion and lethality (Pelclova et al., 2006). Due to its abundance within the liver, activation of the AHR has been shown to cause the onset of hepatic steatosis (Boverhof et al., 2006; Li et al., 1994; Niittynen et al., 2007). The AHR has been linked to hepatic steatosis through its ability to upregulate CD36 and to activate the PPAR α receptor (see PPAR α section). CD36 maintains uptake and intracellular trafficking of fatty acids and is also essential for the esterification of fatty acids into triglycerides. Up-regulation of CD36 via the AHR receptor causes increased fatty acid influx from the peripheral tissues (Figure 2) leading to the accumulation of triglycerides within the liver causing the formation of micro and macrovesicular fat droplets (He et al., 2011; Kawano et al., 2010). The accumulation of micro and macrovesicular fat droplets, as previously discussed, induces nucleus distortion, mitochondrial disruption and endoplasmic reticulum stress causing fatty liver which leads to hepatic steatosis when the lipid concentration reaches 5-10% of total liver weight (Dentin et al., 2006; Miquilena-Colina et al., 2011).

Constitutive Androstane Receptor (CAR)

The CAR, also called the nuclear receptor subfamily 1, group I, member 3 (NR1I3) (Base et al., 1994), is an orphan nuclear receptor that is an essential regulator of drug metabolising enzymes (Yamamoto et al., 2004). Recent studies have highlighted its importance for the control of enzymes such as CYP450s (Honkakoski et al., 1998; Sueyoshi et al., 1999), multidrug-resistant proteins (MRPs) (Cherrington et al., 2002; Kast et al., 2002) and UDP-glucuronosyltransferase (UGT) (Sugatani et al., 2001; Xie et al., 2003). Investigations have determined that CAR is

involved in the regulation of both bile acid (Saini et al., 2004) and bilirubin (Huang et al., 2004; Moreau et al., 2008; Xie et al., 2003) induced liver injury.

CAR forms a heterodimer with RXR upon ligand binding; this mediates transcriptional up-regulation of target genes (Suino et al., 2004). Analogous to PXR, CAR has a highly conserved DNA Binding Domain (DBD) and a moderately conserved Ligand Binding Domain (LBD) (Suino et al., 2004). Similar to the PXR NR, CAR activation can be induced via a wide range of chemicals such as the antiemetic chlorpromazine, the anti-inflammatory drug acetaminophen and the barbiturate phenobarbital (Moreau et al., 2008). Studies have shown that PXR and CAR share some chemical ligands with results showing them to have different ligand binding and activational properties (Maglich et al., 2009; Osabe et al., 2008). Investigations revealed that most CAR agonists, instead of inducing activation by direct binding, induce CAR translocation from the cytoplasm in to the nucleus (Maglich et al., 2009; Osabe et al., 2008). The predominant difference between CAR and other ligand-dependant nuclear receptors is that CAR is constitutionally active (Suino et al., 2004).

CAR activation has been shown to affect lipid homeostasis (Moreau et al., 2008). Studies have demonstrated that upon agonist binding and activation, CAR facilitates fat accumulation and leads to enhanced hepatic steatosis *in vivo* (Xie et al., 2003). The activation of CAR leads to increased expression of PPAR γ (see this section) and Sterol Regulatory Element Binding Protein-1c (SREBP-1c) (Moreau et al., 2008; Wada et al., 2009). SREBP-1c expression leads to up-regulation of lipogenic enzymes essential for *de novo* lipogenesis (lipid production within the cell). Its activation via the CAR receptor leads to increased *de novo* synthesis (Figure 2) and causes lipid accumulation within hepatocytes (Azzout-Marniche et al., 2000 Foretz et al., 1999; Nguyen et al., 2007). Lipid accumulation can lead to the production of micro and macrovesicular fat droplets within hepatocytes if the CAR agonist is not removed, resulting in

a fatty liver which leads to the onset of hepatic steatosis (Azzout-Marniche et al., 2000 Foretz et al., 1999; Nguyen et al., 2007).

oEstrogen Receptor (ER)

The ER, which is also known as the nuclear receptor subfamily 3, group A, member 1/2 (NR3A1/2), is essential for the normal development and maintenance of the sexual and reproductive functions in both males and females (Heldring et al., 2007). ER signalling is linked to a wide range of physiological effects in biological systems such as the immune, cardiovascular, musculoskeletal and the central nervous system in both men and women (Gustafsson, 2005). The ER has two receptor subtypes ER α (NR3A1) and ER β (NR3A2), which share 53% homology of their LBD (Gougelet et al., 2005). Although they share this homology, the two receptor subtypes have noticeable differences in their binding specificities (Gougelet et al., 2005). However, both ER receptor subtypes have strong affinity for 17 β -estradiol with the main difference in binding being the strong affinity ER β has for phytoestrogens (Kuiper et al., 1997).

ER α blockers/ antagonists are used therapeutically for the treatment of breast cancer. ER agonists have been linked to the development of hepatotoxicity such as hepatic steatosis due to their ability to inhibit respiration leading to increased accumulation of triglycerides within the liver (Bandyopadhyay et al., 2006; Foster, 2012; Lelliott et al., 2005; Moya et al., 2010; Shimizu et al., 2007).

Studies have shown that both ER- α agonists and antagonists can lead to the onset of hepatic steatosis, although the mechanisms behind this are not yet fully understood (Lelliott et al., 2005; Moya et al., 2010). It was determined that ER α activation causes increased malonyl-CoA levels within the cell (Bandyopadhyay et al., 2006; Lelliott et al., 2005). Malonyl-CoA is essential for energy homeostasis; its regulation determines the rate of cellular fatty acid β oxidation within

the mitochondria and fatty acid synthesis within the cytosol (Foster, 2012). At lower concentrations, malonyl-CoA acts as a substrate for the enzyme malonyl coenzyme A-acyl carrier protein transacylase during fatty acid synthesis, a rate limiting step for the elongation of fatty acids. When the concentration of malonyl-CoA within the cytosol is high, it acts to inhibit the carnitine acyltransferase 1 transporter which prevents the transport of fatty acids into the mitochondrial matrix and therefore inhibits mitochondrial β oxidation (Foster, 2012; Lelliot et al., 2005). ER α activation causes increased concentrations of malonyl-CoA causing the inhibition of mitochondrial β oxidation and thus preventing the breakdown of fatty acids. This leads to the accumulation of triglycerides which, as previously discussed, can result in the onset of hepatic steatosis (Bandypadhyay et al., 2006; Foster, 2012; Lelliot et al., 2005; Moya et al., 2010)

Farnesoid X receptor (FXR)

The FXR, which is also known as the nuclear receptor subfamily 1, group H, member 4/5 (NR1H4/5) (Forman et al., 1995), is an essential modulator of lipid and glucose homeostasis (Claudel et al., 2005). The FXR has two receptor subtypes FXR α (NR1H4) and FXR β (NR1H5). The endogenous ligand for the FXR receptor is bile acids (Laffitte et al., 2002). After FXR ligand binding has occurred, FXR binds to DNA segments termed FXR response elements (FXREs), which can elicit activation or repression of the FXREs. FXR can act as a monomer or can form a heterodimer with RXR (Claudel et al., 2003). Upon activation via bile acids, the FXR regulates bile acid synthesis, bile acid transport, bile acid conjugation and also impacts on lipid and glucose metabolism (Claudel et al., 2005). The FXR is highly expressed within the liver and intestine (Howard et al., 2000). Studies have demonstrated that the FXR can also be activated via triterpenoids (e.g. forskolin) (Zhao et al., 2004) and polyunsaturated fatty acids (e.g. arachidonic acid) (Iser et al., 1975). The weak FXR agonist ursodeoxycholic acid is the only bile acid used therapeutically. Ursodeoxycholic acid is used for the treatment of gallstone

disease and cholestatic liver diseases (Claudel et al., 2005). As the FXR plays a pivotal role in energy and bile acid metabolism, it has been identified as a promising target for the treatment of dyslipidaemia and liver disorders (Claudel et al., 2005). However, this would require the development of highly specific FXR modulators and will require extensive research before it becomes a reality (Claudel et al., 2005). Chenodeoxycholic acid, demonstrated to be the most naturally potent ligand of the FXR, was taken to clinical trials to be used as a potential therapy for the treatment of gallstone disease and/ or hypertriglyceridemia (Biddie et al., 2002). However, the adverse side effects such as diarrhoea and liver toxicity such as hepatic steatosis lead to the withdrawal of this drug as a therapy and has shown the need for further understanding of the effects FXR activation induces (Biddie et al., 2002). FXR activation is linked to increased expression of PPAR- α (see this section) causing increased accumulation of triglycerides within the liver culminating in hepatic steatosis (Figure 2) (Pineda Torra et al., 2003).

Glucocorticoid Receptor (GR)

GR, also known as the nuclear receptor subfamily 3, group C, member 1(NR3C1) is essential to the maintenance of various metabolic and homeostatic functions within the body (Anbalagan et al., 2012). Cortisol is the naturally occurring ligand for the GR within mammals, it is produced via the adrenal gland and is a cholesterol-derived steroid hormone (Lewis-Tuffin et al., 2007). Upon ligand binding the GR undergoes a conformational change that triggers its translocation to the nucleus, where it induces transcriptional activation of target genes (Buttgereit et al., 2012). Post-translational modifications can also occur (such as phosphorylation, acetylation and ubiquitination), these can alter the function of the GR target genes (Vilasco et al., 2011).

Therapeutically GR agonists were first used in 1940 for the treatment of rheumatoid arthritis (cortisone) (Anbalagan et al., 2012). Since then they have been developed and used for many

other chronic inflammatory conditions such as eye infections, asthma (budesonide), skin infections (dexamethasone) and for immunosuppression (prednisone) in transplant patients. GR agonists have also been developed for the treatment of certain cancers due to their antiproliferative and antiangiogenic properties (prednisone) (Chourbaji et al., 2008).

Upon agonistic binding to the GR anti-angiogenic, anti-inflammatory, proapoptotic and anti-proliferative effects are induced in the musculoskeletal (Bultink et al., 2013; Weinstein et al., 2010), nervous (Tronche et al., 1999), visual (Edelman, 2010; Kiernan et al., 2009), cardiovascular (Fardet et al., 2012), immune (Silverman et al., 2012; Zen et al., 2011), reproductive (Harris et al., 2011), integumentary (Coenraads et al., 2012; Sevilla et al., 2012), respiratory systems (Hakim et al., 2012), and on glucose/ liver metabolism (Rose et al., 2010) within the body.

Studies have shown that when under stress the GR signalling induces glycogenolysis and gluconeogenesis within the liver to replenish glucose levels (Biddie et al., 2012; Jia et al., 2009). This is an adverse effect of GR agonists and has been linked to the onset of Cushing's disease, Addison's disease and to the development of hepatic steatosis (Jia et al., 2009; Letteron et al., 1997). GR agonist have been shown to induce hepatic steatosis due to their ability to both inhibit mitochondrial fatty acid β oxidation and to upregulate CAR expression (see CAR section), both actions lead to increased accumulation of triglycerides within the liver which can lead to hepatic steatosis (Figure 2) (Jia et al., 2009; Letteron et al., 1997)

Liver X Receptor (LXR)

The LXR, also known as the nuclear receptor subfamily 1, group H, member 2/3 (NR1H2/3), is an adopted orphan NR that upon activation forms a heterodimer with RXR to induce transcriptional activation of its target genes (Janowski et al., 1996). The LXR became an adopted NR when it was discovered that endogenous oxysterols serve as LXR ligands (Schultz

et al., 2000). LXR has two isoforms LXR α (NR1H3) and LXR β (NR1H2), which are both essential for the regulation of cholesterol and lipid metabolism (Rippa et al., 2000). Studies have determined that LXR α is expressed mainly in the liver and small intestine whereas LXR β is expressed ubiquitously, with only low expression within the liver (Rippa et al., 2000). Investigations have shown that upon LXR activation, intestinal cholesterol absorption (Rippa et al., 2002) and hepatic cholesterol synthesis (Rippa et al., 2000) are reduced, whereas the expression of genes involved in cholesterol mobilisation (Peet et al., 1998), bile acid synthesis (Zaghini et al., 2002), reverse cholesterol transport (Peet et al., 1998) and cholesterol excretion into the bile (Grefhorst et al., 2002) are increased. Treatments with LXR agonists have shown to be preventative against atherosclerosis during *in vivo* studies. This has led to the creation of a potential therapeutic drug target against atherosclerosis (Rippa et al., 2000). However, LXR activation has been linked to the development of hepatic steatosis (Janowski et al., 1996). LXR activation is shown to cause up-regulation of: carbohydrate responsive element binding protein (ChREBP), SREBP-1c, fatty acid synthase (FAS) and SCD1. ChREBP is crucial for mediating the body's response to glucose on glycolytic and lipogenic genes, it is required for the induction of FAS (responsible for transporting triglycerides into cells) and acetyl-CoA carboxylase (ACC) (essential enzyme involved in *de novo* synthesis). Therefore, the induction of ChREBP via LXR agonists leads to increased accumulation of lipids inside hepatocytes as more lipids are transport into the cell via FAS induction and more lipids are produced via ACC induction (Dentin et al., 2006). Similarly, increased SREBP-1c mRNA expression also leads to up-regulation of lipogenic enzymes ACC and FAS (Nguyen et al., 2007). Up-regulation of the SCD1 enzyme leads to increased fatty acid production as it is the rate limiting step in the production of unsaturated fatty acids (Zhang et al., 1999). In summary, LXR activation induces all the above mentioned pathways, these collectively cause the accumulation of *de novo* fatty acids within the liver resulting in the production of micro and macrovesicular fat droplets within

hepatocytes, thus creating a fatty liver and ultimately leading to hepatic steatosis (Kawano et al., 2010; Rosen et al., 2008; Zafrani, 2004).

Peroxisome Proliferator-Activated Receptor (PPAR)

The PPARs, also known as Nuclear receptor subfamily 1, group C, member 1-3 (NR1C1-3), are orphan receptors that form heterodimers with RXR upon ligand-induced activation, leading to transcriptional regulation of target genes (Issemann et al., 1990). There are three PPAR isoforms, namely PPAR α (NR1C1), PPAR β (NR1C2), and PPAR γ (NR1C3) (Kliewer et al., 1994). PPAR α was the first to be discovered 20 years ago (Lee et al., 1995). PPAR α mediates increased hepatic peroxisome expression causing increased concentration and density of peroxisomes within the liver (Peters et al., 2005). PPAR β and PPAR γ share a sequence homology to PPAR α however they are not involved in peroxisome proliferation (Issemann et al., 1990). Studies have shown that PPAR β and PPAR γ can interfere with NF- κ B and API signalling through protein-protein interactions (Peters et al., 1997; Satoh et al., 2013). All PPAR isoforms are encoded for by separate genes, have different expression levels within various tissues and are expressed by all mammalian species (Dreyer et al., 1992). PPAR α is essential for the down regulation of apolipoprotein CII mRNA expression However; the mechanism behind this pathway is not understood (Misrahi et al., 1987).

PPAR γ agonists are used to regulate glycaemia in type II diabetics. PPAR α agonists are used to treat atherosclerosis. PPAR γ and PPAR α dual agonists are used to improve glycaemia and the condition of the cardiovascular system in patients with type II diabetes. PPAR α and PPAR γ ligands have been shown to cause hepatic steatosis due to lipid accumulation (Honkakoski et al., 1998; Misrahi et al., 1987). The mechanisms behind PPAR α and PPAR γ induced hepatic steatosis differ:

PPAR α

The production of energy (in the form of ATP) in the liver is controlled by both mitochondrial and peroxisomal β -oxidation (Nguyen et al., 2007; Reddy and Roa, 2005). Peroxisomal β -oxidation involves the oxidation of long or very long chain fatty acids and results in less ATP production than mitochondrial β -oxidation. One of its key enzymes is the fatty acyl-CoA oxidase (AOX) which catabolises the dehydrogenation of fatty acids into acetyl -CoA (Hashimoto et al., 1999).

The PPAR- α receptor acts as a sensor for fatty acids and is responsible for the transcriptional activation of the AOX enzyme in the liver. Upon antagonistic binding to the PPAR α , down regulation of the AOX enzyme follows leading to the inhibition of microsomal β -oxidation (Figure 2). This causes an accumulation of fatty acids within the liver, if PPAR α antagonism continues, the build-up of triglycerides will also continue which will eventually lead to the production of micro and macrovesicular intra-cytoplasmic fat droplets within hepatocytes. As discussed previously, micro and macrovesicular fat droplets induce nucleus distortion, mitochondrial disruption and endoplasmic reticulum stress (Figure 2) (Nguyen et al., 2007; Reddy and Roa, 2005; Zafrani, 2004). These effects result in a fatty liver which becomes diagnosable as hepatic steatosis when lipid weight is between 5-10% of the total liver weight (Kawano et al., 2010; Rosen et al., 2008; Zafrani, 2004).

PPAR γ

PPAR γ activation leads to increased expression of SREBP-1c, increased expression of fatty acid translocase CD36 (CD36), and cellular differentiation (Al Sharif et al., 2014; Miquilena-Colina et al., 2011; Rippla et al., 2000) (Figure 2). Increased SREBP-1c mRNA expression leads to up-regulation of lipogenic enzymes such as ACC (catalyses elongation of the fatty acid chain in *de novo* lipogenesis) and FAS (multifunctional enzyme responsible for catalysing many steps in *de novo* lipogenesis) (Azzout-Marniche et al., 2000 Foretz et al., 1999; Nguyen

et al., 2007). CD36 maintains uptake and intracellular trafficking of fatty acids and is also essential for the esterification of fatty acids into triglycerides. The increased expression of SERBP-1c and CD36 are known to induce accumulation of *de novo* fatty acids. *De novo* lipogenesis is the process of lipid production inside the cytosol of a cell and is necessary for energy homeostasis within mammals (Nguyen et al., 2007). Increased production of *de novo* fatty acids has been demonstrated to cause accumulation of triglycerides within the liver which ultimately results in the onset of hepatic steatosis (Figure 2) (Postic and Girard, 2008; Rippla et al., 2000; Peraza et al., 2006).

Pregnane X Receptor (PXR)

The PXR is also known as the steroid Nuclear receptor subfamily 1, group I, member 2 (NR1I2) (Venkatesh et al., 2011). PXR is an adopted orphan nuclear receptor that upon ligand binding forms a heterodimer with RXR, this mediates transcriptional up-regulation of target genes (Gonzalez et al., 1991).

PXR can be activated by both small (e.g. estradiol 268Da) and large molecules (e.g. rifampicin 823 Da). The naturally occurring ligands for PXR are steroids pregnenolone and progesterone. The synthetic ligands that have been shown to bind to PXR are glucocorticoid agonists and antagonists (Bertilsson et al., 1998). The PXR is known to bind a large range of structurally unrelated chemicals. X-ray crystal structures of the PXR-LBD have shown it to have a larger LBD compared to other NRs. This may account for its promiscuous binding (Watkins et al., 2001). Therefore, PXR has low substrate specificity and can be activated by many different chemicals.

Research has demonstrated that the PXR is essential for the modulation of hepatic drug metabolism (Li et al., 2012). Metabolic clearance especially via the phase I cytochrome p450 enzymes (CYP450), is vital for endogenous (bile acids) and exogenous (xenobiotics)

detoxification and essential to the survival of an organism (Waxman et al., 1999). Both the CYP3A and CYP2B isoenzymes mediate the metabolism of many clinical drugs and can be induced by substrate binding, therefore they are up-regulated to meet hepatic requirements (Smith et al., 1993). The PXR and CYP3As are both highly expressed within the liver (Bertilsson et al., 1998). Studies have determined that PXR up-regulates hepatic CYP3A expression along with other CYP450 enzymes. This was demonstrated via the use of PXR Knock Out (KO) mice. The mice were both viable and fertile, however upon dosing with xenobiotics the PXR KO mice developed severe hepatotoxicity as CYP3A could not be up-regulated. These studies established that PXR is the central mediator of CYP3A induction (Hoekstra et al., 2009; Smith et al., 1993; Xie et al., 2000). Studies showed that upon ligand binding PXR forms a heterodimer with RXR and binds to the Direct Repeat 3 (DR3) site on the CYP3A promoter this mediates transcriptional activation (Bertilsson et al., 1998). The mechanisms underlying PXR activation are still unknown.

It has been shown that PXR agonists induce a decrease in plasma LDL cholesterol levels (Bitter et al., 2014) and the onset of hepatic steatosis (Bitter et al., 2014; Kaisimanickam et al., 2013) *in vivo*. Research has shown that PXR agonists increase the expression of PPAR α (see PPAR section) which leads to increased triglyceride accumulation within the liver which can lead to hepatic steatosis (Figure 2) (Moreau et al., 2007).

Retinoic Acid Receptor (RAR)

The RAR, also called nuclear receptor subfamily 1, group b, member 1-3 (NR1B1-3), is divided into three subtypes: RAR α (NR1B1), RAR β (NR1B2) and RAR γ (NR1B3). Bound together with the retinoid X receptor (RXR) as a heterodimer, RAR regulates genetic expression as transcriptional repressors in absence of ligands. RAR is important for regulating of cellular proliferation and differentiation (Liu et al., 2014). All three subtypes of the RAR are

activated by *all-trans* retinoic acid and *9-cis* retinoic acid, which are derivatives of vitamin A (Liu et al., 2014). Ligands are used in pharmacological treatment of dermal diseases, such as *Acne vulgaris*, *Psoriasis vulgaris*, *Keratosis pilaris* and specific types of cancer, e.g. acute promyelocytic leukemia (Alizadeh et al., 2014; Leyden et al., 2005). Effects of RAR agonists include changes in lipid metabolism, which may cause hepatic steatosis and eventually liver inflammation, fibrosis and finally liver failure. Teratogenic effects and neural disorders, such as nausea and headache, have been also reported from retinoids (Biesalski et al., 1989; Moya et al., 2010; Shalita et al., 1988). Moya et al (2010) demonstrated that the retinol forms of RAR agonists are able to induce steatosis in hepatocytes, however, the retinoic acid forms were not able to induce this same effect. The difference in these toxic responses are not yet understood but the mechanistic route for the onset of steatosis is suggested to be via the up regulation of lipid synthesis leading to increased accumulation of triglycerides within the liver (Figure 2) (Moya et al., 2010).

AOP Formulation

All the findings from the NR literature are summarised in Table 2 below. As can be seen the AHR, AR, CAR, ER, FXR, GR, LXR, PPAR, PXR, RAR, RXR and THR are all associated with hepatic injury, however, only AHR, CAR, ER, FXR, GR, LXR, PPAR, PXR, RAR and RXR (Table 2 - highlighted with *) have been shown to induce hepatic steatosis upon binding. Upon studying the mechanistic information gathered it was noted that activation of those NR associated with the onset of hepatic steatosis followed a similar chain of events. Activation of these NRs leads to triglyceride accumulation within the hepatocytes which then ultimately results in hepatic steatosis. Using the evidence found within the literature and the AOPs found on the AOP wiki, an AOP pathway for NR induced hepatic steatosis is proposed (Figure 2).

TABLE 2 HERE

FIGURE 2 HERE

The AOP pathway reported in Figure 2 has been adapted from the diagram available on the AOP wiki website (AOP, May 2015). The AOP pathways 1-7 (blue arrows) are extracted from the original figure and AOPs 7-10 (green and orange arrows) have been added. The AOPs 1-6 that have been highlighted green and the arrows that are green show the NR pathways that have been studied within this critical review and further strengthen the evidence for these AOP pathways. For all AOPs within Figure 2, the chemical is the NR agonist and the MIE is the binding of these agonists within the NR binding pocket to initiate activation of the NR. The key event is triglyceride accumulation and the adverse outcome is hepatic steatosis. The key events of this AOP (Figure 2) are affirmed via the literature search carried out within this review; however, the molecular processes between the MIE and the key events should be explored further.

Conclusion

The links between the chemical structure of NR ligands and the biological responses of the NR they bind to have been elucidated, this has provided knowledge of the toxicological pathways linked to these NR, specifically to the adverse outcome of hepatic steatosis. However, more work will need to be carried out to form this AOP, with the next essential step being the formation of structural alerts. This critical review has given an example of how current AOPs can be extended and developed via the use of literature review in order to bring together different (*in vitro*, *in vivo* and *in silico*) existing mechanistic / toxicological knowledge for a particular AOP. To conclude, this study has highlighted the importance of the promotion of the AOP concept to the scientific community to gain wider participation and has shown the way in which AOP formation can be developed and extended using current knowledge.

Acknowledgement

The research leading to these results has received funding from the European Community's Seventh Framework Program (FP7/2007-2013) under grant agreement n° 266835 and from Cosmetics Europe. More information is available at www.cosmostox.eu.

Declaration of Interest

The authors declare that there are no conflicts of interest

References

- Adler S, Basketter D, Creton S, Pelkonen O, Benthem J, Valérie Zuang V, Andersen K.E, Angers-Loustau A, Aptula A, Bal-Price A, Benfenati E, Bernauer U, Bessems J, Bois F.Y, Boobis A, Brandon E, Bremer S, Broschard T, Casati S, Coecke S, Corvi R, Cronin M.T.D, Daston G, Dekant W, Felton S, Grignard E, Gundert-Remy U, Heinonen T, Kimber I, Kleinjans J, Komulainen H, Kreiling R, Kreysa J, Leite S.B, Loizou G, Maxwell G, Mazzatorta P, Munn S, Pfuhler S, Phrakonkham P, Piersma A, Poth A, Prieto P, Repetto G, Rogiers V, Schoeters G, Schwarz M, Serafimova R, Ta'hti H, Testai E, van Delft J, van Loveren H, Vinken M, Worth A and Zaldivar J-M (2011). Alternative (non-animal) methods for cosmetics testing: current status and future prospects – 2010. *Arch Toxicol* 85, 367-485.
- Alizadeh F, Bolhassani A, Khavari A, Bathaie S.Z, Naji T and Bidgoli S.A (2014). Retinoids and their biological effects against cancer. *Int Immunopharmacol* 18 (2014), 43-49
- Al Sharif M, Alov P, Vitcheva V, Pajeva I and Tsakovska I (2014). Modes-of-action related to repeated dose toxicity: from PPARgamma ligand-dependent dysregulation to non-alcoholic fatty liver disease. *PPAR Research*, Special issue, "PPARs and Metabolic Syndrome", Volume 2014, Article ID 432647
- Anbalagan M, Huderson B, Murphy L and Rowan B.G (2012). Post-translational modifications of nuclear receptors and human disease. *Nucl. Recept. Signal.* 10, e001
- Ankley, G.T., Bennett, R.S., Erickson, R.J., Hoff, D.J., Hornung, M.W., Johnson, R.D., Mount, D.R., Nichols, J.W., Russom, C.L., Schmieder, P.K., Serrano, P.K., Tietge, J.E and Villeneuve, D.L (2010). Adverse outcome pathways: a conceptual framework to support ecotoxicology research and risk assessment. *Environ. Toxicol. Chem.* 29, 730 – 741
- AOP wiki (26th April 2015). The term hepatic steatosis has been searched for within the AOP wiki search function. <https://aopkb.org/aopwiki/index.php?search=hepatic+steatosis&title=Special%3ASearch&fulltext=Search> .Accessed on 26th April 2015
- AOP wiki NR linked to hepatic steatosis (5th May 2015) <https://aopkb.org/aopwiki/index.php/Aop:34>. Accessed on 5th May 2015
- AOP-KB (2015). The OECD webpage describing AOPs. <http://www.oecd.org/chemicalsafety/testing/adverse-outcome-pathways-molecular-screening-and-toxicogenomics.htm>. Accessed on 26th April 2015
- Apriletti J.W, Ribeiro R.C, Wagner R.L, Feng W, Webb P, Kushner P.J, West B.L, Nilsson S, Scanlan T.S, Fletterick R.J and Baxter J.D (1998). Molecular and structural biology of thyroid hormone receptors. *Clin Exp Pharmacol Physiol Suppl.* 1998 Nov;25:S2-11.
- Arnett M.G, Kolber B.J, Boyle M.P and Muglia L.J (2011). Behavioral insights from mouse models of forebrain–and amygdala-specific glucocorticoid receptor genetic disruption. *Mol. Cell. Endocrinol.* 336, 2–5
- Azzout-Marniche D, Bécard D, Guichard C, Foretz M, Ferré P and Foufelle F (2000). Insulin effects on sterol regulatory-element-binding protein-1c (SREBP-1c) transcriptional activity in rat hepatocytes. *Biochem J.* 2000 Sep 1;350 Pt 2:389-93.
- Baes M, Gulick T, Choi HS, Martinoli MG, Simha D and Moore D.D (1994). A new orphan member of the nuclear hormone receptor superfamily that interacts with a subset of retinoic acid response element. *Mol. Cell. Biol.* 14 (3): 1544–52.
- Bahn R.S, Burch H.S, Cooper D.S, Garber J.R, Greenlee C.M, Klein I.L, Laurberg P, McDougall IR, Rivkees S.A, Ross D, Sosa J.A and Stan M.N (2009). The Role of Propylthiouracil in the Management of Graves' Disease in Adults: report of a meeting jointly sponsored by the American Thyroid Association and the Food and Drug Administration. *Thyroid : official journal of the American Thyroid Association* 19 (7): 673–4
- Bandyopadhyay G.K, Yu J.G, Ofrecio J, and Olefsky J.M (2006). Increased Malonyl-CoA Levels in Muscle From Obese and Type 2 Diabetic Subjects Lead to Decreased Fatty Acid Oxidation and Increased Lipogenesis; Thiazolidinedione Treatment Reverses These Defects. *DIABETES*, VOL. 55, AUGUST 2006
- Barral R.L and Gold M.A (2014) Progestin-Only Contraception. *Contraception for Adolescent and Young Adult Women.* 2014, pp 25-44

- Baulieu E and Schumacher M (2000). Progesterone as a neuroactive neurosteroid, with special reference to the effect of progesterone on myelination. *Steroids* 65 (10-11): 605–12
- Bertilsson G, Heidrich J, Svensson K, Asman M, Jendeberg L, Sydow-Bäckman M, Ohlsson R, Postlind H, Blomquist P and Berkenstam A (1998). Identification of a human nuclear receptor defines a new signaling pathway for CYP3A induction. *Proc Natl Acad Sci USA* 1998;95:12208-13
- Biddie S.C, Conway-Campbell B.L and Lightman S.L (2012). Dynamic regulation of glucocorticoid signalling in health and disease. *Rheumatology (Oxford)* 51, 403–412
- Biesalski H.K (1989). Comparative assessment of the toxicology of vitamin A and retinoids in man. *Toxicology* 57 (1989), 117-161
- Bitter A, Rümmele P, Klein K, Kandel B.A, Rieger J.K, Nüssler A.K, Zanger U.M, Trauner M, Schwab M and Burk O (2014). Pregnane x receptor activation and silencing promote steatosis of human hepatic cells by distinct lipogenic mechanisms. *Arch Toxicol.* 2014 Sep 3
- Boverhof D.R, Burgoon L.D, Tashiro C, Sharratt B, Chittim B, Harkema J.R, Mendrick D.L and Zacharewski T.R (2006). Comparative Toxicogenomic Analysis of the Hepatotoxic Effects of TCDD in Sprague Dawley Rats and C57BL/6 Mice *Toxicol. Sci.* 94 (2006) 398–416
- Bultink I.E, Baden M and Lems W.F. (2013). Glucocorticoid-induced osteoporosis: an update on current pharmacotherapy and future directions. *Expert Opin. Pharmacother.* 14, 185–197
- Buttgereit F (2012). A fresh look at glucocorticoids how to use an old ally more effectively. *Bull. NYU Hosp. Jt. Dis.* 70 (Suppl. 1), 26–29
- Chambellan-Tison C, Horen B, Plat-Wilson G and Moulin P (2007). Severe hypercalcemia due to vitamin D intoxication. *Arch Pediatr.* 2007 Nov; 14(11):1328-32
- Cherrington N.J, Hartley D.P, Li N, Johnson D.R and Klaassen C.D (2002). Organ distribution of multidrug resistance proteins 1, 2, and 3 (Mrp1, 2, and 3) mRNA and hepatic induction of Mrp3 by constitutive androstane receptor activators in rats. *J Pharmacol Exp Ther* 2002; 300: 97-104
- Chourbaji S and Gass P (2008). Glucocorticoid receptor transgenic mice as models for depression. *Brain Res. Rev.* 57, 554–560
- Christin-Maitre S, Bouchard P and Spitz I.M (2000). Medical termination of pregnancy. *N Engl J Med* 2000;342(13):946-56
- Claudel T, Inoue Y, Barbier O, Duran-Sandoval D, Kosykh V, Fruchart J, Fruchart J.C, Gonzalez F.J and Staels B (2003). Farnesoid X receptor agonists suppress hepatic apolipoprotein CIII expression. *Gastroenterology.* 2003;125:544 –555.
- Claudel T, Staels B and Kuipers F (2005). The Farnesoid X receptor: a molecular link between bile acid and lipid and glucose metabolism. *Arterioscler. Thromb. Vasc. Biol.* 25:2020–2030.
- Coenraads P.J. (2012) Hand eczema. *N. Engl. J. Med.* 367, 1829–1837
- Cohen Hubal E.A, Richard A, Aylward L, Edwards S, Gallagher J, Goldsmith M.R, Isukapalli S, Tornero-Velez R, Weber E and Kavlock R (2010). Advancing 1169 exposure characterization for chemical evaluation and risk assessment. *J.Toxicol. Environ. Health B Crit. Rev.* 13, 299 - 313.
- Couse J.F and Korach K.S (1999). Estrogen receptor null mice: what have we learned and where will they lead us. *Endocr Rev* 20: 358–417, 1999
- Crescioli C (2014). Vitamin D Receptor Agonists: Suitable Candidates as Novel Therapeutic Options in Autoimmune Inflammatory Myopathy. *BioMed Research International Volume 2014 (2014), Article ID 949730, 10 pages*
- Cronin M.T.D and Livingston D.J (2004). Predicting Chemical Toxicity and Fate. Chapter 18, pp392-393.
- Dalakas M.C (2010). Immunotherapy of myositis: issues, concerns and future prospects. *Nature Reviews Rheumatology*, vol. 6, no. 3, pp. 129–137, 2010
- DeManno D, Elger W, Garg R, Lee R, Schneider B, Hess-Stumpp H, Schubert G and Chwalisz K (2003). Asoprisnil (J867): a selective progesterone receptor modulator for gynecological therapy. *Steroids* 68 (10-13): 1019–32
- Dentin, R.; Denechaud, P. D.; Benhamed, F.; Girard, J.; Postic, C., 2006: Hepatic gene regulation by glucose and polyunsaturated fatty acids: a role for ChREBP. *Journal of Nutrition* **136**, 1145–1149.
- Dreyer C, Krey G, Keller H, Givel F, Helftenbein G and Wahli W (1992). Control of the peroxisomal beta-oxidation pathway by a novel family of nuclear hormone receptors. *Cell* 68, 879–887.
- Edelman J.L. (2010). Differentiating intraocular glucocorticoids. *Ophthalmologica* 224 (Suppl. 1), 25–30
- Elferink C.J, Gasiewicz T.A and Whitlock J.P Jr (1990). Protein-DNA interactions at a dioxin-responsive enhancer. Evidence that the transformed Ah receptor is heteromeric. *J. Biol. Chem.* 265 (1990) 20708–20712.
- Engelken S.F and Eaton R.P (1981). The effects of altered thyroid status on lipid metabolism in the genetic hyperlipemic Zucker rat. *Atherosclerosis* 1981, 38, 177-188.
- Fardet L and Nazareth I (2012). Risk of cardiovascular events in people prescribed glucocorticoids with iatrogenic Cushing’s syndrome: cohort study. *BMJ* 345, e4928

- Foretz M, Guichard C, Ferre P and Foufelle F (1999). Sterol regulatory element binding protein-1c is a major mediator of insulin action on the hepatic expression of glucokinase and lipogenesis-related genes. *Proc. Natl. Acad. Sci. U. S. A.* 96, 12737–12742.
- Forman B.M, Goode E, Chen J, Oro A.E, Bradley D.J, Perlmann T, Noonan D.J, Burka L.T, McMorris T, Lamph W.W, Evans R.M and Weinberger C (1995). Identification of a nuclear receptor that is activated by farnesol metabolites. *Cell* 81 (5): 687–93.
- Forrest D and Vennström B (2000). Functions of thyroid hormone receptors in mice. *Thyroid* 2000, 10, 41-52.
- Förster C, Mäkela S, Wärrä A, Kietz S, Becker D, Hulthenby K, Warner M and Gustafsson J.Å (2002). Involvement of estrogen receptor β in terminal differentiation of mammary gland epithelium. *Proc Natl Acad Sci USA* 99: 15578–15583, 2002
- Foster D.W (2012). Malonyl-CoA: the regulator of fatty acid synthesis and oxidation. *The Journal of Clinical Investigation*. Vol. 122, No.6, pp1958-1959.
- Gellersen B, Fernandes M.S, and Brosens J.J (2009). Non-genomic progesterone actions in female reproduction *Hum. Reprod. Update* (2009) 15 (1): 119-138
- Gelmann EP (2002). Molecular biology of the androgen receptor. *J Clin Oncol*. 2002 Jul 1;20(13):3001-15.
- Gizzo S, Andrisani A, Esposito F, Noventa M, Di Gangi S, Angioni S, Litta P, Gangemi M, and Nardelli G.B (2014). Which luteal phase support is better for each IVF stimulation protocol to achieve the highest pregnancy rate? A superiority randomized clinical trial. *Gynecol Endocrinol, Early Online*: 1–7.2014
- Gocht T, Berggren E, Ahr H.J, Cotgreave I, Cronin M.T.D, Daston G, Hardy B, Heinzle E, Hescheler J, Knight D.J, Mahony C, Peschanski M, Schwarz M, Thomas R.S, Verfaillie C, White A and Whelan M (2015). The SEURAT-1 approach towards animal free human safety assessment *ALTEX* 32(1), 2015
- Gonzalez F.J and Gelboin H.V (1991). Human cytochromes P450: evolution, catalytic activities and interindividual variations in expression. *Prog Clin Biol Res* 1991;372:11-20
- Gougelet A, Bouclier C, Marsaud V, Maillard S, Mueller S.O, Korach K.S and Renoir J.M (2005). Estrogen receptor α and β subtype expression and transactivation capacity are differentially affected by receptor-, hsp90- and immunophilin-ligands in human breast cancer cells. *J Steroid Biochem Mol Biol*. 2005 Feb;94(1-3):71-81. Epub 2005 Feb 23
- Grefhorst A, Elzinga B.M, Voshol P.J, Plösch T, Kok T, Bloks V.W, van der Sluijs F.H, Havekes L.H, Romijn J.A, Verkade H.J and Kuipers F (2002). Stimulation of lipogenesis by pharmacological activation of the liver X receptor leads to production of large, triglyceride-rich very low density lipoprotein particles. *J. Biol. Chem.* 277 (2002) 34182–34190.
- Groh K.J, Carvalho R.N, Chipman J.K, Denslow N.D, Halder M, Murphy C.A, Roelofs D, Rolaki A, Schirmer K and Watanabe K.H (2015). Development and application of the adverse outcome pathway framework for understanding and predicting chronic toxicity: I. Challenges and research needs in ecotoxicology. *Chemosphere* 120 (2015) 764 - 777
- Gustafsson JA (2003). What pharmacologists can learn from recent advances in estrogen signalling. *Trends Pharmacol Sci* 24: 479–485, 2003
- Haas D.M and Ramsey P.S (2008). Progestogen for preventing miscarriage. *Cochrane Database of Systematic Reviews* 2008; Issue 2. Art. No.: CD003511; DOI: 10.1002/14651858.
- Hakim A, Adcock I.M and Usmani O.S (2012). Corticosteroid resistance and novel anti-inflammatory therapies in chronic obstructive pulmonary disease: current evidence and future direction. *Drugs* 72, 1299–1312
- Hansson P, Valdemarsson S and Nilsson-Ehle P (1983). Experimental hyperthyroidism in man: Effects on plasma lipoproteins, lipoprotein lipase and hepatic lipase. *Horm. Metab. Res.* 1983, 15, 449-452.
- Harris W.P, Mostaghel E.A, Nelson P.S and Montgomery B (2009). Androgen deprivation therapy: progress in understanding mechanisms of resistance and optimizing androgen depletion. *Nat Clin Pract Urol* 2009; 6: 76-85
- Hartung, T and McBride M (2011). Food for thought on mapping the human toxome. *ALTEX* 28, 83-93.
- Hashimoto T, Fujita T, Usuda N, Cook W, Qi C, Peters J.M, Gonzalez F.J, Yeldandi A.V, Rao M.S, and Reddy J.K (1999). Peroxisomal and Mitochondrial Fatty Acid β -Oxidation in Mice Nullizygous for Both Peroxisome Proliferator-activated Receptor α and Peroxisomal Fatty Acyl-CoA Oxidase. *The Journal Of Biological Chemistry* Vol. 274, No. 27, pp. 19228–19236
- Haussler M.R, Haussler C.A, Bartik L, Whitfield G.K, Hsieh J, Slater S, and Jurutka P.W (2008). Vitamin D receptor: molecular signaling and actions of nutritional ligands in disease prevention. *Nutr Rev.* 2008 Oct;66(10 Suppl 2):S98-112.
- He J, Lee J.H, Febbraio and Xie W (2011). The emerging roles of fatty acid translocase/CD36 and the aryl hydrocarbon receptor in fatty liver disease. *Exp Biol Med*, 2011 236: 1116
- Heldring N, Pike A, Andersson S, Matthews J, Cheng G, Hartman J, Tujague M, Ström A, Treuter E, Warner M and Gustafsson J.A (2007). Estrogen Receptors: How Do They Signal and What Are Their Targets. *Physiological Reviews*. 2007 Jul;87(3):905-31.
- Heyman R.A, Mangelsdorf D.J, Dyck J.A, Stein R.B, Eichele G, Evans R.M and Thaller (1992). C. 9-cis retinoic acid is a high affinity ligand for the retinoid X receptor. *Cell* 68 (1992), 397-406
- Hoekstra M, Lammers B, Out R, Li Z, Van Eck M and Van Berkel T.J (2009). Activation of the nuclear receptor PXR decreases plasma LDL-cholesterol levels and induces hepatic steatosis in LDL receptor knockout mice. *Mol Pharm.* 2009 Jan Feb;6(1):182-9.

- Honkakoski P, Moore R, Washburn K and Negishi M (1998). Activation by diverse xenochemicals of the 51-base pair phenobarbital-responsive enhancer module in the CYP2B10 gene. *Mol Pharmacol* 1998; 53: 597-601
- Horwitz K.B, Tung L and Takimoto G.S (1996). In *Hormones and Cancer*; Vedeckis, W. V., Ed.; Birkhaeuser: Boston, 1996; p 283.
- Howard W.R, Pospisil J.A, Njolito E and Noonan D.J (2000). Catabolites of cholesterol synthesis pathways and forskolin as activators of the farnesoid X-activated nuclear receptor. *Toxicol Appl Pharmacol*. 2000;163: 195–202.
- Huang W, Zhang J and Moore D.D (2004). A traditional herbal medicine enhances bilirubin clearance by activating the nuclear receptor CAR. *J Clin Invest* 2004; 113: 137-143
- Ihunnah C.A, Jiang M and Xie W (2011). Nuclear receptor PXR, transcriptional circuits and metabolic relevance. *Biochim Biophys Acta* 1812 (2011), 956-963
- Iser J.H, Dowling H, Mok H.Y and Bell G.D (1975). Chenodeoxycholic acid treatment of gallstones. A follow-up report and analysis of factors influencing response to therapy. *N Engl J Med*. 1975;293:378 –383. 145.
- Issemann I and Green S (1990). Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature* 347, 645–650.
- Janowski B.A, Willy P.J, Devi T.R, Falck J.R and Mangelsdorf D.J (1996). An oxysterol signalling pathway mediated by the nuclear receptor LXR alpha. *Nature* 383 (6602): 728–31.
- Jia Y, Viswakarma N, Fu T, Yu S, Rao M.S, Borensztajn J, Reddy J.K (2009). Conditional ablation of mediator subunit MED1 (MED1/PPARBP) gene in mouse liver attenuates glucocorticoid receptor agonist dexamethasone-induced hepatic steatosis. *Gene Expr*. 2009;14(5):291-306.
- Jung U. J, Millman P.N, Tall A.R and Deckelbaum R.J (2011). n–3 Fatty acids ameliorate hepatic steatosis and dysfunction after LXR agonist ingestion in mice. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids* 1811(9): 491-497.
- Karas R.H, Eickels M.V, Lydon J.P, Roddy S, Kwoun M, Aronovitz M, Baur W.E, Conneely O, O'Malley B.W and Mendelsohn M.E (2001). A Complex role for the progesterone receptor in the response to vascular injury. (SCOR) *J Clin Invest* 2001;108(4):611-618
- Kasimanickam V.R, Kasimanickam R.K and Rogers H.A (2013). Immunolocalization of retinoic acid receptor-alpha, -beta, and -gamma, in bovine and canine sperm. *Theriogenology* 79 (2013) 1010-1018
- Kast H.R, Goodwin B, Tarr P.T, Jones S.A, Anisfeld A.M, Stoltz C.M, Tontonoz P, Kliewer S, Willson T.M and Edwards P.A (2002). Regulation of multidrug resistance-associated protein 2 (ABCC2) by the nuclear receptors pregnane X receptor, farnesoid X-activated receptor, and constitutive androstane receptor. *J Biol Chem* 2002; 277: 2908-2915
- Kawaguchi M, Mitsuhashi Y and Kondo S.J (2003). Iatrogenic hypercalcemia due to vitamin D3 ointment (1,24(OH)2D3) combined with thiazide diuretics in a case of psoriasis. *Dermatol*. 2003 Nov; 30(11):801-4
- Kawano Y, Nishiumi S, Tanaka S, Nobutani K, Miki A, Yano Y, Seo Y, Kutsumi H, Ashida H, Azuma T and Yoshida M. (2010). Activation of the aryl hydrocarbon receptor induces hepatic steatosis via the upregulation of fatty acid transport. *Arch Biochem Biophys* 504: 221-227.
- Kettel L.M, Murphy A.A, Mortola J.F, Liu J.H, Ulmann A and Yen S.S (1992). Fertil. Steril. Br J Obstet Gynaecol. Induction of abortion with mifepristone and misoprostol in early pregnancy. 1992 Dec;99(12):1004-7.
- Kiernan D.F and Mieler W.F (2009). The use of intraocular corticosteroids. *Expert Opin. Pharmacother.* 10, 2511–2525
- King T.L and Brucker M.C (2010). *Pharmacology for Women's Health*. Jones & Bartlett Publishers. pp. 372–373.
- Kliewer S.A, Forman B.M, Blumberg B, Ong E.S, Borgmeyer U, Mangelsdorf D.J, Umesono K and Evans R.M (1994). Differential expression and activation of a family of murine peroxisome proliferator- activated receptors. *Proc. Natl. Acad. Sci. U.S.A.* 91, 7355–7359.
- Kliewer S.A, Moore J.T, Wade L, Staudinger J.L, Watson M.A, Jones S.A, McKee D.D, Oliver B.B, Willson T.M, Zetterström R.H, Perlmann T and Lehmann J.M (1998). An orphan nuclear receptor activated by pregnanes defines a novel steroid signaling pathway. *Cell* 1998;92:73-82
- Kuiper G.G, Carlsson B, Grandien K, Enmark E, Haggblad J, Nilsson S and Gustafsson J.A (1997). Comparison of the ligand binding specificity and transcript tissue distribution of estrogen receptors alpha and beta. *Endocrinology* 138: 863–870, 1997
- Laffitte B.A, Kast H.R, Nguyen C.M, Zavacki A.M, Moore D.D and Edwards P.A (2000). Identification of the DNA binding specificity and potential target genes for the farnesoid X-activated receptor. *J Biol Chem*. 2000;275: 10638–10647.
- Lazar M.A (1993). Thyroid hormone receptors: Multiple forms, multiple possibilities. *Endocr. Rev.* 1993, 14, 184-193.
- Lee S.S, Pineau T, Drago J, Lee E.J, Owens J.W, Kroetz D.L, Fernandez-Salguero P.M, Westphal H and Gonzalez F.J (1995). Targeted disruption of the alpha isoform of the peroxisome proliferator-activated receptor gene in mice results in abolishment of the pleiotropic effects of peroxisome proliferators. *Mol. Cell. Biol.* 15, 3012–3022.
- Lelliott C.J, López M, Curtis R.K, Parker N, Laudes M, Yeo G, Jimenez-Liñan M, Grosse J, Saha A.K, Wiggins D, Hauton D, Brand M.D, O'Rahilly S, Griffin J.L, Gibbons G.F and Vidal-Puig A. (2005). Transcript and metabolite analysis of the effects of tamoxifen in rat liver reveals inhibition of fatty acid synthesis in the presence of hepatic steatosis. *FASEB J* 2005 Jul;19(9):1108-19.

- Letteron P, Brahimi-Bourouina N, Robin M-A, Moreau A, Feldman G and Pessayre D (1997). Glucocorticoids inhibit mitochondrial matrix acyl-CoA dehydrogenases and fatty acid β -oxidation. *Am. J. Physiol Gastrointest. Liver Physiol.* 272:G1141–G1150; 1997.
- Lewis-Tuffin L.J, Jewell C.M, Bienstock R.J, Collins J.B and Cidlowski J.A (2007). Human glucocorticoid receptor beta binds RU-486 and is transcriptionally active. *Mol. Cell. Biol.* 27, 2266–2282
- Leyden J.J, Shalita A, Thiboutot D, Washenik K and Webster G (2005). Topical retinoids in inflammatory acne: A retrospective, investigator-blinded, vehicle-controlled, photographic assessment. *Clin Ther* 27 (2005), 216-224
- Li T, Yu R.T, Atkins A.R, Downes M, Tukey R.H and Evans R.M (2012). Targeting the pregnane X receptor in liver injury. *Expert Opinion on Therapeutic Targets* 16(11): 1075.
- Li W, Donat S, Dohr O, Unfried K and Abel J (1994). Ah receptor in different tissues of C57BL/6J and DBA/2J mice: Use of competitive polymerase chain reaction to measure Ah-receptor mRNA expression. *Arch. Biochem. Biophys.* 1994;315:279–284. doi: 10.1006/abbi.1994.1501
- Lin J.Z, Martagón A.J, Hsueh W.A, Baxter J.D, Gustafsson J.Å, Webb P and Phillips K.J (2012). Thyroid hormone receptor agonists reduce serum cholesterol independent of the LDL receptor. *J Endocrinology.* 2012 Dec;153(12):6136-44.
- Liu Z, Hu Q and Rosenfeld M.G (2014). Complexity of the RAR-mediated transcriptional regulatory programs. *Subcell Biochem* 70 (2014), 203-225
- Loireau A, Autissier N, Dumas P, Michael O, Jorgensen E.C and Michael R (1986). Comparative effects of 3,5-dimethyl-3'-isopropyl-L-thyronine (DIMIT) and 3,5-diiodo-3'-isopropylthyoacetic acid (IpTA2) on body weight gain and lipid metabolism in genetically obese Zucker rats. *Biochem. Pharmacol.* 1986, 35,1691-1696.
- Love J.D (2006). Nuclear Receptors and Disease. eLS.
- Lu N.Z, Wardell S.E, Burnstein K.L, Defranco D, Fuller P.J, Giguere V, Hochberg R.B, McKay L, Renoir J.M, Weigel N.L, Wilson E.M, McDonnell D.P and Cidlowski J.A (2006). "International Union of Pharmacology. LXV. The pharmacology and classification of the nuclear receptor superfamily: glucocorticoid, mineralocorticoid, progesterone, and androgen receptors". *Pharmacol. Rev.* 58 (4): 782–97.
- Ma WL, Lai HC, Yeh S, Cai X and Chang C (2014). Androgen receptor roles in hepatocellular carcinoma, fatty liver, cirrhosis and hepatitis Endocrine-Related Cancer. 2014;21(3):R165-R182.
- Maglich J.M, Lobe D.C and Moore J.T (2009). The nuclear receptor CAR (NR1I3) regulates serum triglyceride levels under conditions of metabolic stress. *J. Lipid Res.* 50 (2009) 439–445.
- Makishima M, Okamoto A.Y, Repa J.J, Tu H, Learned R.M, Luk A, Hull M.V, Lustig K.D, Mangelsdorf D.J and Shan B (1999). Identification of a nuclear receptor for bile acids. *Science.* 1999;284:1362–1365.
- Miquilena-Colina M.E, Lima-Cabello E, Sanchez-Campos S, Garcia-Mediavilla M.V, Fernandez-Bermejo M, Lozano-Rodriguez T, Vargas-Castrillon J, Buque X, Ochoa B, Aspichueta P, Gonzalez-Gallego J and Garcia-Monzon C (2012). Hepatic fatty acid translocase CD36 upregulation is associated with insulin resistance, hyperinsulinaemia and increased steatosis in non-alcoholic steatohepatitis and chronic hepatitis C. *Gut* 2011;60:1394-1402.
- Misrahi M, Atger M, d'Auriol L, Loosfelt H, Meriel C, Fridlansky F, Guiochon-Mantel A, Galibert F and Milgrom E (1987). Complete amino acid sequence of the human progesterone receptor deduced from cloned cDNA. *Biochem. Biophys. Res. Commun.* 143 (2): 740–8.
- Morani A.B.R, Imamov O, Hultenby K, Arner A, Warner M and Gustafsson J.A (2006). Lung dysfunction causes systemic hypoxia in estrogen receptor beta knockout (ERbeta^{-/-}) mice. *Proc Natl Acad Sci USA* 103: 7165–7169, 2006
- Moreau A, Vilarem M.J, Maurel P and Pascussi J.M (2008). CAR and PXR activation and consequences on lipid metabolism, glucose homeostasis, and inflammatory response. *Mol. Pharm.* 5 (2008) 35–41.
- Moya M, Gómez-Lechón M.J, Castell J.V and Jover R (2010). Enhanced steatosis by nuclear receptor ligands: A study in cultured human hepatocytes and hepatoma cells with a characterized nuclear receptor expression profile. *Chem Biol Interact* 184 (2010), 376-387
- Murphy A.A, Kettel L.M, Morales A.J, Roberts V.J and Yen S.S (1993). Regression of uterine leiomyomata in response to the antiprogestone RU 486. *J. Clin. Endo. Metab.* 1993, 76, 513.
- Naguyen P, Leray V, Diez M, Serisier S, Le Bloch J and Dumon H (2008). Liver lipid metabolism. *Journal of Animal Physiology and Animal Nutrition* 92(2008) 272-283.
- Naik M.A, Banday K.A, Najar M.S, Reshi A.R and Bhat M.A (2008). Vitamin D intoxication presenting as acute renal failure. *Indian J Nephrol.* Jul 2008; 18(3): 125–126.
- National Research Council (NRC) (2007). Toxicity Testing in the 21st Century: A 1321 Vision and a Strategy. National Academies Press, Washington, DC.
- Nebert D.W, Dalton T.P, Okey A.B and Gonzalez F.J (2004). Role of Aryl Hydrocarbon Receptor-mediated Induction of the CYP1 Enzymes in Environmental Toxicity and Cancer *J. Biol. Chem.* 279 (2004) 23847–23850.

- Niittynen M, Simanainen U, Syrjälä P, Pohjanvirta R, Viluksela M, Tuomisto J and Tuomisto J.T (2007). Differences in acute toxicity syndromes of 2,3,7,8-tetrachlorodibenzo-p-dioxin and 1,2,3,4,7,8-hexachlorodibenzo-p-dioxin in rats. *Toxicology* 235 (2007) 39–51.
- Norman A.W (2006). Minireview: vitamin D receptor: new assignments for an already busy receptor. *Endocrinology* 2006;147:5542–8.
- O'Bryant C.L, Flaig T.W and Utz K.J (2008). Bicalutamide-associated fulminant hepatotoxicity. *Pharmacotherapy*. 2008 Aug;28(8):1071-5.
- Osabe M, Sugatani J, Fukuyama T, Ikushiro S, Ikari A and Miwa M (2008). Expression of hepatic UDP-glucuronosyltransferase 1A1 and 1A6 correlated with increased expression of the nuclear constitutive androstane receptor and peroxisome proliferator-activated receptor alpha in male rats fed a high-fat and highsucrose diet. *Drug Metab. Dispos.* 36 (2008) 294–302.
- Parks D.J, Blanchard S.G, Bledsoe R.K, Chandra G, Consler T.G, Kliewer S.A, Stimmel J.B, Willson T.M, Zavacki A.M, Moore D.D and Lehmann J.M (1999). Bile acids: natural ligands for an orphan nuclear receptor. *Science*. 1999;284:1365–1368.
- Patlewicz G, Simon T, Rowlands J.C, Budinsky R.A, and Becker R.A (2015). Proposing a scientific confidence framework to help support the application of adverse outcome pathways for regulatory purposes. *Regul Toxicol Pharmacol*. 2015 Feb 20;71(3):463-477.
- Peet D.J, Turley S.D, Ma W, Janowski B.A, Lobaccaro J.M, Hammer R.E and Mangelsdorf D.J (1998). Cholesterol and bile acid metabolism are impaired in mice lacking the nuclear oxysterol receptor LXR alpha. *Cell* 93 (1998) 693–704.
- Pelclová D, Urban P, Preiss J, Lukás E, Fenclová Z, Navrátil T, Dubská Z and Senholdová Z (2006). Adverse health effects in humans exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). *Rev. Environ. Health* 21 (2006) 119–138
- Peraza M.A, Burdick A.D, Marin H.E, Gonzalez F.J, Peters J.M (2006). The toxicology of ligands for peroxisome proliferator-activated receptors (PPAR). *Toxicol Sci* 90(2): 269-295.
- Pérez E, Bourguet W, Gronemeyer H and de Lera A.R (2012). Modulation of RXR function through ligand design. *Biochim Biophys Acta* 1821 (2012), 57-69
- Peters J.M, Cheung C and Gonzalez F.J (2005). Peroxisome proliferator activated receptor-alpha and liver cancer: Where do we stand? *J. Mol. Med.* 83, 774–785.
- Peters J.M, Hennuyer N, Staels B, Fruchart J.C, Fievet C, Gonzalez F.J and Auwerx J (1997). Alterations in lipoprotein metabolism in peroxisome proliferator-activated receptor alpha-deficient mice. *J. Biol. Chem.* 272, 27307–27312.
- Petitti D.B (2003). Combination Estrogen–Progestin Oral Contraceptives. *N Engl J Med* 2003; 349:1443–1450 October 9, 2003 DOI: 10.1056/NEJMcp030751
- Pineda Torra I, Claudel T, Duval C, Kosykh V, Fruchart J.C and Staels B (2003). Bile acids induce the expression of the human peroxisome proliferator-activated receptor α gene via activation of the farnesoid X receptor. *Mol. Endocrinol.* 2003;17:259–272.
- Postic C and Girard J (2008). Contribution of de novo fatty acid synthesis to hepatic steatosis and insulin resistance: lessons from genetically engineered mice. *J Clin Invest.* 2008 Mar;118(3):829-38.
- Raparti G, Jain S, Ramteke K, Murthy M, Ghanghas R, Ramanand S and Ramanand J (2013). Selective thyroid hormone receptor modulators. *Indian J Endocrinol Metab.* 2013 Mar-Apr; 17(2): 211–218.
- Reddy J.K and Roa S.M (2006). Lipid metabolism and liver inflammation II. Fatty liver disease and fatty oxidation. *Am J Physiol Gastrointest Liver Physiol* 290:G853-G858
- Repa J.J, Berge K.E, Pomajzl C, Richardson J.A, Hobbs H and Mangelsdorf D.J (2002). Regulation of ATP-binding cassette sterol transporters ABCG5 and ABCG8 by the liver X receptors alpha and beta. *J. Biol. Chem.* 277 (2002) 18793–18800.
- Repa J.J, Turley S.D, Lobaccaro J.A, Medina J, Li L, Lustig K, Shan B, Heyman R.A, Dietschy J.M and Mangelsdorf D.J (2000). Regulation of absorption and ABC1-mediated Efflux of cholesterol by RXR heterodimers. *Science* 289 (2000), 1524-1529
- Rosano G.M.C and Fini M (2001). Comparative cardiovascular effects of different progestins in menopause. *INT J F W M*, 46(5), 2001, pp. 248-256
- Rose A.J, Vegiopoulos A and Herzig S (2010). Role of glucocorticoids and the glucocorticoid receptor in metabolism: insights from genetic manipulations. *J. Steroid Biochem. Mol. Biol.* 122, 10–20
- Rosen M.B, Abbott B.D, Wolf D.C, Corton J.C, Wood C.R, Schmid J.E, Das K.P, Zehr R.D, Blair E.T and Lau C (2008). Gene profiling in the livers of wild-type and PPARalpha-null mice exposed to perfluorooctanoic acid. *Toxicol. Pathol.* 36: 592-607.
- Rossier MF (2006). T channels and steroid biosynthesis: in search of a link with mitochondria. *Cell Calcium.* 40 (2): 155–64.
- Roy AK, Lavrovsky Y, Song CS, Chen S, Jung MH, Velu NK, Bi BY, Chatterjee B (1999). "Regulation of androgen action". *Vitam. Horm. Vitamins & Hormones* 55: 309–52.
- Saini S.P, Sonoda J, Xu L, Toma D, Uppal H, Mu Y, Ren S, Moore D.D, Evans R.M and Xie W (2004). A novel constitutive androstane receptor-mediated and CYP3A-independent pathway of bile acid detoxification. *Mol Pharmacol* 2004; 65: 292-300
- Satoh H, Ide N, Kagawa Y and Maeda T (2013). Hepatic steatosis with relation to increased expression of peroxisome proliferator-activated receptor- γ in insulin resistant mice. *Biol Pharm Bull.* 2013;36(4):616-23. Epub 2013 Feb 2.

- Schultz J.R, Tu H, Luk A, Pepa J.J., Medina J.C, Li L, Schwendner S, Wang S, Thoolen M, Mangelsdorf D.J, Lustig K.D and Shan B. (2000) Role of LXR α in control of lipogenesis, *Genes Dev.* 14 (2000) 2831–2838.
- Sevilla L.M, Latorre V, Sanchis A and Pérez P (2012). Epidermal inactivation of the glucocorticoid receptor triggers skin barrier defects and cutaneous inflammation. *J. Invest. Dermatol.* 133, 361–370
- Shalita A.R (1988). Lipid and teratogenic effects of retinoids. *J Am Acad Dermatol* 19 (1988), 197-198
- Shimizu T, Yu H.P, Hsieh Y.C, Choudhry M.A, Suzuki T, Bland K.I and Chaudry I.H (2007). Flutamide Attenuates Pro-inflammatory Cytokine Production and Hepatic Injury Following Trauma-Hemorrhage via Estrogen Receptor-related Pathway. *Ann Surg.* 2007 Feb;245(2):297-304
- Silverman M.N. and Sternberg, E.M. (2012) Glucocorticoid regulation of inflammation and its functional correlates: from HPA axis to glucocorticoid receptor dysfunction. *Ann. N. Y. Acad. Sci.* 1261, 55–63
- Singer E.A and Srinivasan R (2012). Intravenous therapies for castration-resistant prostate cancer: toxicities and adverse events. *Urol Oncol* 2012; 30: S15-9.
- Sitruk-Ware R and Spitz I.M (2003). Pharmacological properties of mifepristone: toxicology and safety in animal and human studies. *Contraception* 2003;68(6):409-20 .Spitz I.M. Progesterone receptor antagonists. *Curr Opin Investig Drugs.* 2006 Oct;7(10):882-90
- Skouby S.O and Petersen K.R (1991). Clinical experience with the recently developed progestogens. *Int. J. Fertility* 1991;36 (Suppl. 1), 32.
- Smith G, Henderson C.J, Parker M.G White R, Bars R.G and Wolf C.R (1993). 1,4-Bis[2-(3,5-dichloropyridyloxy)] benzene, an extremely potent modulator of mouse hepatic cytochrome P-450 gene expression. *Biochem J* 1993;289:807-13
- Sueyoshi T, Kawamoto T, Zelko I, Honkakoski P and Negishi M (1999). The repressed nuclear receptor CAR responds to phenobar-bital in activating the human CYP2B6 gene. *J Biol Chem* 1999;274: 6043-6046
- Sugatani J, Kojima H, Ueda A, Kakizaki S, Yoshinari K, Gong Q.H, Owens I.S, Negishi M and Sueyoshi T (2001). The phenobar-bital response enhancer module in the human bilirubin UDP- glucuronosyltransferase UGT1A1 gene and regulation by the nuclear receptor CAR. *Hepatology* 2001; 33: 1232-1238
- Suino K, Peng L, Reynolds R, Li Y, Cha J.Y, Repa J.J, Kliewer S.A and Xu H.E. (2004). The Nuclear Xenobiotic Receptor CAR: Structural Determinants of Constitutive Activation and Heterodimerization. *Molecular Cell* (Vol. 16): 893–905.
- Takeda K, Sakurai A, DeGroot L.J and Refetoff S (1992). Recessive inheritance of thyroid hormone resistance caused by complete deletion of the protein-coding region of the thyroid hormone receptor- α gene. *J. Clin. Endocrin., Metab.* 1992, 74, 49-55.
- Thomin A, Keller V, Daraï E and Chabbert-Buffet N (2014). Consequences of emergency contraceptives: the adverse effects. July 2014, Vol. 13, No. 7, Pages 893-902 (doi:10.1517/14740338.2014.921678)
- Tronche F, Kellendonk C, Kretz O, Gass P, Anlag K, Orban P.C, Bock R, Klein R and Schütz G. (1999). Disruption of the glucocorticoid receptor gene in the nervous system results in reduced anxiety. *Nat. Genet.* 23, 99–103
- Vatner D.F, Weismann D, Beddow S.A, Kumashiro N, Erion D.M, Liao X, Grover G.J, Webb P, Phillips K.J, Weiss R.E, Bogan J.S, Baxter J, Shulman G.I and Samuel V.T (2013). Thyroid hormone receptor- β agonists prevent hepatic steatosis in fat-fed rats but impair insulin sensitivity via discrete pathways. *American Journal of Physiology - Endocrinology and Metabolism.* Published 1 July 2013 Vol. 305 no. 1
- Venkatesh M, Wang H, Cayer J, Leroux M, Salvail D, Das B, Wrobel J.E and Mani S (2011). In Vivo and In Vitro Characterization of a First-in-Class Novel Azole Analog That Targets Pregnane X Receptor Activation. *Molecular Pharmacology* 80(1): 124.
- Vilasco M, Communal L, Mourra N, Courtin A, Forgez P and Gompel A (2011). Glucocorticoid receptor and breast cancer. *Breast Cancer Res. Treat.* 130, 1–10
- Vinken M (2013). The adverse outcome pathway concept: A pragmatic tool in toxicology. *Toxicology* 312 (2013) 158– 165
- Wada T, Gao J and Xie W (2009). PXR and CAR in energy metabolism. *Trends in Endocrinology & Metabolism* 20(6): 273-279.
- Wang L, Andersson S, Warner M and Gustafsson J.A (2001). Morphological abnormalities in the brains of estrogen receptor beta knockout mice. *Proc Natl Acad Sci USA* 98: 2792–2796, 2001.
- Watkins R.E, Wisely G.B, Moore L.B, Collins J.L, Lambert M.H, Williams S.P, Willson T.M, Kliewer S.A, Redinbo M.R (2001). The human nuclear xenobiotic receptor PXR: structural determinants of directed promiscuity. *Science* 2001;292:2329-33
- Waxman D.J (1999). P450 gene induction by structurally diverse xenochemicals: central role of nuclear receptors CAR, PXR, and PPAR. *Arch Biochem Biophys* 1999;369:11-23
- Weihua Z, Makela S, Andersson L.C, Salmi S, Saji S, Webster J.I, Jensen E.V, Nilsson S, Warner M and Gustafsson J.A (2001). A role for estrogen receptor beta in the regulation of growth of the ventral prostate. *Proc Natl Acad Sci USA* 98: 6330–6335, 2001
- Weinstein R.S, Wan C, Liu Q, Wang Y, Almeida M, O'Brien C.A, Thostenson J, Roberson P.K, Boskey A.L, Clemens T.L and Manolagas S.C (2010). Endogenous glucocorticoids decrease skeletal angiogenesis, vascularity, hydration, and strength in aged mice. *Aging Cell* 9, 147–161

- Wellington K and Perry C.M (2002). Estradiol valerate/dienogest. *Drugs* 2002;62(3):491-50
- Xie W, Barwick JL, Downes M, et al. Humanized xenobiotic response in mice expressing nuclear receptor SXR. *Nature* 2000;406:435-9
- Xie W, Yeuh M.F, Radominska-Pandya A, Saini S.P, Negishi Y, Bottroff B.S, Cabrera G.Y, Tukey R.H and Evans R.M (2003). Control of steroid, heme, and carcinogen metabolism by nuclear pregnane X receptor and constitutive androstane receptor. *Proc Natl Acad Sci USA* 2003; 100: 4150-4155
- Yaghmaei S, Roberts C, Ai R, Mizwicki M.T and Chang C.A (2013). Agonist and antagonist binding to the nuclear vitamin D receptor: dynamics, mutation effects and functional implications. *In Silico Pharmacology* 2013, 1:2
- Yamamoto Y, Moore R, Goldsworthy T.L, Negishi M and Maronpot R.R (2004). The orphan nuclear receptor constitutive active/androstane receptor is essential for liver tumor promotion by phenobarbital in mice. *Cancer Res* 64(20): 7197-7200.
- Ye L, Li Y.L, Mellström K, Mellin C, Bladh L.G, Koehler K, Garg N, Garcia Collazo A.M, Litten C, Husman B, Persson K, Ljunggren J, Grover G, Sleph P.G, George R and Malm J(2003). Thyroid receptor ligands. 1. Agonist ligands selective for the thyroid receptor beta1. *Med Chem.* 2003 Apr 24;46(9):1580-8.
- Yin Y, Yu Z, Xia M, Luo X, Lu X and Ling W (2012). Vitamin D attenuates high fat diet-induced hepatic steatosis in rats by modulating lipid metabolism. Guangdong Provincial Key Laboratory of Food, Nutrition and Health, Department of Nutrition, School of Public Health, Sun Yat-sen University, Guangzhou, China. *European journal of clinical investigation* 07/2012; 42(11):1189-96.
- Zafrani E (2004). Non-alcoholic fatty liver disease: an emerging pathological spectrum. *Virchows Arch*, 2004,444:3-12
- Zaghini I, Landrier J.F, Grober J, Krief S, Jones S.A, Monnot M.C, Lefrère I, Watson M.A, Collins J.L, Fujii H and Besnard P (2002). Sterol regulatory element-binding protein-1c is responsible for cholesterol regulation of ileal bile acid-binding protein gene in vivo. Possible involvement of liver-X-receptor. *J. Biol. Chem.* 277 (2002) 1324–1331.
- Zambrana J, Zambrana F, Neto F, Gonçalves A, Zambrana F and Ushirohira J (2005). Agranulocytosis with tonsillitis associated with methimazole therapy. *Brazilian journal of otorhinolaryngology* 71 (3): 374–377.
- Zen M, Canova M, Campana C, Bettio S, Nalotto L, Rampudda M, Ramonda R, Iaccarino L and Doria A (2011). The kaleidoscope of glucocorticoid effects on immune system. *Autoimmun. Rev.* 10, 305–310
- Zhang, L.; Ge, L.; Parimoo, S.; Stenn, K.; Prouty, S. M. (1999). "Human stearoyl-CoA desaturase: Alternative transcripts generated from a single gene by usage of tandem polyadenylation sites". *The Biochemical journal.* 340 (Pt 1) (Pt 1): 255–264.
- Zhao A, Yu J, Lew J.L, Huang L, Wright S.D and Cui J (2004). Polyunsaturated fatty acids are FXR ligands and differentially regulate expression of FXR targets. *DNA Cell Biol.* 2004;23:519 –526.

Table 1: **Nuclear Receptors associated with liver injury and their abbreviations**

| Nuclear receptor name | Abbreviation |
|--|--------------|
| Aryl Hydrocarbon Receptor | AHR |
| Constitutive Androstane Receptor | CAR |
| oEstrogen Receptor | ER |
| Farnesoid X Receptor | FXR |
| Glucocorticoid Receptor | GR |
| Liver X Receptor | LXR |
| Peroxisome Proliferator-Activated Receptor | PPAR |
| Pregnane X Receptor | PXR |
| Retinoic Acid Receptor | RAR |

Table 2: Summary of the effects on the liver following activation of Nuclear Receptors

| Nuclear receptor | Agonist effect on liver | Antagonist effect on liver |
|------------------|--|----------------------------|
| AHR | Induces hepatic steatosis* | - |
| CAR | Induces hepatic steatosis* | - |
| ER | Induces hepatic steatosis* | Induces hepatotoxicity |
| FXR | Induces hepatic steatosis* | - |
| GR | Induces hepatic steatosis* | |
| LXR | Induces hepatic steatosis* | - |
| PPAR | PPAR γ Induces hepatic steatosis* | PPAR α |
| PXR | Induces hepatic steatosis* | - |
| RAR | Induces hepatic steatosis* | - |
| RXR | Induces hepatic steatosis* | - |

Figure Titles

Figure 1. Structure of an AOP adapted from Ankley et al (2010)

Figure 2. Updated AOP for steatosis, developing that presented by Vinken (2013)




Figure 1




Figure 2