

1 **Patterns of philopatry and longevity contribute to the evolution of post-reproductive**
2 **lifespan in mammals**

3 Nichols H.J.*⁺¹, Zecherle L.^{1*}, Arbuckle, K.²

4 1. Liverpool John Moores University, School of Natural Science and Psychology, Liverpool, UK
5 L3 3AF.

6 2. University of Liverpool, Institute of Integrative Biology, Liverpool, UK L69 7ZB.

7 * These authors contributed equally to the manuscript

8 † Corresponding author: h.j.nichols@ljmu.ac.uk

9 **Abstract**

10 While menopause has long been known as a characteristic trait of human reproduction, evidence for
11 post-reproductive lifespan (PRLS) has recently been found in other mammals. Adaptive and non-
12 adaptive hypotheses have been proposed to explain the evolution of PRLS, but formal tests of these
13 are rare. We use a phylogenetic approach to evaluate hypotheses for the evolution of PRLS among
14 mammals. In contrast to theoretical models predicting that PRLS may be promoted by male
15 philopatry (which increases relatedness between a female and her group in old age), we find little
16 evidence that male philopatry led to the evolution of a post-reproductive period. However, the
17 proportion of life spent post-reproductive was related to lifespan and patterns of philopatry,
18 suggesting that the duration of PRLS may be impacted by both non-adaptive and adaptive processes.
19 Finally, the proportion of females experiencing PRLS was higher in species with male philopatry and
20 larger groups, in accordance with adaptive models of PRLS. We suggest that the origin of PRLS
21 primarily follows the non-adaptive 'mismatch' scenario, but that patterns of philopatry may
22 subsequently confer adaptive benefits of late-life helping.

23 **Key Words**

24 Menopause, life-history evolution, kin-selection, dispersal, primate, cetacean

25

26 **Introduction**

27 Menopause has long been known as a characteristic of human reproduction [1] but the
28 existence of post-reproductive lifespan (PRLS) in other mammalian species has been recognised
29 relatively recently [2, 3]. Post-reproductive periods of 20+ years (similar to that observed in humans)
30 have been found in two long-lived cetacean species [4]. Shorter periods of PRLS have been identified
31 in a wide range of mammalian species, including cetaceans, primates and rodents [2]. However, the

32 existence of post-reproductive lifespan is an evolutionary paradox; natural selection would be
33 expected to disfavour the premature cessation of reproduction. Why then is PRLS so widespread?

34 PRLS may have an adaptive value. For example, menopause could be favoured if mothers
35 provide help to adult offspring, thereby increasing the production of grand-offspring [5]. Lahdenperä
36 et al. [6] showed that human grandmothers are able to boost the reproductive success of their
37 children. Similarly, in killer whales, post-reproductive females appear to extend the longevity of their
38 adult offspring [7]. This adaptive 'grandmother hypothesis' for the evolution of PRLS depends on the
39 presence of kin towards whom help can be directed.

40 Johnstone and Cant [8] developed a model to explain the involvement of kin selection in the
41 evolution of prolonged PRLS in cetaceans and humans which suggested parallel routes for the
42 evolution of PRLS. While early humans are believed to have been male-philopatric, in killer whales
43 and short-finned pilot whales neither sex disperses and mating occurs outside the family group.
44 These dispersal strategies are predicted to lead to an increase in relatedness to other group
45 members throughout the lifetime of a female. In both situations, females begin their reproductive
46 life away from their father and other paternal relatives (either because she has dispersed or because
47 she was the product of an extra-group mating). However, her sons remain within the social group,
48 and hence relatedness between the female and local males (and therefore average relatedness to
49 group members) increases over the female's lifespan, thereby leading to the evolution of an
50 adaptive period of post-reproductive helping behaviour [8].

51 While the literature focuses on PRLS as an adaptive trait, it could simply be a non-adaptive
52 by-product of other life history traits. PRLS is of short duration in most mammals, leading to the
53 proposal that PRLS is a consequence of mismatch in somatic versus reproductive senescence [2].
54 This mismatch may be more likely to occur in long-lived species as the associated variability in
55 maximum lifespan leads to an increased probability that some individuals exceed the age by which
56 oocytes are depleted.

57 Attempts to test the predictions of adaptive versus non-adaptive hypotheses for the
58 evolution of PRLS are lacking, despite a theoretical framework for both classes of explanation [2, 8].
59 In this study, we use a comparative approach to investigate whether natural history traits can
60 predict the existence and extent of PRLS among mammals. If PRLS has arisen adaptively due to kin
61 selection, we expect sex-specific dispersal dynamics to be important in the evolution of PRLS [8].
62 Alternatively, if PRLS arises primarily due to a mismatch between somatic and reproductive ageing,
63 then we would expect PRLS to be seen in longer-lived species.

64 **Methods**

65 ***Data Collection***

66 A literature search was conducted to identify all mammalian species for which reliable PRLS
67 data is available (see supplementary material for our strategy for categorising PRLS including
68 caveats, and Table S1 for the data obtained). We recorded the presence or absence of PRLS, the
69 duration of PRLS, and the frequency with which PRLS is experienced in the population. For species
70 which we had data on the presence or absence of PRLS we continued our literature search to obtain
71 data on natural history variables likely to influence local relatedness (male philopatry, female
72 philopatry, and group-size), and lifespan (in years), which could influence the mismatch between
73 somatic and reproductive ageing. Only data from wild populations was used since captivity can alter
74 the incidence and details of PRLS [9] and therefore arguably adds no information to evolutionary
75 studies of the trait. A dated phylogenetic tree of mammals was obtained from the literature [10] and
76 pruned in Mesquite [11] to leave the 26 mammal species for which we had PRLS data (note that we
77 included three populations of humans in some analyses so some sample sizes were greater than 26).
78 This pruned tree was used for all comparative analyses conducted.

79 ***Statistical analysis***

80 ***What influences the presence of PRLS?***

81 We fit generalised linear mixed models (GLMMs) with a binomial error structure using
82 Markov chain Monte Carlo (MCMC) with an inverse gamma hyperprior to investigate whether each
83 natural history variable (male philopatry, female philopatry, group size and lifespan) was a predictor
84 of the presence of PRLS. We coded the absence or presence of PRLS as having states 0 and 1
85 respectively and used this as our response variable. The phylogeny was included as a random effect
86 to account for evolutionary history and these models were run in the MCMCglmm package in R [12].
87 To avoid over-parameterisation each model contained only one explanatory variable. Each MCMC
88 GLMM was run for 15 million generations, the first 500,000 of which were conservatively discarded
89 as burnin. The chain was sampled every 10,000 generations, giving 1,450 posterior samples for each
90 model.

91 For significant predictors of the presence/absence of PRLS we also reconstructed ancestral
92 states to further assess how the traits evolved with respect to each other. Ancestral state
93 reconstruction was conducted using Bayesian stochastic mapping in phytools [13] and inference
94 made based on 10,000 simulations.

95 ***What influences the duration of PRLS?***

96 We measured relative duration of PRLS as the proportion of maximum lifespan spent post-
97 reproductive (Table S1). We tested for effects of each natural history variable (male philopatry,
98 female philopatry, group-size, and lifespan) individually on the relative duration of PRLS using
99 generalised estimating equations (GEEs), which were fitted in the ape package in R [14]. The
100 variance-covariance matrix for the GEEs was specified based on the phylogeny, which controls for
101 phylogenetic relationships between species by including this information within the model.

102 ***What influences the frequency of PRLS?***

103 To investigate which factors influence the proportion of individuals that experience PRLS, we
104 modelled this variable as a function of each natural history variable (male philopatry, female

105 philopatry, group-size, and lifespan). We used GEEs to control for any influence of phylogeny which
106 were fit as described in the preceding section.

107

108 **Results and Discussion**

109 We took a phylogenetic approach to investigate natural history factors influencing post-
110 reproductive lifespan in mammals with the aim of assessing whether adaptive or non-adaptive
111 scenarios best explain its evolution. In accordance with theoretical work by Johnstone and Cant [8],
112 we found a significant association between the presence of PRLS and male philopatry (MCMC
113 GLMM: $\beta=340.52$, $P=0.018$, Table S2). However, Johnstone and Cant's model predicts that male
114 philopatry is a key (but not the only) evolutionary driver of PRLS, which was not supported by our
115 results. All 5 species with confirmed male philopatry exhibited PRLS, but ancestral state
116 reconstructions suggest that PRLS evolves first, followed by male philopatry (at least in primates)
117 (Figure 1). Furthermore, 50% of the 18 species with dispersing males also exhibited PRLS, again
118 suggesting that male philopatry is unlikely to explain the origin of PRLS in mammals.

119 If PRLS is typically of short duration then it is possible that patterns of philopatry are
120 important in the evolution of an extended period of PRLS due to their influence on kinship [4].
121 Supporting this, we found that species with female philopatry had significantly shorter periods of
122 PRLS (GEE: $\beta\pm SE=-1.573\pm 0.681$, $t_{1,20}=-2.308$, $P=0.048$, Table S3, Figure 2a), and species with male-
123 philopatry had a (non-significant) trend for increased periods of PRLS (GEE: $\beta\pm SE=1.394\pm 0.676$,
124 $t_{1,20}=2.063$, $P=0.071$, Table S3). However we note that we cannot rule out the possibility that male
125 philopatry is associated with PRLS via its positive relationship with lifespan (pGLS: $t=4.06$, $df=1,20$,
126 $P=0.001$). Furthermore, factors other than dispersal patterns are expected to influence the adaptive
127 evolution of PRLS, such as the opportunity for late-life helping and competition (which we did not
128 fully investigate here), but Johnstone and Cant [8] propose dispersal as an important driver.

129 We found that the relative duration of PRLS was greater in longer-lived species (GEE:
130 $\beta \pm SE = 0.038 \pm 0.011$, $t_{1,23} = 3.482$, $P = 0.007$, Table S3, Figure 2b). While a previous study [15] found a
131 relationship between time spent postreproductive and lifespan, they modelled the total duration of
132 PRLS, rather than the relative duration (as we have calculated here). Our measure is unlikely to be
133 inherently associated with lifespan, and therefore suggests that the relationship between PRLS and
134 lifespan is not simply an artefact of longer living species spending more months/years post-
135 reproductive. Instead, our results are consistent with the idea that extended PRLS can occur due to
136 mismatching of somatic and reproductive ageing [2]. It could also be related to selection on
137 increased male lifespan, which could in turn lead to extended female lifespan via intersexual genetic
138 correlations, even if they are not reproductive [2, 16, 17]. Disentangling these alternative non-
139 adaptive scenarios would require detailed investigation into the evolutionary genetic constraints on
140 lifespan across a wide range of mammals.

141 The proportion of females experiencing PRLS was higher in male philopatric species (GEE:
142 $\beta \pm SE = 1.900 \pm 0.786$, $t_{1,15} = 2.418$, $P = 0.047$, Table S3, Figure 2c), suggesting that male philopatry could
143 drive the evolution of widespread PRLS in a species [7]. We also found that PRLS was more prevalent
144 in larger groups (GEE: $\beta \pm SE = 0.051 \pm 0.014$, $t_{1,15} = 3.762$, $P = 0.0075$, Table S3, Figure 2d), possibly
145 because these contain more philopatric young, and hence greater opportunities for helping.
146 However, there was significant covariation between male philopatry and group size (GEE:
147 $\beta \pm SE = 43.680 \pm 18.437$, $t_{1,20} = 2.369$, $P = 0.045$), making it difficult to distinguish between their effects,
148 especially with the limited number of species for which data is currently available. Future models of
149 the evolution of PRLS may benefit from exploring these relationships further by investigating the
150 impacts of both group size and philopatry on kinship dynamics.

151

152 **Conclusions**

153 We tested the predictions of the most common adaptive model for the evolution of PRLS [8]
154 and our results provide mixed support for such a model. We suggest that adaptive models such as
155 that by Johnstone and Cant [8] may be important in explaining prolonged periods of PRLS (as they
156 were intended to do), but do not provide a good explanation for the occurrence of shorter periods
157 of PRLS that seem to be prevalent across mammals [2, 5]. Rather, the evolutionary origin of PRLS
158 appears to primarily follow a non-adaptive scenario such as the 'mismatch' hypothesis [2, 18].
159 Patterns of philopatry may subsequently confer adaptive benefits of late-life helping which extends
160 the duration and frequency of PRLS [5, 7, 8]. Under this scenario we suggest that the prolonged
161 periods of PRLS found in a few species such as humans and cetaceans are a consequence of non-
162 adaptive origins followed by adaptive evolutionary 'tinkering'. Our results also demonstrate that for
163 some analyses, it may be important to consider different components of PRLS separately, rather
164 than combined in a single measure such as PrR [15]. Different factors are likely to govern the
165 evolution of the presence, absence and duration of PRLS and conflation of these elements in a single
166 index limits our ability to evaluate many ideas.

167

168 **Data Accessibility**

169 Data are provided in Table S1.

170

171 **Ethical Statement**

172 This study did not use human or animal subjects and therefore there were no ethical concerns which
173 required evaluation.

174

175 **Competing Interests**

176 The authors declare that there are no competing interests.

177

178 **Author Contributions**

179 HJN and KA devised the study. LZ collected the majority of the data. KA and LZ analysed the data. KA
180 and HJN wrote the first draft of the manuscript. All authors commented on the manuscript,
181 approved the final version for publication, and accept accountability for all aspects of the work.

182

183 **Funding Statement**

184 No funding was received for the work presented herein.

185

186 **Figure Legends**

187 Figure 1. Summary of ancestral state reconstructions for PRLS (left) and male philopatry (right).

188 Posterior probabilities (PP) of state 1 (trait is present) are represented by a greyscale gradient.

189 Figure 2. Relationships between the duration of PRLS and (a) female philopatry, (b) maximum
190 lifespan, and between the prevalence of PRLS and (c) male philopatry and (d) group size.

191

192 **References**

- 193 1. Blurton-Jones, N.G., Howkes, K., O'Connell, J.F. 2002 Antiquity of post-reproductive life: are
194 there modern impacts on hunter-gatherer postreproductive lifespans? *American Journal of Human*
195 *Biology* **14**,184-205.
- 196 2. Cohen, A.A. 2004 Female post-reproductive lifespan: a general mammalian trait. *Biological*
197 *Reviews* **79**,733-750.
- 198 3. Packer, C., Tatar, M., Collins, A. 1998 Reproductive cessation in female mammals. *Nature*
199 **392**,807-811.

200 4. Croft, D.P., Brent, L.J.N., Franks, D.W., Cant, M.A. 2015 The evolution of prolonged life after
201 reproduction. *Trends in Ecology & Evolution* **30**,407-416.

202 5. Shanley, D.P., Kirkwood, T.B. 2001 Evolution of the human menopause. *Bioessays* **23**,282-
203 287.

204 6. Lahdenperä, M., Lummaa, V., Helle, S., Tremblay, M., Russell, A.F. 2004 Fitness benefits of
205 prolonged post-reproductive lifespan in women. *Nature* **428**,178-181.

206 7. Foster, E.A., Franks, D.W., Mazzi, S., Darden, S.K., K.C., B., Ford, J.K.B., Croft, D.P. 2012
207 Adaptive prolonged postreproductive life span in killer whales. *Science* **337**,1313.

208 8. Johnstone, R.A., Cant, M.A. 2010 The evolution of menopause in cetaceans and humans: the
209 role of demography. *Proceedings of the Royal Society of London Series B, Biological Sciences*
210 **277**,3765-3771.

211 9. Levitis, D.A., Burger, O., Lackey, L.B. 2013 The human post-fertile lifespan in comparative
212 evolutionary context. *Evolutionary Anthropology: Issues, News, and Reviews* **22**,66-79.

213 10. Bininda-Emonds, O.R.P., Cardillo, M., Jones, K.E., MacPhee, R.D.E., Beck, R.M.D., Grenyer, R.,
214 Price, S.A., Vos, R.A., Gittleman, J.L., Purvis, A. 2007 The delayed rise of present-day mammals.
215 *Nature* **446**,507 - 512.

216 11. Maddison, W.P., Maddison, D.R. Mesquite: a modular system for evolutionary analysis.
217 Version 2.75. <http://mesquiteproject.org>2011.

218 12. Hadfield, J.D. 2010 MCMC Methods for Multi-Response Generalized Linear Mixed Models:
219 The MCMCglmm R Package. *Journal of Statistical Software* **33**,1-22.

220 13. Revel, L.J. 2012 Phytools: an R package for phylogenetic comparative biology (and other
221 things). *Methods in Ecology and Evolution* **3**,217-223.

222 14. Paradis, E., J., C. 2002 Analysis of comparative data using generalized estimating equations.
223 *Journal of Theoretical Biology*, **218**,175-185.

224 15. Levitis, D.A., Lackey, L.B. 2011 A measure for describing and comparing postreproductive life
225 span as a population trait. *Methods in Ecology and Evolution* **2**,446-453.

226 16. Bonduriansky, R., Chenoweth, S.F. 2009 Intralocus sexual conflict. *Trends in Ecology and*
227 *Evolution* **24**,280-288.

228 17. Tuljapurkar, S.D., Puleston, C.O., Gurven, M.D. 2007 Why Men Matter: Mating Patterns Drive
229 Evolution of Human Lifespan. *PLoS ONE* **2**,e785.

230 18. Ricklefs, R.E., Scheuerlein, A., Cohen, A. 2003 Age-related patterns of fertility in captive
231 populations of birds and mammals. *Experimental Gerontology* **38**,714-745.