

# **THERMOREGULATION AND FLUID BALANCE DURING A 30 KM MARCH IN 60 VERSUS 80-YEAR OLD SUBJECTS**

RUNNING PAGE HEADLINE: Temperature and fluid balance in elderly

COEN C.W.G. BONGERS<sup>1</sup>

THIJS M.H. EIJVOGELS<sup>1,2</sup>

JEAN NYAKAYIRU<sup>1</sup>

MATTHIJS T.W. VELTMEIJER<sup>1</sup>

DICK H.J. THIJSEN<sup>1,3</sup>

MARIA T.E. HOPMAN<sup>1</sup>

<sup>1</sup>Department of Physiology, Radboud university medical center, Nijmegen, The Netherlands

<sup>2</sup>Henry Low Heart Center, Department of Cardiology, Hartford Hospital, Hartford, Connecticut, USA

<sup>3</sup>Liverpool John Moores University, Liverpool, United Kingdom

**Total Word Count:** 4.542

**Word Count Abstract:** 228

**Total Number of Tables:** 2

**Total Number of Figures:** 2

**Conflicts of interest:** None

**Disclosure statement funding:** T.M.H.E is financially supported by the Netherlands Organization for Scientific Research (Rubicon Grant 825.12.016), D.H.J.T received an E. Dekker stipend of the Dutch Heart Foundation (2009T064).

**Author for correspondence:**

Prof. Dr. Maria Hopman, Department of Physiology (392), Radboud university medical center,  
P.O. Box 9101, 6500 HB Nijmegen, The Netherlands, Tel +31 24 36 14 200, Fax +31 24 36  
16413, E-mail: [Maria.Hopman@radboudumc.nl](mailto:Maria.Hopman@radboudumc.nl)

## ABSTRACT

**BACKGROUND.** The presence of impaired thermoregulatory and fluid balance responses to exercise in older individuals is well established. To improve our understanding on thermoregulation and fluid balance during exercise in older individuals, we compared thermoregulatory and fluid balance responses between sexagenarians and octogenarians during prolonged exercise. **METHODS.** 40 sexagenarians ( $60\pm 1$  year) and 36 octogenarians ( $81\pm 2$  year) volunteered to participate in a 30 km march at a self-selected pace. Intestinal temperature ( $T_{in}$ ) and heart rate were recorded every 5 km. Subjects reported fluid intake, whilst urine output was measured and sweat rate was calculated. **RESULTS.** Octogenarians demonstrated a lower baseline  $T_{in}$  and a larger exercise-induced increase in  $T_{in}$  compared to sexagenarians ( $1.2\pm 0.5^{\circ}\text{C}$  *versus*  $0.7\pm 0.4^{\circ}\text{C}$ ,  $p<0.01$ ), whilst maximum  $T_{in}$  tended to be higher in octogenarians ( $38.4\pm 0.4^{\circ}\text{C}$  *versus*  $38.2\pm 0.3^{\circ}\text{C}$ ,  $p=0.09$ ). Exercise intensity ( $70\pm 11\%$  *versus*  $70\pm 9\%$ ) and exercise duration ( $7\text{h}45\text{min}\pm 0\text{h}57\text{min}$  *versus*  $7\text{h}24\text{min}\pm 0\text{h}58\text{min}$ ) were not different between octogenarians and sexagenarians. Octogenarians demonstrated lower fluid intake ( $251\pm 97$  mL/h *versus*  $325\pm 125$  mL/h,  $p=0.01$ ) and urine output ( $28\pm 22$  mL/h *versus*  $52\pm 40$  mL/h,  $p<0.01$ ) compared to sexagenarians. Furthermore, the sweat rate tended to be lower ( $294\pm 150$  mL/h *versus*  $364\pm 148$  mL/h,  $p=0.07$ ) in the octogenarian group. Sodium levels and plasma volume changes were not different between sexagenarians and octogenarians (all  $p>0.05$ ). **CONCLUSIONS.** These results suggest that thermoregulatory responses deteriorate with advancing age, while fluid balance is regulated appropriately during a 30 km walking march under moderate ambient conditions.

**KEYWORDS:** Walking exercise, hyperthermia, dehydration, aging, endurance exercise

## INTRODUCTION

The increased metabolic heat production during exercise leads to a rise in core body temperature (Bouchama and Knochel 2002; Kenefick et al. 2007) and may ultimately lead to the development of heat-related illnesses (Tattersson et al. 2000). The primary heat dissipating mechanism during exercise is evaporation through sweating, accounting for 80% of total heat loss (Bar-Or 1998; Sawka et al. 2007). However, the production of sweat in combination with insufficient fluid replacement may lead to the development of dehydration (Cheuvront et al. 2010) and could impair exercise performance and increase the risk to develop hyperthermia (Sawka 1992).

Advanced age is associated with a negative impact on thermoregulatory and fluid balance responses during exercise (Blatteis 2012; Chester and Rudolph 2011). Previous studies, which compared young *versus* old subjects, showed that elderly  $\geq 60$  years have a lower baseline core body temperature, a reduced skin vasodilatory capacity, a less effective sweat response, and a decreased sensitivity of the thermal receptors (Blatteis 2012; Chester and Rudolph 2011; Guergova and Dufour 2011). Furthermore, advanced age is associated with a decreased total body water (Davis 1994; Schoeller 1989), decreased thirst sensation (Mack et al. 1994; Phillips et al. 1984), decline in kidney functioning (Davis 1994; Lindeman et al. 1985) and a reduced plasma vasopressin regulation at rest and during dehydration (Miller and Shock 1953; Sheehy et al. 1999). These characteristics make older humans a vulnerable population for the development of hyperthermia and dehydration (Begum 2010; Blatteis 2012; Chester and Rudolph 2011; Montes 2006; Sheehy et al. 1999). Most of this knowledge is derived from cross-sectional observations between cohorts of young (usually  $<30$  years) and older humans (usually  $>60$

years). Accordingly, it is unknown whether thermoregulation and fluid balance control deteriorates further with aging or plateaus at some point.

The purpose of our study was, therefore, to compare thermoregulatory responses and fluid balance during prolonged, moderate-intensity exercise between sexagenarians and octogenarians. We hypothesized that exercise-induced thermoregulatory responses and fluid balance control are further deteriorated in octogenarians compared to sexagenarians.

## **MATERIALS and METHODS**

### **Subjects**

A total of 79 subjects volunteered to participate in this study (Table 1). Subjects were recruited via a newsletter, which was distributed by the organisation of the Nijmegen Four Days Marches (an annual walking event in The Netherlands). Based on the age classification of previous publications, subjects were divided into the group of sexagenarians ( $60\pm 1$  year, 24 men and 16 women) or octogenarians ( $81\pm 2$  year, 25 men and 14 women). The following subjects were excluded from participation: I) body mass < 36.5 kg, II) implanted electro-medical device, III) a history of gastro-intestinal disease or abdominal surgery, and IV) a scheduled MRI-scan within 1 week after participating in the present study (all based on the use of the temperature pill). The study was approved by the Medical Ethical Committee of the Radboud university medical center, and all subjects gave written informed consent prior to participation. This study was conducted in accordance with the Declaration of Helsinki.

## **Experimental design**

Baseline measurements, including subject characteristics, body composition, urine specific gravity and blood levels of sodium, hemoglobin and hematocrit, were performed in our temperature controlled laboratory one or two days prior to the exercise bout. The blood pressure was measured in supine position after lying on a bed for 5 minutes, whilst a blood sample was taken in sitting position. Immediately before the start of exercise, subjects underwent assessment of body mass, heart rate (HR), and intestinal temperature ( $T_{in}$ ). Thereafter, subjects participated in the first day of the Nijmegen Four Days marches and walked 30 km at a self-selected pace, all starting at 7.00 AM. The exercise bout consisted of a 30 km self-paces march on a flat course over tarmac roads. During exercise,  $T_{in}$  and HR were recorded after every 5 km, while urine excretion was collected using collecting bags and all subjects registered their fluid intake using a diary. Directly after finishing, subjects reported to our laboratory to determine post-exercise body mass, HR,  $T_{in}$ , fluid balance (urine and blood analysis).

## **Measurements**

*Subject characteristics.* Body mass (Seca 888 scale, Hamburg, Germany) and body height were measured and body mass index (BMI) was calculated. A four-point skin fold thickness measurement (biceps, triceps, sub-scapular, supra-iliac) was performed in order to calculate the lean body mass (Durnin and Womersley 1974). Waist circumference was measured midway between the lower rib margin and iliac crest. Hip circumference was measured at the level of widest circumference over greater trochanters. Waist-to-hip ratio was calculated as waist circumference divided by hip circumference. Thereafter, resting heart rate and blood pressure

were measured twice using an automated sphygmomanometer (M5-1 intellisense, Omron Healthcare, Hoofddorp, the Netherlands) after 5 minutes supine rest. Finally, all subjects completed a questionnaire concerning their physical activity and health status, in which subjects were asked for medical history, medication usage, daily activity (hours of sport participating per week) and training status (walking-specific training history in the year preceding the walking march).

### **Thermoregulation**

*Intestinal temperature ( $T_{in}$ ).* A portable telemetry system (CoreTemp™ system, HQ Inc, Palmetto, USA) was used to measure  $T_{in}$ , which is a safe and reliable method to examine  $T_{in}$  at rest and during exercise (Byrne and Lim 2007; Gant et al. 2006). Subjects ingested an individually calibrated telemetric temperature pill on the evening preceding the exercise bout, to avoid interaction with fluid ingestion during testing (Wilkinson et al. 2008). Baseline  $T_{in}$  was determined as the average of 3 consecutive measurements. This procedure was repeated every 5 km along the route. The highest value of these measurements was presented as peak  $T_{in}$  and increment in  $T_{in}$  ( $\Delta T_{in}$ ) was calculated as peak  $T_{in}$  minus baseline  $T_{in}$ .

*Heart rate (HR).* Heart rate was measured simultaneously with  $T_{in}$  (i.e. 3 consecutive measurements every 5 km) using a 2-channel ECG chest band system (Polar Electro Oy, Kempele, Finland). Mean heart rate during exercise was calculated as the average heart rate, excluding the values measured directly before the start and after the finish. Exercise intensity was calculated by dividing the mean heart rate during exercise by the maximal predicted heart rate ( $208-0.7*\text{age}$ ) (Tanaka et al. 2001).

### **Fluid balance**

*Fluid intake.* All subjects received written and individual oral instructions regarding the registration of their fluid intake. During exercise subjects were allowed to drink *ad libitum*, while they registered the time (in blocks of 1 hour), amount (using standard sized cups (125 mL), cans (330 mL) and bottles (500 mL)), and type ('water', 'sports drink' or 'other') of their individual fluid intake in a diary. Directly post-exercise, the fluid intake diary was checked by the research team, and clarifications were provided by the subjects if necessary. The relative change in body mass (in %) between the measurement immediately before the start and directly after finishing was calculated.

*Sweat rate.* Sweat rate (mL/h) was calculated as a combination of body mass, fluid intake and urine output data using the following formula:  $\text{sweat rate (mL/h)} = (\text{pre-exercise body mass} - \text{post exercise body mass} + \text{fluid intake} - \text{urine output}) / \text{exercise duration}$  (Casa et al. 2000).

*Blood sodium levels + plasma volume.* Subjects were seated for 5 minutes after which a 2 mL blood sample was drawn from the antecubital vein. Blood samples were directly analyzed for plasma levels of sodium, hemoglobin and hematocrit (Rapidpoint® 400, Siemens Healthcare Diagnostics Inc., Tarrytown, New York, USA). Hyponatremia and hypernatremia were defined as a plasma sodium concentration of  $\leq 135$  and  $\geq 145$  mmol/L, respectively (Adroque and Madias 2000; Hew-Butler et al. 2005). Relative changes in plasma volume were calculated from blood hematocrit and hemoglobin concentrations using Dill and Costill's equation (Dill and Costill 1974).

*Urine analysis.* To determine urine specific gravity, a 5 mL urine sample was provided by all subjects and directly analyzed (Clinitek Status® Analyzer, Siemens Healthcare Diagnostics Inc., Tarrytown, New York, USA). A urine specific gravity of  $\geq 1.020$  g/mL is indicative for

dehydration (Casa et al. 2000). To determine the total amount of urine output, subjects were instructed to exclusively urinate into a specialized collecting bag (Roadbag/Ladybag, KETs GmbH, Köln, Germany) during the entire exercise bout. Bags were collected after every 5 km and weighted at the laboratory within 0.1 g accuracy (PT 1400, Sartorius AG, Göttingen, Germany).

*Ambient conditions.* During the experiment, dry bulb, wet bulb and globe temperatures were measured every 30 minutes using a portable climate monitoring device (Davis instruments Inc., Hayward, USA), which is positioned at the start/finish area. The wet bulb globe temperature index (WBGT) was calculated using the formula:  $WBGT = 0.1 (T_{dry\ bulb}) + 0.7 (T_{wet\ bulb}) + 0.2 (T_{globe})$  (Armstrong et al. 2007).

### **Statistical analysis**

All values are presented as mean±standard deviation, unless indicated otherwise. Statistical analyses were performed using Statistical Package for Social Sciences 20.0 (IBM SPSS version 20.0, Armonk, New York, USA) and the level of significance was set at  $p < 0.05$ . The Kolmogorov-Smirnov test was used to examine the normality of the data distribution. When data demonstrated a non-Gaussian distribution, Ln-transformation was applied. Comparisons of baseline measurements between groups were assessed using an unpaired Student's t-test. To assess differences in fluid balance between the sexagenarians and the octogenarians, an unpaired Student's t-test was used. A linear mixed model analysis was used to determine whether changes in  $T_{in}$  and exercise intensity across the 30-km exercise differed between both groups. A two-way (2X2) repeated measures ANOVA was used to examine whether exercise altered plasma sodium,



hemoglobin and hematocrit concentration ('exercise': pre *versus* post), and whether these changes differed between groups ('group': sexagenarians *versus* octogenarians). Finally, a Pearson's Chi-square test was applied to assess differences in the prevalence of dehydration, hypo- or hypernatremia and high urine specific gravities between groups. Results of the Pearson's Chi-square test were presented as relative risk (RR) and their 95% confidence intervals.

## **RESULTS**

Three octogenarian subjects did not finish the 30 km walking exercise bout due to a fractured wrist and severe muscle pains and fatigue, and were excluded from further analysis. The average walking distance in the year prior to the march was  $501\pm 423$  km and  $990\pm 723$  km for the sexagenarians and octogenarians, respectively ( $p<0.01$ ). Octogenarians had a lower height, body mass, body mass index (BMI), body fat, lean body mass and abdominal and waist circumference compared to sexagenarians (all  $p<0.05$ , Table 1), whilst no differences were found for mean arterial pressure and heart rate. No differences in prevalence of cardiovascular diseases, hypertension, hypercholesterolemia and diabetes were reported, additionally we found no differences in medication usage (Table 1, all  $p>0.05$ ).

### **Exercise characteristics**

Relative humidity and wet bulb globe temperature (WBGT) were 95% and 12°C at the start of the exercise bout (i.e. 7.00 AM) and gradually changed to 43% and 24°C at 5.00 PM. Exercise

duration and walking speed were comparable between both groups (Table 2). Exercise intensity increased during the exercise bout to a similar extent in both groups ( $p=0.88$ , Figure 1A).

### **Intestinal temperature**

Baseline intestinal temperature ( $T_{in}$ ) was  $0.3^{\circ}\text{C}$  lower in the octogenarians ( $37.2\pm 0.3$ ) compared to the sexagenarians ( $37.5\pm 0.4$ ,  $p<0.01$ ).  $T_{in}$  increased significantly during exercise ( $p<0.01$ ), whilst the increase in  $T_{in}$  was significantly larger in octogenarians compared to sexagenarians ( $1.2\pm 0.5^{\circ}\text{C}$  versus  $0.7\pm 0.4^{\circ}\text{C}$ ,  $p<0.01$ ). In contrary, maximum  $T_{in}$  tended to be higher in the octogenarians ( $38.4\pm 0.4^{\circ}\text{C}$  versus  $38.2\pm 0.3^{\circ}\text{C}$ ;  $p=0.09$ ).

### **Fluid balance**

Average fluid intake during exercise was lower for the octogenarians ( $251\pm 97$  mL/hour) compared to the sexagenarians ( $325\pm 125$  mL/hour,  $p=0.01$ ) (Figure 2A). Furthermore, octogenarian subjects demonstrated a significantly lower urine output compared to sexagenarians ( $p<0.01$ ), and the sweat rate tended to be lower in the octogenarian group ( $p=0.07$ ) (Figure 2B-C). Post-exercise plasma volume, plasma sodium, hemoglobin and hematocrit concentration, and incidence of urine specific gravity levels  $\geq 1.020$  g/mL were not different between groups (all  $p>0.05$ , Table 2). Both groups demonstrated an exercise-induced increase in hemoglobin and hematocrit concentration ( $p<0.01$ ), whilst the exercise-induced changes in plasma sodium concentration were not different in both groups ( $p>0.05$ ). Both groups demonstrated a decrease in body mass after exercise. The relative decrease in body mass was significantly larger for the

octogenarians ( $p=0.04$ , Figure 2D), whilst no differences between groups were observed in dehydration status (Table 2).

## DISCUSSION

The purpose of our study was to compare thermoregulatory responses and fluid balance during prolonged, moderate-intensity exercise between sexagenarians and octogenarians. We found that octogenarians demonstrate a lower baseline  $T_{in}$  and a larger rise in  $T_{in}$  during prolonged walking exercise compared to sexagenarians. As workload and exercise intensity were similar between groups, these results may suggest that the thermoregulatory control declines with advanced age. Moreover, octogenarians reported a lower fluid intake and urine output in combination with a larger loss of body mass compared to sexagenarians, whilst other fluid balance parameters were not different between groups. Thus, in moderate ambient conditions and with *ad libitum* fluid intake the fluid balance regulation of both groups was adequate to avoid development of sodium disturbances or severe dehydration.

In accordance with literature, baseline  $T_{in}$  was  $0.3^{\circ}\text{C}$  lower in octogenarians compared to sexagenarians (Howell 1948). The most likely explanations for this observation is the smaller muscle mass and, concomitant, lower metabolic heat production in the oldest group (Blatteis 2012; Howell 1948). Furthermore, higher age was associated with a significantly higher exercise-induced increase in  $T_{in}$ . Since exercise characteristics were similar between groups (i.e. duration and intensity), we propose 3 possible explanations for the higher increase in  $T_{in}$  in the octogenarians. First, the sweat rate tended to be lower in aged subjects due to a decrease in individual sweat gland output as well as a reduced number of activated sweat glands by heat (Blatteis 2012;

Shibasaki and Crandall 2010). Presumably, when aging progresses the sweat rate response is further impaired, which may result in a larger increment of  $T_{in}$  during prolonged exercise. Second, the initiation of the thermoregulatory responses to heat exposure occurs by altered signaling from thermosensors located both in the periphery and the core of the body (Blatteis 2012). With advanced age, there is a higher peripheral fiber loss and a lower conduction velocity of the nervous system, which may delay vascular responses to heat. This results in a decreased peripheral thermal sensation, a delayed response to heat exposure (Blatteis 2012; Guergova and Dufour 2011), and subsequently a larger exercise-induced increase in  $T_{in}$ . A third possible mechanism relates to the attenuated cutaneous vasodilator response to heat in aged people (Holowatz et al. 2010; Holowatz et al. 2007). Both, a delay in the initiation of the vasodilatory as well as a decreased vasodilatory capacity of the skin are known to limit heat dissipation (Holowatz et al. 2010; Holowatz et al. 2007). The above mentioned factors limit the human skin to lose heat, thereby causing  $T_{in}$  to rise and potentially increase the risk to develop heat related illnesses. In parallel, a trend ( $p=0.09$ ) for differences in maximum  $T_{in}$  between octogenarians and sexagenarians was observed. Whilst the present study was performed in moderate ambient conditions and maximum  $T_{in}$  of both groups was within the normal physiological range, differences in maximum  $T_{in}$  may be more pronounced under hot and humid ambient conditions. In summary, our data may suggest a progressive age-related impairment of thermoregulatory responses, with larger exercise-induced  $T_{in}$  increases in octogenarians compared to sexagenarians.

Octogenarians demonstrated a larger relative body mass loss compared to sexagenarians, suggesting a deficiency in fluid intake and/or a higher fluid loss. Indeed, the self-reported fluid balance diaries demonstrated that octogenarians consumed 74 mL/h less fluid compared to

sexagenarians. This observation is in line with Phillips *et al.* (Phillips *et al.* 1984) who demonstrated that older subjects (67-75 years) reported a lower thirst sensitivity compared to young controls (20-31 years) after a 24-h period of water deprivation. The blunted thirst response in older humans was later confirmed by others, who compared young and old subjects (Kenney and Chiu 2001; Menten 2006). Next to the differences in fluid intake, the octogenarians also reported a lower urine output. It may be assumed that there is a causal relationship between the lower fluid intake and the lower urine output. The lower total body water content may enhance water reabsorption in the kidneys and hence reduce the urine production. Despite the lower fluid intake and urine output, no differences were observed between groups for changes in body mass, blood sodium concentration and plasma volume. These findings suggest that octogenarians and sexagenarians are both capable to regulate their fluid balance appropriately during prolonged walking exercise.

*Clinical relevance.* Health problems like heat stroke, dehydration and hyponatremia are frequently reported during prolonged exercise. Although octogenarians reported a larger  $T_{in}$  increase and lower fluid intake compared to sexagenarians, medical problems did not occur in both groups. Risk assessment must be revised when these vulnerable groups perform exercise in hot and humid ambient conditions. Particularly since recent fluid intake guidelines advice athletes to drink according their thirst sensation to avoid dehydration (Dion *et al.* 2013; Goulet 2011) and previous studies noted a decreased thirst sensitivity in elderly (Kenney and Chiu 2001; Mack *et al.* 1994; Phillips *et al.* 1984). For that reason, the “drink according to thirst” regimen may be inappropriate in the older individuals under severe ambient conditions. Therefore, it is

recommended that future fluid replacement guidelines should include specific information on rehydration in elderly.

*Limitations.* This study includes two unique groups of advanced aged subjects that performed prolonged walking exercise. However, one might argue that these individuals do not represent the general advanced age population, since these subjects have a highly active lifestyle. Consequently, there may be selection bias in the current study. Fact is that medical advances stimulate healthy aging and subsequent longevity in modern society. As a direct consequence, physical activity is increasingly popular in this population, which is reinforced by the growing number of older humans that participate in endurance exercise events. Therefore, insight into the physiological responses of these individuals is necessary for a safe exercise environment with appropriate recommendations. Secondly, sexagenarians reported a significantly higher BMI and lean body mass compared to octogenarians. As these characteristics increase the risk for fluid disturbances and the development of dehydration in sexagenarians (Eijsvogels et al. 2011), we may have underestimated the differences in thermoregulatory and fluid balance responses between both groups. A third limitation of this study is the used estimation of the maximum heart rate to calculate the exercise intensity. Since we used the formula by Tanaka et al. (Tanaka et al. 2001), a widely used method to calculate the predicted maximum heart rate, we believe that the consecutively calculated exercise intensities properly reflect the real exercise intensities. Finally, clothing was not reported during the study, while it plays an important role in the capacity to dissipate heat. However, the majority of the subjects wore normal walking clothes (t-shirt+shorts), and therefore, we strongly believe that clothing did not influence our results.

In conclusion, octogenarians demonstrate an impaired thermoregulatory control compared to sexagenarians during prolonged moderate-intensity exercise under moderate ambient conditions. In contrast, fluid balance was well controlled in both groups and did not deteriorate with aging. This enabled sexagenarians as well as the octogenarians to successfully complete a 30 km walking march without heat or fluid related health problems.

### **Acknowledgements**

This work was supported by the Netherlands Organization for Scientific Research (Rubicon Grant 825.12.016, T.M.H.E) and the Dutch Heart Foundation (E-Dekker stipend 2009T064, D.H.J.T). None of the authors reported a conflict of interest. We recognize the excellent help of the organization of the Nijmegen Four Days Marches and the Diagnostic Division of Siemens, The Netherlands. Furthermore, we acknowledge the valuable assistance of Leonie Prins, Martijn Maessen, Nathalie Benda, Joost Seeger, Tim Schreuder, Bregina Hijmans-Kersten and Elvira van Vlijmen.

### **Conflict of Interest**

None.

### **REFERENCES**

- Adroge HJ, Madias NE (2000) Hyponatremia The New England journal of medicine 342:1493-1499 doi:10.1056/NEJM200005183422006
- Armstrong LE, Casa DJ, Millard-Stafford M, Moran DS, Pyne SW, Roberts WO (2007) American College of Sports Medicine position stand. Exertional heat illness during training and competition Medicine and science in sports and exercise 39:556-572 doi:10.1249/MSS.0b013e31802fa199
- Bar-Or O (1998) Effects of age and gender on sweating pattern during exercise International journal of sports medicine 19 Suppl 2:S106-107 doi:10.1055/s-2007-971970

- Begum MNJ, C.S. (2010) A review of the literature on dehydration in the institutionalized elderly The European e-Journal of Clinical Nutrition and Metabolism 5:47-53  
doi:<http://dx.doi.org/10.1016/j.eclnm.2009.10.007>
- Blatteis CM (2012) Age-dependent changes in temperature regulation - a mini review Gerontology 58:289-295 doi:10.1159/000333148
- Bouchama A, Knochel JP (2002) Heat stroke The New England journal of medicine 346:1978-1988  
doi:10.1056/NEJMra011089
- Byrne C, Lim CL (2007) The ingestible telemetric body core temperature sensor: a review of validity and exercise applications British journal of sports medicine 41:126-133  
doi:10.1136/bjism.2006.026344
- Casa DJ et al. (2000) National athletic trainers' association position statement: fluid replacement for athletes Journal of athletic training 35:212-224
- Chester JG, Rudolph JL (2011) Vital signs in older patients: age-related changes Journal of the American Medical Directors Association 12:337-343 doi:10.1016/j.jamda.2010.04.009
- Cheuvront SN, Kenefick RW, Montain SJ, Sawka MN (2010) Mechanisms of aerobic performance impairment with heat stress and dehydration Journal of applied physiology 109:1989-1995  
doi:10.1152/jappphysiol.00367.2010
- Criswell D, Renshler K, Powers SK, Tulley R, Cicale M, Wheeler K (1992) Fluid replacement beverages and maintenance of plasma volume during exercise: role of aldosterone and vasopressin European journal of applied physiology and occupational physiology 65:445-451
- Davis KM, KL (1994) Disorders of fluid balance: Dehydration and hyponatremia. In: Hazard WB, El; Blass, JP; Ettinger, WH; Halter, JB (ed) Principles of geriatric medicine and gerontology. 3 edn. McGraw Hill, New York, pp 1182-1190
- Dill DB, Costill DL (1974) Calculation of percentage changes in volumes of blood, plasma, and red cells in dehydration J Appl Physiol 37:247-248
- Dion T, Savoie FA, Asselin A, Garipey C, Goulet ED (2013) Half-marathon running performance is not improved by a rate of fluid intake above that dictated by thirst sensation in trained distance runners European journal of applied physiology 113:3011-3020 doi:10.1007/s00421-013-2730-8
- Durnin JV, Womersley J (1974) Body fat assessed from total body density and its estimation from skinfold thickness: measurements on 481 men and women aged from 16 to 72 years The British journal of nutrition 32:77-97
- Eijvogels TM, Veltmeijer MT, Schreuder TH, Poelkens F, Thijssen DH, Hopman MT (2011) The impact of obesity on physiological responses during prolonged exercise International journal of obesity 35:1404-1412 doi:10.1038/ijo.2010.277
- Gant N, Atkinson G, Williams C (2006) The validity and reliability of intestinal temperature during intermittent running Medicine and science in sports and exercise 38:1926-1931  
doi:10.1249/01.mss.0000233800.69776.ef
- Goulet ED (2011) Effect of exercise-induced dehydration on time-trial exercise performance: a meta-analysis British journal of sports medicine 45:1149-1156 doi:10.1136/bjism.2010.077966
- Guergova S, Dufour A (2011) Thermal sensitivity in the elderly: a review Ageing research reviews 10:80-92 doi:10.1016/j.arr.2010.04.009
- Hew-Butler T et al. (2005) Consensus statement of the 1st International Exercise-Associated Hyponatremia Consensus Development Conference, Cape Town, South Africa 2005 Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine 15:208-213
- Holowatz LA, Thompson-Torgerson C, Kenney WL (2010) Aging and the control of human skin blood flow Front Biosci (Landmark Ed) 15:718-739



Holowatz LA, Thompson-Torgerson CS, Kenney WL (2007) Altered mechanisms of vasodilation in aged human skin *Exercise and sport sciences reviews* 35:119-125 doi:10.1097/jes.0b013e3180a02f85

Howell TH (1948) Normal temperatures in old age *Lancet* 1:517

Kenefick RW, Cheuvront SN, Sawka MN (2007) Thermoregulatory function during the marathon *Sports medicine* 37:312-315

Kenney WL, Chiu P (2001) Influence of age on thirst and fluid intake *Medicine and science in sports and exercise* 33:1524-1532

Lindeman RD, Tobin J, Shock NW (1985) Longitudinal studies on the rate of decline in renal function with age *Journal of the American Geriatrics Society* 33:278-285

Mack GW, Weseman CA, Langhans GW, Scherzer H, Gillen CM, Nadel ER (1994) Body fluid balance in dehydrated healthy older men: thirst and renal osmoregulation *Journal of applied physiology* 76:1615-1623

Mentes J (2006) Oral hydration in older adults: greater awareness is needed in preventing, recognizing, and treating dehydration *The American journal of nursing* 106:40-49; quiz 50

Miller JH, Shock NW (1953) Age differences in the renal tubular response to antidiuretic hormone *Journal of gerontology* 8:446-450

Phillips PA, Rolls BJ, Ledingham JG, Forsling ML, Morton JJ, Crowe MJ, Wollner L (1984) Reduced thirst after water deprivation in healthy elderly men *The New England journal of medicine* 311:753-759 doi:10.1056/NEJM198409203111202

Sawka MN (1992) Physiological consequences of hypohydration: exercise performance and thermoregulation *Medicine and science in sports and exercise* 24:657-670

Sawka MN, Burke LM, Eichner ER, Maughan RJ, Montain SJ, Stachenfeld NS (2007) American College of Sports Medicine position stand. Exercise and fluid replacement *Medicine and science in sports and exercise* 39:377-390

Sawka MN, Noakes TD (2007) Does dehydration impair exercise performance? *Medicine and science in sports and exercise* 39:1209-1217 doi:DOI 10.1249/mss.0b013e318124a664

Schoeller DA (1989) Changes in total body water with age *The American journal of clinical nutrition* 50:1176-1181; discussion 1231-1175

Sheehy CM, Perry PA, Cromwell SL (1999) Dehydration: biological considerations, age-related changes, and risk factors in older adults *Biological research for nursing* 1:30-37

Shibasaki M, Crandall CG (2010) Mechanisms and controllers of eccrine sweating in humans *Frontiers in bioscience* 2:685-696

Tanaka H, Monahan KD, Seals DR (2001) Age-predicted maximal heart rate revisited *Journal of the American College of Cardiology* 37:153-156

Tatterson AJ, Hahn AG, Martin DT, Febbraio MA (2000) Effects of heat stress on physiological responses and exercise performance in elite cyclists *Journal of science and medicine in sport / Sports Medicine Australia* 3:186-193

Wilkinson DM, Carter JM, Richmond VL, Blacker SD, Rayson MP (2008) The effect of cool water ingestion on gastrointestinal pill temperature *Medicine and science in sports and exercise* 40:523-528 doi:10.1249/MSS.0b013e31815cc43e

## FIGURE LEGENDS

**Fig. 1** Time course of a) exercise intensity (% of HRmax) and b) intestinal temperature (°C) in sexagenarians (solid line) and octogenarians (dashed line). While, octogenarians demonstrated a larger increase in intestinal temperature compared to sexagenarians ( $p < 0.01$ ), no differences were observed in the change in exercise intensity between the groups ( $p = 0.88$ ). Data is presented as mean  $\pm$  SEM

**Fig. 2** Fluid balance parameters (a= Fluid intake, b= urine output, c= sweat rate and d= body mass loss) presented for both sexagenarians (black bars) and octogenarians (white bars). Sexagenarians demonstrated a higher fluid intake and sweat rate and subsequently a lower body mass loss. Data were presented as mean  $\pm$  SEM. \* Represents a significant difference between both groups.

## TABLE LEGENDS

**Table 1.** Subject characteristics

**Table 2.** Exercise characteristics and fluid balance presented per age group