




N-Benzyl-5-methoxytryptamines as Potent Serotonin 5-HT₂ Receptor Family Agonists and Comparison with a Series of Phenethylamine Analogues

David E. Nichols,^{*,†} M. Flori Sassano,[†] Adam L. Halberstadt,[‡] Landon M. Klein,[§] Simon D. Brandt,^{||} Simon P. Elliott,[⊥] and Wolfgang J. Fiedler[#][†]Division of Chemical Biology and Medicinal Chemistry, University of North Carolina, Chapel Hill, North Carolina 27599, United States[‡]Department of Psychiatry, and [§]Department of Neurosciences, University of California San Diego, La Jolla, California 92093, United States^{||}School of Pharmacy and Biomolecular Sciences, Liverpool John Moores University, Byrom Street, Liverpool L3 3AF, U.K.[⊥]ROAR Forensics, Malvern Hills Science Park, Geraldine Road, Malvern, Worcestershire WR14 3SZ, U.K.[#]Hurststrasse 6a, D-69514 Laudenbach, Germany

ABSTRACT: A series of *N*-benzylated-5-methoxytryptamine analogues was prepared and investigated, with special emphasis on substituents in the meta position of the benzyl group. A parallel series of several *N*-benzylated analogues of 2,5-dimethoxy-4-iodophenethylamine (2C-I) also was included for comparison of the two major templates (i.e., tryptamine and phenethylamine). A broad affinity screen at serotonin receptors showed that most of the compounds had the highest affinity at the 5-HT₂ family receptors. Substitution at the para position of the benzyl group resulted in reduced affinity, whereas substitution in either the ortho or the meta position enhanced affinity. In general, introduction of a large lipophilic group improved affinity, whereas functional activity often followed the opposite trend. Tests of the compounds for functional activity utilized intracellular Ca²⁺ mobilization. Function was measured at the human 5-HT_{2A}, 5-HT_{2B}, and 5-HT_{2C} receptors, as well as at the rat 5-HT_{2A} and 5-HT_{2C} receptors. There was no general correlation between affinity and function. Several of the tryptamine congeners were very potent functionally (EC₅₀ values from 7.6 to 63 nM), but most were partial agonists. Tests in the mouse head twitch assay revealed that many of the compounds induced the head twitch and that there was a significant correlation between this behavior and functional potency at the rat 5-HT_{2A} receptor.

KEYWORDS: Serotonin, 5-HT₂ receptors, 5-HT_{2A} agonist, phenethylamine, 5-methoxytryptamine, mouse head twitch



INTRODUCTION

Recently, an extremely potent hallucinogenic phenethylamine, 25I-NBOMe (*N*-(2-methoxybenzyl)-2,5-dimethoxy-4-iodophenethylamine; “smiles”) **1** has been available on the illicit drug market.¹ For purposes of enforcement, it is presently considered by the Drug Enforcement Administration (DEA) to be an analogue of 2C-I (**2**), which is currently a Schedule I controlled substance. The procedure to classify **1** as a Schedule I substance has been initiated, and it has been placed temporarily into Schedule I.² Unfortunately, several deaths have been associated with the use of **1**,^{3–5} but it is not clear whether the deaths resulted from the ingestion of lethal amounts of pure solid drug, or whether the drug has some inherent toxicity that is not normally associated with other hallucinogens.


There has been increasing global interest in **1** and closely related analogues. For example, the European Monitoring

Centre for Drugs and Drug Addiction (EMCDDA) has received a range of notifications from EU Member States about analytically confirmed nonfatal and fatal intoxications associated with **1**. This was then followed by a risk assessment conducted by the Scientific Committee of the EMCDDA in order to assess health and social risks associated with this particular analogue.⁶ In addition, the World Health Organization’s Expert Committee on Drug Dependence reviewed the status of a range of new substances for its 36th meeting in June 2014, which included **1** and its 4-bromo and 4-chloro analogues.⁷ In September 2014, the Council of the European Union decided to subject **1** to control measures and criminal penalties throughout the European Union.⁸

Special Issue: Serotonin Research


Received: November 10, 2014

Published: December 29, 2014



Typically, simple *N*-alkylation dramatically attenuates or abolishes hallucinogenic activity in phenethylamines.^{9,10} The *N*-benzyl moiety, however, confers exceptionally high potency onto the molecule,^{11–15} and we have presented evidence that the *N*-benzyl may engage F339 in the human 5-HT_{2A} receptor.¹⁴ We also examined various *N*-arylmethyl substituents and found that a variety of aryl groups were effective in enhancing potency.^{16,17} In addition, the presence of a polar substituent at the ortho position of the aryl ring (a possible hydrogen bond acceptor) further enhances activity.¹⁸ Silva et al.¹⁸ also have reported that in an *in vitro* cylindrical rat tail artery strip **1** had a pEC₅₀ of 10.09 and an *E*_{max} of 30%.

Two decades ago, Glennon et al.¹⁹ reported that the affinities of the *N*-benzyl compound **3a**, as well as the 4-bromo- and 4-iodo-*N*-benzyl compounds, **3b** and **3c**, respectively, were 2–3 times higher than that of the parent primary amine. There have been no further reports on these compounds, and in our own work, we had never examined 3- or 4-substituted benzyl substituents in the phenethylamine series.




3a X = H, **3b** X = 4-Br, **3c** X = 4-I

In addition to the phenethylamine type 5-HT_{2A} agonists, certain simple tryptamines possess similar pharmacology, particularly 4- or 5-oxygenated molecules. In the report by Glennon et al., placing an *N*-benzyl moiety on the amine of 5-methoxytryptamine had essentially no effect on affinity. Interestingly, *N*-benzyl-5-methoxytryptamine previously had been reported to be an antagonist of serotonin-induced contraction in the rat stomach fundus, the isolated guinea pig uterus, and the isolated guinea pig taenia cecum.²⁰ In addition, Leff et al.²¹ had shown that *N*-benzyl-5-methoxytryptamine had only weak partial agonist activity at 5-HT₂ type receptors in rabbit aorta and rat jugular vein.


Surprisingly, however, in the Glennon report,¹⁹ a 5-HT_{2A} receptor affinity of 0.1 nM was reported for the *N*-4-bromobenzyl compound (compound **33** in the Glennon report, numbered here as **5f**), with 1000-fold selectivity for 5-HT_{2A} over 5-HT_{2C} receptors. We found these data particularly intriguing. This degree of selectivity was overestimated, however, because affinity at the 5-HT_{2A} receptor was measured by displacement of an agonist radioligand, whereas affinity at the 5-HT_{2C} receptor was measured by displacement of an antagonist radioligand. Nonetheless, no specific 5-HT_{2A}-selective agonist has been available, although such a compound would be very valuable for serotonin neuroscience research.

Although it was reported¹⁹ that 4-bromo compound **5f** had 0.1 nM affinity at the human 5-HT_{2A} receptor, the 4-fluoro-, 4-chloro-, and 4-iodo-substituted benzyl congeners had reported affinities of 40, 105, and 120 nM, respectively, in that same report. We found this discontinuity in the structure–activity

relationship (SAR) puzzling, where the 4-bromo compound would be such an outlier in the family of halogen-substituted benzyls. Further investigation by Jensen, however, revealed that the authentic 4-bromo compound **5f** actually had relatively low affinity for the 5-HT_{2A} receptor, more consistent with the reported affinities of the other halogenated compounds.²² Although spectroscopic data were not reported by Glennon et al.¹⁹ that might explain the basis for this discrepancy, their publication indicated elemental analysis data to be consistent with the proposed structure. If the elemental analysis data were correct, the mostly likely explanation for the discordant biological data therefore seemed to be that **5f** might have been an isomer other than the 4-substituted compound.



1, X = 2-OCH₃
4a, X = 3-OCH₃
4b, X = 4-OCH₃
4c, X = 2-Br
4d, X = 3-Br
4e, X = 4-Br



5a, X = 2-OCH₃
5b, X = 3-OCH₃
5c, X = 4-OCH₃
5d, X = 2-Br
5e, X = 3-Br
5f, X = 4-Br
5g, X = 3-F
5h, X = 3-Cl
5i, X = 3-I
5j, X = 3-CH₃
5k, X = 3-CF₃
5l, X = 3-SCH₃

On the basis of the hypothesis that the original data were associated with an isomer other than the 4-bromo compound, we subsequently discovered that *N*-3-bromobenzyl compound **5e** did have higher affinity for the 5-HT_{2A} receptor (*K*_i 1.48 nM), compared to that of the 4-bromo congener **5f** (*K*_i 11.2 nM). Further, the effect of an ortho-oxygenated *N*-benzyl appeared not to be significant for affinity in the tryptamine series, suggesting perhaps different binding orientations of the *N*-benzyltryptamines versus the *N*-benzylphenethylamines within the receptor. That is, compound **5a** has been reported to have agonist potency (pEC₅₀ 7.08) in a rat tail artery assay not significantly different from the compound with an unsubstituted *N*-benzyl moiety (pEC₅₀ 7.00), although the *E*_{max} was slightly higher for the 2'-methoxy compound.¹⁸ These findings prompted us to synthesize a small series of structurally related congeners to determine whether other substitutions might have even greater affinity and/or selectivity for the 5-HT_{2A} receptor.

Thus, in this article we describe the facile synthesis of compounds **1**, **4a–4e**, and **5a–5l**, preliminary screening at a variety of 5HT family receptors, and more detailed testing at human 5-HT_{2A}, 5-HT_{2B}, and 5-HT_{2C} receptors, including affinity measurements using displacement of the agonist radioligand [¹²⁵I]-DOI and functional effects in elevating intracellular calcium. We also present behavioral data for the mouse head twitch response (HTR) as a measure of *in vivo* 5-HT_{2A} receptor activation.²³

Compound **1** has been previously reported,²⁴ and the NMR and electron ionization mass spectra of **4a** and **4b** have been reported but without any biological data.²⁵ We thus decided to compare all of the series members at the same time to elucidate a consistent SAR.

CHEMISTRY

All of the compounds were most easily prepared using a modification of the facile method first reported by Abdel-Magid et al.²⁶ The free base of **2** was stirred in 3 mL of MeOH for 30 min with the appropriate aldehyde, followed by reduction of the intermediate enamine with NaBH₄. Following appropriate workup, the bases were converted to their HCl or maleate salts and crystallized in good to excellent yields.

PHARMACOLOGY

Affinities at a panel of 5-HT receptors were determined by the NIMH-sponsored PDSP program (<http://pdsp.med.unc.edu/kidb.php>). Affinities at both the human and rat 5-HT_{2A} and 5-HT_{2C} receptors also were determined, using both agonist and antagonist radioligands. As a measure of functional potency and efficacy, changes in intracellular Ca²⁺ levels were measured using a fluorometric imaging plate reader (FLIPR^{TETRA}, Molecular Devices), at the human 5-HT_{2A}, 5-HT_{2B}, and 5-HT_{2C} receptors, and at the rat 5-HT_{2A} and 5-HT_{2C} receptors. Finally, as a measure of in vivo 5-HT_{2A} receptor activation, we assessed the ability of all compounds to induce the mouse HTR.²³ We hypothesized that functional potency at the rat 5-HT_{2A} receptor might correlate best with the mouse head twitch behavioral data because ligand affinities at the rat 5-HT_{2A} receptor correlate with the mouse 5-HT_{2A} receptor but not with the human 5-HT_{2A} receptor.²⁷

RESULTS

Further exploration of a small library of 3-substituted *N*-benzyl tryptamines allowed us to develop a tentative SAR for this series, and it is clear that substituents on the *N*-benzyl 3-position do modulate affinity in the tryptamine series. In the broad screening of 5-HT receptor types, all of the compounds had the highest affinity at the 5-HT₂ family of receptors (Tables 1 and 2).

At the 5-HT_{2A} and 5-HT_{2C} receptors, the highest affinity was observed in the competition displacements with [¹²⁵I]-DOI. Except for **5c** and **5f**, all of the tryptamine compounds had low nanomolar or subnanomolar affinity for the human 5-HT_{2A} receptor. The known phenethylamine **1** had by far the highest affinity at 5-HT_{2A/2C} receptors, with subnanomolar affinity at both subtypes. We have previously reported an affinity for **1** at the human 5-HT_{2A} receptor of 0.04 nM.¹⁴ Of the tryptamines, only the 3-iodobenzyl compound **5i**, had subnanomolar affinity at the 5-HT_{2A} receptor, although all of the tryptamines had high affinity at this receptor. It should be noted that *N*-methylation of **5e** completely abolished affinity at the 5-HT_{2A} receptor (*K_i* > 10 μM; data not shown), indicating that tertiary amines are not tolerated in the *N*-benzyltryptamines.

The rank order of affinity of all compounds at the [¹²⁵I]-DOI-labeled h5-HT_{2C} receptor generally paralleled that measured at the 5-HT_{2A} receptor, although the affinities tended to be somewhat lower. Again, among the tryptamines studied **5i** had the highest affinity at this receptor, as well as at the 5-HT_{2B} receptor. Affinities measured at the [¹²⁵I]-DOI site tended to be on the order of 5–10 times higher than that at the antagonist labeled sites at both receptors.

Functional potencies at the rat and human 5-HT_{2A} and 5-HT_{2C} receptors and the human 5-HT_{2B} receptor are shown in Table 3. Compound **1** was a nearly full agonist at both receptor types, with a 4.2 nM EC₅₀ at the human 5-HT_{2A} receptor and 11 nM EC₅₀ at the rat 5-HT_{2A} receptor. The most potent

Table 1. Affinities of New Compounds for the Human 5-HT_{2A} and 5-HT_{2C} Receptors Using Both Agonist and Antagonist Radioligands^a

compd	h5-HT _{2A} p <i>K_i</i> ± SEM (<i>K_i</i> nM)		h5-HT _{2C} p <i>K_i</i> ± SEM (<i>K_i</i> nM)	
	[³ H]ketanserin	[¹²⁵ I]DOI	[³ H]mesulergine	[¹²⁵ I]DOI
1	9.28 ± 0.11 (0.52)	9.80 ± 0.15 (0.16)	9.16 ± 0.09 (0.69)	9.30 ± 0.16 (0.50)
4a	8.81 ± 0.17 (1.5)	9.57 ± 0.09 (0.27)	8.38 ± 0.01 (4.17)	9.90 ± 0.07 (0.13)
4b	7.93 ± 0.13 (11.7)	9.15 ± 0.16 (0.70)	7.85 ± 0.02 (14.1)	8.44 ± 0.14 (3.63)
4c	8.63 ± 0.18 (2.34)	9.42 ± 0.09 (0.38)	8.06 ± 0.07 (8.71)	8.99 ± 0.18 (1.02)
4d	8.40 ± 0.04 (3.98)	9.24 ± 0.12 (0.57)	8.12 ± 0.02 (7.59)	8.79 ± 0.08 (1.62)
4e	7.28 ± 0.14 (52.5)	8.49 ± 0.09 (3.24)	7.34 ± 0.02 (45.7)	8.48 ± 0.25 (3.31)
5a	7.78 ± 0.05 (16.6)	8.82 ± 0.19 (1.51)	7.49 ± 0.14 (32.4)	8.47 ± 0.10 (3.39)
5b	8.11 ± 0.10 (7.76)	8.98 ± 0.14 (1.05)	7.42 ± 0.12 (38.0)	8.23 ± 0.09 (5.89)
5c	7.16 ± 0.16 (69.2)	7.98 ± 0.04 (10.5)	6.90 ± 0.03 (126)	7.85 ± 0.13 (14.1)
5d	7.60 ± 0.12 (25.1)	8.63 ± 0.19 (2.34)	7.00 ± 0.01 (100)	7.85 ± 0.10 (14.1)
5e	8.17 ± 0.11 (6.76)	8.83 ± 0.10 (1.48)	7.58 ± 0.05 (26.3)	8.25 ± 0.11 (5.62)
5f	6.37 ± 0.12 (427)	7.95 ± 0.22 (11.2)	6.60 ± 0.15 (251)	7.54 ± 0.19 (28.8)
5g	7.67 ± 0.04 (21.4)	8.58 ± 0.17 (2.63)	7.32 ± 0.09 (47.9)	8.06 ± 0.14 (8.71)
5h	8.28 ± 0.08 (5.25)	8.98 ± 0.10 (1.05)	7.55 ± 0.06 (28.2)	8.37 ± 0.05 (4.27)
5i	8.46 ± 0.09 (3.47)	9.21 ± 0.16 (0.62)	8.19 ± 0.09 (6.46)	8.98 ± 0.08 (1.05)
5j	8.32 ± 0.17 (4.79)	8.93 ± 0.11 (1.17)	7.65 ± 0.03 (22.4)	8.47 ± 0.08 (3.39)
5k	7.55 ± 0.05 (28.2)	8.53 ± 0.19 (2.95)	6.99 ± 0.06 (102)	7.83 ± 0.26 (14.8)
5l	8.05 ± 0.15 (8.91)	8.51 ± 0.17 (3.09)	7.88 ± 0.23 (13.2)	8.68 ± 0.30 (2.09)

^ap*K_i* ± SEM (affinities in nM); *n* = 3–5 separate displacement curves.

compound was **5a**, with an EC₅₀ of 1.9 nM and 85% efficacy at the h5-HT_{2A}. Notably, this compound has the *N*-2-methoxybenzyl substituent, the same as the most potent phenethylamine **1**, suggesting that it may be optimal for activation of the 5-HT_{2A} receptor when placed at the 2-position of the *N*-benzyl moiety. Efficacies of the tryptamines at the rat and human 5-HT_{2A} receptors and human 5-HT_{2C} receptor varied from about 40% to 80%, with a few compounds that were full agonists (e.g., **5a** and **5c**), whereas at the rat 5-HT_{2C} receptor all of the compounds were full agonists.

It is noteworthy that the functional potencies in the rat and human 5-HT_{2A} receptors are essentially identical for phenethylamine compounds **1**, and **4a–4e**, yet the potencies for tryptamine compounds **5a–5l** are 4–10-fold higher at the human 5-HT_{2A} receptor than at the rat 5-HT_{2A} receptor. This finding may reflect the single amino acid difference in the orthosteric binding site of these two receptors at position 5.46. In the rat or mouse 5-HT_{2A} receptor, residue 5.46 is an alanine, whereas in the human receptor it is a serine. We have previously shown that mutation of this residue in the human receptor from serine to alanine has little effect on affinity or function for phenethylamine 5-HT_{2A} agonists but does have a significant effect for tryptamines.²⁸ One might infer, therefore, from these potency differences that the indole NH in the

Table 2. PDSP Screening Affinities for All Compounds at Other Human Serotonin Receptor Types^a

compd	S-HT _{2B}	S-HT _{1A}	S-HT _{1B}	S-HT _{1D}	S-ht1e	S-HT ₃	S-ht5a	S-HT ₆	S-HT ₇
1	8.86 ± 0.03 (1.4)	5.99 ± 0.05 (1033)	5.23 ± 0.06 (5886)	6.27 ± 0.05 (533)	>10,000	>10,000	5.55 ± 0.07 (2795)	7.5 ± 0.06 (32)	5.81 ± 0.06 (1542)
4a	8.34 ± 0.03 (4.6)	6.03 ± 0.05 (925)	5.49 ± 0.05 (3232)	6.36 ± 0.05 (439)	5.77 ± 0.05 (1707)	>10,000	7.24 ± 0.06 (57)	7.17 ± 0.06 (67)	6.23 ± 0.06 (583)
4b	7.78 ± 0.03 (17)	5.97 ± 0.05 (1064)	5.8 ± 0.05 (1592)	6.49 ± 0.05 (325)	5.89 ± 0.05 (1285)	>10,000	5.99 ± 0.06 (1020)	7.12 ± 0.03 (75)	5.8 ± 0.06 (1575)
4c	7.7 ± 0.04 (20)	5.94 ± 0.06 (1155)	>10,000	6.37 ± 0.05 (423)	>10,000	>10,000	5.64 ± 0.09 (2290)	6.59 ± 0.06 (257)	5.59 ± 0.05 (2547)
4d	7.89 ± 0.04 (13)	6.17 ± 0.06 (670)	5.80 ± 0.05 (1568)	6.79 ± 0.05 (162)	6.10 ± 0.04 (792)	>10,000	6 ± 0.08 (1009)	6.76 ± 0.06 (175)	6.45 ± 0.05 (355)
4e	7.17 ± 0.04 (68)	6.19 ± 0.06 (649)	5.22 ± 0.05 (5093)	6.51 ± 0.05 (311)	>10,000	5.61 ± 0.05 (2460)	5.73 ± 0.06 (1848)	6.46 ± 0.05 (350)	6.19 ± 0.05 (641)
5a	8.04 ± 0.03 (9)	6.64 ± 0.05 (231)	>10,000	5.89 ± 0.05 (1292)	>10,000	>10,000	>10,000	7.06 ± 0.03 (87)	5.75 ± 0.06 (1770)
5b	8.6 ± 0.03 (2.5)	6.48 ± 0.05 (335)	>10,000	6.48 ± 0.06 (334)	>10,000	>10,000	5.9 ± 0.06 (1261)	7.6 ± 0.03 (25)	6.39 ± 0.05 (406)
5c	7.49 ± 0.03 (33)	7.12 ± 0.06 (76)	5.97 ± 0.04 (1060)	6.79 ± 0.06 (161)	>10,000	>10,000	5.62 ± 0.09 (2388)	6.45 ± 0.03 (353)	7.44 ± 0.05 (37)
5d	7.62 ± 0.03 (24)	6.54 ± 0.05 (286)	>10,000	6.11 ± 0.05 (782)	>10,000	5.21 ± 0.07 (6169)	>10,000	6.69 ± 0.05 (203)	5.96 ± 0.05 (1086)
5e	8.45 ± 0.03 (3.6)	6.81 ± 0.05 (155)	5.19 ± 0.06 (6433)	6.42 ± 0.05 (381)	>10,000	>10,000	6.21 ± 0.06 (612)	7.34 ± 0.03 (45)	6.93 ± 0.06 (116)
5f	6.83 ± 0.03 (150)	7.11 ± 0.05 (78)	5.35 ± 0.05 (4374)	6.57 ± 0.05 (271)	>10,000	>10,000	5.99 ± 0.08 (1034)	6.25 ± 0.03 (566)	6.45 ± 0.06 (358)
5g	7.66 ± 0.03 (22)	6.53 ± 0.04 (295)	5.57 ± 0.06 (2674)	6.50 ± 0.06 (319)	>10,000	>10,000	5.61 ± 0.05 (2450)	7.23 ± 0.03 (59)	6.62 ± 0.05 (242)
5h	8.16 ± 0.02 (6.6)	6.71 ± 0.05 (195)	5.36 ± 0.06 (4392)	6.55 ± 0.06 (282)	>10,000	>10,000	5.64 ± 0.06 (2310)	7.30 ± 0.03 (50)	6.55 ± 0.05 (281)
5i	9.12 ± 0.03 (0.76)	6.91 ± 0.05 (122)	5.53 ± 0.05 (2963)	6.70 ± 0.06 (199)	>10,000	>10,000	5.81 ± 0.05 (1536)	7.58 ± 0.03 (27)	7.66 ± 0.05 (22)
5j	8.71 ± 0.03 (1.9)	6.57 ± 0.04 (271)	5.37 ± 0.07 (4241)	6.55 ± 0.06 (283)	5.41 ± 0.05 (3876)	>10,000	5.41 ± 0.06 (3852)	7.21 ± 0.03 (62)	6.67 ± 0.05 (212)
5k	7.56 ± 0.02 (28)	6.62 ± 0.05 (240)	>10,000	6.56 ± 0.06 (278)	>10,000	>10,000	5.51 ± 0.06 (3091)	7.06 ± 0.03 (87)	6.58 ± 0.05 (262)
5l	8.39 ± 0.04 (4.1)	6.90 ± 0.05 (127)	>10,000	6.18 ± 0.05 (659)	>10,000	>10,000	6.08 ± 0.08 (841)	8.01 ± 0.06 (9.7)	6.87 ± 0.05 (136)

^apK_i ± SEM, (affinity in nM).

present series also engages this serine in the human receptor but not the alanine in the rat receptor, consistent with mutagenesis studies reported by others.^{29,30}

Figure 1 shows an illustrative dose–response curve for compound **5h** in the mouse HTR. HTR data for all compounds are given in Table 4. Although some of the compounds failed to induce the HTR at doses up to 30 mg/kg, most of the “inactive” compounds displayed relatively low potency at S-HT_{2A} (see Figure 2), so it is possible that they would induce the HTR if tested at higher doses. Importantly, for the subset of compounds that induced the HTR, behavioral potency was significantly correlated with functional potency at the r5-HT_{2A} receptor ($r = 0.69$, $p < 0.03$; Figure 2), but there was no correlation with functional EC₅₀ values at the r5-HT_{2C} receptor ($r = 0.17$, $p > 0.1$). Despite the overall correlation between mouse HTR and r5-HT_{2A} potency, the relationship was not always orderly for individual compounds. Compound **1** was by far the most potent compound in that assay, with an ED₅₀ of 0.078 mg/kg (data taken from Halberstadt and Geyer³¹). It is not clear why **1** should be so much more potent than any other compound because, for example, **4d** is inactive but appears nearly comparable functionally, with an EC₅₀ of 14 nM and efficacy of 69%, compared with an EC₅₀ of 11 nM for **1** with an efficacy of 79%. The next most potent compounds in the mouse HTR are **4c** and **5j**, with identical ED₅₀s of 2.31 mg/kg, about 300-fold less potent than **1**. Although they have similar functional EC₅₀ values (36 and 26 nM), nothing in the

functional or binding data can explain their lower potency compared to that of **1**. Further, compounds **5a**, **5b**, and **5g** have virtually identical ED₅₀ values in the mouse HTR, yet their functional EC₅₀s at the rat 5-HT_{2A} receptor are 21, 34, and 80 nM, respectively.

With the exception of **5k** and **5l**, which had relatively low functional potencies at the r5-HT_{2A} (EC₅₀ values of 770 and 120 nM, respectively), all of the meta-substituted *N*-benzyl derivatives of 5-methoxytryptamine induced the HTR. That included the 3-methyl (**5j**; ED₅₀ = 2.31 mg/kg), 3-methoxy (**5b**; ED₅₀ = 3.28 mg/kg), 3-fluoro (**5g**; ED₅₀ = 3.33 mg/kg), 3-chloro (**5h**; ED₅₀ = 4.43 mg/kg), 3-bromo (**5e**; ED₅₀ = 5.18 mg/kg), and 3-iodo (**5i**; ED₅₀ = 7.77 mg/kg) compounds.

The HTR produced by compounds **5b** and **5j** showed a biphasic bell-shaped dose–response function (the response peaked at 10 mg/kg and 30 mg/kg was inactive). Other 5-HT_{2A} agonists, including DOI, DOM, 2C-T-7, and 5-MeO-DIPT, have been shown to produce similar nonmonotonic responses.^{32–34} Fantegrossi et al.³⁴ have argued that the descending arm of the biphasic HTR dose–response is a consequence of 5-HT_{2C} activation, which attenuates the response to 5-HT_{2A} activation. Recently, however, it was reported that *N*-(2-hydroxybenzyl)-2,5-dimethoxy-4-cyanophenethylamine (25CN-NBOH), a 5-HT_{2A} agonist with 100-fold selectivity over 5-HT_{2C}, also induces the HTR with a biphasic dose–response.³⁵ The fact that the descending arm of the response to 25CN-NBOH was not affected by a 5-HT_{2C}

Table 3. Functional Data for New Compounds in Rat and Human 5-HT_{2A} and 5-HT_{2C} and Human 5-HT_{2B} Receptors^a

cmpd	r5-HT _{2A}		h5-HT _{2A}		h5-HT _{2B}		r5-HT _{2C}		h5HT _{2C}	
	pEC ₅₀ (EC ₅₀ nM)	E _{max} %	pEC ₅₀ (EC ₅₀ nM)	E _{max} %	pEC ₅₀ (EC ₅₀ nM)	E _{max} %	pEC ₅₀ (EC ₅₀ nM)	E _{max} %	pEC ₅₀ (EC ₅₀ nM)	E _{max} %
5-HT	8.3 ± 0.04 (5.4)	100 ± 1.5	8.7 ± 0.05 (2.0)	100 ± 1.6	9.31 ± 0.04 (0.49)	99.9 ± 1.1	9.70 ± 0.03 (0.20)	99.6 ± 0.73	9.52 ± 0.08 (0.30)	98.5 ± 2.4
1	8.0 ± 0.04 (11)	79.4 ± 1.1	8.4 ± 0.05 (4.2)	86.4 ± 1.4	7.81 ± 0.09 (15)	65 ± 2	7.02 ± 0.05 (95)	104 ± 2	7.38 ± 0.12 (41.7)	92 ± 0
4a	7.6 ± 0.04 (27)	51.7 ± 0.9	7.6 ± 0.03 (28)	71.6 ± 0.9	7.4 ± 0.3 (38)	NA ^b	6.88 ± 0.06 (133)	91 ± 3	7.47 ± 0.36 (33.8)	41 ± 6
4b	7.3 ± 0.04 (50)	53.8 ± 0.9	7.2 ± 0.03 (60)	74.1 ± 0.8	7.1 ± 0.1 (87)	38 ± 2	6.88 ± 0.05 (132)	97 ± 2	7.36 ± 0.31 (43.2)	50 ± 7
4c	7.4 ± 0.06 (36)	65.6 ± 1.6	7.4 ± 0.04 (42)	88.0 ± 1.5	6.82 ± 0.07 (134)	83 ± 3	6.98 ± 0.02 (105)	104 ± 1	7.24 ± 0.13 (57.6)	87 ± 5
4d	7.8 ± 0.03 (14)	68.5 ± 0.9	7.8 ± 0.04 (17)	87.5 ± 1.3	7.05 ± 0.05 (85)	90 ± 2	7.44 ± 0.05 (36)	101 ± 2	7.28 ± 0.17 (57.6)	74 ± 5
4e	6.8 ± 0.03 (150)	67.3 ± 0.9	6.8 ± 0.03 (170)	88.0 ± 1.4	6.21 ± 0.04 (610)	90 ± 2	6.54 ± 0.04 (290)	105 ± 2	6.66 ± 0.14 (200)	77 ± 5
5a	7.7 ± 0.03 (21)	80.9 ± 1.1	8.7 ± 0.05 (19)	85.2 ± 1.4	8.2 ± 0.1 (6.7)	52 ± 2	7.79 ± 0.04 (16)	102 ± 2	7.24 ± 0.12 (57.1)	119 ± 6
5b	7.5 ± 0.04 (34)	52.2 ± 0.9	8.2 ± 0.04 (6.2)	70.0 ± 1.0	6.0 ± 0.4 (949)	NA ^b	6.78 ± 0.05 (168)	102 ± 2	6.75 ± 0.15 (178)	65 ± 5
5c	6.7 ± 0.03 (190)	75.0 ± 1.3	7.4 ± 0.04 (42)	84.1 ± 1.3	7.64 ± 0.04 (23)	81 ± 1	7.73 ± 0.04 (19)	102 ± 2	7.12 ± 0.11 (75.1)	112 ± 5
5d	6.3 ± 0.04 (450)	49.7 ± 1.2	7.5 ± 0.05 (30)	74.7 ± 1.5	6.8 ± 0.3 (168)	NA ^b	6.05 ± 0.05 (898)	104 ± 3	6.36 ± 0.09 (439)	94 ± 5
5e	6.9 ± 0.03 (130)	65.5 ± 0.8	7.9 ± 0.04 (13)	73.8 ± 1.1	7.5 ± 0.2 (29)	20 ± 2	6.38 ± 0.04 (422)	112 ± 3	6.49 ± 0.23 (321)	64 ± 8
5f	5.8 ± 0.04 (1500)	77.6 ± 2.4	6.4 ± 0.02 (430)	90.3 ± 1.2	6.54 ± 0.05 (290)	90 ± 2	6.69 ± 0.03 (204)	108 ± 2	6.28 ± 0.14 (529)	83 ± 7
5g	7.1 ± 0.04 (80)	69.1 ± 1.3	8.0 ± 0.1 (10)	89.3 ± 1.1	7.42 ± 0.08 (38)	37 ± 1	7.34 ± 0.07 (46)	100 ± 3	6.72 ± 0.13 (192)	83 ± 5
5h	7.1 ± 0.03 (83)	70.1 ± 1.0	7.9 ± 0.04 (14)	81.2 ± 1.3	7.3 ± 0.2 (50)	NA ^b	6.54 ± 0.04 (286)	105 ± 2	6.50 ± 0.13 (316)	85 ± 6
5i	6.9 ± 0.04 (120)	73.4 ± 1.4	7.8 ± 0.04 (16)	79.0 ± 1.1	7.4 ± 0.2 (43)	31 ± 2	6.51 ± 0.05 (313)	110 ± 3	6.35 ± 0.09 (445)	94 ± 5
5j	7.6 ± 0.04 (26)	56.2 ± 0.9	8.2 ± 0.04 (6.5)	73.3 ± 1.0	NA ^b		6.72 ± 0.04 (192)	104 ± 2	6.54 ± 0.10 (289)	75 ± 4
5k	6.1 ± 0.03 (770)	69.6 ± 1.4	7.1 ± 0.04 (87)	75.5 ± 1.2	6.97 ± 0.07 (107)	51 ± 2	6.79 ± 0.03 (162)	104 ± 2	6.29 ± 0.11 (512)	75 ± 5
5l	6.9 ± 0.05 (120)	32.0 ± 0.7	7.5 ± 0.04 (32)	46.9 ± 0.8	NA ^b		6.69 ± 0.05 (205)	101 ± 2	6.55 ± 0.11 (283)	60 ± 4

^aValues are pEC₅₀ ± SEM, with (EC₅₀) values in nM and E_{max} given in percentage of the maximum response to 5-HT. ^bNA, not active; E_{max} ≤ 15%.




Figure 1. Representative dose–response plot in the mouse head twitch assay for compound 5h. **p* < 0.05 versus vehicle (Tukey's test).

antagonist³⁵ demonstrates that the inhibition of the HTR at high doses does not necessarily result from competing activity at 5-HT_{2C}. One potential alternative explanation for the biphasic HTR is that high levels of 5-HT_{2A} activation may produce competing behaviors that interfere with expression of head shaking. Along those lines, it has been reported that high doses of quipazine, 5-MeO-DMT, and (+)-LSD produce

stereotypic behaviors that preclude head shakes and wet dog shakes in rats.^{36,37}

DISCUSSION

Unfortunately, despite the report by Glennon et al.,¹⁹ compound 5e was not selective for the h5-HT_{2A} receptor versus the h5-HT_{2C} receptor. Using affinity at the [¹²⁵I]-DOI-labeled receptors, the selectivity of 5e was slightly less than 4-fold. Even using affinity at the [¹²⁵I]-DOI-labeled h5-HT_{2A} receptor and the [³H]-mesulergine-labeled h5-HT_{2C} receptor, “selectivity” was only about 18-fold. The most selective compound in the entire series, with respect to affinity, was 5d, but with only 6-fold selectivity.

With respect to selectivity in function at the h5-HT_{2A} vs h5-HT_{2C}, the most selective tryptamine was 5j, with 44-fold selectivity and less than a 3-fold difference in affinity at the agonist-labeled receptors. Indeed, we were disappointed that none of the compounds had high selectivity for the h5-HT_{2A} receptor.

Overall, with the exception of compound 1, none of the compounds was particularly potent in producing the HTR. This low potency is somewhat surprising, given that many known hallucinogens with high affinity for the 5-HT_{2A} receptor, such

Table 4. Activity of New Compounds in Producing the Mouse Head Twitch

	ED ₅₀ mg/kg (95% CI)	test duration (min)	N	dose range	active doses (mg/kg)	max counts	maximally effective dose (mg/kg)	magnitude of peak effect × vehicle
1	0.078 (0.055–0.111)	30	5	0.03–1.0	0.1, 0.3, 1	102.6	1	16.0
4a	4.34 (1.41–13.32)	10	10	0.3–30	3, 10, 30	11.4	30	5.7
4b	inactive		5	0.3–30				
4c	2.31 (1.41–3.77)	20	5	0.3–30	3, 10, 30	23.2	10	3.0
4d	inactive		5–7	0.3–10				
4e	inactive		6	1–30				
5a	3.15 (1.94–5.12)	20	10	0.3–30	10, 30	25.4	10	3.9
5b	3.28 (1.53–7.04)	10	5–6	1–30	10	9.2	10	3.7
5c	inactive		5	30				
5d	inactive		5	0.3–30				
5e	5.18 (2.35–11.38)	10	5–6	1–30	10, 30	14.2	30	4.4
5f	inactive		5	0.3–30				
5g	3.33 (2.25–4.93)	10	6	1–30	10, 30	14.5	10	7.3
5h	4.43 (2.03–9.69)	10	5–6	1–30	10, 30	10.6	30	8.0
5i	7.77 (3.40–17.53)	10	6	1–30	10, 30	20.2	30	3.4
5j	2.31 (0.82–6.51)	10	5	0.3–30	10	14.6	10	3.5
5k	inactive		5	30				
5l	inactive		4–5	30				




Figure 2. Plots of active and inactive compounds as a function of potency and efficacy at the rat 5-HT_{2A} receptor (panel A) and the human 5-HT_{2A} receptor (panel B).

as 2,5-dimethoxy-4-iodoamphetamine (DOI), *R*(-)-2,5-dimethoxy-4-methylamphetamine (*R*-DOM), *R*(-)-2,5-dimethoxy-4-bromoamphetamine (*R*-DOB), 2,5-dimethoxy-4-propylthiophenethylamine (2C-T-7), psilocin, and 5-MeO-*N,N*-diisopropyltryptamine (5-MeO-DIPT) produce the head twitch in mice at doses of ≤ 1 mg/kg.^{32,33,38–40} However, certain tryptamine hallucinogens, including 5-MeO-*N,N*-dimethyltryptamine (5-MeO-DMT) and α -methyltryptamine, are active within the same dose range (3–30 mg/kg) as the *N*-benzyltryptamines tested herein.^{40–42} It is unlikely that the low *in vivo* potencies of the compounds studied here are related to the use of an automated HTR detection system because we have confirmed that the results obtained using this system are consistent with published data based on visual scoring.²³ For example, the potency of LSD measured using the automated system (ED₅₀ = 0.13 μ mol/kg)²³ is almost exactly the same as the potency assessed using direct observation (ED₅₀ = 0.14 μ mol/kg).⁴¹ One possible explanation for the low potencies might be rapid first pass metabolism of *N*-benzyl-analogues in general⁴³ combined with a slow release from subcutaneous tissue due to the highly hydrophobic nature of the compounds.

Substitution on the *N*-benzyl ring has different effects, depending on whether the phenethylamines or the tryptamines are being studied. For example, *ortho*-bromo-substituted tryptamine congener **5d** failed to induce the HTR when tested

at doses up to 30 mg/kg (~ 60 μ mol/kg), yet *N*-3-bromobenzyl **5e** is active. By contrast, *N*-2-bromobenzyl phenethylamine **4c** is active, whereas *N*-3-bromobenzyl **4d** is inactive in the HTR assay.

None of the phenethylamines or tryptamines with 4-substituted *N*-benzyl groups, **4b**, **4e**, **5c**, or **5f**, was active in the HTR. All of these compounds were partial agonists with relatively low potency in the r5-HT_{2A} functional assay. Although **5e**, with a 3-substituted *N*-benzyl, has an EC₅₀ and *E*_{max} virtually identical to **4e**, it is active in the HTR assay. It is possible that differences in pharmacokinetics or metabolic lability could explain these data. Nevertheless, if only the compounds active in the mouse HTR assay are compared, one finds a significant correlation between potency in the rat 5-HT_{2A} receptor and potency in the HTR assay, as shown in Figure 3.

Taken together, these data show that for *N*-benzylphenethylamines the highest *in vivo* potency in mice is associated with an *ortho*-substituent on the benzyl group, whereas the *N*-benzyltryptamines are more active *in vivo* when a meta-




Figure 3. Regression analysis of pED₅₀ for the mouse head twitch response on the pEC₅₀ for function for active compounds at the rat 5-HT_{2A} receptor; *n* = 10.

substituent is present. Hence, there are SAR differences between the *N*-benzyltryptamines and the *N*-benzylphenethylamines for the induction of the HTR, which likely reflect different binding orientations in the 5-HT_{2A} receptor. Obviously, the indole system is larger than a simple phenyl ring, something that would clearly affect the binding modes for the two different series at the orthosteric site. For example, the distance from the indole C(3) atom to the 5-oxygen atom is 4.94 Å, whereas the corresponding distance from the 5-methoxy oxygen to C(1) of the aryl ring is only 3.70 Å. Even the distance of 4.85 Å from C(1) of the aryl ring to the 4-iodo atom of the phenethylamines is less than the 4.94 Å distance measured from C(3) of the indole to the 5-methoxy.

One exception is that for both the *N*-benzyltryptamines and *N*-benzylphenethylamines, oxygenated substituents are tolerated at the ortho- and meta-positions of the benzyl moiety. For example, **1**, **4a**, **5a**, and **5b** are all active in the HTR assay, whereas **4d** and **5d** are inactive over a range of doses. This observation again would be consistent with some structural feature in the 5-HT_{2A} receptor that could engage a polar oxygen atom at the ortho-position of the *N*-benzyl moiety. There has been speculation, based on virtual docking studies with phenethylamines and tryptamines, that an oxygen atom in the ortho-position of the *N*-benzyl moiety may interact with a hydrogen bond donor (possibly the OH of Tyr 370^(7,43) in the h5-HT_{2A} receptor.^{14,18} It is conceivable that an oxygen atom at the meta-position in *N*-benzyltryptamines also could form a hydrogen bond with Tyr 370, possibly involving a water molecule.

Unfortunately, a 5-HT_{2A} selective agonist did not emerge from this small library of compounds. There are now only two selective 5-HT_{2A} agonists reported,^{44,45} but they have not been available for extensive study. Thus, research on 5-HT_{2A} receptor function has been forced to employ either a mixed 5-HT_{2A/2C} agonist such as DOI in combination with a specific 5-HT_{2C} antagonist, or to administer antagonists alone, the latter paradigm really being appropriate to study receptor function only when there are high levels of endogenous receptor activation or constitutive activity of the receptors. Genetic knockout mice have not revealed particular behavioral phenotypes and have served primarily to demonstrate that a particular drug depends on the presence of 5-HT_{2A} or 5-HT_{2C} receptors for its effect. Hence, the psychopharmacology of a “pure” 5-HT_{2A} agonist remains completely unknown. Furthermore, the tremendous present interest in the role of the 5-HT_{2A} receptor in normal brain function makes it imperative that scientists in the field gain access to a 5-HT_{2A} specific agonist so that research into the roles of the 5-HT_{2A} receptor can be more fully elucidated.

METHODS

Chemistry. General Methods. Reagents were purchased from Sigma-Aldrich Co. (St. Louis, MO) or Alfa Aesar (Ward Hill, MA) and used as delivered, unless otherwise specified. Thin layer chromatography was carried out using J. T. Baker flexible sheets (silica gel IB2-F) with fluorescent indicator, visualizing with UV light at 254 nm or iodine stain. Melting points were determined using a Mel-Temp apparatus and are uncorrected. NMR experiments were carried out using a Bruker Advance 300 MHz instrument, and the chemical shift (δ) values are in parts per million (ppm) relative to tetramethylsilane at 0.00 ppm. The solvent was CD₃OD. NMR samples were dissolved in MeOD. Ph = aromatic protons/carbons of benzyl group; In = aromatic protons/carbons of the indole nucleus; Ar = either phenyl or indole resonances, or phenyl in the case of compounds **1**–**4f**.

Coupling constants (*J*) are presented in Hertz. Abbreviations used in the reporting of NMR spectra include: br = broad, s = singlet, d = doublet, t = triplet, q = quartet, and quint = quintuplet.

Mass spectra were performed by high resolution LC-QTOF-MS on protonated molecules [M + H]⁺. UHPLC-Q-TOF-MS conditions for UHPLC separation employed a mobile phase consisting of 100% MeCN that included 1% formic acid (organic phase) and an aqueous solution of 1% formic acid (aqueous phase). The column was maintained at 40 °C with a 0.6 mL/min flow rate and 5.5 min acquisition time. The elution was a 5–70% MeCN gradient ramp over 3.5 min, then up to 95% MeCN in 1 min and held for 0.5 min before returning to 5% MeCN in 0.5 min. Q-TOF-MS data were acquired in positive mode scanning from 100 to 1000 *m/z* with and without auto MS/MS fragmentation. Ionization was achieved with an Agilent JetStream electrospray source and infused internal reference masses. Agilent 6540 Q-TOF-MS parameters: gas temperature, 325 °C; drying gas, 10 L/min; and sheath gas temperature, 400 °C. Internal reference masses of 121.05087 and 922.00979 *m/z* were used.

For compounds **1** and **4a–4e**, 0.5 mmol of the free base of 4-iodo-2,5-dimethoxyphenethylamine^{10,46} was stirred for 30 min at room temperature with 0.55 mmol of the appropriate aldehyde in 3 mL of methanol. The reaction was then placed on an ice bath, and 48 mg (1.25 mmol) of NaBH₄ was added in three portions over 15 min. The ice bath was removed and the reaction allowed to stir for an additional 15 min. The reaction was then transferred to a separatory funnel with 50 mL of EtOAc. The organic phase was washed three times with saturated NaCl, then dried overnight over Na₂SO₄. The drying agent was removed by suction filtration, and the filtrate was concentrated under reduced pressure. EtOH (1 mL) was added to the amber residue, and the HCl salt was prepared by acidification with 0.5 mL of 1 N HCl/EtOH. Dilution with EtOAc or diethyl ether then led to crystallization of the HCl salts, generally in good yields. In most cases, the supernatant was simply decanted from the crystalline product, followed by resuspension of the crystals in Et₂O and decantation, then air drying to afford the products as white to off-white fine needles. No attempt was made to optimize the yields, but in one case the supernatant was reduced to dryness and the residue crystallized from EtOH/Et₂O to afford an additional 6% of product. This small additional recovery was not deemed sufficient to warrant the extra effort. Thus, all reported yields are those obtained after the first crystallization.

The synthesis of tryptamines **5a–5l** followed essentially the same procedure, except that maleate salts were prepared. As an example, 1.0 mmol of 5-methoxytryptamine free base (Aldrich) was stirred for 30 min with 1.10 mmol of the appropriate aldehyde in 5 mL of methanol. The reaction was then placed on an ice bath, and 96 mg (2.5 mmol) of NaBH₄ was added in three portions over 15 min. The ice bath was removed and the reaction allowed to stir for an additional 15 min. The reaction was then transferred to a separatory funnel with 50 mL of EtOAc and was washed three times with saturated NaCl. The organic phase was dried overnight over Na₂SO₄, then filtered and concentrated under reduced pressure. Maleic acid (116 mg, 1 mmol) and 1.0 mL of acetone were then added to the residual amber oil, and the solution swirled until all of the maleic acid had dissolved. The reaction was then diluted with 10 mL of EtOAc, and Et₂O was added nearly to the cloud point. In most cases, crystallization occurred rapidly and spontaneously, and the product solution was stored overnight in a cold room. Crystalline products were collected by suction filtration, washed on the filter with EtOAc, and then air-dried to afford white to off-white fine needles.

N-(2-Methoxybenzyl)-2-(4-iodo-2,5-dimethoxyphenyl)ethan-1-amine Hydrochloride (**1**). Obtained as needles following crystallization from acetone/EtOAc/Et₂O; yield 86%; mp 168–170 °C, Lit²⁴ mp 162–166 °C, 166.¹³ ¹H NMR (300 MHz, CD₃OD) δ ppm 7.46 (1H, td, *J* = 8.2, 1.7 Hz, Ar–H), 7.37 (1H, dd, *J* = 7.6, 1.6 Hz, Ar–H), 7.35 (1H, s, Ar–H), 7.09 (1H, d, *J* = 8.3 Hz, Ar–H), 7.02 (1H, td, *J* = 7.5, 1.0 Hz, Ar–H), 6.86 (1H, s, Ar–H), 4.24 (2H, s, NB–CH₂), 3.88 (3H, s, OCH₃), 3.81 (3H, s, OCH₃), 3.78 (3H, s, OCH₃), 3.20–3.25 (2H, m, α -CH₂), 3.03–2.98 (2H, m, β -CH₂). ¹³C NMR (CD₃OD): δ ppm 159.37 (Ar–Cq), 154.44 (Ar–Cq), 153.60 (Ar–Cq), 132.81

(Ar-CH), 132.73 (Ar-CH), 126.99 (Ar-Cq), 123.19 (Ar-CH), 122.13 (Ar-CH), 120.29 (Ar-Cq), 114.98 (Ar-CH), 112.16 (Ar-CH), 85.04 (Ar-Cq-iodine), 57.59 (OCH₃), 56.71 (OCH₃), 56.24 (OCH₃), 48.1 (NB-CH₂), 48.0 (α -CH₂), 28.49 (β -CH₂). HRMS calculated for C₁₈H₂₃INO₃ [M + H]⁺, 428.07171; observed [M + H]⁺, 428.07239. The EI mass spectrum also has been reported by Casale and Hays.²⁵

N-(3-Methoxybenzyl)-2-(4-iodo-2,5-dimethoxyphenyl)ethan-1-amine Hydrochloride (**4a**). Obtained as needles following crystallization from acetone/EtOAc/Et₂O; yield 85%; mp 171–2 °C. ¹H NMR (300 MHz, CD₃OD) δ ppm 7.38 (1H, t, *J* = 7.7 Hz, Ar-H), 7.34 (1H, s, Ar-H), 6.98–7.10 (3H, m, Ar-H), 6.86 (1H, s, Ar-H), 4.19 (2H, s, NB-CH₂), 3.83 (3H, s, OCH₃), 3.81 (3H, s, OCH₃), 3.79 (3H, s, OCH₃), 3.22–3.27 (2H, m, α -CH₂), 2.99–3.04 (2H, m, β -CH₂). ¹³C NMR (CD₃OD): δ ppm 161.77 (Ar-Cq), 154.43 (Ar-Cq), 153.63 (Ar-Cq), 133.82 (Ar-Cq), 131.48 (Ar-CH), 127.01 (Ar-Cq), 123.14 (Ar-CH), 122.92 (Ar-CH), 116.53 (Ar-CH), 116.13 (Ar-CH), 114.95 (Ar-CH), 85.00 (Ar-Cq-iodine), 57.59 (OCH₃), 56.68 (OCH₃), 55.93 (OCH₃), 52.23 (NB-CH₂), 48.1 (α -CH₂), 28.65 (β -CH₂). HRMS calculated for C₁₈H₂₃INO₃ [M + H]⁺, 428.07171; observed [M + H]⁺, 428.07319. The EI mass spectrum has also been reported by Casale and Hays.²⁵

N-(4-Methoxybenzyl)-2-(4-iodo-2,5-dimethoxyphenyl)ethan-1-amine Hydrochloride (**4b**). This particular compound was extremely difficult to crystallize, providing unfilterable gels upon attempts to crystallize it from EtOH, EtOH/Et₂O, or MeOH/Et₂O. It was finally obtained by dissolving in a minimum amount of boiling acetonitrile and allowing the solution to cool. Upon cooling, the solution also took on a gel-like appearance, but unlike other attempts, this material could be collected by vacuum filtration through a sintered glass filter funnel. The voluminous white solid was washed on the filter with a small amount of cold acetonitrile, then left on the funnel with suction until dry; yield 72%; mp 180–182 °C. ¹H NMR (300 MHz, CD₃OD): δ ppm 7.41 (2H, d, *J* = 8.7 Hz, 2 x Ar-H), 7.34 (1H, s, Ar-H), 7.00 (2H, d, *J* = 8.5 Hz, 2 x Ar-H), 6.85 (1H, s, Ar-H), 4.15 (2H, s, NB-CH₂), 3.82 (3H, s, OCH₃), 3.81 (3H, s, OCH₃), 3.79 (3H, s, OCH₃), 3.18–3.23 (2H, m, α -CH₂), 2.96–3.01 (2H, m, β -CH₂). ¹³C NMR (CD₃OD): δ ppm 162.27 (Ar-Cq), 154.42 (Ar-Cq), 153.64 (Ar-Cq), 132.59 (2 x Ar-CH), 127.05 (Ar-Cq), 124.22 (Ar-Cq), 123.16 (Ar-CH), 115.64 (2 x Ar-CH), 114.94 (Ar-CH), 84.98 (Ar-Cq-iodine), 57.59 (OCH₃), 56.68 (OCH₃), 55.90 (OCH₃), 51.90 (NB-CH₂), 47.8 (α -CH₂), 28.67 (β -CH₂). HRMS calculated for C₁₈H₂₃INO₃ [M + H]⁺, 428.07171; observed [M + H]⁺, 428.07320. The EI mass spectrum has also been reported by Casale and Hays.²⁵

N-(2-Bromobenzyl)-2-(4-iodo-2,5-dimethoxyphenyl)ethan-1-amine Hydrochloride (**4c**). Obtained as needles following crystallization from acetone/EtOAc/Et₂O; yield 79%; mp 170–1 °C. ¹H NMR (300 MHz, CD₃OD): δ ppm 7.74 (1H, dd, *J* = 7.9, 1.3 Hz, Ar-H), 7.61 (1H, dd, *J* = 7.7, 1.7 Hz, Ar-H), 7.49 (1H, td, *J* = 7.5, 1.3 Hz), 7.39 (1H, td, *J* = 7.9, 1.9 Hz), 7.35 (1H, s, Ar-H), 6.89 (1H, s, Ar-H), 4.42 (2H, s, NB-CH₂), 3.82 (3H, s, OCH₃), 3.81 (3H, s, OCH₃), 3.31–3.36 (2H, m, α -CH₂), 3.03–3.08 (2H, m, β -CH₂). ¹³C NMR (CD₃OD): δ ppm 154.46 (Ar-Cq), 153.62 (Ar-Cq), 134.74 (Ar-CH), 133.03 (Ar-CH), 132.81 (Ar-CH), 132.32 (Ar-Cq), 129.70 (Ar-CH), 126.87 (Ar-Cq), 125.94 (Ar-Cq), 123.19 (Ar-CH), 114.98 (Ar-CH), 85.08 (Ar-Cq-iodine), 57.60 (OCH₃), 56.73 (OCH₃), 51.99 (NB-CH₂), 48.7 (α -CH₂), 28.62 (β -CH₂). HRMS calculated for C₁₇H₂₀BrINO₂ [M + H]⁺, 475.97166; observed [M + H]⁺, 475.97212.

N-(3-Bromobenzyl)-2-(4-iodo-2,5-dimethoxyphenyl)ethan-1-amine Hydrochloride (**4d**). Obtained as needles following crystallization from acetone/EtOAc/Et₂O; yield 89%; mp 199–201 °C. ¹H NMR (300 MHz, CD₃OD): δ ppm 7.69–7.74 (1H, m, Ar-H), 7.60–7.66 (1H, m, Ar-H), 7.45–7.51 (1H, m, Ar-H), 7.41 (1H, d, *J* = 7.7 Hz, Ar-H), 7.35 (1H, s, Ar-H), 6.86 (1H, s, Ar-H), 4.22 (2H, s, NB-CH₂), 3.81 (3H, s, OCH₃), 3.79 (3H, s, OCH₃), 3.22–3.27 (2H, m, α -CH₂), 2.98–3.03 (2H, m, β -CH₂). ¹³C NMR (CD₃OD): δ ppm 154.44 (Ar-Cq), 153.63 (Ar-Cq), 134.97 (Ar-Cq), 134.03 (Ar-CH), 133.87 (Ar-CH), 132.12 (Ar-CH), 129.89 (Ar-CH), 126.93 (Ar-Cq), 124.00 (Ar-Cq), 123.18 (Ar-CH), 114.95 (Ar-CH),

85.06 (Ar-Cq-iodine), 57.59 (OCH₃), 56.70 (OCH₃), 51.51 (NB-CH₂), 48.3 (α -CH₂), 28.68 (β -CH₂). HRMS calculated for C₁₇H₂₀BrINO₂ [M + H]⁺, 475.97166; observed [M + H]⁺, 475.97281.

N-(4-Bromobenzyl)-2-(4-iodo-2,5-dimethoxyphenyl)ethan-1-amine Hydrochloride (**4e**). Obtained as needles following crystallization from acetone/EtOAc/Et₂O; yield 81%; mp 196–7 °C. ¹H NMR (300 MHz, CD₃OD) δ ppm 7.64 (2H, d, *J* = 8.7 Hz, 2 x Ar-H), 7.42 (2H, d, *J* = 8.5 Hz, 2 x Ar-H), 7.34 (1H, s, Ar-H), 6.86 (1H, s, Ar-H), 4.21 (2H, s, NB-CH₂), 3.81 (3H, s, OCH₃), 3.79 (3 H, s, OCH₃), 3.22–3.27 (2H, m, α -CH₂), 2.98–3.03 (2H, m, β -CH₂). ¹³C NMR (CD₃OD): δ ppm 154.43 (Ar-Cq), 153.62 (Ar-Cq), 133.50 (2 x Ar-CH), 133.00 (2 x Ar-CH), 131.71 (Ar-Cq), 126.92 (Ar-Cq), 124.92 (Ar-Cq), 123.16 (Ar-CH), 114.94 (Ar-CH), 85.02 (Ar-Cq-iodine), 57.60 (OCH₃), 56.68 (OCH₃), 51.58 (NB-CH₂), 48.2 (α -CH₂), 28.66 (β -CH₂). HRMS calculated for C₁₇H₂₀BrINO₂ [M + H]⁺, 475.97166; observed [M + H]⁺, 475.97268.

N-(2-Methoxybenzyl)-2-(5-methoxy-1*H*-indol-3-yl)ethan-1-amine Hydrochloride (**5a**). Obtained as needles following crystallization from EtOH/EtOAc; yield 91%; mp 232–4 °C. ¹H NMR (300 MHz, CD₃OD): δ ppm 7.42 (1H, td, *J* = 7.9, 1.7 Hz, Ph-H), 7.33 (1H, dd, *J* = 7.4, 1.6 Hz, Ph-H), 7.28 (1H, dd, *J* = 8.9, 0.6 Hz, In-H), 7.16 (1H, s, In-H), 6.97–7.03 (2H, m, Ph-H), 6.95 (1H, d, *J* = 2.4 Hz, In-H), 6.81 (1H, dd, *J* = 8.8, 2.4 Hz, In-H), 4.23 (2H, s, NB-CH₂), 3.78 (3H, s, OCH₃), 3.67 (3H, s, OCH₃), 3.28–3.33 (2H, m, α -CH₂, overlapping with solvent), 3.12–3.17 (2H, m, β -CH₂). ¹³C NMR (CD₃OD): δ ppm 159.25 (Ph-Cq), 155.42 (In-Cq), 133.67 (Ar-Cq), 132.77 (Ph-CH), 132.68 (Ph-CH), 128.34 (Ar-Cq), 125.35 (In-CH), 122.12 (In-CH), 120.13 (Ar-Cq), 113.41 (Ph-CH), 113.21 (In-CH), 112.06 (Ph-CH), 109.51 (Ar-Cq), 101.00 (In-CH), 56.35 (OCH₃), 55.93 (OCH₃), 48.90 (α -CH₂), 48.3 (NB-CH₂), 23.21 (β -CH₂). HRMS calculated for C₁₉H₂₃N₂O₂ [M + H]⁺, 311.17540; observed [M + H]⁺, 311.17548.

N-(3-Methoxybenzyl)-2-(5-methoxy-1*H*-indol-3-yl)ethan-1-amine Maleate (**5b**). Obtained as needles following crystallization from acetone/EtOAc/Et₂O; yield 84%; mp 124–5 °C. ¹H NMR (300 MHz, CD₃OD): δ ppm 7.33–7.39 (1H, m, Ph-H), 7.26 (1H, dd, *J* = 8.9, 0.6 Hz, In-H), 7.13 (1H, s, In-H), 6.99–7.01 (4H, m, overlapping 3 x Ph-H, 1 x In-H), 6.80 (1H, dd, *J* = 8.8, 2.4 Hz, In-H), 6.24 (2H, s, maleate), 4.18 (2H, s, NB-CH₂), 3.81 (3H, s, OCH₃), 3.80 (3H, s, OCH₃), 3.28–3.35 (2H, α -CH₂, overlapping with solvent), 3.11–3.16 (2H, m, β -CH₂). ¹³C NMR (CD₃OD): δ ppm 170.89 (maleate), 161.79 (Ph-Cq), 155.39 (In-Cq), 136.79 (maleate), 133.89 (Ar-Cq), 133.60 (Ar-Cq), 131.50 (Ph-CH), 128.44 (Ar-Cq), 125.00 (In-CH), 122.87 (Ph-CH), 116.40 (Ph-CH), 116.15 (Ph-CH), 113.35 (In-CH), 113.07 (In-CH), 109.84 (Ar-Cq), 101.04 (In-CH), 56.39 (OCH₃), 55.88 (OCH₃), 52.16 (NB-CH₂), 49.0 (α -CH₂), 23.36 (β -CH₂). HRMS calculated for C₁₉H₂₃N₂O₂ [M + H]⁺, 311.17540; observed [M + H]⁺, 311.17572.

N-(4-Methoxybenzyl)-2-(5-methoxy-1*H*-indol-3-yl)ethan-1-amine Maleate (**5c**). Obtained as needles following crystallization from acetone/EtOAc/Et₂O; yield 82%; mp 172–3 °C. ¹H NMR (300 MHz, CD₃OD): δ ppm 7.36 (2H, d, *J* = 8.0 Hz, 2 x Ph-H), 7.26 (1 H, dd, *J* = 8.8, 0.5 Hz, In-H), 7.12 (1H, s, In-H), 6.99 (1H, d, *J* = 2.5 Hz, In-H), 6.97 (2H, d, *J* = 6.6 Hz, 2 x Ph-H), 6.80 (1H, dd, *J* = 8.9, 2.4 Hz, In-H), 6.24 (2H, s, maleate), 4.15 (2H, s, NB-CH₂), 3.81 (3H, s, OCH₃), 3.80 (3H, s, OCH₃), 3.27–3.32 (2H, m, α -CH₂, overlapping with solvent), 3.09–3.14 (2H, m, β -CH₂). ¹³C NMR (CD₃OD): δ ppm 170.90 (maleate), 162.24 (Ph-Cq), 155.37 (In-Cq), 136.78 (maleate), 133.60 (Ar-Cq), 132.52 (2 x Ph-CH), 128.45 (Ar-Cq), 124.97 (In-CH), 124.27 (Ar-Cq), 115.64 (2 x Ph-CH), 113.34 (In-CH), 113.06 (In-CH), 109.89 (Ar-Cq), 101.06 (In-CH), 56.39 (OCH₃), 55.89 (OCH₃), 51.78 (NB-CH₂), 48.5 (α -CH₂), 23.38 (β -CH₂). HRMS calculated for C₁₉H₂₃N₂O₂ [M + H]⁺, 311.17540; observed [M + H]⁺, 311.17632.

N-(2-Bromobenzyl)-2-(5-methoxy-1*H*-indol-3-yl)ethan-1-amine Maleate (**5d**). Obtained as needles following crystallization from acetone/EtOAc/Et₂O; yield 72%; mp 93–5 °C. ¹H NMR (300 MHz, CD₃OD): δ ppm 7.70 (1H, dd, *J* = 7.9, 1.3 Hz, Ph-H), 7.54 (1H, dd, *J* = 7.7, 1.9 Hz, Ph-H), 7.45 (1H, td, *J* = 7.5, 1.4 Hz, Ph-H), 7.36 (1H, td, *J* = 7.8, 1.8 Hz, Ph-H), 7.26 (1H, dd, *J* = 8.9, 0.6 Hz, In-H), 7.16

(1H, s, In-H), 7.02 (1H, d, $J = 2.3$ Hz, In-H), 6.80 (1H, dd, $J = 8.9$, 2.4 Hz, In-H), 6.24 (2H, s, maleate), 4.41 (2H, s, NB-CH₂), 3.82 (3H, s, OCH₃), 3.40–3.45 (2H, m, α -CH₂), 3.16–3.21 (2H, m, β -CH₂). ¹³C NMR (CD₃OD): δ ppm 170.89 (maleate), 155.41 (In-Cq), 136.75 (maleate), 134.71 (Ph-CH), 133.64 (Ar-Cq), 133.04 (Ph-CH), 132.76 (Ph-CH), 132.40 (Ar-Cq), 129.65 (Ph-CH), 128.44 (Ar-Cq), 125.94 (Ar-Cq), 125.12 (In-CH), 113.38 (In-CH), 113.10 (In-CH), 109.67 (Ar-Cq), 101.06 (In-CH), 56.40 (OCH₃), 51.90 (NB-CH₂), 49.3 (α -CH₂), 23.32 (β -CH₂). HRMS calculated for C₁₈H₂₀BrN₂O [M + H]⁺, 359.07535; observed [M + H]⁺, 359.07581.

N-(3-Bromobenzyl)-2-(5-methoxy-1H-indol-3-yl)ethan-1-amine Maleate (5e). Obtained as needles following crystallization from acetone/EtOAc/Et₂O; yield 86%; mp 137–8 °C. ¹H NMR (300 MHz, CD₃OD): δ ppm 7.67–7.68 (1H, m, Ph-H), 7.61 (1H, dt, $J = 7.7$, 1.6 Hz, Ph-H), 7.34–7.45 (2H, m, Ph-H), 7.27 (1H, d, $J = 8.7$ Hz, In-H), 7.14 (1H, s, In-H), 7.01 (1H, d, $J = 2.3$ Hz, In-H), 6.80 (1H, dd, $J = 8.9$, 2.4 Hz, In-H), 6.24 (2H, s, maleate), 4.21 (2H, s, NB-CH₂), 3.82 (3H, s, OCH₃), 3.31–3.36 (2H, m, α -CH₂, overlapping with solvent), 3.11–3.16 (2H, m, β -CH₂). ¹³C NMR (CD₃OD): δ ppm 170.92 (maleate), 155.40 (In-Cq), 136.78 (maleate), 135.10 (Ar-Cq), 134.00 (Ph-CH), 133.82 (Ph-CH), 133.61 (Ar-Cq), 132.10 (Ph-CH), 129.81 (Ph-CH), 128.45 (Ar-Cq), 125.01 (In-CH), 124.03 (Ar-Cq), 113.37 (In-CH), 113.08 (In-CH), 109.82 (Ar-Cq), 101.06 (In-CH), 56.42 (OCH₃), 51.52 (NB-CH₂), 49.1 (α -CH₂), 23.40 (β -CH₂). HRMS calculated for C₁₈H₂₀BrN₂O [M + H]⁺, 359.07535; observed [M + H]⁺, 359.07547.

N-(4-Bromobenzyl)-2-(5-methoxy-1H-indol-3-yl)ethan-1-amine Maleate (5f). Obtained as needles following crystallization from acetone/EtOAc/Et₂O; yield 75%; mp 181–3 °C. ¹H NMR (CD₃OD): δ ppm 7.60 (2H, d, $J = 8.5$ Hz, 2 x Ph-H), 7.37 (2H, d, $J = 8.5$ Hz, 2 x Ph-H), 7.26 (1H, d, $J = 8.9$ Hz, In-H), 7.13 (1H, s, In-H), 6.99 (1H, d, $J = 2.3$ Hz, In-H), 6.80 (1H, dd, $J = 8.9$, 2.4 Hz, In-H), 6.24 (2H, s, maleate), 4.20 (2H, s, NB-CH₂), 3.81 (3H, s, OCH₃), 3.31–3.36 (2H, m, α -CH₂, overlapping with solvent), 3.11–3.16 (2H, m, β -CH₂). ¹³C NMR (CD₃OD): δ ppm 170.89 (maleate), 155.38 (In-Cq), 136.75 (maleate), 133.61 (Ar-Cq), 133.51 (2 x Ph-CH), 132.92 (2 x Ph-CH), 131.77 (Ar-Cq), 128.43 (Ar-Cq), 125.02 (In-CH), 124.90 (Ar-Cq), 113.36 (In-CH), 113.06 (In-CH), 109.76 (Ar-Cq), 101.06 (In-CH), 56.41 (OCH₃), 51.50 (NB-CH₂), 48.90 (α -CH₂), 23.40 (β -CH₂). HRMS calculated for C₁₈H₂₀BrN₂O [M + H]⁺, 359.07535; observed [M + H]⁺, 359.07597.

N-(3-Fluorobenzyl)-2-(5-methoxy-1H-indol-3-yl)ethan-1-amine Maleate (5g). Obtained as needles following crystallization from acetone/EtOAc/Et₂O; yield 78%; mp 150–2 °C. ¹H NMR (300 MHz, CD₃OD): δ ppm 7.44–7.51 (1H, m, Ph-H), 7.16–7.29 (4H, m, overlapping 3 x Ph-H, 1 x In-H), 7.14 (1H, s, In-H), 7.01 (1H, d, $J = 2.4$ Hz, In-H), 6.80 (1H, dd, $J = 8.9$, 2.4 Hz, In-H), 6.24 (2H, s, maleate), 4.24 (2H, s, NB-CH₂), 3.81 (3H, s, OCH₃), 3.31–3.37 (2H, m, α -CH₂, overlapping with solvent), 3.12–3.17 (2H, m, β -CH₂). ¹³C NMR (CD₃OD): δ ppm 170.89 (maleate), 164.38 (Ph-Cq-3', d, $J = 246.2$ Hz), 155.40 (In-Cq), 136.74 (maleate), 135.08 (Ph-Cq-1', d, $J = 7.5$ Hz), 133.61 (In-Cq), 132.31 (Ph-C-5', d, $J = 8.3$ Hz), 128.45 (In-Cq), 126.88 (Ph-C-6', d, $J = 3.0$ Hz), 125.00 (In-CH), 117.79 (Ph-C-2', d, $J = 22.5$ Hz), 117.60 (Ph-C-4', d, $J = 21.8$ Hz), 113.37 (In-CH), 113.08 (In-CH), 109.79 (In-Cq), 101.06 (In-CH), 56.41 (OCH₃), 51.58 (NB-CH₂, $J = 1.5$ Hz), 49.1 (α -CH₂), 23.38 (β -CH₂). HRMS calculated for C₁₈H₂₀FN₂O [M + H]⁺, 299.15542; observed [M + H]⁺, 299.15602.

N-(3-Chlorobenzyl)-2-(5-methoxy-1H-indol-3-yl)ethan-1-amine Maleate (5h). Obtained as needles following crystallization from acetone/EtOAc/Et₂O; yield 79%; mp 116–8 °C. ¹H NMR (300 MHz, CD₃OD): δ ppm 7.52 (1H, br, s, Ph-H), 7.34–7.49 (3H, m, Ph-H), 7.26 (1H, d, $J = 8.9$ Hz, In-H), 7.14 (1H, s, In-H), 7.01 (1H, d, $J = 2.4$ Hz, In-H), 6.80 (1H, dd, $J = 8.9$, 2.4 Hz, In-H), 6.24 (2H, s, maleate), 4.22 (2H, s, NB-CH₂), 3.82 (3H, s, OCH₃), 3.31–3.37 (2H, m, α -CH₂, overlapping with solvent), 3.12–3.17 (2H, m, β -CH₂). ¹³C NMR (CD₃OD): δ ppm 155.41 (In-Cq), 136.76 (maleate), 136.10 (Ar-Cq), 134.82 (Ar-Cq), 133.60 (Ar-Cq), 131.89 (Ph-CH), 131.04 (Ph-CH), 130.84 (Ph-CH), 129.38 (Ph-CH), 128.45 (Ar-Cq), 125.01 (In-CH), 113.36 (In-CH), 113.08 (In-CH), 109.78 (In-

Cq), 101.04 (In-CH), 56.40 (OCH₃), 51.54 (NB-CH₂), 49.1 (α -CH₂), 23.39 (β -CH₂). HRMS calculated for C₁₈H₂₀ClN₂O [M + H]⁺, 315.12587; observed [M + H]⁺, 315.12666.

N-(3-Iodobenzyl)-2-(5-methoxy-1H-indol-3-yl)ethan-1-amine Maleate (5i). Obtained as needles following crystallization from acetone/EtOAc/Et₂O; yield 84%; mp 131–2 °C. ¹H NMR (300 MHz, CD₃OD): δ ppm 7.87 (1H, brs, Ph-H), 7.81 (1H, d, $J = 7.9$ Hz, Ph-H), 7.45 (1H, d, $J = 7.7$ Hz, Ph-H), 7.27 (1H, d, $J = 8.3$ Hz, In-H), 7.21 (1H, t, $J = 7.8$ Hz, Ph-H), 7.13 (1H, s, In-H), 7.01 (1H, d, $J = 2.3$ Hz, In-H), 6.80 (1H, dd, $J = 8.9$, 2.3 Hz, In-H), 6.24 (2H, s, maleate), 4.18 (2H, s, NB-CH₂), 3.82 (3H, s, OCH₃), 3.31–3.36 (2H, m, α -CH₂, overlapping with solvent), 3.11–3.16 (2H, m, β -CH₂). ¹³C NMR (CD₃OD): δ ppm 155.39 (In-Cq), 139.98 (Ph-CH), 139.88 (Ph-CH), 136.76 (maleate), 134.98 (Ar-Cq), 133.58 (Ar-Cq), 132.03 (Ph-CH), 130.31 (Ph-CH), 128.46 (Ar-Cq), 124.99 (In-CH), 113.36 (In-CH), 113.08 (In-CH), 109.80 (Ar-Cq), 101.03 (In-CH), 95.42 (Ar-Cq-iodine), 56.42 (OCH₃), 51.41 (NB-CH₂), 49.1 (α -CH₂), 23.37 (β -CH₂). HRMS calculated for C₁₈H₂₀I₂N₂O [M + H]⁺, 407.06148; observed [M + H]⁺, 407.06188.

N-(3-Methylbenzyl)-2-(5-methoxy-1H-indol-3-yl)ethan-1-amine Maleate (5j). Obtained as needles following crystallization from acetone/EtOAc/Et₂O; yield 78%; mp 125–7 °C. ¹H NMR (300 MHz, CD₃OD): δ ppm 7.22–7.35 (5H, m, overlapping 4 x Ph-H and 1 x In-H), 7.13 (1H, s, In-H), 6.99 (1H, d, $J = 2.3$ Hz, In-H), 6.80 (1H, dd, $J = 8.9$, 2.4 Hz, In-H), 6.24 (2H, s, maleate), 4.17 (2H, s, NB-CH₂), 3.82 (3H, s, OCH₃), 3.29–3.35 (2H, m, α -CH₂, overlapping with solvent), 3.10–3.15 (2H, m, β -CH₂), 2.36 (3H, s, CH₃). ¹³C NMR (CD₃OD): δ ppm 170.90 (maleate), 155.38 (In-Cq), 140.47 (Ar-Cq), 136.80 (maleate), 133.61 (Ar-Cq), 132.46 (Ar-Cq), 131.49 (Ph-CH), 131.40 (Ph-CH), 130.27 (Ph-CH), 128.46 (Ar-Cq), 127.93 (Ph-CH), 125.00 (In-CH), 113.35 (In-CH), 113.06 (In-CH), 109.87 (Ar-Cq), 101.07 (In-CH), 56.40 (OCH₃), 52.24 (NB-CH₂), 48.9 (α -CH₂), 23.37 (β -CH₂), 21.36 (CH₃). HRMS calculated for C₁₉H₂₃N₂O [M + H]⁺, 295.18049; observed [M + H]⁺, 295.18090.

N-(3-Methylthiobenzyl)-2-(5-methoxy-1H-indol-3-yl)ethan-1-amine Maleate (5j). Obtained as needles following crystallization from acetone/EtOAc/Et₂O; yield 80%; mp 151–2 °C. ¹H NMR (300 MHz, CD₃OD): δ ppm 7.31–7.39 (3H, m, Ph-H), 7.26 (1H, d, $J = 8.9$ Hz, In-H), 7.19 (1H, dt, $J = 7.0$, 1.9 Hz, Ph-H), 7.13 (1H, s, In-H), 7.00 (1H, d, $J = 2.4$ Hz, In-H), 6.80 (1H, dd, $J = 8.9$, 2.4 Hz, In-H), 6.24 (2H, s, maleate), 4.19 (2H, s, NB-CH₂), 3.81 (3H, s, OCH₃), 3.30–3.35 (2H, m, α -CH₂, overlapping with solvent), 3.11–3.16 (2H, m, β -CH₂), 2.48 (3H, s, CH₃). ¹³C NMR (CD₃OD): δ ppm 170.91 (maleate), 155.40 (Ar-Cq), 141.95 (Ar-Cq), 136.78 (maleate), 133.60 (Ar-Cq), 133.34 (Ar-Cq), 130.74 (Ph-CH), 128.46 (Ar-Cq), 128.40 (Ph-CH), 128.30 (Ph-CH), 127.20 (Ph-CH), 125.00 (In-CH), 113.36 (In-CH), 113.08 (In-CH), 109.84 (Ar-Cq), 101.04 (In-CH), 56.41 (OCH₃), 52.05 (NB-CH₂), 49.1 (α -CH₂), 23.38 (β -CH₂), 15.37 (CH₃). HRMS calculated for C₁₉H₂₃N₂OS [M + H]⁺, 327.15256; observed [M + H]⁺, 327.15362.

N-(3-Trifluoromethylbenzyl)-2-(5-methoxy-1H-indol-3-yl)ethan-1-amine Maleate (5k). Obtained as needles following crystallization from acetone/EtOAc/Et₂O; yield 62%; mp 161–2 °C. ¹H NMR (300 MHz, CD₃OD): δ ppm 7.84 (1H, brs, Ph-H), 7.62–7.78 (3H, m, Ph-H), 7.26 (1H, d, $J = 8.8$ Hz, In-H), 7.14 (1H, s, In-H), 7.02 (1H, d, $J = 2.1$ Hz, In-H), 6.80 (1H, dd, $J = 8.9$, 2.3 Hz, In-H), 6.24 (2H, s, maleate), 4.32 (2H, s, NB-CH₂), 3.81 (3H, s, OCH₃), 3.35–3.40 (2H, m, α -CH₂), 3.13–3.18 (2H, m, β -CH₂). ¹³C NMR (CD₃OD): δ ppm 170.91 (maleate), 155.41 (In-Cq), 136.74 (maleate), 134.85 (Ph-CH), 134.04 (Ph-Cq), 133.60 (In-Cq), 132.56 (Ph-Cq, d, $J = 32.3$ Hz), 131.24 (Ph-CH), 128.46 (In-Cq), 127.84 (Ph-CH, q, $J = 4.0$ Hz), 127.46 (Ph-CH, q, $J = 3.8$ Hz), 125.4 (CF₃, q, $J = 272$ Hz), 125.01 (In-CH), 113.36 (In-CH), 113.06 (In-CH), 109.79 (In-Cq), 101.06 (In-CH), 56.39 (OCH₃), 51.63 (NB-CH₂), 49.20 (α -CH₂), 23.41 (β -CH₂). HRMS calculated for C₁₉H₂₀F₃N₂O [M + H]⁺, 349.15222; observed [M + H]⁺, 349.15259.

Pharmacology. Receptor Affinity. Receptor affinity values for a panel of human serotonin receptors were obtained for all compounds through the NIMH-sponsored PDSP program (www.pdsp.med.unc.edu). Affinity data from screening are reported in Table 1. Following

the initial screen, more detailed values were obtained for affinity at the human 5-HT_{2A} and 5-HT_{2C} receptors using both an antagonist radioligand ([³H]ketanserin for 5-HT_{2A}) and ([³H]mesulergine for 5-HT_{2C}) and an agonist radioligand ([³H]-DOI) for both receptors. Those data are reported in Table 2.

Receptor Efficacy and Potency in the Ca²⁺ Mobilization Assay. Changes in intracellular Ca²⁺ levels were measured using a Fluorometric Imaging plate reader (FLIPR^{TETRA}, Molecular Devices), essentially as described in the PDSP (NIMH Psychoactive Drug Screening Program) Assay Protocol Book (www.pdsp.med.unc.edu). PO1C cells stably transfected with r5-HT_{2C} or r5-HT_{2A} receptors, and HEK 293 cells stably transfected with h5-HT_{2A}, h5-HT_{2B}, or h5-HT_{2C} receptors were plated (20,000 cells/well) into poly-L-lysine coated 394-well clear-bottom black-walled microplates (Greiner Bio-one) with 50 μ L of media (DMEM media supplemented with 500 μ g/mL Geneticin sulfate (G-418), 10% dialyzed fetal bovine serum, and 50 U of penicillin/50 μ g of streptomycin) and incubated overnight (37 °C, 5% CO₂). The following day, media were replaced with 20 μ L of FLIPR Calcium 4 Assay Kit (Molecular Devices) diluted in assay buffer (HBSS, 2.5 mM probenecid, and 20 mM HEPES, pH 7.4–7.8) and incubated for 45 min at 37 °C and 15 min at room temperature. Compounds were initially dissolved in DMSO. The 16-point curves were prepared as 3 \times serial dilutions for each compound with final concentrations ranging from 10 μ M to 0.003 nM. Basal fluorescence was measured for 10 s, then 10 μ L of test or control compounds was added followed by continued fluorescence measurement for an additional 120 s. Raw data were normalized to baseline fluorescence (0%) and 5HT at 10 μ M (100%), expressed as percent activation, and plotted as a function of molar concentration of test compound using Prism 5.0 (GraphPad Software). These data are reported in Table 3.

Mouse Head Twitch Response. Animals. Male C57BL/6J mice (6–8 weeks old) were obtained from Jackson Laboratories (Bar Harbor, ME, USA) and housed in a vivarium at the University of California, San Diego, an AAALAC-approved animal facility that meets Federal and State requirements for the care and treatment of laboratory animals. Mice were housed up to four per cage in a climate-controlled room with a reversed light-cycle (lights on at 1900 h, off at 0700 h). Food and water were provided ad libitum, except during behavioral testing. Testing was performed between 1000 and 1830 h. Experiments were conducted in accord with NIH guidelines and were approved by the UCSD animal care committee.

Procedures. The HTR was assessed using a head-mounted magnet and a magnetometer detection coil. Mice were anesthetized (100 mg/kg ketamine, 3 mg/kg acepromazine, and 20 mg/kg xylazine, IP), and a neodymium magnet (4.57 \times 4.57 \times 2.03 mm, 375 mg) was attached to the skull using dental cement. The magnet was positioned so that the N–S axis was parallel to the dorsoventral plane of the head. Mice were allowed to recover for 2 weeks after surgery. HTR experiments were conducted in a well-lit room. Test compounds were dissolved in water containing 5% Tween-80 and administered SC (5 or 10 mL/kg). Mice were injected with drug or vehicle and placed in a glass cylinder surrounded by a magnetometer coil. Head movements were recorded and analyzed for HTR as described previously.^{23,31} Coil voltage was low-pass filtered (5–10 kHz), amplified, and digitized (40 kHz sampling rate) using a Powerlab/8SP with LabChart v 7.3.2 (ADInstruments, Colorado Springs, CO, USA). The data were filtered off-line (40–200 Hz band-pass), and HTRs were identified by manually searching for sinusoidal wavelets possessing at least two bipolar peaks, spectrum in the 40–160 Hz range, amplitude exceeding the background noise level, and duration <0.15 s, with stable coil voltage during the period immediately before and after each response.

Analysis. HTR counts were analyzed using one-way analyses of variance (ANOVAs). Post-hoc comparisons were made using Tukey's studentized range method. Significance was demonstrated by surpassing an α -level of 0.05. ED₅₀ values and 95% confidence limits were calculated using nonlinear regression. These data are reported in Table 4.

AUTHOR INFORMATION

Corresponding Author

*Division of Chemical Biology and Medicinal Chemistry, Genetic Medicine Building, 2078, 120 Mason Farm Road, University of North Carolina, Chapel Hill, NC 27599. Phone: (765) 404-0350. E-mail: denichol@email.unc.edu.

Author Contributions

D.N. directed the project, synthesized all of the compounds, supervised the integration of the various studies, and was responsible for the writing and final editing of the manuscript, F.S. carried out the calcium mobilization functional assays, A.H. supervised the mouse head twitch assays, L.M.K. assisted with the mouse assays, S.D.B. and S.P.E. carried out the analytical chemistry assays, and W.F. made key suggestions for the project and edited the manuscript.

Funding

This work was supported by the NIMH Psychoactive Drug Screening Program, NIMH K01 MH100644, NIDA R01 DA002925, and the Brain and Behavior Research Foundation.

Notes

The authors declare no competing financial interest.

REFERENCES

- (1) Lawn, W., Barratt, M., Williams, M., Horne, A., and Winstock, A. (2014) The NBOMe hallucinogenic drug series: Patterns of use, characteristics of users and self-reported effects in a large international sample. *J. Psychopharmacol.* 28, 780–788.
- (2) Drug Enforcement Administration, D. o. J. Schedules of Controlled Substances: Temporary Placement of Three Synthetic Phenethylamines into Schedule I, 21 CFR Part 1308, 2013, [Docket NO. DEA-382].
- (3) Poklis, J. L., Devers, K. G., Arbefeville, E. F., Pearson, J. M., Houston, E., and Poklis, A. (2014) Postmortem detection of 25I-NBOMe [2-(4-iodo-2,5-dimethoxyphenyl)-N-[(2-methoxyphenyl)methyl]ethanamine] in fluids and tissues determined by high performance liquid chromatography with tandem mass spectrometry from a traumatic death. *Forensic Sci. Int.* 234, e14–e20.
- (4) Walterscheid, J. P., Phillips, G. T., Lopez, A. E., Gonsoulin, M. L., Chen, H. H., and Sanchez, L. A. (2014) Pathological findings in 2 cases of fatal 25I-NBOMe toxicity. *Am. J. Forensic Med. Pathol.* 35, 20–25.
- (5) Nikolaou, P., Papoutsis, I., Stefanidou, M., Spiliopoulou, C., and Athanasielis, S. (2015) 2C-I-NBOMe, an “N-bomb” that kills with “Smiles”. Toxicological and legislative aspects. *Drug Chem. Toxicol.* 38, 113–119.
- (6) European Monitoring Centre for Drugs and Drug Addiction (2014) *Risk Assessment Report of a New Psychoactive Substance: 2-(4-Iodo-2,5-dimethoxyphenyl)-N-(2-methoxybenzyl)ethanamine (25I-NBOMe)*, EMCDDA, Lisbon, Portugal.
- (7) World Health Organization (2014) *Thirty-Sixth Meeting of the Expert Committee on Drug Dependence*, World Health Organization, Geneva, Switzerland.
- (8) Council of the European Union (2014) 2014/688/EU: Council implementing decision of 25 December 2014 on subjecting –4-iodo-2,5-dimethoxy-N-(2-methoxybenzyl)phenethylamine (25I-NBOMe), 3,4-dichloro-N-[[1-(dimethylamino)cyclohexyl]methyl]benzamide (AH-7921), 3,4-methylenedioxypropylvalerone (MDPV) and 2-(3-methoxyphenyl)-2-(ethylamino)cyclohexanone (methoxetamine) to control measures. *Off. J. Eur. Union* L287, 22.
- (9) Shulgin, A. T. (1978) Psychotomimetic Drugs: Structure-Activity Relationships, in *Handbook of Psychopharmacology*, Chapter 6, pp 243–333, Plenum Press, New York.
- (10) Shulgin, A., and Shulgin, A. (1991) *PIHKAL: A Chemical Love Story*, Transform Press, Berkeley, CA.
- (11) Heim, R., Pertz, H. H., and Elz, S. (1999) Preparation and in vitro pharmacology of novel secondary amine-type 5-HT_{2A} receptor

agonists: from submillimolar to subnanomolar activity. *Arch. Pharm. Pharm. Med. Chem.* 332, 34.

(12) Elz, S., Klass, T. H. R., Wamke, U., and Pertz, H. H. (2002) Development of highly potent partial agonists and chiral antagonists as tools for the study of 5-HT_{2A}-receptor mediated functions. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 365 (Suppl 1), R29.

(13) Heim, R. (2003) Synthese und Pharmakologie potenter 5-HT_{2A}-Rezeptoragonisten mit N-2-Methoxybenzyl-Partialstruktur Entwicklung eines neuen Struktur-Wirkungskonzepts, Ph.D. Thesis, Freie Universität Berlin, Berlin, Germany.

(14) Braden, M. R., Parrish, J. C., Naylor, J. C., and Nichols, D. E. (2006) Molecular interaction of serotonin 5-HT_{2A} receptor residues Phe339(6.51) and Phe340(6.52) with superpotent N-benzyl phenethylamine agonists. *Mol. Pharmacol.* 70, 1956–1964.

(15) Heim, R., and Elz, S. (2000) Novel extremely potent partial 5-HT_{2A} receptor agonists: successful application of a new structure-activity concept. *Arch. Pharm. Pharm. Med. Chem.* 333, 39.

(16) Nichols, D. E. (2012) Structure-activity relationships of serotonin 5-HT_{2A} agonists. *WIREs Membr. Transp. Signaling* 1, 559–579.

(17) Braden, M. R. (2007) Towards a Biophysical Understanding of Hallucinogen Action, Ph.D. Thesis, Purdue University, West Lafayette, IN.

(18) Silva, M. E., Heim, R., Strasser, A., Elz, S., and Dove, S. (2011) Theoretical studies on the interaction of partial agonists with the 5-HT_{2A} receptor. *J. Comput.-Aided Mol. Des.* 25, 51–66.

(19) Glennon, R. A., Dukat, M., el Bermawy, M., Law, H., De Los Angeles, J., Teitler, M., King, A., and Herrick-Davis, K. (1994) Influence of amine substituents on 5-HT_{2A} versus 5-HT_{2C} binding of phenylalkyl- and indolylalkylamines. *J. Med. Chem.* 37, 1929–1935.

(20) Takagi, K., Takayanagi, I., Irikura, T., Nishino, K., and Ito, M. (1969) A potent competitive inhibitor of 5-hydroxytryptamine: 3-(2'-benzylaminoethyl)-5-methoxyindol hydrochloride. *Jpn. J. Pharmacol.* 19, 234–239.

(21) Leff, P., Martin, G. R., and Morse, J. M. (1986) The classification of peripheral 5-HT₂-like receptors using tryptamine agonist and antagonist analogues. *Br. J. Pharmacol.* 89, 493–499.

(22) Jensen, N. (2004) Tryptamines as Ligands and Modulators of the Serotonin 5-HT_{2A} Receptor and the Isolation of Aerginascins from the Hallucinogenic Mushroom *Inocybe aeruginascens*, Ph.D. Thesis, Georg-August-Universität zu Göttingen, Göttingen, Germany.

(23) Halberstadt, A. L., and Geyer, M. A. (2013) Characterization of the head-twitch response induced by hallucinogens in mice: Detection of the behavior based on the dynamics of head movement. *Psychopharmacology* 227, 727–739.

(24) Hansen, M. (2010) Design and Synthesis of Selective Serotonin Receptor Agonists for Positron Emission Tomography Imaging of the Brain, Ph.D. Thesis, University of Copenhagen, Copenhagen, Denmark.

(25) Casale, J. F., and Hays, P. A. (2012) Characterization of eleven 2,5-dimethoxy-N-(2-methoxybenzyl)phenethylamine (NBOMe) derivatives and differentiation from their 3- and 4-methoxybenzyl analogues - Part I. *Microgram J.* 9, 84–109.

(26) Abdel-Magid, A. F., Carson, K. G., Harris, B. D., Maryanoff, C. A., and Shah, R. D. (1996) Reductive amination of aldehydes and ketones with sodium triacetoxyborohydride. Studies on direct and indirect reductive amination procedures. *J. Org. Chem.* 61, 3849–3862.

(27) Dougherty, J. P., and Aloyo, V. J. (2011) Pharmacological and behavioral characterization of the 5-HT_{2A} receptor in C57BL/6N mice. *Psychopharmacology* 215, 581–593.

(28) Braden, M. R., and Nichols, D. E. (2007) Assessment of the roles of serines 5.43(239) and 5.46(242) for binding and potency of agonist ligands at the human serotonin 5-HT_{2A} receptor. *Mol. Pharmacol.* 72, 1200–1209.

(29) Johnson, M. P., Loncharich, R. J., Baez, M., and Nelson, D. L. (1994) Species variations in transmembrane region V of the 5-hydroxytryptamine type 2A receptor alter the structure-activity relationship of certain ergolines and tryptamines. *Mol. Pharmacol.* 45, 277–286.

(30) Johnson, M. P., Audia, J. E., Nissen, J. S., and Nelson, D. L. (1993) N(1)-substituted ergolines and tryptamines show species differences for the agonist-labeled 5-HT₂ receptor. *Eur. J. Pharmacol.* 239, 111–118.

(31) Halberstadt, A. L., and Geyer, M. A. (2014) Effects of the hallucinogen 2,5-dimethoxy-4-iodophenethylamine (2C-I) and superpotent N-benzyl derivatives on the head twitch response. *Neuropharmacology* 77, 200–207.

(32) Fantegrossi, W. E., Harrington, A. W., Eckler, J. R., Arshad, S., Rabin, R. A., Winter, J. C., Coop, A., Rice, K. C., and Woods, J. H. (2005) Hallucinogen-like actions of 2,5-dimethoxy-4-(n)-propylthiophenethylamine (2C-T-7) in mice and rats. *Psychopharmacology* 181, 496–503.

(33) Fantegrossi, W. E., Harrington, A. W., Kiessel, C. L., Eckler, J. R., Rabin, R. A., Winter, J. C., Coop, A., Rice, K. C., and Woods, J. H. (2006) Hallucinogen-like actions of 5-methoxy-N,N-diisopropyltryptamine in mice and rats. *Pharmacol., Biochem. Behav.* 83, 122–129.

(34) Fantegrossi, W. E., Simoneau, J., Cohen, M. S., Zimmerman, S. M., Henson, C. M., Rice, K. C., and Woods, J. H. (2010) Interaction of 5-HT_{2A} and 5-HT_{2C} receptors in R(-)-2,5-dimethoxy-4-iodoamphetamine-elicited head twitch behavior in mice. *J. Pharmacol. Exp. Ther.* 335, 728–734.

(35) Fantegrossi, W. E., Gray, B. W., Bailey, J. M., Smith, D. A., Hansen, M., Kristensen, J. L., and Woods, J. H. (2014) Hallucinogen-like effects of 2-([2-(4-cyano-2,5-dimethoxyphenyl) ethylamino]-methyl)phenol (2SCN-NBOH), a novel N-benzylphenethylamine with 100-fold selectivity for 5-HT receptors, in mice. *Psychopharmacology (Berl.)*.

(36) Bedard, P., and Pycocock, C. J. (1977) "Wet-dog" shake behaviour in the rat: a possible quantitative model of central 5-hydroxytryptamine activity. *Neuropharmacology* 16, 663–670.

(37) Matthews, W. D., and Smith, C. D. (1980) Pharmacological profile of a model for central serotonin receptor activation. *Life Sci.* 26, 1397–1403.

(38) Benneyworth, M. A., Xiang, Z., Smith, R. L., Garcia, E. E., Conn, P. J., and Sanders-Bush, E. (2007) A selective positive allosteric modulator of metabotropic glutamate receptor subtype 2 blocks a hallucinogenic drug model of psychosis. *Mol. Pharmacol.* 72, 477–484.

(39) Canal, C. E., Olaghere da Silva, U. B., Gresch, P. J., Watt, E. E., Sanders-Bush, E., and Airey, D. C. (2010) The serotonin 2C receptor potently modulates the head-twitch response in mice induced by a phenethylamine hallucinogen. *Psychopharmacology* 209, 163–174.

(40) Halberstadt, A. L., Koedood, L., Powell, S. B., and Geyer, M. A. (2011) Differential contributions of serotonin receptors to the behavioral effects of indoleamine hallucinogens in mice. *J. Psychopharmacol.* 25, 1548–1561.

(41) Corne, S. J., and Pickering, R. W. (1967) A possible correlation between drug-induced hallucinations in man and a behavioural response in mice. *Psychopharmacologia* 11, 65–78.

(42) Darmani, N. A., Martin, B. R., and Glennon, R. A. (1990) Withdrawal from chronic treatment with (±)-DOI causes supersensitivity to 5-HT₂ receptor-induced head-twitch behaviour in mice. *Eur. J. Pharmacol.* 186, 115–118.

(43) Leth-Petersen, S., Bundgaard, C., Hansen, M., Carnerup, M. A., Kehler, J., and Kristensen, J. L. (2014) Correlating the metabolic stability of psychedelic 5-HT agonists with anecdotal reports of human oral bioavailability. *Neurochem. Res.* 39, 2018–2023.

(44) Juncosa, J. I., Jr., Hansen, M., Bonner, L. A., Cueva, J. P., Maglathlin, R., McCorvy, J. D., Marona-Lewicka, D., Lill, M. A., and Nichols, D. E. (2013) Extensive rigid analogue design maps the binding conformation of potent N-benzylphenethylamine 5-HT_{2A} serotonin receptor agonist ligands. *ACS Chem. Neurosci.* 4, 96–109.

(45) Hansen, M., Phonekeo, K., Paine, J. S., Leth-Petersen, S., Begtrup, M., Brauner-Osborne, H., and Kristensen, J. L. (2014) Synthesis and structure-activity relationships of N-benzyl phenethylamines as 5-HT_{2A/2C} agonists. *ACS Chem. Neurosci.* 5, 243–249.

(46) Braun, U., Shulgin, A. T., Braun, G., and Sargent, T., III. (1977) Synthesis and body distribution of several iodine-131 labeled centrally acting drugs. *J. Med. Chem.* 20, 1543–1546.