



Accepted but uncorrected

Identification and characterization of an imidazolium by-product formed during the synthesis of 4-methylmethcathinone (mephedrone).

Journal:	<i>Drug Testing and Analysis</i>
Manuscript ID:	DTA-15-0016.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	14-Feb-2015
Complete List of Authors:	Power, John; Trinity College, Pharmacology and Therapeutics Kavanagh, Pierce; Trinity College, Pharmacology and Therapeutics Brandt, Simon; Liverpool John Moores University, School of Pharmacy and Pharmaceutical Sciences O'Briern, John; Trinity College, School of Chemistry Talbot, Brian; Trinity College, School of Pharmacy and Pharmaceutical Sciences Dowling, Geraldine; State Labotatory, Barry, Michael; Trinity College, Pharmacology and Therapeutics Twamley, Brian; Trinity College, School of Chemistry McLaughlin, Gavin; Trinity College, Pharmacology and Therapeutics
Keywords:	forensic, mephedrone, cathinones, synthesis, imidazolium, by-product, TMMPI
Abstract:	<p>Abstract</p> <p>4-Methylmethcathinone (2-methylamino-1-(4-methylphenyl)propan-1-one, mephedrone) is a psychoactive substance that has been associated with recreational use worldwide. Analytical data related to mephedrone are abundantly available but the characterization of by-products obtained during organic synthesis remains to be explored. This study presents the identification of a 1,2,3,5-tetramethyl-4-(4-methylphenyl)-1H-imidazol-3-ium salt (TMMPI), which was formed during the synthesis of mephedrone. When diethyl ether was added to the crude reaction product, solid material precipitated from the solution. Analytical characterization of TMMPI employed a range of analytical techniques including chromatographic analysis in combination with various mass spectrometric detection methods, nuclear magnetic resonance spectroscopy and crystal structure analysis. Additional confirmation was obtained from organic synthesis of the imidazolium by-product. When TMMPI was subjected to analysis by gas chromatography-mass spectrometry (GC-MS), isomerization and degradation into two distinct compounds were observed, which pointed toward thermal instability under GC conditions. An LC-MS based investigation into a micro scale synthesis of mephedrone and three additional analogues revealed that the corresponding TMMPI analogue was formed. Interestingly, storage of mephedrone freebase in a number of</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	organic solvents also gave rise to TMMPI and it appeared that its formation during storage was significantly reduced in the absence of air. The present study was aimed to support clandestine forensic investigations by employing analytical strategies that are applicable to manufacturing sites. The imidazolium salts will most likely be found amongst the waste products of any clandestine lab site under investigation rather than with the desired product.

SCHOLARONE™
Manuscripts

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

**Identification and characterization of
an imidazolium by-product formed during
the synthesis of 4-methylmethcathinone (mephedrone).**

John D. Power,^{a, b,*} Pierce Kavanagh,^a Gavin McLaughlin,^a John O'Brien,^c Brian Talbot,^d Michael Barry,^a Brendan Twamley,^c Geraldine Dowling^c and Simon D. Brandt^f

- a. Department of Pharmacology and Therapeutics, School of Medicine, Trinity Centre for Health Sciences, St. James Hospital, Dublin 8*
- b. Forensic Science Ireland, Garda HQ, Dublin 8, Ireland*
- c. School of Chemistry, Trinity College, Dublin 2.*
- d. School of Pharmacy and Pharmaceutical Sciences, Trinity College Dublin 2, Ireland.*
- e. State Laboratory, Backweston Campus, Co.Kildare, Ireland.*
- f. School of Pharmacy and Biomolecular Sciences, Liverpool John Moores University, Byrom Street, L3 3AF, Liverpool, UK*

**Corresponding author. E-Mail: jdpower@tcd.ie, phone: +353 879793483*

Abstract

4-Methylmethcathinone (2-methylamino-1-(4-methylphenyl)propan-1-one, mephedrone) is a psychoactive substance that has been associated with recreational use worldwide. Analytical data related to mephedrone are abundantly available but the characterization of by-products obtained during organic synthesis remains to be explored. This study presents the identification of a 1,2,3,5-tetramethyl-4-(4-methylphenyl)-1*H*-imidazol-3-ium salt (TMMPI), which was formed during the synthesis of mephedrone. When diethyl ether was added to the crude reaction product, solid material precipitated from the solution. Analytical characterization of TMMPI employed a range of analytical techniques including chromatographic analysis in combination with various mass spectrometric detection methods, nuclear magnetic resonance spectroscopy and crystal structure analysis. Additional confirmation was obtained from organic synthesis of the imidazolium by-product. When TMMPI was subjected to analysis by gas chromatography-mass spectrometry (GC-MS), isomerization and degradation into two distinct compounds were observed, which pointed toward thermal instability under GC conditions. An LC-MS based investigation into a micro scale synthesis of mephedrone and three additional analogues revealed that the corresponding TMMPI analogue was formed. Interestingly, storage of mephedrone freebase in a number of organic solvents also gave rise to TMMPI and it appeared that its formation during storage was significantly reduced in the absence of air. The present study was aimed to support clandestine forensic investigations by employing analytical strategies that are applicable to manufacturing sites. The imidazolium salts will most likely be found amongst the waste products of any clandestine lab site under investigation rather than with the desired product.

1
2
3
4 **Keywords:** forensic; mephedrone; cathinones; synthesis; imidazolium, by-product;
5
6 TMMPI.
7
8
9

10 **Introduction**

11
12 The cathinone molecular scaffold gives rise to a large range of biologically active
13 compounds and a considerable number of substituted cathinones has gained
14 popularity amongst drug users. These substances were often accessible to the general
15 public, for example, via specialized street shops (e.g. head shops/smart shops), and
16 the Internet that included retailers operating in the ‘surface’ net or hidden ‘dark’ net
17 environment. The rapid emergence of these largely un-tested new psychoactive
18 substances (NPS) has raised concerns worldwide from emergency personnel,
19 clinicians, policy makers, law enforcement and the general public.^[1,2,3] Amongst the
20 more frequently encountered substituted cathinones, which showed some form of
21 diffusion into specific subpopulations was 4-methylmethcathinone (mephedrone).
22
23 When mephedrone was controlled under the Irish misuse of drugs legislation, it was
24 encountered on the Irish illicit market where it was typically sold either as a powder
25 purporting to be cocaine or in a tablet form purporting to represent 3,4-
26 methylenedioxymethamphetamine (MDMA).^[4,5]
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 The synthesis of mephedrone was first published in 1929 by Saem de Burnaga
49 Sanchez.^[6] The route involved the bromination of 4-methylpropiophenone to yield 4-
50 methyl-2-bromopropiophenone which, when reacted with methylamine hydrochloride
51 and triethylamine as an acidic scavenger, gave mephedrone. The reaction was
52 quenched with hydrogen chloride (aqueous or gaseous) that provided the racemic
53 hydrochloride mephedrone salt. A Friedel-Crafts acylation route, starting with toluene
54
55
56
57
58
59
60

1
2
3
4 and propanoyl chloride in the presence of a suitable Lewis acid as a catalyst, may also
5
6 be used to produce the 4-methylpropiophenone intermediate.^[7] Another potential
7
8 route to mephedrone synthesis involves the oxidation of 4-methylephedrine in a
9
10 simple one step reaction process or a Grignard reaction using 4-methylbenzaldehyde
11
12 as the starting material.^[8]
13
14
15
16

17 Detailed published information on possible synthetic mephedrone impurities and
18
19 route-specific by-products are currently unavailable although it has been recognized
20
21 that the formation of iso-cathinones may be relevant for the chemistry of synthetic
22
23 cathinones.^[9-11] Mephedrone pyrolysis has also been reported to yield a range of
24
25 degradation products.^[12] In addition, GC-based heat-induced formation of artifacts or
26
27 decomposition products may be observed during the analysis of substituted
28
29 cathinones, which deserves further study.^[13-18]
30
31
32
33
34
35

36 As part of the authors' ongoing collective research efforts regarding the systematic
37
38 preparation and forensically relevant characterization of targeted NPS, the present
39
40 study reports on the formation of a by-product during mephedrone synthesis. During a
41
42 series of experiments involving the use of a 33% methylamine solution and a number
43
44 of different solvents containing the α -bromo-4-methylpropiophenone, one
45
46 experiment included the addition of diethyl ether to the crude mephedrone product (in
47
48 preparation for forming a mephedrone hydrochloride salt). A solid product, however,
49
50 was observed to precipitate from the solution, initially, this solid was presumed to be
51
52 mephedrone freebase and accordingly filtered, dried and stored in a freezer for further
53
54 work.
55
56
57
58
59
60

1
2
3 This product was subsequently identified as a 1,2,3,5-tetramethyl-4-(4-
4 methylphenyl)-1*H*-imidazol-3-ium salt (TMMPI). Analytical characterization
5 included chromatographic techniques coupled with various forms of mass
6 spectrometry, nuclear magnetic resonance spectroscopy, crystal structure analysis and
7 additional confirmation by synthesis of this imidazolium by-product. The presence of
8 TMMPI was also observed in the crude reaction mixtures of subsequent syntheses
9 along with mephedrone and a mephedrone dimer. Given the polar salt nature of
10 TMMPI, the data presented is also aimed to inform forensic investigators and
11 researchers involved in the examination of waste products derived from clandestine
12 manufacturing facilities suspected of synthesizing substituted cathinones.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

33 **Experimental**

34 *Reagents and standards*

35 All reagents and dry solvents used in the syntheses were obtained from Sigma Aldrich
36 Ltd. (Arklow, Ireland). LC-MS grade solvents were obtained from Fisher Scientific
37 (Dublin, Ireland).
38
39
40
41
42
43
44
45

46 *Isolation of the imidazolium by-product from a mephedrone reaction mixture.*

47
48
49
50 A solution of α -bromo-4-methylpropiophenone (5.68 g, 25 mmol) and ethanolic
51 methylamine (6.25 mL, 8 M, 50 mmol) in absolute ethanol (50 mL) was stirred
52 overnight at room temperature. The mixture was evaporated to a small volume under
53 vacuum at room temperature and diethyl ether (150 mL) was added. After vigorous
54 mixing, the mixture was filtered. The residue was extracted with acetonitrile (30 mL)
55
56
57
58
59
60

1
2
3
4 and filtered. The filtrate was evaporated to a small volume to afford a precipitate. The
5
6 remaining solvent was removed with a pipette and the solid was washed with
7
8 acetonitrile (3 x 1 mL). The residue was dissolved in dichloromethane (15 mL),
9
10 filtered and the filtrate was evaporated to dryness under vacuum to give a colorless
11
12 solid (172 mg). A sample was recrystallized (slow evaporation) from dichloromethane
13
14 to afford colorless crystals (m. pt. 224 - 226 °C uncorrected): ¹H NMR (DMSO) δ
15
16 7.42 (d, *J*=7.5 Hz, 2H, CH Ar), 7.42 (d, *J*=7.5 Hz, 2H, CH Ar), 3.74 (s, 3H, CH₃),
17
18 3.74 (s, 3H, CH₃), 2.68 (s, 3H, CH₃), 2.42 (s, 3H, CH₃), 2.22 (s, 3H, CH₃);
19
20
21 ¹³C NMR (d₆DMSO) δ 144.2 (C imidazolium), 140.1 (C phenyl), 130.7 (C phenyl),
22
23 129.5 (C imidazolium), 129.3 (C phenyl), 123.4 (C phenyl), 33.2 (C imidazolium),
24
25 32.6 (C imidazolium), 21.4 (C phenyl methyl), 10.5 (C imidazolium), 9.2 (C
26
27 imidazolium. HR-ESIMS found 215.1548 (theory M⁺: C₁₄H₁₉N₂, 215.1548)
28
29
30
31
32
33
34
35
36
37
38
39
40

1,2,3,5-Tetramethyl-4-(4-methylphenyl)-1H-imidazol-3-ium iodide synthesis

41
42 A mixture of α-bromo-4-methylpropiophenone (1.22 g, 5.4 mmol), acetamidine
43
44 hydrochloride (1.04 g, 10.9 mmol) and anhydrous potassium carbonate (2.20 g) in
45
46 acetonitrile (120 mL) was refluxed for 15 hr. The mixture was cooled, filtered and
47
48 evaporated to dryness to afford a light yellow powder (729 mg), which was dissolved
49
50 in DMF (3 mL). A suspension of sodium hydride (115 mg, 4.8 mmol) in DMF (3 mL)
51
52 was added and the mixture was stirred at room temperature for 2 h. Iodomethane (2
53
54 mL) was added and the mixture was stirred for 3 h at room temperature. More
55
56 iodomethane (2 mL) was added and the mixture was stirred overnight at room
57
58 temperature. The mixture was diluted with acetonitrile (100 mL), filtered and the
59
60

1
2
3
4 volatiles were removed under vacuum. The residue was dissolved in diethyl ether
5
6 (100 mL) and the precipitated solid was collected by filtration. The solid was mixed
7
8 with dichloromethane (20 mL), filtered and the filtrate was concentrated to give a
9
10 light yellow solid (234 mg). This was recrystallized from dichloromethane/*tert*-butyl
11
12 methyl ether to afford a colorless solid (92 mg, m.p. 220 - 222 °C uncorrected): ¹H
13
14 NMR (DMSO) δ 7.40 (d, *J*=7.5 Hz, 2H, CH Ar), 7.34 (d, *J*=7.5 Hz, 2H, CH Ar),
15
16 3.73 (s, 3H, CH₃), 3.56 (s, 3H, CH₃), 2.68 (s, 3H, CH₃), 2.40 (s, 3H, CH₃), 2.21 (s,
17
18 3H, CH₃) ; ¹³C NMR (d₆ DMSO) δ 143.6 (C imidazolium), 139.5 (C phenyl), 130.2
19
20 (C phenyl), 129.7 (C imidazolium), 129.0 (C phenyl), 126.6 (C phenyl), 33.7 (C
21
22 imidazolium), 31.9 (C imidazolium), 20.9 (C phenyl methyl), 10.1 (C imidazolium),
23
24 9.7 (C imidazolium. HR-ESIMS found 215.1548 (theory M⁺: C₁₄H₁₉N₂, 215.1543)
25
26
27
28
29
30
31
32
33

34 ***Micro scale reactions for LC-MS analysis***

35
36 A solution of the α-bromo ketone (0.25 mmol) and the amine (0.5 mmol) in the
37
38 specified solvent (500 μL) was allowed to stand at room temperature overnight. This
39
40 solution was diluted with acetonitrile/water (1/1, containing 0.1 % formic acid, 1.5
41
42 mL), an aliquot (8 μL) was further diluted with acetonitrile/water (1/1, containing
43
44 0.1 % formic acid, 1 mL) and analyzed by LC-MS.
45
46
47
48
49

50 ***Micro scale formation of TMMPI from freebase mephedrone***

51
52 A solution of freebase mephedrone in acetonitrile (0.5 mmol/500 μL acetonitrile, 100
53
54 μL) and acetonitrile (100 μL) was allowed to stand overnight (in the presence or
55
56 absence of air) at room temperature. This solution was diluted with acetonitrile/water
57
58 (1/1, containing 0.1 % formic acid, 600 μL), an aliquot (8 μL) was further diluted
59
60

1
2
3 with acetonitrile/water (1/1, containing 0.1 % formic acid, 1 mL) and analyzed by
4
5 LC-MS. The by-product formation was observed in the presence of air, but very
6
7 significantly, this was observed to drop by over 90 % in the absence of air.
8
9

10 11 12 ***Preparative formation of TMPPI from mephedrone freebase.***

13
14 A solution of mephedrone free base (prepared from 2.15 g of the HCl salt, 10 mmol)
15
16 was dissolved in acetonitrile and stirred at room temperature for 48 h. The solution
17
18 was evaporated to dryness and the residual brown oil was washed with diethyl ether
19
20 (4 x 10 mL). The residue was dissolved in dichloromethane and hydrogen chloride in
21
22 2-propanol (5-6 M) was added. The mixture was concentrated to a small volume
23
24 under vacuum, the supernatant was removed and drying was continued under vacuum
25
26 to afford a light brown solid (219 mg). The supernatant was concentrated to give a
27
28 brown oil (178 mg) which was analyzed by NMR and also used for further
29
30 chromatographic analysis.
31
32
33
34
35
36
37
38

39 ***Preparation of crude mephedrone hydrochloride.***

40
41
42 A solution of α -bromo-4-methylpropiofenone (1.13 g, 5 mmol) and ethanolic
43
44 methylamine (1.25 mL, 8 M, 10 mmol) in absolute ethanol (10 mL) was stirred
45
46 overnight at room temperature. The mixture was evaporated to dryness and, taken up
47
48 in dichloromethane and a solution of hydrogen chloride in diethyl ether (2 M) was
49
50 added. Removal of the volatiles afforded a brown powder (1.41 g)
51
52
53
54
55
56
57

58 ***Instrumental analysis***

59 ***Gas chromatography - mass spectrometry***

60

1
2
3
4 Samples were analyzed on an Agilent 6890N gas chromatograph coupled to a 5973
5
6 MSD. A HP-5MS column (30 m × 0.25 mm × 0.25 μm) was used in splitless mode
7
8 with helium carrier gas at a constant flow of 1 mL/min. The injection port and transfer
9
10 line temperatures were set at 250 °C and 280 °C, respectively. The initial oven
11
12 temperature was 90 °C, held for 1 min, then ramped at 15 °C/min to 280 °C and held
13
14 for 5 min, followed by a ramp of 20 °C/min to 300 °C with a final hold time of 1 min.
15
16 The mass spectra data was collected after a 5 minute solvent delay. The ionization
17
18 energy was set at 70 eV and the mass range was m/z 40-550.
19
20
21
22

23 24 *Liquid chromatography - mass spectrometry*

25
26 LC-MS analyses, equipped with an electrospray ionization source, were performed on
27
28 an Agilent 1100 LC system. The column (Allure PFP Propyl, 5 μm, 50 × 2.1 mm)
29
30 was from Restek (Bellefonte, PA, USA) and the aqueous mobile phase A consisted of
31
32 0.05% formic acid in water whereas mobile phase B was prepared from 0.05% formic
33
34 acid in acetonitrile, respectively. The Agilent LC-MSD settings were as follows:
35
36 positive electrospray mode, capillary voltage 3000 V, drying gas (N₂) 12 L/min at
37
38 350 °C, nebulizer gas (N₂) pressure 60 psi, m/z 50-500, fragmentor voltage either 50
39
40 V or 120 V. Samples for LC-MS analysis (2 μL injection volume) were dissolved in
41
42 acetonitrile/water (1:1, containing 0.1% formic acid) at a concentration of 5 μg/mL.
43
44 The following gradient elution program was used: 0-5 min 12% A and then increased
45
46 to 35% over 30 min using a linear gradient. The flow rate was 1 mL/min and the
47
48 column temperature was 30 °C.
49
50
51
52
53
54
55
56
57
58
59
60

Nuclear magnetic resonance spectroscopy

1
2
3 Samples were prepared in DMSO- d_6 . ^1H (600 MHz) and ^{13}C (150 MHz) NMR spectra
4
5 were recorded on a Bruker AV600 NMR spectrometer using a 5 mm TCI cryoprobe.
6
7
8 ^1H NMR spectra were referenced to an external TMS reference at $\delta = 0$ ppm.
9

10 11 12 *High-resolution electrospray mass spectrometry*

13
14
15
16 HR-ESI mass spectra were recorded by direct injection into a LTQ Orbitrap
17
18 Discovery (Thermo Fisher, UK). Samples were dissolved in acetonitrile/water (1:1,
19
20 containing 0.1% formic acid) and infused at a rate of 5 $\mu\text{L}/\text{min}$. Full accurate high-
21
22 resolution (30000) mass scans were performed in positive electrospray mode.
23
24 Measured accurate masses were within ± 5 ppm of the theoretical masses. The
25
26 following conditions were used: drying gas (N_2) 10000 mL/min, capillary temperature
27
28 310 $^\circ\text{C}$, spray voltage 4 V, capillary voltage 22 V and tube lens 77 V.
29
30
31

32 33 *X-ray crystallography*

34
35
36
37 Data for the by-product $\text{C}_{14}\text{H}_{19}\text{BrN}_2$ (colourless needle, 0.060 mm x 0.090 mm
38
39 x 0.220 mm) were collected on a Bruker APEX DUO using Mo $\text{K}\alpha$ radiation ($\lambda =$
40
41 0.71073 \AA) using a MiTeGen micromount and at 100(2) K (Oxford Cobra
42
43 Cryosystem). Bruker APEX2 software was used to collect and reduce data, determine
44
45 the space group, solve and refine the structure.^[19] A total of 2942 frames were
46
47 collected. The total exposure time was 12.26 hours. Absorption corrections were
48
49 applied using SADABS 2012.^[20] All final refinements were performed with SHELXL
50
51 2014/3.^[21] All non-hydrogen atoms were refined anisotropically. Hydrogen atoms
52
53
54
55
56 were assigned to calculated positions using a riding model.
57
58
59
60

1
2
3
4 **Crystal Data:** C₁₄H₁₉BrN₂, *M* = 295.22, Monoclinic, *a* = 11.2256(3), *b* = 7.4874(2), *c*
5 = 16.0597(5) Å, β = 99.3230(10)°, *U* = 1332.00(7) Å³, *T* = 100 K, space group P2₁/c,
6 *Z* = 4, μ (Mo Kα) = 3.067 mm⁻¹, ρ = 1.472 Mg/cm³, 86862 reflections collected, 4895
7 independent (*R*_{int} = 0.0263), ^a*R*_I = 0.0205, w*R*₂ = 0.0517 (*I* > 2σ(*I*)), *S* = 1.045.

8
9
10
11
12
13
$$^aR_1 = \frac{\sum ||F_o| - |F_c||}{\sum |F_o|}, wR_2 = [\frac{\sum w(F_o^2 - F_c^2)^2}{\sum w(F_o^2)^2}]^{1/2}.$$

14 (CCDC 1033669 contains the supplementary crystallographic data for this paper.

15
16
17 These data can be obtained free of charge from The Cambridge Crystallographic Data
18
19
20 Centre via www.ccdc.cam.ac.uk/data_request/cif.)
21
22

23 24 25 26 27 **Results and discussion**

28
29
30
31 The analytical characterization of synthetic cathinones is a relatively recent research
32
33 field. Typically, the unique determination of substances in most forensic and law
34
35 enforcement laboratories rely on various separation technologies linked to mass
36
37 spectrometric detection. In the analysis of submitted samples where cathinone
38
39 structures are suspected, particular attention is paid to the base peak that is generally
40
41 formed from whatever side chain substitution pattern is present in the cathinone. The
42
43 molecular ion for substituted cathinones in EI mass spectrometry (70 eV) may not be
44
45 easily identifiable; hence molecular weights may not easily be ascertained. Typical
46
47 cathinone fragmentation patterns are dominated by alpha-cleavage and formation of
48
49 immonium ions. In cases where charge retention is observed following alternative
50
51 alpha cleavage, substituted benzoyl ions are then detected.^[22] Although analytical data
52
53 on mephedrone are abundantly available, published information on synthetic routes,
54
55 possible by-products or possible route specific compounds are not available.
56
57
58
59
60

1
2
3
4 The present investigation reports on the identification and characterization of 1,2,3,5-
5 tetramethyl-4-(4-methylphenyl)-1*H*-imidazol-3-ium salt (TMMPI) obtained as a by-
6 product during the synthesis of mephedrone. As previously stated, when diethyl ether
7 was added to synthesised crude mephedrone product (in preparation of forming a
8 mephedrone salt), a solid product precipitated out of solution that was filtered, dried
9 and stored at -20 °C for further work. Initially, GC-MS analysis was performed on
10 this product and the chromatogram obtained which surprisingly showed two distinct
11 peaks at 15.58 min and 17.54 min, respectively (see supplemental 1a). The GC
12 retention time and mass spectra associated with these two peaks were not consistent
13 with the retention time or mass spectra of the expected synthesis product mephedrone.
14
15 The GC-MS spectra of the synthesis by-product were similar to each other each
16 having a base peak at m/z 200 but some differences were also noted. The
17 fragmentation obtained for the compound at 17.54 min showed the second largest
18 peak at m/z 185 following the loss of a methyl group from the base peak at m/z 200.
19 This m/z 185 was much less abundant in the compound eluting at 15.58 min.
20 Interestingly, a relatively large m/z 132 present in the species at 15.58 min was
21 observed (suggesting loss of a neutral C_4H_6N moiety from the base peak) but this was
22 not detected in the chromatographic peak at 17.54 min. A potential explanation is
23 that the formation of the m/z 132 ion would require the breakage of two double bonds
24 in the imidazolium structure proposed for the by-product, which is energetically less
25 favoured compared to only one double bond breakage that could account for the m/z
26 132 ion formation associated with the compound eluting first. Two interconverting
27 structures could give rise to the mass spectra obtained via loss of either C_4H_6N or
28 C_5H_9N and subsequent rearrangement of the imidazole ring (see supplemental 1b).
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

An LC-MS analysis was performed on the mephedrone synthesis product and the

1
2
3 chromatogram consisted of a single peak, which supported the suggestion that the
4 presence of two GC chromatographic peaks were formed artificially under GC-MS
5 conditions. The electrospray ionization mass spectrum (180 V) showed a base peak at
6 m/z 215 ($C_{14}H_{19}N_2^+$) and the second most abundant product ions was observed at m/z
7 132, presumably due to the loss of C_5H_9N from the base peak and even-electron
8 molecular mass (Figure 1a). Ions observed at m/z 200, m/z 199 and m/z 185 might
9 have reflected the loss of CH_3 , CH_4 and C_2H_6 , respectively. The imidazolium ring
10 may also undergo opening with the loss of C_3H_5N or C_3H_6N to yield m/z 160 and m/z
11 159, and further losses from m/z 159 (H or CH_3) gave m/z 158 and m/z 144
12 respectively. A suggested mechanism is provided in Figure 1b. The difference
13 between the GC-MS base peak and the LC-MS base peak also indicated possible
14 thermal degradation during GC-MS analysis (loss of a methyl moiety) since ions
15 above the m/z 200 base peak in the GC-MS spectrum were not detected.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The solid state crystal structure of the mephedrone by-product was also obtained. The structure of the mephedrone by-product was found to be a bromide salt and shown to consist of a *para*-methyl substituted phenyl ring attached to a tetra-methyl substituted five-membered heterocyclic imidazole ring (Figure 2). The bromine atom was only weakly coordinated to the organic moiety via $CH\dots Br$ interactions (3.75 - 3.98 Å) and each Br^- linked four organic moieties together. The twist between the rings (torsion angle N7-C9-C10-C16, 133.6(1) Å) allowed the alternating packing of the imidazolium/phenyl system into a stack parallel to the b-axis (supplemental 2). This arrangement allowed the rings to overlap and align with an intermolecular centroid (C13-C15-C16-C10-C11-C12) to centroid (N3-C5-N7-C9-C2) distance of ca. 3.6 Å. The characterization of the mephedrone by-product added to the few structurally

1
2
3 known tetramethylimidazolium- $R^+ X^-$ compounds ($R = \text{Me}$, $X = \text{I}$; $R = \text{CH}_2\text{I}$, $X = \text{I}$; R
4
5 = Me , $X = (\text{SCN})_4\text{Crphenanthroline}$; $R = \text{Me}$, $X = (\text{SCN})_4\text{Cr}(\text{NH}_3)_2$; $R = \text{Me}$, $X =$
6
7 $\text{N}(\text{SO}_2\text{CF}_3)_2$).^[22-27]
8
9

10
11
12 A mechanism of formation is proposed to account for the formation of TMMPI during
13
14 mephedrone synthesis (Figure 3). The carbonyl group on mephedrone was thought to
15
16 undergo a reduction to the alcohol followed by dimerization and subsequent loss of
17
18 water. This intermediate in which the carbonyl functional group is directly attached to
19
20 the 4-methylphenyl moiety was proposed to undergo losses of 4-methylbenzaldehyde
21
22 and a hydride ion to yield the imidazolium by-product.
23
24
25

26
27
28 Confirmation of the mechanisms proposed by-products was obtained when a series of
29
30 micro scale experiments were performed using cathinone substituents in which the
31
32 alpha-alkyl group was either H, CH_3 , CH_2CH_3 and $\text{CH}_2\text{CH}_2\text{CH}_3$. The corresponding
33
34 substituted by-products were observed when the reaction solvent was methanol,
35
36 ethanol, acetonitrile and tetrahydrofuran. It is significant and noteworthy that in the
37
38 absence of air during these micro experiments, a considerable reduction $> 90\%$ was
39
40 observed in the formation of TMMPI by-product. This reduction may be due to the
41
42 absence of an efficient mechanism to regenerate the hydride ion in the proposed
43
44 mechanism in the absence of air and hence a corresponding reduction in by-product
45
46 formation. The regeneration of the hydride ion may have some analogous mechanism
47
48 to those found in biological processes. For example, dihydrobenzo-imidazoles have
49
50 been the subject of extensive studies for hydride transfer mechanisms which might
51
52 make the phenyl-1*H*-imidazol-3-ium by-products observed in this study useful as
53
54 potential organic hydride donors.^[28]
55
56
57
58
59
60

1
2
3
4 The results of the micro scale experiments are presented in Figure 4A-D where the
5
6 TMMPI product was shifted in mass based on the nature of the alpha-alkyl group (R)
7
8 during LC-MS analysis, i.e. at m/z values of 187, 215, 243 and 271, respectively.
9
10
11 When R was an ethyl-substituted cathinone, a peak was observed in the expected LC
12
13 window of the R as a methyl substituted cathinone at m/z 215. Co-injection into the
14
15 LC-MS of the CH_3 and CH_2CH_3 micro experiments, clearly show that the second
16
17 peak observed in the m/z 215 target window is an artifact and not the mephedrone by-
18
19 product (see figure 4 B, C. co-injected).
20
21

22
23 Additional confirmation for the unambiguous identification of the mephedrone by-
24
25 product as TMMPI was obtained from its organic synthesis via the reaction of 4-
26
27 methyl-1-phenyl-2-bromo-propan-1-one with acetamidine and potassium carbonate
28
29 (Figure 5). The synthesized compound and the mephedrone by-product was then
30
31 compared using GC-MS, LC-MS and NMR and found to be identical (supplemental
32
33 data 3, NMR data).
34
35
36
37
38
39

40 Analysis of the reaction products derived from in-house synthesis of mephedrone
41
42 showed that the imidazolium by-product was observed in the crude reaction mixture
43
44 along with mephedrone and a mephedrone dimer. GC-MS data clearly showed the
45
46 presence of two peaks at the expected retention times (supplemental data 4a). LC-MS
47
48 data also showed the presence of a single peak at the corresponding retention time
49
50 expected for TMMPI in addition to the expected ESI mass spectrum (supplemental
51
52 data 4b).
53
54
55
56
57
58

59 Mephedrone freebase was placed in acetonitrile and allowed stand for 48 hours,
60
which resulted in the formation of both an oil and a solid. Mephedrone was not

1
2
3
4 detected in both forms but analysis of these reaction products indicated that along
5
6 with the formation of TMMPI an additional compound was detected. Proton NMR of
7
8 the solid indicated the presence of extra peaks in addition to those expected for
9
10 TMMPI, thus suggesting that a para substituted aromatic compound with a methyl
11
12 substituent was also present in the solid (supplemental data 5a).

13
14
15 LC-MS (negative electrospray ionization) analysis of both the oil and solid indicated
16
17 that this additional compound had a deprotonated $[M-H]^-$ m/z 135 (supplemental data
18
19 5b). GC-MS analysis of both forms provided indication that it was consistent with 4-
20
21 methyl benzoic acid (supplemental data 5c). The scheme outlining the formation of
22
23 the TMMPI (Figure 3) showed the loss of 4-methylbenzaldehyde from an
24
25 intermediate and its subsequent oxidation could have been consistent with the
26
27 formation of 4-methyl benzoic acid or it could have been formed by another oxidative
28
29 pathway.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

In the investigation of sites where the manufacture of mephedrone (or other substituted cathinones) is suspected, it is important that identification of substances that can be related to a synthesis route are fully explored. The analytical data presented here will be beneficial to such investigations, as any imidazolium by-product found will give an indication as to what substituted cathinone was synthesized. The imidazolium salts will most likely be found amongst the waste products at a clandestine lab site. It is often the case, that waste products are either dumped in remote locations or are the only items remaining at such illicit synthesis sites when enforcement personnel actually get to examine a premises suspected of having been used in illicit clandestine manufacture. This may be the case were significant by-products are identifiable and associated with a particular class of

1
2
3
4 cathinones. The typical forensic analytical approach taken when examining items
5
6 seized at a suspected clandestine laboratory facility or of suspected waste material
7
8 generated at such a manufacturing site and dumped at a different location is to
9
10 initially screen such samples by GC-MS. If this is the sole screening methodology,
11
12 then challenges may be encountered when attempting to identify the imidazolium
13
14 salts due to GC-induced isomerization and thermal degradation. It is vital that
15
16 analytical strategies utilized for clandestine laboratory investigations are fit for
17
18 purpose and investigators should be aware that some traditional screening
19
20 technologies may have unforeseen limitations.
21
22
23
24
25
26
27

28 **Conclusion**

29
30
31
32
33 The majority of forensic examinations of drug submissions involve samples with
34
35 questionable origin. Verifiable data for samples derived from clandestine
36
37 manufacturing conditions remain illusive unless such operations are found while in
38
39 production and with products being available. The ability to determine a potential link
40
41 to clandestine production with the identification of a by-product linkable to a known
42
43 synthesis route would be a very valuable forensic option. This study explored the
44
45 relationship between a synthetic route commonly used for mephedrone production
46
47 and the formation of a specific imidazolium by-product identified as a 1,2,3,5-
48
49 tetramethyl-4-(4-methylphenyl)-1*H*-imidazol-3-ium salt. Furthermore, it was also
50
51 established that mephedrone freebase was unstable in several organic solvents. Given
52
53 the polar nature of the imidazolium salt formed during a mephedrone synthesis, it is
54
55 conceivable that it might be found amongst the waste products generated at such
56
57
58
59
60

1
2
3 clandestine manufacturing sites or it may be formed post synthesis and as in our
4
5 initial belief presumed to be mephedrone. The present study was aimed to support
6
7 clandestine forensic investigations by targeting specific compounds and the analytical
8
9 strategies which are applicable to the analysis of waste products that are generally
10
11 either dumped or remain abandoned at sites after production has ceased.
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

31 **List of figures**

- 32
33 **Figure 1a:** LC-MS chromatogram of imidazolium by-product.
34
35 **Figure 1b:** LC-MS proposed fragmentation of imidazolium by-product.
36
37 **Figure 2:** X-ray data. Structure of the mephedrone by-product with
38
39 thermal displacement shown at 50%. Hydrogen atoms drawn as
40
41 arbitrary spheres and are unlabelled for clarity.
42
43
44 **Figure 3:** General scheme for the formation of imidazolium by-
45
46 products during the synthesis of substituted cathinones.
47
48
49 **Figure 4:** Microscale LC-MS single ion chromatograms, A,B,C,D and
50
51 BC co-injected.
52
53
54 **Figure 5:** Synthetic route to 1,2,3,5-tetramethyl-4-(4-methylphenyl)-
55
56 1H-imidazol-3-ium (TMMPI)
57
58
59
60

1
2
3
4
5
6
7
8
9
10 **Supplemental data**
11
12
13
14


- 15 **1A GC-MS of imidazolium by-product.**
16
17 **1B Two interconverting structures for TMMPI in GC inj port**
18
19
20 **2 X-Ray crystallography supplemental 2.**
21
22 2. Packing diagram of the mephedrone by-product viewed
23 down the b-axis. (A) hydrogen atoms shown with dashed lines
24 indicating weak CH...Br interactions.(B) Hydrogen atoms
25 removed for clarity to show the alternating head-to-tail packing
26 assembly of the organic moiety.
27
28
29
30
31 **3 TMPPI synthesized and the mephedrone by-product NMR**
32 **4 Crude mephedrone mixture analysis by GC-MS (4a), LC-**
33 **MS (4b)**
34
35
36
37 **5 Mephedrone freebase mixture analysis by NMR (5a), LC-**
38 **MS (5b) and GC-MS (5c)**
39
40
41
42
43
44
45
46
47

48 **References:**
49
50

- 51 [1] S.D. Brandt, L.A. King, M. Evans-Brown. The new drug phenomenon. *Drug*
52 *Test. Anal.* **2014**, *6*, 587.
53 [2] United Nations Office on Drugs and Crime (UNODC). The challenge of new
54 psychoactive substances. A Report from the Global SMART Programme
55 March **2013**. United Nations Publication, Vienna, 2013.
56 [3] Vardakou I, Pistos C, Spiliopoulou Ch. Drugs for youth via Internet and the
57 example of mephedrone. *Toxicol Lett.* **2011**, *201*, *3*, 191.
58
59
60

- 1
2
3
4 [4] K.McElrath, C.O'Neill. Experiences with mephedrone pre and post
5 legislative controls, perceptions of safety and sources of supply.
6 *International Journal of Drug Policy*, **2011**, 22, 2, 120.
- 7 [5] M.C.VanHout, R Brennan. Plant food for thought A quantitative study of
8 mephedrone in Ireland. *Drugs, Education, Prevention and policy*, **2011**,
9 18, 5, 371.
- 10 [6] M.J. Saem de Burnaga Sanchez. Sur un homologue de l'éphédrine. *Bull. Soc.*
11 *Chim. Fr.* **1929**, 45, 284.
- 12 [7] P. Tamilselvan, Y. B. Basavaraju, R. Murugesan, E. Sampathkumar.
13 Cobalt(II) acetylacetonate catalyzed FriedelCrafts acylation of anisole,
14 thioanisole, and toluene. *Catalysis Communications*, **2008**, 10, 3, 300.
- 15 [8] J.D. Power, P. McGlynn, K. Clarke, S.D. McDermott, P. Kavanagh, J. O'Brien.
16 The analysis of substituted cathinones. Part 1: chemical analysis of 2-, 3-
17 and 4-methylmethcathinone. *Forensic Sci. Int.* **2011**, 212, 6.
- 18 [9] S.D. Brandt, S. Freeman, H.R. Sumnall, F. Measham, J. Cole. Analysis of NRG
19 'legal highs' in the UK: identification and formation of novel cathinones.
20 *Drug Test. Anal.* **2011**, 3, 569.
- 21 [10] F. Westphal, T. Junge, U. Girreser, W. Greibl, C. Doering. Mass, NMR and IR
22 spectroscopic characterization of pentedrone and pentylone and
23 identification of their isocathinone by-products. *Forensic Sci. Int.* **2012**,
24 217, 157.
- 25 [11] S.D. McDermott, J.D. Power, P. Kavanagh, J. O'Brien. The analysis of
26 substituted cathinones. Part 2: an investigation into the phenylacetone
27 based isomers of 4-methylmethcathinone and N-ethylcathinone. *Forensic*
28 *Sci. Int.* **2011**, 212, 13.
- 29 [12] P. Kavanagh, J. O'Brien, J.D. Power, B. Talbot, S.D. McDermott. 'Smoking'
30 mephedrone: the identification of the pyrolysis products of 4-
31 methylmethcathinone hydrochloride. *Drug Test. Anal.* **2013**, 5, 291.
- 32 [13] R.P. Archer. Fluoromethcathinone, a new substance of abuse. *Forensic*
33 *Sci.Int.* **2009**, 185, 10.
- 34 [14] S.D. Brandt, P.F. Daley, N.V. Cozzi. Analytical characterization of three
35 trifluoromethyl-substituted methcathinone isomers. *Drug Test. Anal.*
36 **2012**, 4, 525.
- 37 [15] P. Jacob III, A.T. Shulgin, Preparation of novel N-substituted-2-amino-3',4'-
38 methylenedioxypropiofenones as anti-depressant and anti-
39 Parkinsonism agents. US Patent No. WO9639133(A1), **1996**.
- 40 [16] S.D. Brandt, R.C.R. Wootton, G. De Paoli, S. Freeman. The naphyrone story:
41 the alpha or beta-naphthyl isomer? *Drug Test. Anal.* **2010**, 2, 496.
- 42 [17] Noggle FT, DeRuiter J, Valaer A, Clark CR. GC-MS analysis of
43 methcathinone and its major decomposition product. *Microgram* **1994**,
44 27, 4, 106.
- 45 [18] Tsujikawa K, Mikuma T, Kuwayama K, Miyaguchi H, Kanamori T, Iwata
46 YT, Inoue H. Degradation pathways of 4-methylmethcathinone in alkaline
47 solution and stability of methcathinone analogs in various pH solutions.
48 *Forensic Sci Int.* **2012**, 10; 220, 1-3,103.
- 49 [19] Bruker APEX v2012.12-0, Bruker AXS Inc., Madison, Wisconsin, USA.
- 50 [20] SADABS (2012) Bruker AXS Inc., Madison, Wisconsin, USA; Sheldrick, G.
51 M. University of Göttingen, Germany.
- 52 [21] Sheldrick, G. M., SHELXL 2014/3, University of Göttingen, Germany, 2014.
- 53
54
55
56
57
58
59
60


- 1
2
3
4 [22] Power, J., McDermott, S., Talbot, B., O'Brien, J., Kavanagh P. The
5 analysis of amphetamine-like cathinone derivatives using positive
6 electrospray ionization with in-source collision-induced
7 dissociation. *R.C.M.*, **2012**, 30;26, 22, 2601.
8
9 [23] Kuhn, N., Henkel, G., Kreutzberg, J. *Z. Naturforsch., B: Chem. Sci.*, **1991**, 46,
10 1706.
11 [24] Kuhn, N., Gohner, M., Steimann, M. *Z. Anorg. Allg. Chem.*, **2001**, 627, 2048.
12 [25] Peppel, T., Thiele, P., Kockerling, M. *Russ. J. Coord. Chem.*, **2012**, 38, 207.
13 [26] Peppel, T., Schmidt, C., Kockerling, M. *Z. Anorg. Allg. Chem.*, **2011**, 637
14 1314.
15 [27] Roth, C., Peppel, T., Fumino, K., Kockerling, M., Ludwig, R. *Angew. Chem.,*
16 *Int. Ed.* **2010**, 49, 10221.
17 [28] Xiao-Qing, Z., Ming-Tian, Z., Ao, Y., Chun-Hua, W., Jin-Pei, C. Hydride,
18 hydrogen atom, proton, and electron transfer driving forces of various five-
19 membered heterocyclic organic hydrides and their reaction intermediates in
20 acetonitrile. *JACS* 03, **2008**; 130, 8, 2501.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




254x190mm (300 x 300 DPI)

review


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



271x184mm (300 x 300 DPI)





145x102mm (300 x 300 DPI)




259x169mm (300 x 300 DPI)

Review




166x223mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



166x227mm (300 x 300 DPI)


B and C co-injected



166x89mm (300 x 300 DPI)

Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



262x47mm (300 x 300 DPI)

For Peer Review