

**Beilschlabrines A and B: Two new bioactive phenanthrene alkaloids from the stem bark of *Beilschmiedia glabra***

Wan Mohd Nuzul Hakimi Wan Salleh<sup>a</sup>, Farediah Ahmad<sup>a,\*</sup>, Khong Heng Yen<sup>b</sup>, Razauden Mohamed Zulkifli<sup>c</sup>, Jih-Jung Chen<sup>d</sup>, Lutfun Nahar<sup>e</sup>, Jean Duplex Wansi<sup>e</sup>, Satyajit Dey Sarker<sup>e</sup>

<sup>a</sup>*Department of Chemistry, Faculty of Science, Universiti Teknologi Malaysia (UTM), 81310 Skudai, Johor, Malaysia*

<sup>b</sup>*School of Chemistry and Environment Studies, Faculty of Applied Sciences, Universiti Teknologi MARA (UiTM) Sarawak, Jalan Meranek, 94300 Kota Samarahan, Sarawak, Malaysia*

<sup>c</sup>*Department of Bioscience and Health Sciences, Faculty of Biosciences and Medical Engineering, Universiti Teknologi Malaysia (UTM), 81310 Skudai, Johor, Malaysia*

<sup>d</sup>*Department of Pharmacy & Graduate Institute of Pharmaceutical Technology, Tajen University, Pingtung 907, Taiwan*

<sup>e</sup>*Medicinal Chemistry and Natural Products Research Group, School of Pharmacy and Biomolecular Sciences, Faculty of Science, Liverpool John Moores University, James Parsons Building, Byrom Street, Liverpool L3 3AF, United Kingdom*

\*Corresponding author: [farediah@kimia.fs.utm.my](mailto:farediah@kimia.fs.utm.my)

Tel: +607-5534137; Fax: +607-5566162

## ABSTRACT

Two new phenanthrene alkaloids, beilschglabrin A (**1**) and B (**2**) were isolated from the stem bark of *Beilschmiedia glabra*, together with lupeol, taraxerol, and 24-methylenelanosta-7,9-diene-3 $\beta$ -15 $\alpha$ -diol. The structures of the isolated compounds were elucidated by extensive spectroscopic data analysis and comparison with respective literature data. The compounds were tested for DPPH radical scavenging, acetylcholinesterase and lipoxygenase inhibitory activities. Compound **1** displayed considerable activity in the acetylcholinesterase (IC<sub>50</sub> 50.4  $\mu$ M), the DPPH radical scavenging (IC<sub>50</sub> 115.9  $\mu$ M) and the lipoxygenase (IC<sub>50</sub> 32.8  $\mu$ M) assays.

*Keywords:*

*Beilschmiedia glabra*

Lauraceae

Phenanthrene alkaloid

DPPH radical scavenging activity

Acetylcholinesterase

Anti-inflammatory

## 1. Introduction

*Beilschmiedia* is a genus of the family Lauraceae with 250 species, which are distributed in tropical and subtropical Australia, New Zealand, North America, South America and Asia (Nishida, 1999). The chemical analyses of *Beilschmiedia* species have revealed the presence of alkaloids, endiandric acid derivatives, essential oils, fatty acids, epoxyfuranoid lignans, flavonoids and terpenoids, which have been shown to possess anti-oxidant, antibacterial, cytotoxic, antimalarial and antitubercular activities (Chen et al., 2007; Lenta et al., 2009; Chouna et al., 2010; Salleh et al., 2016a).

*Beilschmiedia glabra* Kosterm, locally known as *kayau temblouh* or *kayuh tefuluh*, is distributed in Peninsular Malaysia, Sumatra and Borneo. It can be found in mixed dipterocarp forests up to 600 m altitude, often along rivers. In secondary forests, it is usually present as a pre-disturbance remnant tree. The leaves are alternate to opposite, simple, penni-veined and glabrous, while the fruits are rusty brown and fleshy drupes (Nishida, 2008). We have recently reported the chemical composition and biological activity of the leaf and bark oil (Salleh et al., 2015). The leaf and bark oil contained mainly  $\beta$ -eudesmol (15.4-19.3%) and  $\beta$ -selinene (12.2-16.9%). The leaf oil had the highest phenolic content (233.4 mg GA/g), while the bark oil showed potent activity in the  $\beta$ -carotene/linoleic acid bleaching assay (77.6%). Both the leaf and bark oils were active against *Candida glabrata* and *Saccharomyces cerevisiae* with MIC values of 31.3 and 62.5  $\mu$ g/mL, respectively. Tyrosinase (leaf 73.7%; bark 76.0%) and acetylcholinesterase (leaf 48.1%; bark 45.2%) inhibitory activities, and anti-inflammatory (leaf 59.7%; bark 48.9%) property were also identified (Salleh et al., 2015).

In continuation of phytochemical and bioactivity studies on this genus (Salleh et al., 2016b), herein we report the isolation, identification of phenanthrene alkaloids, named beilschglabrines A (1) and B (2) (Fig. 1) from the stem bark of *B. glabra* and assessment of

their potential free radical scavenging, acetylcholinesterase inhibitory and anti-inflammatory activities.

## 2. Results and discussion

Compound **1** was obtained as a yellow amorphous powder. The HREIMS spectrum revealed the molecular ion peak at  $m/z$  421.1517 (calculated for 421.1525) and the molecular formula was deduced as  $C_{24}H_{23}O_6N$ . The IR spectrum showed the absorption bands at 3367, 1637, 1600 and  $1462\text{ cm}^{-1}$  ascribable to hydroxyl, conjugated double bonds, and aromatic functional groups, respectively, while the peak assignable to the carbonyl group of an aldehyde functionality was observed at  $1727\text{ cm}^{-1}$ . The UV spectrum exhibited absorption bands at 212 and 260 nm, typical of a phenanthrene skeleton (Castro et al., 1986). The  $^1\text{H}$  NMR spectrum (**Table 1**) showed signals for two AB systems for four aromatic protons, at  $\delta_{\text{H}}$  7.58 (1H,  $J = 9.2$  Hz), 8.03 (1H,  $J = 9.2$  Hz), 6.97 (1H,  $J = 4.0$  Hz) and 6.20 (1H,  $J = 4.0$  Hz), corresponding to the aromatic protons of H-9, H-10, H-2' and H-3', respectively. Three singlets resonating at  $\delta_{\text{H}}$  8.96 (1H), 7.36 (1H) and 7.14 (1H) were assigned to aromatic protons H-5, H-8, and H-2, respectively. In addition, each of the two triplets centered at  $\delta_{\text{H}}$  3.51 ( $J = 8.4$  Hz) and 4.67 ( $J = 7.6$  Hz) integrating for two protons were assigned to H-11 and H-12, respectively, and those are typical signals for a phenanthrene alkaloid (Shamma, 1972). A singlet resonating at  $\delta_{\text{H}}$  4.35 (1H) was attributed to H-5' of a cyclopenta-1,3-diene group. Two sharp singlets appeared at  $\delta_{\text{H}}$  3.87 (3H) and 4.11 (3H), and were assigned to two methoxyl groups. The presence of hydroxyl groups was revealed by a broad signal at  $\delta_{\text{H}}$  6.02. Besides, a singlet at  $\delta_{\text{H}}$  9.65 was characteristic for the proton of an aldehyde group. The COSY spectrum exhibited correlations between H-2' ( $\delta_{\text{H}}$  6.97) and H-3' ( $\delta_{\text{H}}$  6.20), and between H-9 ( $\delta_{\text{H}}$  7.58) and H-10 ( $\delta_{\text{H}}$  8.03). Furthermore, H-12 ( $\delta_{\text{H}}$  4.67) was found to correlate with H-11 ( $\delta_{\text{H}}$  3.51). The  $^{13}\text{C}$  NMR spectrum exhibited 24 signals corresponding to twenty four carbons. The DEPT spectrum showed the presence of two

methoxyls, seven methines, three methylenes, one carbonyl and eleven quaternary carbons. The HMBC long-range correlations (**Fig. 2**) furnished correlations of H-9 ( $\delta_{\text{H}}$  7.58) to C-8/C-8a/C-4b/C-10a; H-10 ( $\delta_{\text{H}}$  8.03) to C-4b/C-4a/C-10a/C-1'; H-5 ( $\delta_{\text{H}}$  8.97) to C-4b/C-4a/C-8a/C-7/C-6; H-8 ( $\delta_{\text{H}}$  7.36) to C-4b/C-7/C-6; H-2 ( $\delta_{\text{H}}$  7.14) to C-11/C-10a/C-4/C-3; H-12 ( $\delta_{\text{H}}$  4.67) to C-11/C-1/C-4'; H-11 ( $\delta_{\text{H}}$  3.51) to C-2/C-10a/C-1/C-12, and H-5' ( $\delta_{\text{H}}$  4.35) to C-3'/C-4'. The positions of two methoxyl groups were confirmed by a NOESY experiment, in which a series of correlations were observed between H-5/4-OCH<sub>3</sub> and H-5/6-OCH<sub>3</sub>, in addition to H-11–H-12/2/10, H-5'/H-12, H-12–H-5'/H-10, H-2–H-11, and H-8/H-9. On the basis of the above findings and detailed spectral analyses, the structure of compound **1** was established as 4'-((12-(3,7-dihydroxy-4,6-dimethoxyphenanthren-1-yl)ethyl)(hydroxy)amino)cyclopenta-1',3'-dienecarbaldehyde, and given a trivial name of beilschglabrine A.

Compound **2** was obtained as a yellow amorphous powder. The HREIMS spectrum revealed the molecular formula C<sub>25</sub>H<sub>25</sub>O<sub>6</sub>N, corresponding to the molecular ion peak at  $m/z$  435.1672 (calculated 435.1681). The <sup>1</sup>H NMR and COSY spectra (**Table 1**) displayed similar profile with those of compound **1** except for one of the hydroxyl group position which was replaced by a methoxyl group. This was consistent with the addition of another singlet signal at  $\delta_{\text{H}}$  4.08 (3H) in the <sup>1</sup>H NMR spectrum. The presence of hydroxyl groups was revealed by a broad signal at  $\delta_{\text{H}}$  6.16. The <sup>13</sup>C NMR and DEPT spectra indicated the presence of twenty five carbons, with one carbonyl, eleven quaternary, seven methines, three methylenes, and three methoxyls carbons. The location of the third methoxyl group was confirmed at C-7 by NOESY spectrum (**Fig. 3**) in which good correlation was observed between 7-OCH<sub>3</sub> with H-8. Thus, the structure of compound **2** was established as 4'-(hydroxy(12-(3-hydroxy-4,6,7-trimethoxyphenanthren-1-yl)ethyl)amino)cyclopenta-1',3'-dienecarbaldehyde, and given a trivial name of beilschglabrine B. The known compounds, lupeol (Yang et al., 2009), taraxerol

(Chen et al. 1998) and 24-methylenelanosta-7,9-diene-3 $\beta$ -15 $\alpha$ -diol (Lue et al., 1998), were identified by comparison of their spectroscopic data with published data.

*Beilschmiedia* like other genera in the Lauraceae, is an important source of isoquinoline alkaloids (e.g., benzyloisoquinoline, aporphine) which are widely distributed in the plant kingdom, and more rarely, phenanthrene alkaloids. Some phenanthrene alkaloids possess various biological properties such as antiplatelet (Jantan et al., 2006), anti-inflammatory (Estellés et al., 2005), antimicrobial (Okwu and Nnamdi, 2011), and acetylcholinesterase inhibitory activity (Chiou et al., 1998). In this study, the isolated phenanthrene alkaloids (**1** and **2**) were tested for free radical scavenging, acetylcholinesterase inhibitory and anti-inflammatory activities. The results are shown in **Table 2**. The free radical scavenging activity was evaluated by the DPPH assay with a series of different concentrations (Chima et al., 2014). Compounds **1** and **2** showed a moderate level of activity as compared to that of ascorbic acid with IC<sub>50</sub> values of 115.9 and 134.4  $\mu$ M, respectively. The compounds were also screened for AChE inhibitory activity using the Ellman's colorimetric method in a 96-well plate (Ellman et al., 1961). Both **1** and **2** displayed strong activity against AChE with IC<sub>50</sub> values of 50.4 and 84.6  $\mu$ M, respectively. In the LOX assay (Salleh et al., 2016), compounds **1** and **2** exhibited significant activity with IC<sub>50</sub> values of 32.8 and 40.4  $\mu$ M, respectively. Compound **1** showed slightly higher activity than that of **2** in all assays.

### 3. Experimental

#### 3.1. General experimental procedures

The mass spectra were recorded on the EPSRC UK National Mass Spectrometry Facility (Swansea University). The UV spectra were obtained in methanol on a Shimadzu UV 1601PC spectrophotometer. The IR spectra were obtained on a Perkin-Elmer 1600 FTIR spectrophotometer. The 1D and 2D NMR spectra were recorded in deuterated chloroform on a

Bruker Avance 400 MHz spectrometer, chemical shifts ( $\delta$ ) are reported in ppm on  $\delta$  scale, and the coupling constants ( $J$ ) are given in Hz. Vacuum liquid chromatography (VLC) was performed using Merck silica gel 230-400 mesh, while column chromatography (CC) was performed using Merck silica gel 70-230 mesh. Preparative thin layer chromatography (PTLC) was prepared using silica gel 60 PF<sub>254</sub>. Thin layer chromatography (TLC) aluminum sheets pre-coated with silica gel 60 F<sub>254</sub> (0.2 mm thickness) was used to detect and monitor components presence in the crude samples or fractions. The TLC and PTLC spots were visualized under UV light (254 and 366 nm) followed by spraying with 5% H<sub>2</sub>SO<sub>4</sub> and 1% vanillin in MeOH and heating at 120°C for 5 min. All solvents were AR grade.

### 3.2. *Plant material*

The stem bark of *B. glabra* was collected from Kluang, Johor in October 2014. The species was identified by Dr. Shamsul Khamis from the Institute of Bioscience (IBS), UPM and the voucher specimen (SK2570/14) deposited at the Herbarium of IBS.

### 3.3. *Extraction and isolation*

Cold extraction of the powdered stem bark (2 kg) of *B. glabra*, sequentially, with *n*-hexane, EtOAc and MeOH yielded the crude extracts, *n*-hexane (BGBH: 15.4 g, 0.77%), EtOAc (BGBE: 18.9 g, 0.95%) and MeOH (BGBM: 25.3 g, 1.27%). The MeOH extract (BGBM) (20.0 g) was subjected to VLC on silica gel eluted with *n*-hexane:CHCl<sub>3</sub>:EtOAc to afford 15 major fractions. Fraction BGBM5-10 were combined and purified by multiple CC on silica gel and preparative TLC to yield (1) (10.2 mg) and (2) (11.0 mg). The EtOAc extract (15.0 g) was fractionated by VLC on silica gel, eluted with *n*-hexane:CHCl<sub>3</sub>:EtOAc to afford eight major fractions (BGBE1-8). Fractions BGBE2-5 and BGBE6-8 were purified by CC on silica gel to obtain taraxerol (25.5 mg) and 24-methylenelanosta-7,9-diene-3 $\beta$ -15 $\alpha$ -diol (14.5 mg), respectively. Similar VLC (on silica gel) purification of the *n*-hexane extract (15.0 g) eluted

with *n*-hexane:CHCl<sub>3</sub>:EtOAc afforded 10 major fractions (BGBH1-10). The combined fractions BMBH1-5 was subjected to multiple CC on silica gel to yield lupeol (5.6 mg).

### 3.3.1 *Beilschglabrine A (1)*

Yellow amorphous powder; IR (KBr): 3367, 2917, 2849, 1727, 1637, 1600, 1515, 1462, 1399, 1273, 1088 cm<sup>-1</sup>; UV<sub>max</sub> (MeOH) nm (log ε) 212 (2.15), 264 (2.85); HREIMS *m/z* 421.1517 (calculated 421.1525 for C<sub>24</sub>H<sub>23</sub>O<sub>6</sub>N); <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz), <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100 MHz), COSY, NOESY and HMBC data see **Table 1**.

### 3.3.2 *Beilschglabrine B (2)*

Yellow amorphous powder; IR (KBr): 3312, 2925, 2867, 1715, 1636, 1602, 1514, 1454, 1388, 1223, 1009 cm<sup>-1</sup>; UV<sub>max</sub> (MeOH) nm (log ε) 208 (1.55), 263 (2.62); HREIMS *m/z* 435.1672 (calculated 435.1681 for C<sub>25</sub>H<sub>25</sub>O<sub>6</sub>N); <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz), <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100 MHz), COSY, NOESY and HMBC data see **Table 1**.

## 3.4. *Biological activities*

### 3.4.1. *Solvents and chemicals*

**Anti-oxidant:** 1,1-Diphenyl-2-picrylhydrazyl (DPPH), and ascorbic acid were obtained from Sigma-Aldrich (Germany). Analytical grade ethanol (EtOAc), methanol (MeOH), ethyl acetate (EtOAc), *n*-hexane and dimethylsulfoxide (DMSO) were purchased from Merck (Germany). **Acetylcholinesterase:** AChE enzyme (Type-VI-S, EC3.1.1.7), acetylthiocholine iodide (AChI), 5,5'-dithio-bis(2-nitrobenzoic) acid (DTNB), and galantamine were purchased from Sigma-Aldrich (Germany). **Anti-inflammatory:** Lipoxigenase inhibitor screening assay kit, Cayman Chemicals Co. (Item No. 760700) was purchased from Cayman Chemical Company (Ann Arbor, MI, USA)

### 3.4.2. *Free radical scavenging activity*

The anti-oxidant activity was determined using DPPH free radical scavenging assay as described previously with minor modifications (Shimada et al., 1992; Chima et al., 2014).



Briefly, 0.1 mM DPPH (1 mL) dissolved in EtOH was added to an EtOH solution (3 mL) of **the test samples or ascorbic acid** at different concentrations. An equal volume of EtOH was added in the control test. The mixture was shaken vigorously and allowed to stand at room temperature for 30 min. Then the absorbance at 517 nm was measured with a UV–vis spectrophotometer. The percentage of scavenging of DPPH was calculated using the following equation:

$$I\% = [ A_{\text{blank}} - A_{\text{sample}} / A_{\text{blank}} ] \times 100$$

where  $A_{\text{blank}}$  is the absorbance value of the control reaction (containing all reagents except the test compound) and  $A_{\text{sample}}$  is the absorbance value of the test compound. The sample concentration providing 50% inhibition ( $IC_{50}$ ) was calculated by plotting inhibition percentages against concentrations of the sample. All tests were carried out in triplicate and inhibition values were reported as means  $\pm$  SD of triplicates.

### 3.4.3 Acetylcholinesterase (AChE) inhibitory activity

AChE inhibitory activity of the compounds was measured by slightly modifying the spectrophotometric method (Ellman et al., 1961; Salleh et al., 2016b). Electric eel AChE was used, while acetylthiocholine iodide (AChI) was employed as the substrate of the reaction. DTNB acid was used for the measurement of the AChE activity. Briefly, 140  $\mu$ L of sodium phosphate buffer (pH 8.0), 20  $\mu$ L of DTNB, 20  $\mu$ L of test compounds and 20  $\mu$ L of AChE solution were added by multichannel automatic pipette in a 96-well microplate and incubated for 15 min at 25°C. The reaction was then initiated with the addition of 10  $\mu$ L of AChI. Hydrolysis of AChI was monitored by the formation of the yellow 5-thio-2-nitrobenzoate anion as a result of the reaction of DTNB with thiocholines, catalysed by enzymes at 412 nm utilizing a 96-well microplate reader (Epoch Micro-Volume Spectrophotometer, USA). Percentage of inhibition (I%) of AChE was determined by comparison of rates of reaction of samples relative to blank sample (EtOH in phosphate buffer pH = 8) using the formula:

$$I\% = [ E - S / E ] \times 100$$

where  $E$  is the activity of enzyme without test sample and  $S$  is the activity of enzyme with test sample. The experiments were done in triplicate. Galantamine was used as the positive controls.

#### 3.4.4 Anti-inflammatory activity

The reagents were prepared according to the standard protocol (Lipoxygenase inhibitor screening assay kit, Item No. 760700). Stock solutions of samples were prepared so as to obtain concentrations of 100 - 6.25  $\mu\text{M}$  in the respective wells. The prepared solutions were then introduced onto 96 well plates where the cells were distributed as blanks 1A-2A-1D (triplicate), positive control 1B-2B (duplicate), and 100% initial activity wells 1C-2C-2D (triplicate). The remaining wells were designated for inhibitor (compounds) solutions in duplicate. The addition of the reagents was done according to the standard protocol, according to which, 100  $\mu\text{L}$  of assay buffer was added to the blank wells and 90  $\mu\text{L}$  of lipoxygenase (5-LOX) enzyme and 10  $\mu\text{L}$  of assay buffer were added to positive control wells. For the 100% initial activity wells, 90  $\mu\text{L}$  of lipoxygenase enzyme and 10  $\mu\text{L}$  of solvent (DMSO) were added. The inhibitor (compounds) wells were charged with 90  $\mu\text{L}$  of lipoxygenase enzyme and 10  $\mu\text{L}$  of respective stock (compounds) solution. The reaction was initiated by adding 10  $\mu\text{L}$  of the substrate (AA) to all wells. The plate was then shaken for 5 min on an orbital shaker. Ultimately, 100  $\mu\text{L}$  of chromogen solution (prepared according to standard protocol) was added to each well to stop the enzyme catalysis. The plate was incubated for half an hour and was read at 500 nm. The percentage inhibitions ( $I\%$ ) of the compounds were calculated using the following equation:

$$I\% = [ A_{\text{initial activity}} - A_{\text{inhibitor}} / A_{\text{initial activity}} ] \times 100$$

where  $A_{\text{initial activity}}$  is the absorbance of 100% initial activity wells without sample and  $A_{\text{inhibitor}}$  is the absorbance of compounds/reference. Analyses were expressed as means  $\pm$ SD of triplicates.

#### 3.5 Statistical analysis

Data obtained from biological activities were expressed as mean values. **The statistical analyses were carried out using one way ANOVA (p<0.05).** A statistical package (SPSS version 16.0) was used for the data analysis.

### **Appendix A. Supplementary data**

Supplementary data including IR, 1D/2D NMR, UV and MS for compound (1-2) are available as supporting information.

### **Acknowledgments**

The authors are grateful to Research University Grant (GUP) for financial support under vote Q.J130000.2526.03H93 and the Faculty of Science, Universiti Teknologi Malaysia for research facilities.

### **References**

- Castro, O., López, J., Stermitz, F.R., 1986. New aporphine alkaloids from *Phoebe valeriana*. J. Nat. Prod. 49, 1036-1040.
- Chen, K.S., Chang, F.R., Chia, Y.C., Wu, T.S., Wu, Y.C., 1998. Chemical constituents of *Neolitsea parvigemma* and *Neolitsea konishii*. J. Chinese Chem. Soc. 45, 103-110.
- Chen, J.J., Chou, E.T., Peng, C.F., Chen, I.S., Yang, S.Z., Huang, H.Y., 2007. Novel epoxyfuranoid lignans and antitubercular constituents from the leaves of *Beilschmiedia tsangii*. Planta Med. 73, 567-571.
- Chima, N.K., Nahar, L., Majinda, R.R.T., Celik, S., Sarker, S.D., 2014. Assessment of free radical scavenging activity of the extracts and fractions of *Gypsophila pilulifera*: Assay-guided isolation of the active component. Braz. J. Pharmacogn. 24, 38-43.

- Chiou, C.M., Kang, J.J., Lee, S.S., 1998. Litebamine N-homologues: preparation and antiacetylcholinesterase activity. *J. Nat. Prod.* 61, 46-50.
- Chouna, J.R., Nkeng-Efouet, P.A., Lenta, B.N., Wansi, J.D., Kimbu, S.F., Sewald, N., 2010. Endiandric acid derivatives from the stem bark of *Beilschmiedia anacardioides*. *Phytochem. Lett.* 3, 13-16.
- Ellman, G.L., Courtney, K.D., Andres, V., Featherstone, R.M., 1961. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem. Pharmacol.* 7, 88-95.
- Estellés, R., Milian, L., Nabah, Y.N., Mateo, T., Cerdá-Nicolás, M., Losada, M., Ivorra, M.D., Issekutz, A.C., Cortijo, J., Morcillo, E.J., Blázquez, M.A., Sanz, M.J., 2005. Effect of boldine, secoboldine, and boldine methine on angiotensin II-induced neutrophil recruitment in vivo. *J. Leukoc. Biol.* 78, 696-704.
- Jantan, I., Raweh, S.M., Yasin, Y.H.M., Murad, S., 2006. Antiplatelet activity of aporphine and phenanthrenoid alkaloids from *Aromadendron elegans* Blume. *Phytother. Res.* 20, 493-496.
- Lenta, B.N., Tantangmo, F., Devkota, K.P., Wansi, J.D., Chouna, J.R., Soh, R.C.F., Neumann, B., Stammer, H.G., Tsamo, E., Sewald, N., 2009. Bioactive constituents of the stem bark of *Beilschmiedia zenkeri*. *J. Nat. Prod.* 72, 2130-2134.
- Lue, Y.P., Mu, Q., Zheng, H.L., Li, C.M., 1998. 24-Methylene tetracyclic triterpenes from *Polyalthia lancilimba*. *Phytochemistry* 49, 2053-2056.
- Nishida, W.S., 1999. Revision of *Beilschmiedia* (Lauraceae) in the Neotropics. *Ann. Missouri Bot. Gard.* 86, 657-701.
- Nishida, W.S., 2008. Taxonomic revision of *Beilschmiedia* (Lauraceae) in Borneo. *Blumea* 53, 345-383.

- Okwu, D.E., Nnamdi, F.U., 2011. A novel antimicrobial phenanthrene alkaloid from *Bryopyllum pinnatum*. E-J Chem. 8, 1456-1461.
- Salleh, W.M.N.H.W., Ahmad, F., Khong, H.Y., Zulkifli, R.M., 2015. Chemical compositions and biological activities of essential oils of *Beilschmiedia glabra*. Nat. Prod. Commun. 10, 1297-1300.
- Salleh, W.M.N.H.W., Ahmad, F., Khong, H.Y., Zulkifli, R.M., 2016a. Comparative study of the essential oils of three *Beilschmiedia* species and their biological activities. Int. J. Food Sci. Tech. 51, 240-249.
- Salleh, W.M.N.H.W., Ahmad, F., Khong, H.Y., Zulkifli, R.M., Sarker, S.D., 2016b. Madangones A and B: Two new neolignans from the stem bark of *Beilschmiedia madang* and their bioactivities. Phytochem. Lett. 15, 168-173.
- Shamma, M., 1972. *The Isoquinoline Alkaloids*, Academic Press, New York and Londres, Verlag Chemie, Weinheim, pp. 192.
- Shimada, K., Fujikawa, K., Yahara, K., Nakamura, T., 1992. Antioxidative properties of xanthin on autoxidation of soybean oil in cyclodextrin emulsion. J. Agric. Food Chem. 40, 945-948.
- Yang, P.S., Cheng, M.J., Peng, C.F., Chen, J.J., Chen, I.S., 2009. Endiandric acid analogues from the roots of *Beilschmiedia erythrophloia*. J. Nat. Prod. 72, 53-58.