

Article (refereed) - postprint

Vestergard, Mette; Reinsch, Sabine; Bengtson, Per; Ambus, Per; Christensen, Soren. 2016. **Enhanced priming of old, not new soil carbon at elevated atmospheric CO₂**. *Soil Biology & Biochemistry*, 100. 140-148.
[10.1016/j.soilbio.2016.06.010](https://doi.org/10.1016/j.soilbio.2016.06.010)

© 2016 Elsevier Ltd

This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>



This version available <http://nora.nerc.ac.uk/513847/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

NOTICE: this is the author's version of a work that was accepted for publication in *Soil Biology & Biochemistry*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Soil Biology & Biochemistry*, 100. 140-148. [10.1016/j.soilbio.2016.06.010](https://doi.org/10.1016/j.soilbio.2016.06.010)

www.elsevier.com/

Contact CEH NORA team at
noraceh@ceh.ac.uk

1 Title: Enhanced priming of old, not new soil carbon at elevated atmospheric CO₂

2

3 Authors: Mette Vestergård^{1*}, Sabine Reinsch^{2a}, Per Bengtson³, Per Ambus^{2b}, Søren Christensen¹

4 ¹Terrestrial Ecology Section, Department of Biology, University of Copenhagen,

5 Universitetsparken 15, DK-2100 Copenhagen Ø, Denmark

6 ² Technical University of Denmark, Department for Chemical and Biochemical

7 Engineering, Frederiksborgvej 399, DK-4000 Roskilde, Denmark

8 ³ Department of Biology, Microbial Ecology, Lund University, Sölvegatan 37, 223 62

9 Lund, Sweden

10

11 *Corresponding author: Mette Vestergård, Terrestrial Ecology Section, Department of Biology,

12 University of Copenhagen, Universitetsparken 15, DK-2100 Copenhagen Ø, Denmark.

13 Phone: +45 51827047. E-mail: mevmadsen@bio.ku.dk

14

15 Key-words: Carbon-13; Drought; FACE; Global change; Heathland; Warming

16

17

18

19

20

21

22

^a Present address: Centre for Ecology and Hydrology, Environment Centre Wales, Bangor, LL57 2UW, UK

^b Present address: Department of Geosciences and Natural Resource Management, University of Copenhagen, 1350, Copenhagen, Denmark.

23 **Abstract**

24 Rising atmospheric CO₂ concentrations accompanied by global warming and altered precipitation
25 patterns calls for assessment of long-term effects of these global changes on carbon (C) dynamics in
26 terrestrial ecosystems, as changes in net C exchange between soil and atmosphere will impact the
27 atmospheric CO₂ concentration profoundly. In many ecosystems, including the heath/grassland
28 system studied here, increased plant production at elevated CO₂ increase fresh C input from litter
29 and root exudates to the soil and concurrently decrease soil N availability. Supply of labile C to the
30 soil may accelerate the decomposition of soil organic C (SOC), a phenomenon termed ‘the priming
31 effect’, and the priming effect is most pronounced at low soil N availability. Hence, we
32 hypothesized that priming of SOC decomposition in response to labile C addition would increase in
33 soil exposed to long-term elevated CO₂ exposure. Further, we hypothesized that long-term warming
34 would enhance SOC priming rates, whereas drought would decrease the priming response.
35 We incubated soil from a long-term, full-factorial climate change field experiment, with the factors
36 elevated atmospheric CO₂ concentration, warming and prolonged summer drought with either labile
37 C (sucrose) or water to assess the impact of labile C on SOC dynamics. We used sucrose with a
38 ¹³C/¹²C signature that is distinct from that of the native SOC, which allowed us to assess the
39 contribution of these two C sources to the CO₂ evolved. Sucrose induced priming of SOC, and the
40 priming response was higher in soil exposed to long-term elevated CO₂ treatment. Drought tended
41 to decrease the priming response, whereas long-term warming did not affect the level of priming
42 significantly.

43 We were also able to assess whether SOC-derived primed C in elevated CO₂ soil was assimilated
44 before or after the initiation of the CO₂ treatment 8 years prior to sampling, because CO₂
45 concentrations were raised by fumigating the experimental plots with pure CO₂ that was ¹³C-
46 depleted compared to ambient CO₂. Surprisingly, we conclude that sucrose addition primed

47 decomposition of relatively old SOC fractions, i.e. SOC assimilated more than 8 years before
48 sampling.

49

50 **1. Introduction**

51 The global terrestrial soil organic carbon (SOC) pool is the largest terrestrial carbon (C) pool and
52 constitutes a C stock that is more than twice the size of the atmospheric CO₂-C pool (IPCC, 2013).

53 Therefore, even relatively moderate fluctuations in net C exchange between soil and atmosphere
54 will impact the CO₂ concentration in the atmosphere profoundly. Faced by rising atmospheric CO₂
55 levels and the anticipated climatic changes that will result from this rise, we need to better
56 understand how such changes will influence SOC decomposition and CO₂-release from terrestrial
57 organic C pools.

58 At least two factors that can potentially alter SOC decomposition, namely nitrogen (N) availability
59 and input of fresh plant C, are expected to change with rising CO₂ levels. Supply of fresh plant
60 derived C into the soil matrix may accelerate the decomposition of SOC and decrease soil C stocks
61 (Fontaine et al., 2004); a phenomenon termed ‘the priming effect’. Priming effects induced by root
62 exudation and rhizodeposition can cause an up to 350 % increase in SOC decomposition compared
63 to the root free soil (Cheng et al., 2014). Even so, most C-cycling models do not consider the
64 influence of priming and living roots on SOC decomposition rates (Cheng et al., 2014), perhaps as a
65 result of limited knowledge about underlying mechanisms and factors influencing the magnitude of
66 priming. However, the few attempts that have been made to date to represent plant-induced priming
67 of SOC decomposition have resulted in improved model performance, also regarding global change
68 effects (Cheng et al., 2014, Perveen et al., 2014). It is therefore evident that models predicting
69 future climatic conditions and atmospheric CO₂ levels, as well as our means of mitigating the
70 effects of rising CO₂ levels, depend on more in-depth understanding of feedbacks between climate,
71 elevated atmospheric CO₂ levels, and SOC decomposition caused by priming.

72

73 Common plant physiological responses to elevated CO₂ comprise enhanced photoassimilation of C,
74 increased root volume, and increased input of plant C to the soil in the form of root exudates and
75 rhizodeposits (Hungate et al., 1997; Adair et al., 2009; Albert et al., 2011; Arndal et al., 2013,
76 2014). Since the extent of priming seems to depend on the concentration of labile C inputs, with no
77 or low priming at low concentrations (Blagodatskaya & Kuzyakov, 2008; Gude et al., 2012) and
78 gradually increasing priming with increasing concentrations (Blagodatskaya & Kuzyakov, 2008;
79 Paterson & Sim, 2013) up to a point of saturation (Guenet et al., 2010; Xiao et al., 2015), it can be
80 expected that priming will increase at elevated atmospheric CO₂ concentration due to higher inputs
81 of labile plant C to soils (Paterson et al., 2008). Accordingly, elevated atmospheric CO₂ has been
82 shown to increase decomposition of SOC in grasslands (Xie et al., 2005; Niklaus & Falloon, 2006)
83 as well as forests (Phillips et al., 2012). However, there are also examples where input of labile
84 plant C resulted in negative priming (Sullivan & Hart, 2013; Cheng et al., 2014), i.e. inhibited the
85 decomposition of SOC. A possible reason to the contradictory findings is that the magnitude and
86 direction of priming is dependent on the nutrient status of the soil. In fact, it has been suggested that
87 the decomposition of SOC in response to input of labile C is driven by enhanced microbial demand
88 for nutrients retained in soil organic matter (SOM) (Paterson, 2009; Philips et al., 2012).
89 Accordingly, Bengtson et al. (2012) demonstrated a strong link between rhizodeposition, SOM
90 decomposition and gross N mineralization in a coniferous forest soil, while Fontaine et al. (2004,
91 2011) found that soil C losses caused by priming increased when soil microbes are nutrient limited.
92 In line with these findings, a recent meta-analysis concluded that input of labile C enhanced
93 decomposition of native SOM, but only in soils with low nitrogen (N) content (Zhang et al., 2013).
94 Since it has been observed that increased plant N demand at elevated atmospheric CO₂ commonly
95 leads to decreased soil N availability (Luo et al., 2004; Larsen et al., 2011), this can also explain
96 why priming can be expected to increase under elevated atmospheric CO₂ conditions.

97

98 However, in order to fully appreciate how priming will influence the net ecosystem exchange of C
99 in a high CO₂ world we also need to consider climatic parameters, such as temperature and
100 precipitation patterns that are also undergoing changes that are expected to continue for the decades
101 to come (IPCC, 2013). In Northern Europe we expect an annual mean temperature increase between
102 0.75 and 0.1 °C over the coming 20 years and more extreme precipitation patterns, for instance
103 prolonged summer droughts according to the IPCC RCP4.5 scenario (IPCC, 2013). Hence,
104 evaluation of elevated CO₂ impacts on priming of SOC decomposition must consider projected
105 temperature and precipitation scenarios.

106 Low soil moisture generally reduces soil microbial activity (Moyano et al., 2013) and may also
107 reduce rhizosphere priming of SOC decomposition (Dijkstra & Cheng, 2007). It can, therefore, be
108 expected that prolonged summer droughts will decrease priming. However, as plants have better
109 water use efficiency at elevated CO₂ (Field et al., 1995; Ainsworth & Rogers, 2007), drought
110 impacts on microbial activity are in some cases less severe when combined with elevated CO₂
111 (Kassem et al., 2008). The combined effect of elevated CO₂ and drought on the magnitude of
112 priming is to our knowledge not known.

113 Likewise, little is known about temperature effects on priming. The only study to date that has
114 systematically addressed the temperature dependency of priming found the process to be non-
115 responsive to temperature variations (Ghee et al., 2013). In general, even moderate warming
116 enhances the activity of heterotrophic microbial SOM decomposers (Wang et al., 2014). Therefore,
117 in systems with low N availability temperature dependent stimulation of microbial activity could
118 enhance the need for microbial N acquisition through SOM decomposition and increase priming.
119 However, the priming response to warming may very well depend on soil moisture conditions, since
120 warming enhances evaporation. This could potentially exacerbate negative effects of drought on
121 soil microbial activity.

122

123 Previous studies demonstrated that elevated CO₂ changed C turnover dynamics of different fractions
124 of SOM. Elevated CO₂ increased the content of recently assimilated C in both coarse and fine
125 particulate fractions of SOM, but decreased the content of older C in more physically protected, fine
126 particulate organic matter and mineral-associated organic matter (Hofmockel et al., 2011). This
127 suggests that elevated CO₂ elicits priming of older relatively stable rather than recent SOC pools. In
128 the current experiment we are able to test this hypothesis, as we raised the atmospheric CO₂
129 concentration by fumigating with CO₂ that was ¹³C-depleted compared to the naturally occurring
130 atmospheric CO₂. Therefore, C fixed in elevated CO₂ treatments was ¹³C-depleted compared to the
131 C assimilated before CO₂ treatment started and compared to the C pools of ambient CO₂ treatments
132 (Reinsch & Ambus, 2013). A comparison of the isotopic composition of primed SOC-derived CO₂-
133 C from elevated CO₂ and ambient CO₂ treatments can therefore reveal if primed C derives from C
134 fixed before or after the initiation of CO₂ fumigation.

135

136 The aim of this study was to test the effects of long-term elevated CO₂ exposure, warming and
137 annual extended drought events on potential priming in a nutrient-poor temperate heath/grassland,
138 where long-term elevated CO₂ exposure has reduced the relative N content of organic inputs
139 (Larsen et al., 2011; Arndal et al., 2013, 2014; Vestergård et al., 2015). Moderate warming has also
140 prolonged the plant growth season with two weeks in the spring at the site (Kongstad et al., 2012).
141 If an extended growth period also enhances plant N uptake over the season, this could potentially
142 intensify microbial N demand. We hypothesize, in accordance with other reports (van Groenigen et
143 al., 2005; Xie et al., 2005; Niklaus & Falloon, 2006), that potential priming of soil C is enhanced in
144 soil exposed to elevated CO₂, where plant production and hence C input to the soil is enhanced and
145 the relative N availability has declined. We further hypothesize that warming enhances priming in
146 soil exposed to elevated CO₂, because we expect that an earlier onset of spring growth and

147 enhanced microbial activity under warming further reduced N availability. We hypothesize that
148 summer drought, which is expected to reduce microbial activity, will reduce priming in soil exposed
149 to elevated CO₂. Finally, we expect that warming augments this effect of drought. Further, we will
150 clarify if primed soil C is recently fixed or of older origin.

151

152 Addition of labile carbohydrates is a common method to assess and compare potential priming
153 activity between different soils and treatments (Wu et al., 1993; Zyakun & Dilly, 2005; Garcia-
154 Pausas & Paterson, 2011; Paterson & Sim, 2013; Reinsch et al., 2013). If the ¹³C/¹²C signature of
155 the added carbohydrate is distinct from that of native SOC it is possible to assess the contribution of
156 these two C sources to the respiratory CO₂ evolved. In the present study we estimated potential
157 priming by incubating soils with labile sucrose, a common constituent of root exudates (Grayston et
158 al., 1998), with a ¹³C/¹²C ratio that is distinct from the isotopic ratio of the native SOC.

159

160 **2. Materials and Methods**

161

162 *2.1 Field site*

163 Our field experiment was carried out in an unmanaged temperate heath/grassland in North Zealand
164 c. 50 km northwest of Copenhagen, Denmark (55°53'N, 11°58' E). The soil is a Cambic Arenosol
165 (FAO classification) developed on a nutrient-poor sandy deposit. The organic layer is 5-10 cm thick
166 and the pH is around 5. From 1975-2005 the average annual precipitation was 610 mm and the
167 mean annual temperature was 8 °C (Danish Meteorological Institute). From 2005 to 2013 the mean
168 annual precipitation was 742 mm with a range between 648 and 894 mm and the mean annual
169 temperature was 9.7 °C with a range between 7 and 10 °C. The prevailing species are the grass
170 *Deschampsia flexuosa* (c. 70 % coverage) and the dwarf shrub *Calluna vulgaris* (c. 30 % coverage)
171 intermixed with other grasses, herbs, mosses and lichens (Kongstad et al., 2012).

172

173 2.2 Climate manipulation treatments

174 Climate and CO₂ manipulations, aimed to simulate climatic conditions and atmospheric CO₂ levels
175 that are predicted for Denmark in 2075, were initiated in October 2005. The global change factors
176 drought (D), warming (T) and CO₂ concentration (CO₂) were manipulated individually and in all
177 possible combinations. The CO₂ concentration was increased with 120 ppm based on the average of
178 predicted concentrations in 2075 in five atmospheric CO₂ stabilization scenarios (SP450, SP550,
179 SP650, SP750 and SP1000) (IPCC 2007). The temperature increase chosen was the average of
180 predicted temperature responses for Northern Europe (IPCC 2007), and drought manipulations were
181 also based on the predictions in the IPCC 2007 report. The CO₂ concentration was increased *in-situ*
182 via the free-air carbon dioxide enrichment (FACE) technique in octagon shaped plots (octagons)
183 during daytime hours. Extended spring/summer droughts were imposed using moveable curtains to
184 exclude precipitation for a period of ~1 month during spring/early summer each year. Drought
185 curtains reduced precipitation by 7.6 ± 2.1 % (mean \pm SD) annually. In 2013, the drought period
186 was conducted between 29th of April and 27th of May. From 27th of May to sampling, June 5th-6th,
187 the site only received a few mm precipitation, effectively extending the drought period until
188 sampling. Passive night time warming was achieved via moveable curtains that covered the
189 experimental plots during night time hours and prevented heat loss to the atmosphere. The warming
190 effect at 20 cm above ground surface ranged between 0.5 °C and 1.5 °C over the year (Scherber et
191 al., 2013). Warming curtains were withdrawn during rainfall.

192 The experiment is a full-factorial split plot design organized in 6 blocks. One block contains two
193 octagons each of 6.8 m diameter, one exposed to ambient CO₂ concentration and one exposed to
194 elevated CO₂ concentration, respectively. Each octagon is divided into four plots, which amounts to
195 a total of 48 plots. Within each octagon, one plot is subjected to the drought treatment, one is
196 subjected to warming, and a third plot is subjected to the combined drought and warming treatment.

197 The treatment, which is not subjected to any of the global change treatments, represents ambient
198 conditions (A). For further details regarding online measurements, treatments and experimental
199 setup, see Mikkelsen et al. (2008).

200 Characteristics of the soil in the treatments at sampling in June 2013 are shown in Table 1. SOC
201 content in 0-10 cm depth was higher at elevated CO₂ than at ambient CO₂, and at elevated CO₂
202 drought further increased the SOC content.

203

204 2.3 Soil sampling and incubation

205 Within each of the 48 plots, an undisturbed area of 0.5 m × 0.5 m was selected for this experiment.

206 Areas were chosen to contain an approximately equal amount of *C. vulgaris* and grasses (mainly

207 *D. flexuosa*) at initiation of the experimental setup in 2003. Unfortunately, an error in the

208 experimental procedures forced us to discard samples from one of the blocks, and we therefore

209 present data from five blocks ($n = 5$).

210 Soil cores were taken on the 5th-6th of June 2013 with an 8.7 cm diameter cylinder auger to 10 cm

211 depth. The soil was sieved (2 mm) in the field to separate roots from the soil and kept under cool

212 conditions (5 °C) until use. Two weeks after sampling, two 117 mL serum flasks per soil sample

213 were each added 4 g (fw) soil. We added 10 mL water or 10 mL of sucrose solution (4 g L⁻¹) to

214 each of the paired flasks, respectively. Assuming an average bulk density of 1.24 g cm⁻³ in the 10

215 top cm of the soil the sucrose added corresponds to an input of 572 g C m⁻². This is well above the

216 total annual C input to the soil, given that the annual net primary production roughly corresponds to

217 350 g C m⁻² (Chapin III et al., 2002). Hence, during the incubation, the soil was fully water

218 saturated, and microorganisms were at no risk of experiencing C limitation. We used sugar cane-

219 derived sucrose with a ¹³C/¹²C ratio of δ¹³C = -12 ‰, as it is distinct from the δ¹³C value (-28.5 ‰)

220 of the C3 plants in the area (Reinsch & Ambus, 2013). This enables us to distinguish between CO₂

221 derived from the added sucrose and from SOC. To eliminate the initial CO₂-content of the flasks,

222 the flasks were sealed, evacuated (< 10% air remaining) and refilled to atmospheric pressure with
223 CO₂-free atmospheric air (Alphagaz Luft 1, Air Liquide, Denmark). Evacuation and refilling was
224 repeated twice and finally 20 mL extra CO₂-free air was added. We incubated the flasks on a shaker
225 (4 h, 20 °C). Following incubation, we sampled 19 mL headspace gas with a syringe and flushed an
226 evacuated 5.9 ml Exetainer vial (Labco Scientific, High Wycombe, UK) with the gas sample
227 leaving the vial at atmospheric pressure for subsequent analyses of CO₂ concentration and isotopic
228 ¹³C/¹²C ratio. Likewise, we sampled gas from four background control flasks, which only contained
229 10 mL water or 10 mL sucrose solution.

230

231 *2.4 Soil and sucrose analysis*

232 We dried 10 g soil at 103 °C to determine gravimetric soil water content. The total C content and
233 ¹³C/¹²C isotopic ratio of soil and sucrose were measured in dried samples by Dumas combustion
234 (1020 °C) on an elemental analyser (CE 1110, Thermo Electron, Milan, Italy) coupled in continuous
235 flow mode to a Finnigan MAT Delta PLUS isotope ratio mass spectrometer (Thermo Scientific,
236 Bremen, Germany). Homogenized portions of 2 mg (sucrose) or 15 mg (soil) were weighed out into
237 tin combustion cups for elemental analysis. Acetanilide (Merck, Darmstadt, Germany) and soil
238 standards (Elemental Microanalysis, Okehampton, UK) were used for elemental analyser mass
239 calibration. As working standard for isotope ratio analysis we used pure CO₂ gas calibrated against
240 certified reference ¹³C-sucrose (IAEA, Vienna, Austria). Performance of analysis (Qa/Qc) was
241 assessed by the inclusion of reference samples of biological origin (Peach leaves (NIST 1547),
242 National Institute of Standards and Technology, Gaithersburg, MD, USA).

243

244 *2.5 CO₂ concentration and ¹³C/¹²C isotopic ratio*

245 CO₂ concentrations and isotopic ¹³C/¹²C ratio were analysed on a DeltaV Advantage Isotope Ratio
246 Mass Spectrometer (Thermo Scientific, Bremen, Germany) coupled in continuous flow mode to a

247 GasBench II. Two calibration gas mixtures of CO₂ in synthetic air were included in the analytical
 248 runs, viz. 362 ppm CO₂ at δ¹³C = -2.7 ‰ vs. Vienna Pee Dee Belemnite (VPDB) and 356 ppm CO₂
 249 at δ¹³C = -29.3 ‰ (Messer Denmark, Padborg, Denmark). CO₂ concentrations and δ¹³C were
 250 corrected according to the values measured in the background control treatments.

251

252 2.6 Data analysis

253 Because we eliminated the initial CO₂ content of the flasks before the soil incubation, we assessed
 254 respiration rates based on the CO₂ concentration measured after the incubation period. We
 255 calculated the sucrose-induced respiration as the difference between CO₂-C evolved in sucrose-
 256 amended and non-amended flasks.

257

258 We calculated the proportion of SOC-derived respiratory C (P_{SOC}) and sucrose-derived respiratory
 259 C (P_{suc}) in sucrose-amended flasks using a two end-member-mixing-model:

260

261 [Equation 1]
$$P_{\text{suc}} = \frac{\delta^{13}\text{C}_{\text{SOC}} - \delta^{13}\text{C}_{\text{samp}}}{\delta^{13}\text{C}_{\text{SOC}} - \delta^{13}\text{C}_{\text{suc}}}$$

262

263 [Equation 2]
$$P_{\text{SOC}} = 1 - P_{\text{suc}} = 1 - \frac{\delta^{13}\text{C}_{\text{SOC}} - \delta^{13}\text{C}_{\text{samp}}}{\delta^{13}\text{C}_{\text{SOC}} - \delta^{13}\text{C}_{\text{suc}}}$$

264

265 where δ¹³C_{samp} denotes the isotopic value of the CO₂ evolved in flasks amended with sucrose,
 266 δ¹³C_{SOC} denotes the isotopic value of soil C, and δ¹³C_{suc} is the isotopic value of the added sucrose (-
 267 12 ‰).

268

269 By calculating the total SOC-derived CO₂ evolved in sucrose-amended flasks and subtracting the
 270 basal respiratory CO₂ evolved, i.e. the CO₂ produced in non-amended flasks, we can calculate the
 271 amount of SOC primed in response to sucrose addition as follows:

272

273 [Equation 3] $C_{\text{primed}} = P_{\text{SOC}} \times [CO_{2\text{suc}}] - [CO_{2\text{H}_2\text{O}}]$,

274

275 where [CO_{2suc}] and [CO_{2H₂O}] denote the CO₂ evolved in flasks with and without sucrose,
 276 respectively. We present the priming effect as the increase in SOC-derived CO₂ with sucrose-
 277 amendment in relation to the SOC content of individual samples.

278

279 To assess whether primed C for elevated CO₂ octagons derived from C assimilated before or after
 280 the initiation of the FACE in 2003, we calculated the δ¹³C of the primed SOC-derived CO₂,
 281 δ¹³C_{primed}. First, we calculated the δ¹³C of the sucrose-induced respiratory CO₂, i.e. the additional
 282 CO₂ produced from sucrose and from SOC priming in response to sucrose addition, as the
 283 difference in ¹³C respired with and without sucrose divided by the difference in CO₂ produced with
 284 and without sucrose:

285

286 [Equation 4] $\delta^{13}\text{C of sucrose-induced CO}_2\text{-production} = \frac{[\delta^{13}\text{C}_{\text{suc}}] \times [CO_{2\text{suc}}] - [\delta^{13}\text{C}_{\text{H}_2\text{O}}] \times [CO_{2\text{H}_2\text{O}}]}{[CO_{2\text{suc}}] - [CO_{2\text{H}_2\text{O}}]}$,

287

288 where subscripts “suc” and “H₂O” denote CO₂ evolved in flasks with and without sucrose addition,
 289 respectively.

290 At the same time, δ¹³C of sucrose-induced CO₂-production can also be expressed in relation to the
 291 proportional contribution of sucrose and SOC to the sucrose-induced CO₂-production:

292

293 [Equation 5] $\delta^{13}\text{C}$ of sucrose-induced CO_2 -production = $P_{\text{suc}}^* \times \delta^{13}\text{C}_{\text{suc}} + P_{\text{SOC}}^* \times \delta^{13}\text{C}_{\text{SOC}}$,

294

295 where P_{suc}^* and P_{SOC}^* denote the proportions of sucrose-derived and SOC-derived CO_2 in the
 296 sucrose-induced respiration, respectively. We know all the variables of equation 4, and by
 297 combining equations 4 and 5, we can express $\delta^{13}\text{C}_{\text{primed}}$ as follows:

298

299 [Equation 6]

$$300 \delta^{13}\text{C}_{\text{primed}} = \frac{\frac{P_{\text{suc}}^* \times \delta^{13}\text{C}_{\text{suc}} + P_{\text{SOC}}^* \times \delta^{13}\text{C}_{\text{SOC}} - P_{\text{SOC}}^* \times \delta^{13}\text{C}_{\text{primed}}}{P_{\text{suc}}^* - P_{\text{SOC}}^*}}{P_{\text{suc}}^* - P_{\text{SOC}}^*} =$$

$$301 \frac{P_{\text{suc}}^* \times \delta^{13}\text{C}_{\text{suc}} + P_{\text{SOC}}^* \times \delta^{13}\text{C}_{\text{SOC}} - P_{\text{SOC}}^* \times \delta^{13}\text{C}_{\text{primed}}}{P_{\text{suc}}^* - P_{\text{SOC}}^*}$$

302 We know that $\delta^{13}\text{C}_{\text{suc}}$ is -12 ‰, so we must calculate P_{suc}^* . We calculate P_{suc}^* based on the sucrose-
 303 derived CO_2 , $\text{CO}_{2\text{sucrose-derived}}$, and the total sucrose-induced respiration, i.e. the difference between
 304 CO_2 evolved in sucrose-amended and non-amended flasks. This is possible, because we can
 305 calculate $\text{CO}_{2\text{sucrose-derived}}$ as the product of the proportion of the sucrose-derived respiratory CO_2
 306 (P_{suc}) in sucrose-amended flasks and the CO_2 evolved in sucrose-amended flasks:

308

$$309 \text{ [Equation 7] } P_{\text{suc}}^* = \frac{[\text{CO}_{2\text{sucrose-derived}}]}{[\text{CO}_2]_{\text{sucrose-amended}} - [\text{CO}_2]_{\text{non-amended}}} = P_{\text{suc}} \times \frac{[\text{CO}_2]_{\text{sucrose-amended}}}{[\text{CO}_2]_{\text{sucrose-amended}} - [\text{CO}_2]_{\text{non-amended}}}$$

310

311 We know P_{suc} from equation 1; hence:

312

$$313 \text{ [Equation 8] } P_{\text{suc}}^* = \frac{[\text{CO}_2]_{\text{sucrose-amended}} \times P_{\text{suc}}}{[\text{CO}_2]_{\text{sucrose-amended}} - [\text{CO}_2]_{\text{non-amended}}}$$

314

315 and by inserting equation 8 in equation 6, we find that $\delta^{13}\text{C}$ of the primed SOC, $\delta^{13}\text{C}_{\text{primed}}$ can be
 316 calculated accordingly:

317

318 [Equation 9]
$$\delta^{13}\text{C}_{\text{primed}} = \frac{\left(\frac{[\text{C}_{\text{primed}}] \times \delta^{13}\text{C}_{\text{primed}} - [\text{C}_{\text{total}}] \times \delta^{13}\text{C}_{\text{total}}}{[\text{C}_{\text{primed}}] - [\text{C}_{\text{total}}]} \right) \times \left(\frac{[\text{C}_{\text{total}}] \times \delta^{13}\text{C}_{\text{total}} - [\text{C}_{\text{primed}}] \times \delta^{13}\text{C}_{\text{primed}}}{[\text{C}_{\text{total}}] - [\text{C}_{\text{primed}}]} \right)}{\left(\frac{[\text{C}_{\text{total}}] \times \delta^{13}\text{C}_{\text{total}} - [\text{C}_{\text{primed}}] \times \delta^{13}\text{C}_{\text{primed}}}{[\text{C}_{\text{total}}] - [\text{C}_{\text{primed}}]} \right)}$$

319

320 For one of the soil samples from the ambient treatment the $\delta^{13}\text{C}$ value of CO_2 evolved in the flasks
 321 was unexpectedly high, which suggests that the CO_2 partly originated from carbonate C. We
 322 therefore omitted this sample from data analyses. We tested the effects of elevated CO_2 , drought
 323 and warming on all response variables with full factorial three-way ANOVAs. Homogeneity of
 324 variance was assessed with the Brown-Forsythe test. Data for basal respiration rate were log
 325 transformed prior to analysis to obtain homogeneity of variance. All statistical analyses were
 326 executed in Sigma Plot version 13.0.

327

328 3. Results

329 The soil-weight-specific basal respiration and sucrose-induced respiration, i.e. the extra CO_2
 330 produced in sucrose-amended flasks compared to non-amended flasks, was on average c. 50 %
 331 higher in soil from elevated CO_2 plots, and this increase was more pronounced when drought and
 332 elevated CO_2 were combined (Table 2 and 3). This is in line with the higher SOC content at
 333 elevated CO_2 , which was also highest when elevated CO_2 and drought were combined (Table 1). In
 334 contrast, warming did not affect basal or sucrose-induced respiration. The basal decomposition of
 335 SOC, expressed as the respiration activity per g SOC, was independent of treatments (Table 2 and
 336 3). Likewise, the SOC-specific sucrose-induced respiration activity did not differ between
 337 treatments (Table 2 and 3).

338

339 Sucrose addition enhanced decomposition of native SOC (Fig. 1), hence priming occurred. At
340 ambient CO₂, sucrose enhanced the SOC decomposition rate with 35-49 μg C g SOC⁻¹ h⁻¹. The
341 priming effect was significantly higher at elevated CO₂ (Table 3), where sucrose addition enhanced
342 SOC decomposition rate with 43-59 μg C g SOC⁻¹ h⁻¹. There was a tendency towards reduced
343 priming in soils exposed to drought (*P*=0.11), but warming did not affect the level of priming (Fig.
344 1, Table 3).

345

346 As a consequence of the eight years of FACE with pure ¹³C-depleted CO₂, the δ¹³C of the total CO₂
347 efflux from elevated CO₂ soil (Fig. 2a) was significantly 3.0-5.4 ‰ lower than from ambient CO₂
348 soil in flasks without sucrose (Table 3). Addition of sucrose with the higher δ¹³C of -12 ‰ raised
349 the δ¹³C of the CO₂ evolved during the four-hour incubation. Nevertheless, CO₂ from elevated CO₂
350 octagons in the sucrose-amended flasks was still significantly 1.2-2.0 ‰ lower than CO₂ evolved
351 from ambient CO₂ soil (Fig. 2b, Table 3).

352 In contrast to the total CO₂ efflux (Fig. 2), the δ¹³C of primed SOC-derived CO₂-C from elevated
353 CO₂ soil was not lower than the δ¹³C of primed SOC-derived CO₂-C from ambient CO₂ soil
354 (Fig. 3). At ambient CO₂, the δ¹³C of primed CO₂-C released from soils that were not exposed to
355 warming was significantly lower than the δ¹³C of primed C in all other treatments (Fig. 3, Table 3).

356

357 **4. Discussion**

358 The soil C content in the upper 0-10 cm of the soil profile had increased with 12-22 % after eight
359 years of elevated CO₂ exposure in treatments without experimental drought exposure, and drought
360 further increased soil C content at elevated CO₂ (Table 1), which is consistent with the increased
361 root production at elevated CO₂ recorded in 2009-2010 at the same field site (Arndal et al., 2013).
362 This build-up of organic C resulted in larger basal and sucrose-induced respiration activities
363 expressed per soil weight, whereas the SOC-specific respiration did not respond to any of the

364 treatments (Table 2). Stimulating effects of elevated CO₂ on soil respiration rates (Fig. 1a) have
365 been reported at the current site of this investigation (Selsted et al., 2012), and are also well
366 described from other studies (Zak et al., 2000, van Groenigen et al., 2014).

367

368 As we hypothesized, the priming effect was more pronounced in soils exposed to elevated CO₂
369 (Fig. 1). With the increased soil C content and C:N ratios of aboveground (Vestergård et al., 2015)
370 and belowground (Arndal et al., 2013, 2014) organic inputs at elevated CO₂, it is likely that the
371 microbial N demand increased. Consequently, the enhanced priming and mineralization of SOC
372 may be a result of increased microbial N mining (Dijkstra et al. 2013; Chen et al. 2014). In line with
373 this, soils from elevated CO₂ plots at our field site exhibited higher activity of enzymes involved in
374 SOC degradation (Partavian et al., 2015). In a previous laboratory set-up with soil from the current
375 field site, Reinsch et al. (2013) assessed the temporal development of glucose-induced priming over
376 two weeks at 8°C and also found positive priming induced by labile C (glucose), with stronger
377 effects at elevated CO₂. The similar outcomes of the two studies demonstrate that the short-term (4
378 h) immediate priming response to labile C input, i.e. the priming capacity of the inherent microbial
379 community prior to microbial growth on the added labile substrate, is a relevant indicator also of
380 long-term priming effects. While Reinsch et al. (2013) only examined the occurrence of priming in
381 soils from a subset of the field treatments, i.e. the ambient (A), elevated CO₂ (CO₂) and the full
382 combination of all treatment factors (TDCO₂), we assessed the effects of all possible combinations
383 of the global change factors, i.e. elevated CO₂, warming and drought, on priming responses. In the
384 longer-term experiment, the priming effect diminished in soils exposed to elevated CO₂, drought
385 and warming in combination (Reinsch et al., 2013). In the present study we did not find a
386 comparable significant interaction between the three global change factors and potential priming of
387 SOC, although we note that the priming response in soils exposed to drought tended to be lower
388 than in soils that were not subjected to experimental drought.

389 In our investigation, we incubated soil samples from the different field treatments under
390 standardized conditions with respect to moisture, temperature and CO₂. Hence, we address whether
391 the long-term field manipulation of climate accommodated changes in the microbial decomposition
392 of SOC, which could be caused by altered availability and quality of SOC and N and/or altered
393 microbial community activity or composition. We sampled the soil immediately after the annual
394 drought treatment, where the water content in drought plots was still significantly reduced
395 (Table 1), and we hypothesized that reduced microbial activity after the drought would impair
396 priming. However, contrary to our expectation, both basal respiration and sucrose-induced
397 respiration per soil weight were enhanced by drought in combination with elevated CO₂, and
398 drought tended to increase the SOC-specific respiration (Table 2). This probably reflects a high
399 turnover of drought-decimated microorganisms upon re-wetting in the incubation experiment
400 (Groffman & Tiedje, 1988). On the other hand, drought tended to reduce the sucrose-induced
401 priming of SOC (Fig. 1). The stimulation of microbial activity upon re-wetting of soils after a
402 severe drought event thus appears uncoupled from the microbial priming of SOC in response to
403 labile C input.

404 Contrary to our hypothesis, long-term warming did not affect basal respiration, sucrose induced
405 respiration (Table 2), or potential priming (Fig. 1). In the field, warming enhanced microbial
406 abundance (Larsen et al., 2011, Haugwitz et al., 2014) and initiated earlier plant growth in the
407 spring (Kongstad et al., 2012). We expected this to result in decreased N availability, which would
408 be reflected in increased priming (Fontaine et al., 2004, 2011; Zhang et al., 2013), but we found no
409 evidence for this hypothesis. A possible reason is that eight years of warming and earlier onset of
410 spring growth of plants did not decrease soil N availability sufficiently to influence priming.

411 Further, at the field site the warming treatment only raised mean soil temperatures at 5 cm depth by
412 0.1-0.2 °C over the 3 months preceding the soil sampling (Vestergård et al., 2015). This is hardly a
413 temperature increase that would stimulate microbial activity considerably.

414

415 It is remarkable that the $\delta^{13}\text{C}$ of respiratory CO_2 derived from SOC priming in soils exposed to eight
416 years of elevated CO_2 with reduced ^{13}C was not lower than in ambient CO_2 soils (Fig. 3). In contrast,
417 the isotopic composition of total respired CO_2 from soils exposed to elevated CO_2 was ^{13}C -
418 depleted compared to CO_2 evolved from ambient CO_2 soils (Fig. 2). This shows that C assimilated
419 in the elevated CO_2 treatments was indeed decomposed in the soil basal respiration, whereas this
420 pool of newly assimilated C was not subject to primed decomposition; hence its decomposition was
421 apparently not energy limited. This implies that the primed C was assimilated more than eight years
422 before sampling. We can therefore add evidence to support previous statements that elevated CO_2
423 induces decomposition of older soil C (van Groenigen et al., 2005; Xie et al., 2005; Niklaus &
424 Falloon, 2006). Likewise, elevated CO_2 enhanced the formation of coarse particulate SOM (fresh
425 SOM) and decreased the fraction of physically protected SOM (old SOM) in forest soil (Hofmockel
426 et al., 2011) and in prairie soil (Procter et al., 2015). Given that old SOM pools contain significant,
427 yet (to a large extent) physically and chemically protected N stocks, this lends support to the
428 hypothesis that priming in response to labile C supply is a mechanism by which (some)
429 microorganisms gain access to a reservoir of N to meet their enhanced N demand under conditions
430 of ample C supply (Dijkstra et al., 2013; Chen et al., 2014). If enhanced priming at elevated CO_2 is
431 caused by increased microbial N demand, because more SOM with lower relative N content enters
432 the system at elevated CO_2 , it is reasonable that priming should be directed towards SOM pools
433 with a higher N content, i.e. SOM pools incorporated into the system before the elevated CO_2
434 treatment was initiated.

435 Bulk SOC encompasses different pools of SOC of varying age and particle size, and the $\delta^{13}\text{C}$ of
436 these different pools vary considerably (Gerzabek et al., 2001). As expected, the SOC $\delta^{13}\text{C}$ was
437 decreased from -27.8 ‰ in ambient CO_2 soil to -29.3 ‰ in elevated CO_2 soil, whereas drought and
438 warming did not affect the isotopic composition of SOC. The lower $\delta^{13}\text{C}$ of the C primed in the

439 ambient plots and plots subjected to drought as a single factor compared to the other treatments
440 (Fig. 3), therefore suggests that sucrose-amendment primed the decomposition of different SOC
441 pools in the different treatments.

442 It has been argued that short-term incubations as employed in the current study reflects ‘apparent’
443 rather than ‘real’ priming effects. Theoretically, apparent priming is a state, where the initial
444 enhanced respiratory pulse induced by labile C addition, derives from turn-over of microbial
445 biomass C rather than decomposition of SOC, i.e. part of the inherent microbial biomass C pool is
446 substituted by the added labile C. ‘Real priming’, on the other hand, describes the enhanced
447 decomposition of SOC after prolonged incubation with labile C (Blagodatskaya & Kuzyakov, 2008;
448 Blagodatsky et al., 2010). We argue, though, that the finding that the primed C was at least eight
449 years old is strong indication that even in our short-term incubation study, the addition of labile C
450 resulted in real priming; i.e. the enhanced decomposition of SOC. If the sucrose-induced priming
451 did indeed represent apparent priming, it would imply that the pool-substituted microbial biomass C
452 was more than eight years old. Microbial biomass turnover is on average much faster than eight
453 years, and it is quite unlikely that microorganisms grow preferentially on older C pools. Therefore,
454 we find it most plausible that the enhanced soil-derived CO₂-C flux represents real priming of SOC
455 rather than pool substitution of microbial biomass C.

456 It has been suggested that increased primary production at elevated CO₂ will enhance C
457 sequestration in terrestrial ecosystems and thus counteract the rise in atmospheric CO₂
458 concentration (Oren et al., 2001; Jastrow et al., 2005; Houghton, 2007). However, this and other
459 studies (Carney et al., 2007; van Groenigen et al., 2014) demonstrate that ecosystems exposed to
460 elevated CO₂ concentrations will be more prone to SOC decomposition triggered by labile C input.
461 This will thus reduce the anticipated increase in C sequestration. In our heath/grassland system
462 elevated CO₂ did enhance the C input to the system and hence the SOC pool (Table 1). However,
463 we demonstrate that labile C inputs accelerate the turnover of older SOC pools and alter C

464 dynamics of the system under elevated CO₂. Therefore, in the longer term, the net C balance of this
465 and other systems in a high CO₂ world will depend on the extent to which the build up of new
466 organic C will compensate for the increased loss of older organic C pools.

467

468 **Acknowledgements**

469 We thank colleagues from the CLIMAITE project for collaboration during sampling for this study.
470 The CLIMAITE experiment was funded by the Villum Foundation, DONG Energy and Air
471 Liquide. MV was also supported by the Danish Council for Strategic Research (ASHBACK, DSF-
472 12-132655) and the Danish Council for Independent Research (OP-RICE-ING, DFF-4002-00274),
473 and PB was supported by grants from the Swedish Research Council Formas (grant number 2012-
474 1541).

475

476 **References**

477 Adair, E.C., Reich, P.B., Hobbie, S.E., Knops, J.M.H., 2009. Interactive effects of time, CO₂, N,
478 and diversity on total belowground carbon allocation and ecosystem carbon storage in a grassland
479 community. *Ecosystems* 6, 1037-1052.

480

481 Ainsworth, E.A., Rogers, A., 2007. The response of photosynthesis and stomatal conductance to
482 rising [CO₂]: mechanisms and environmental interactions. *Plant, Cell and Environment* 30, 258-
483 270.

484

485 Albert, K.R., Ro-Poulsen, H., Mikkelsen, T.N., Michelsen, A., van der Linden, L., Beier, C., 2011.
486 Effects of elevated CO₂, warming and drought episodes on plant carbon uptake in a temperate heath
487 ecosystem are controlled by soil water status. *Plant, Cell & Environment* 34, 1207-1222.

488

489 Arndal, M.F., Merrild, M.P., Michelsen, A., Schmidt, I.K., Mikkelsen, T.N. & Beier, C., 2013. Net
490 root growth and nutrient acquisition in response to predicted climate change in two contrasting
491 heathland species. *Plant and Soil* 369, 615-629.

492

493 Arndal, M.F., Schmidt, I.K., Kongstad, J., Beier, C. & Michelsen, A., 2014. Root growth and N
494 dynamics in response to multi-year experimental warming, summer drought and elevated CO₂ in a
495 mixed heathland-grass ecosystem. *Functional Plant Biology* 41, 1-10.

496

497 Bengtson, P., Barker, J., Grayston, S.J., 2012. Evidence of a strong coupling between root
498 exudation, C and N availability, and stimulated SOM decomposition caused by rhizosphere priming
499 effects. *Ecology and Evolution* 2, 1843-1852.

500

501 Blagodatskaya, E., Kuzyakov, Y., 2008. Mechanisms of real and apparent priming effects and their
502 dependence on soil microbial biomass and community structure: critical review. *Biology and
503 Fertility of Soils* 45, 115-131.

504

505 Blagodatsky, S., Blagodatskaya, E., Yuyukina, T., Kuzyakov, Y., 2010. Model of apparent and real
506 priming effects: Linking microbial activity with soil organic matter decomposition. *Soil Biology &
507 Biochemistry* 42, 1275-1283.

508

509 Carney, K.M., Hungate, B.A., Drake, B.G., Megonigal, J.P., 2007. Altered soil microbial
510 community at elevated CO₂ leads to loss of soil carbon. *Proceedings of the National Academy of
511 Sciences* 104, 4990–4995.

512

513 Chapin III, F.S., Matson, P.A., Mooney, H.A., 2002. Principles of Terrestrial Ecosystem Ecology.
514 Springer, New York.
515

516 Chen, R.R., Senbayram, M., Blagodatsky, S., Myachina, O., Dittert, K., Lin, X.G., Blagodatskaya,
517 E., Kuzyakov, Y., 2014. Soil C and N availability determine the priming effect: microbial N mining
518 and stoichiometric decomposition theories. *Global Change Biology* 20, 2356-2367.
519

520 Cheng, W.X., Parton, W.J., Gonzalez-Meler, M.A., Phillips, R., Asao, S., McNickle, G.G., Brzostek,
521 E., Jastrow, J.D., 2014. Synthesis and modeling perspectives of rhizosphere priming. *New*
522 *Phytologist* 201, 31-44.
523

524 Dijkstra, F.A., Cheng, W., 2007. Moisture modulates rhizosphere effects on C decomposition in
525 two different soil types. *Soil Biology & Biochemistry* 39, 2264-2274.
526

527 Dijkstra, F.A., Carrillo, Y., Pendall, E., Morgan, J.A., 2013. Rhizosphere priming: a nutrient
528 perspective. *Frontiers in Microbiology* 4, article 216.
529

530 Field, C.B., Jackson, J.A., Vilgalys, R., Jackson, R.B., 1995. Stomatal responses to increased CO₂:
531 implications from the plant to the global scale. *Plant, Cell and Environment* 18, 1214-1225.
532

533 Fontaine, S., Bardoux, G., Abbadie, L., Mariotti, A., 2004. Carbon input to soil may decrease soil
534 carbon content. *Ecology Letters* 7, 314-320.
535

536 Fontaine, S., Henault, C., Aamor, A., Bdioui, N., Bloor, J.M.G., Maire, V., Mary, B., Revalliot, S.,
537 Maron, P.A., 2011. Fungi mediate long term sequestration of carbon and nitrogen in soil through
538 their priming effect. *Soil Biology & Biochemistry* 43, 86-96.
539

540 Garcia-Pausas, J., Paterson, E., 2011. Microbial community abundance and structure are
541 determinants of soil organic matter mineralisation in the presence of labile carbon. *Soil Biology &*
542 *Biochemistry* 43, 1705-1713.
543

544 Gerzabek, M.H., Habernauer, G., 2001. Soil organic matter pools and carbon-13 natural abundances
545 in particle-size fraction of a long-term agricultural field experiment receiving organic amendments.
546 *Soil Science Society of America Journal* 65, 352-358.
547

548 Ghee, C., Neilson, R., Hallett, P.D., Robinson, D., Paterson, E., 2013. Priming of soil organic
549 matter mineralisation is intrinsically insensitive to temperature. *Soil Biology & Biochemistry* 66,
550 20-28.
551

552 Grayston, S.J., Wang, S., Campbell, C.D., Edwards, A.C., 1998. Selective influence of plant species
553 on microbial diversity in the rhizosphere. *Soil Biology & Biochemistry* 30, 369-378.
554

555 Groffman, P.M., Tiedje, J.M., 1988. Denitrification hysteresis during wetting and drying in soil.
556 *Soil Science Society of America Journal* 52, 1626-1629.
557

558 Gude, A., Kandeler, E., Gleixner, G., 2012. Input related microbial carbon dynamic of soil organic
559 matter in particle size fractions. *Soil Biology & Biochemistry* 47, 209-219.
560

561 Guenet, B., Neill, C., Bardoux, G., Abbadie, L., 2010. Is there a linear relationship between priming
562 effect intensity and the amount of organic matter input? *Applied Soil Ecology* 46, 436-442.
563

564 Haugwitz, M.S., Bergmark, L., Priemé, A., Christensen, S., Beier, C., Michelsen, A., 2014. Soil
565 microorganisms respond to five years of climate change manipulations and elevated atmospheric
566 CO₂ in a temperate heath ecosystem. *Plant and Soil* 374, 211-222
567

568 Hofmockel, K.S., Zak, D.R., Moran, K.K., Jastrow, J.D., 2011. Changes in forest soil organic
569 matter pools after a decade of elevated CO₂ and O₃. *Soil Biology & Biochemistry* 43, 1518-1527.
570

571 Houghton, R.A., 2007. Balancing the Global Carbon Budget. *Annual Review of Earth and Planetary*
572 *Sciences* 35, 313–347.
573

574 Hungate, B.A., Holland, E.A., Jackson, R.B., Chapin III, F.S., Mooney, H.A., Field, C.B., 1997.
575 The fate of carbon in grasslands under carbon dioxide enrichment. *Nature* 388, 576–579.
576

577 IPCC, 2007. *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I*
578 *to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.* Cambridge
579 University Press, Cambridge, UK and New York.
580

581 IPCC, 2013. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I*
582 *to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.* Cambridge
583 University Press, Cambridge, UK and New York.
584

585 Jastrow, J.D., Miller, R.M., Matamala, R., Norby, R.J., Boutton, T.W., Rice, C.W., Owensby, C.E.,
586 2005. Elevated atmospheric carbon dioxide increases soil carbon. *Global Change Biology* 11,
587 2057–2064.

588

589 Kassem II, J.P., Sigler, V., Heckathorn, S., Wang, Q., 2008. Effect of elevated CO₂ and drought on
590 soil microbial communities associated with *Andropogon gerardii*. *Journal of Integrative Plant*
591 *Biology* 50, 1406-1415.

592

593 Kongstad, J., Schmidt, I.K., Riis-Nielsen, T., Arndal, M.F., Mikkelsen, T.N., Beier, C., 2012 High
594 resilience in heathland plants to changes in temperature, drought, and CO₂ in combination: Results
595 from the CLIMAITE experiment. *Ecosystems* 15, 269-283.

596

597 Larsen, K.S., Andresen, L.C., Beier, C., Jonasson, S., Albert, K.R., Ambus, P., Arndal, M., Carter,
598 M.S., Christensen, S., Holmstrup, M., Ibrom, A., Kongstad, J., van der Linden, L., Maraldo, K.,
599 Michelsen, A., Mikkelsen, T.N., Pilegaard, K., Priemé, A., Ro-Poulsen, H., Schmidt, I.K., Selsted,
600 M.B., Stevnbak, K., 2011. Reduced N cycling in response to elevated CO₂, warming, and drought
601 in a Danish heathland: Synthesizing results of the CLIMAITE project after two years of treatments.
602 *Global Change Biology* 17, 1884-1899.

603

604 Luo, Y., Su, B., Currie, W.S., Zak, D.R., Dukes, J.S., Finzi, A., Hartwig, U., Hungate, B.,
605 McMurtrie, R.E., Oren, R., Parton, W.J., Pataki, D.E., Shaw, M.R., Field, C.B., 2004. Progressive
606 nitrogen limitation of ecosystem responses to rising atmospheric carbon dioxide. *BioScience* 54,
607 731-739.

608

609 Mikkelsen, T.N., Beier, C., Jonasson, S., Holmstrup, M., Schmidt, I.K., Ambus, P., Pilegaard, K.,
610 Michelsen, A., Albert, K., Andresen, L.C., Arndal, M.F., Bruun, N., Christensen, S., Danbæk, S.,
611 Gundersen, P., Jørgensen, P., Linden, J.G., Kongstad, J., Maraldo, K., Priemé, A., Riis-Nielsen, T., Ro-
612 Poulsen, H., Stevnbak, K., Selsted, M.B., Sørensen, P., Larsen, K.S., Carter, M.S., Ibrom, A.,
613 Martinussen, T., Miglietta, F., Sverdrup, H., 2008. Experimental design of multifactor climate change
614 experiments with elevated CO₂, warming and drought: the CLIMAITE project. *Functional Ecology* 22,
615 185-195.

616

617 Moyano, F.E., Manzoni, S., Chenu, C., 2013. Responses of soil heterotrophic respiration to
618 moisture availability: An exploration of processes and models. *Soil Biology & Biochemistry* 59, 72-
619 85.

620

621 Niklaus, P.A., Falloon, P., 2006. Estimating soil carbon sequestration under elevated CO₂ by
622 combining carbon isotope labelling with soil carbon cycle modelling. *Global Change Biology* 12,
623 1909-1921.

624

625 Oren, R., Ellsworth, D.S., Johnsen, K.H., Phillips, N., Ewers, B.E., Maier, C., Schäfer, K.V.R.,
626 McCarthy, H., Hendrey, G., McNulty, S.G., Katul, G.G., 2001. Soil fertility limits carbon
627 sequestration by forest ecosystems in a CO₂-enriched atmosphere. *Nature* 411, 469–472.

628

629 Partavian, A., Mikkelsen, T.N., Vestergård, M., 2015. Plants increase laccase activity in soil with
630 long-term elevated CO₂ legacy. *European Journal of Soil Biology* 70, 97-103.

631

632 Paterson, E., Thornton, B., Midwood, A.J., Osborne, S.M., Sim, A., Millard, P., 2008. Atmospheric
633 CO₂ enrichment and nutrient additions to planted soil increase mineralisation of soil organic matter,

634 but do not alter microbial utilisation of plant- and soil C-sources. *Soil Biology & Biochemistry* 40,
635 2434-2440.

636

637 Paterson, E., 2009. Comments on the regulatory gate hypothesis and implications for C-cycling in
638 soil. *Soil Biology & Biochemistry* 41, 1352-1354.

639

640 Paterson, E., Sim, A., 2013. Soil-specific response functions of organic matter mineralization to the
641 availability of labile carbon. *Global Change Biology* 19, 1562-1571.

642

643 Perveen, N., Barot, S., Alvarez, G., Klumpp, K., Martin, R., Rapaport, A., Herfurth, D., Louault, F.,
644 Fontaine, S., 2014. Priming effect and microbial diversity in ecosystem functioning and response to
645 global change: a modeling approach using the SYMPHONY model. *Global Change Biology* 20,
646 1174-1190.

647

648 Phillips, R.P., Meier, I.C., Bernhardt, E.S., Grandy, A.S., Wickings, K., Finzi, A.C., 2012. Roots
649 and fungi accelerate carbon and nitrogen cycling in forests exposed to elevated CO₂. *Ecology*
650 *Letters* 15, 1042-1049.

651

652 Procter, A.C., Gill, R.A., Fay, P.A., Polley, H.W., Jackson, R.B., 2015. Soil carbon responses to
653 past and future CO₂ in three Texas prairie soils. *Soil Biology & Biochemistry* 83, 66-75.

654

655 Reinsch, S., Ambus, P., 2013. In situ ¹³CO₂ pulse-labeling in a temperate heathland - Development
656 of a mobile multi-plot field setup. *Rapid Communications in Mass Spectrometry* 27, 1417-1428.

657

658 Reinsch, S., Ambus, P., Thornton, B., Paterson, E., 2013, Impact of future climatic conditions on
659 the potential for soil organic matter priming. *Soil Biology & Biochemistry*, 133-140.
660

661 Scherber, C., Gladbach, D., Stevnbak, K., Karsten, R.J., Schmidt, I.K., Michelsen, A., Albert, K.R.,
662 Larsen, K.S., Mikkelsen, T.N., Beier, C., Christensen, S., 2013. Multi-factor climate change effects
663 on insect herbivore performance. *Ecology and Evolution* 3, 1449-1460.
664

665 Selsted, M.B., van der Linden, L., Ibrom, A., Michelsen, A., Larsen, K.S., Pedersen, J.K.,
666 Mikkelsen, T.N., Pilegaard, K., Beier, C., Ambus, P., 2012. Soil respiration is stimulated by
667 elevated CO₂ and reduced by summer drought: three years of measurements in a multifactor
668 ecosystem manipulation experiment in a temperate heathland (CLIMAITE). *Global Change*
669 *Biology* 18, 1216-1230.
670

671 Sullivan, B.W., Hart, S.C., 2013. Evaluation of mechanisms controlling the priming of soil carbon
672 along a substrate age gradient. *Soil Biology & Biochemistry* 58, 293-301.
673

674 van Groenigen, K.J., Gorissen, A., Six, J., Harris, D., Kuikman, P.J., van Groenigen, J.W., van
675 Kessel, C., 2005. Decomposition of C-14-labeled roots in a pasture soil exposed to 10 years of
676 elevated CO₂. *Soil Biology & Biochemistry* 37, 497-506.
677

678 van Groenigen, K.J., Qi, X., Osenberg, C.W., Luo, Y., Hungate, B.A., 2014. Faster decomposition
679 under increased atmospheric CO₂ limits soil carbon storage. *Science* 344, 508-509.
680

681 Vestergård, M., Dyrnum, K., Michelsen, A., Damgaard, C., Holmstrup, M., 2015. Long-term
682 multifactorial climate change impacts on mesofaunal biomass and nitrogen content. *Applied Soil*
683 *Ecology* 92, 54-63
684

685 Wang, X., Liu, L., Piao, S., Janssens, I.A., Tang, J., Liu, W., Chi, Y., Wang, J., Xu, S., 2014. Soil
686 respiration under climate warming: differential response of heterotrophic and autotrophic
687 respiration. *Global Change Biology* 20, 3229-3237.
688

689 Wu, J., Brookes, P.C., Jenkinson, D.S., 1993. Formation and destruction of microbial biomass
690 during the decomposition of glucose and ryegrass in soil. *Soil Biology & Biochemistry* 25, 1435-
691 1441.
692

693 Xiao, C., Guenet, B., Zhou, Y., Su, J., Janssens, I.A., 2015. Priming of soil organic matter
694 decomposition scales linearly with microbial biomass response to litter input in steppe vegetation.
695 *Oikos* 124, 649-657.
696

697 Xie, Z.B., Cadisch, G., Edwards, G., Baggs, E.M., Blum, H., 2005. Carbon dynamics in a temperate
698 grassland soil after 9 years exposure to elevated CO₂ (Swiss FACE). *Soil Biology & Biochemistry*
699 37, 1387-1395.
700

701 Zak, D.R., Pregitzer, K.S., King, J.S., Holmes, W.E., 2000. Elevated atmospheric CO₂, fine roots
702 and the response of soil microorganisms: a review and hypothesis *New Phytologist* 147, 201-222.
703

704 Zhang, W.D., Wang, X.F., Wang, S.L., 2013. Addition of external organic carbon and native soil
705 organic carbon decomposition: a meta-analysis. *PLoS ONE* 8, e54779.

706

707 Zyakun, A.M., Dilly, O., 2005. Use of carbon isotope composition for characterization of microbial
708 activity in arable soils. *Applied Biochemistry and Microbiology*5, 512-520.

709

710

711

712

713

714

715

716

717

718

719

720

721

722

723

724

725

726

727

728

729

730

731 **Figure legends**

732 Figure 1

733 Mean priming effect assessed as the increase in SOC decomposition rate in soils amended with
734 sucrose compared to soils incubated with water. Soils were collected from field plots in a Danish
735 grass/heathland exposed to 8 years of ambient conditions (A), annually repeated spring/early
736 summer drought (D), warming (T), elevated atmospheric CO₂ (CO₂) and all possible combinations
737 of single factors. Error bars depict SE. *n*=4-5.

738

739 Figure 2

740 Mean $\delta^{13}\text{C}$ values of CO₂-C respired during incubation of soils collected from field plots in a
741 Danish grass/heathland exposed to 8 years of ambient conditions (A), annually repeated
742 spring/early summer drought (D), warming (T), elevated atmospheric CO₂ (CO₂) and all possible
743 combinations of single factors. Soils were incubated with water (a) or with a sucrose solution (b).
744 Error bars depict SE. *n*=4-5.

745

746 Figure 3

747 Mean $\delta^{13}\text{C}$ values of SOC-derived CO₂-C primed by sucrose addition during incubation of soils
748 collected from field plots in a Danish grass/heathland exposed to 8 years of ambient conditions (A),
749 annually repeated spring/early summer drought (D), warming (T), elevated atmospheric CO₂ (CO₂)
750 and all possible combinations of single factors. Treatments A and D, marked with asterisk, are
751 significantly different from the other treatments (Tukey *P*<0.05). Error bars depict SE. *n*=4-5.

752

753

Table 1. Mean water content and C content in soils collected from field plots in a Danish grass/heathland exposed to 8 years of ambient conditions (A), annually repeated spring/early summer drought (D), warming (T), elevated CO₂ (CO₂) and all possible combinations of single factors. SE depicted in parentheses. *n*=4-5.

| | <u>Soil water content (%)</u> | | <u>Soil C content (%)</u> | |
|-------------------|---|--------|--|--------|
| A | 10.30 | (0.42) | 3.15 | (0.29) |
| D | 8.04 | (1.32) | 3.08 | (0.32) |
| T | 10.69 | (1.00) | 3.00 | (0.23) |
| TD | 6.71 | (0.53) | 2.89 | (0.23) |
| CO ₂ | 11.86 | (0.50) | 3.52 | (0.33) |
| DCO ₂ | 9.27 | (0.69) | 4.80 | (0.41) |
| TCO ₂ | 11.68 | (1.16) | 3.65 | (0.60) |
| TDCO ₂ | 11.24 | (2.50) | 5.75 | (1.28) |
| Treatment effects | P _{CO₂} =0.024 P _D =0.012 | | P _{CO₂} =0.002 P _{CO₂xD} =0.038 | |

Table 2. Mean basal and sucrose-induced respiration in relation to soil dry weight and soil organic C (SOC) content during incubation of soils collected from field plots in a Danish grass/heathland exposed to 8 years of ambient conditions (A), annually repeated spring/early summer drought (D), warming (T), elevated atmospheric CO₂ (CO₂) and all possible combinations of single factors. Sucrose-induced respiration is the difference between respiration activity in sucrose-amended and control samples. SE depicted in parentheses. *n*=4-5.

| | Basal respiration ($\mu\text{g CO}_2\text{-C g soil}^{-1} \text{ h}^{-1}$) | | Sucrose-induced respiration ($\mu\text{g CO}_2\text{-C g soil}^{-1} \text{ h}^{-1}$) | | SOC-specific basal respiration ($\mu\text{g CO}_2\text{-C g SOC}^{-1} \text{ h}^{-1}$) | | SOC-specific sucrose- induced respiration ($\mu\text{g CO}_2\text{-C g SOC}^{-1} \text{ h}^{-1}$) | |
|-------------------|---|--------|--|--------|--|---------|---|---------|
| A | 2.05 | (0.38) | 5.07 | (0.51) | 63.44 | (6.45) | 160.51 | (3.57) |
| D | 2.00 | (0.31) | 4.69 | (0.78) | 65.63 | (10.02) | 158.85 | (30.94) |
| T | 1.97 | (0.24) | 5.00 | (0.53) | 65.34 | (4.99) | 166.12 | (9.85) |
| TD | 1.84 | (0.21) | 3.98 | (0.37) | 64.05 | (4.95) | 138.67 | (8.36) |
| CO ₂ | 2.17 | (0.16) | 5.76 | (0.73) | 62.76 | (4.45) | 162.63 | (9.50) |
| DCO ₂ | 3.51 | (0.47) | 8.23 | (1.04) | 72.45 | (5.20) | 169.88 | (11.04) |
| TCO ₂ | 2.27 | (0.49) | 5.59 | (1.03) | 61.22 | (5.32) | 152.83 | (13.68) |
| TDCO ₂ | 4.39 | (0.81) | 8.49 | (1.10) | 80.18 | (6.41) | 162.39 | (17.79) |

Figure 1




Figure 2




Figure 3

