



Al-Ghafari, Ayat B. and Punjaruk, Wiyada and Storer, L.C.D. and Carrier, D.J. and Hussein, D. and Coyle, Beth and Kerr, Ian D. (2016) Long-term exposure to irinotecan reduces cell migration in glioma cells. *Journal of Neuro-Oncology*, 127 (3). pp. 455-462. ISSN 1573-7373

Access from the University of Nottingham repository:

<http://eprints.nottingham.ac.uk/34240/1/Al-Ghafari-final%20draft%20post-refereeing.pdf>

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see:
http://eprints.nottingham.ac.uk/end_user_agreement.pdf

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

[Click here to view linked References](#)**J NeuroOncol**

Long-term exposure to irinotecan reduces cell migration in glioma cells

Al-Ghafari, A.B^{1*}, Punjaruk, W^{2*}, Storer, L², Carrier, D.J.², Hussein, D², Coyle, B^{2#}, Kerr, I.D^{1#}.

¹School of Life Sciences, Queen's Medical Centre, University of Nottingham, Nottingham, NG7 2UH

²School of Medicine, Queen's Medical Centre, University of Nottingham, Nottingham, NG7 2UH

* - these authors contributed equally to data generation.

- authors to whom correspondence may be addressed.

ian.kerr@nottingham.ac.uk tel: 00 44 8230122

Beth.coyle@nottingham.ac.uk tel: 00 44 8230719

Keywords: glioma; topoisomerase; etoposide; irinotecan; ABCB1; invasion; chemotherapy

1
2 **Introduction**
3

4
5 Glioblastoma multiforme (GBM) is the most common malignant brain tumour in adults (annual
6 incidence 3 per 100,000; [1]). It carries a very poor prognosis with mean overall survival of 12-17
7 months. Despite testing a multitude of chemotherapy agents and strategies, very little progress has
8 been made in extending the life expectancy of those diagnosed with a GBM, since the establishment
9 of the Stupp protocol (a combination of radiotherapy and temozolomide) a decade ago [2, 3]. This is
10 believed to be due to a high level of intrinsic drug resistance [4] combined with the fact that these
11 tumours tend to diffusely invade the local brain parenchyma making it difficult to identify the
12 perimeter and effectively remove them surgically [5-7]. An improvement in clinical outcome
13 therefore requires identification of drugs that can inhibit invasion as well as promote cell death.
14
15

16
17 Antineoplastic effects of topoisomerase inhibitors have been observed in glioma cells.
18 Topoisomerases are enzymes involved in cleaving and religating supercoiled DNA. Topoisomerase I
19 does this through making single strand breaks whereas topoisomerase II breaks both strands.
20 Inhibition of either enzyme results in cell death and cancer cells have been shown to be selectively
21 sensitive to topoisomerase inhibitors. Indeed there are a number of ongoing glioma clinical trials
22 exploring the combination of established inhibitors with other marketed drugs e.g. the phase II
23 clinical trial NCT01574092 is assessing irinotecan (topoisomerase I inhibitor) in combination with
24 cisplatin. In addition to which, small molecule inhibitors such as the topoisomerase II inhibitor
25 Banoxantrone (phase I/II) or the topoisomerase I inhibitors Cositecan (phase I) and Gimatecan
26 (phase I) are also currently in clinical trials [8]. Over time, however, studies have shown that many
27 tumours become resistant to topoisomerase II inhibitors due to increased DNA repair or acquired
28 mutations in topoisomerase II [9, 10], although whether this is paralleled by an increase in general
29 drug resistance mechanisms (e.g. multidrug efflux pumps) is unknown.
30
31

32
33 In this study, we set out to determine the cellular effects of long-term exposure to topoisomerase
34 inhibitors in a glioma cell line (C6), focusing on acquired cross-resistance and invasion. The C6
35 glioma cell line was grown out from a glioma induced in an outbred Wistar rat strain by repeated
36 rounds of treatment with N'N'-methylnitrosourea [11, 12]. Injection of the C6 cell line into newborn
37 rats produced a tumour with pathology consistent with a glioblastoma multiforme including a
38 diffuse invading border [13]. Since then it has been used as an experimental model in a number of
39 different studies (reviewed in [14]). C6 cells were continuously exposed to either a topoisomerase I
40 inhibitor (irinotecan) or a topoisomerase II inhibitor (etoposide). In common with other studies we
41 found that continuous treatment with etoposide resulted in increased resistance to this agent, but
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 did not induce cross-resistance to irinotecan. In contrast, even after many rounds of exposure to the
2 irinotecan, sensitivity to this drug was maintained and, importantly, we also observed a reduction in
3 migratory capacity of treated cells. This data argues for the further clinical investigation of
4 topoisomerase I inhibition as a mechanism to limit glioma invasion without incurring increased
5 resistance to chemotherapy.
6
7
8
9

10 11 **Materials and Methods**

12 13 **Cell culture**

14
15 The rat C6 glioma cell line was maintained at 37°C/5% CO₂ in Dulbecco's Modified Eagle Medium
16 (DMEM) (Invitrogen) with 10% foetal bovine serum (FBS). Drug selected cell lines (C6-etoposide and
17 C6-irinotecan) were established by continuous culture over 5 passages at the drug concentration
18 established to cause death of 70% of cells i.e. in the presence of 16 µM etoposide (Sigma) and 10 µM
19 irinotecan (Sigma). In the case of etoposide selection, cells were incubated for 2 hours [15] and then
20 media was changed while for irinotecan selection, cells were incubated for 96 hours then media was
21 changed [16]. Similarly, in experiments with prolonged duration such as clonogenic assays, the drugs
22 were removed after 2 hours and 96 hours for etoposide and irinotecan, respectively.
23
24
25
26
27
28
29
30
31
32
33
34

35 36 **Western blot analysis**

37
38 Cell pellets were resuspended in PBS/glycerol (10% v/v) supplemented with EDTA free protease
39 inhibitor cocktail set III (Calbiochem) and lysed by probe sonication for 3 x 10-20 second bursts.
40 Equal quantities of total protein were resolved on 8-12% w/v polyacrylamide gels and protein was
41 transferred to Hybond™-P PVDF membrane (Amersham Biosciences). Membranes were blocked
42 with 5% w/v non-fat Marvel milk in TBST buffer (25 mM Tris pH 7.6, 0.15 M NaCl, 0.1% v/v Tween-
43 20) at room temperature for an hour followed by incubation with primary antibodies (dilutions given
44 in figure legends) at 4°C overnight. The membranes were then washed with TBST and incubated for
45 1 hour at room temperature with horseradish peroxidase conjugated secondary antibodies.
46 Following washing, the specific proteins were detected using a SuperSignal® West Pico
47 Chemilluminescent Substrate (Thermo Scientific). Where band intensity was quantified, the density
48 of unsaturated exposures was determined with Scion Image software (NIH, USA) after scanning the
49 blots at 600 dpi resolution.
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Quantitative RT- PCR analysis

1
2 RNA was isolated using a *mirVana*[™] miRNA isolation kit (Ambion) and transcribed into cDNA using
3 Superscript III reverse transcriptase (Invitrogen). Quantitative reverse transcription PCR (qPCR)
4 analysis was carried out using a CFX96 real-time PCR machine (Biorad) and iQ SYBR Green Superrmix
5 (Biorad; primers in Supplementary Table 1). Samples were analysed in triplicate and data obtained
6 from three independent experiments were used for analysis of relative gene expression compared to
7 GAPDH and β -Actin [17] with the Pfaffl equation [18].
8
9
10
11
12
13
14
15
16

Clonogenic assay

17 Viability was determined following an established clonogenic assay protocol [19]. Approximately 100
18 cells were seeded in a 2 ml volume into a 6 well plate and were treated with etoposide (5 - 80 μ M),
19 irinotecan (0.5 - 24 μ M) or a vehicle control (DMSO either 0.08% v/v or 0.02% v/v for etoposide and
20 irinotecan respectively). Plates were incubated for 7-10 days until colonies containing approximately
21 50 cells were visible in control wells. Subsequently, media was removed and colonies were fixed
22 with paraformaldehyde (4% (w/v) PFA in PBS, 20 minutes) and stained with crystal violet (0.1% (w/v)
23 in distilled H₂O, 5 minutes), prior to colony counting.
24
25
26
27
28
29
30
31
32
33
34

Flow cytometry

35 Single cell suspensions (2×10^6 cells/ 0.1 ml) in fluorescent activated cell sorting (FACS) buffer (0.5%
36 w/v BSA in phenol red free DMEM) were incubated with rhodamine 123 (2 μ M) and periodically
37 inverted during a 30 minute incubation at 37°C. Cells were washed twice by pelleting (250g, 3
38 minutes, 4°C) and resuspension in FACS buffer. Cellular fluorescence was determined using a
39 Cytomics FC500 (Beckman Coulter) flow cytometer and data analysed with WinMDI (Scripps
40 Institute).
41
42
43
44
45
46
47
48
49
50
51

Reactive oxygen species (ROS) assay

52 Black clear-bottom 96-well plates (Greiner Bio-one) were seeded with 10,000 cells per well. After
53 overnight incubation, media was replaced by Hanks Balanced Salt Solution (HBSS) supplemented
54 with 50 μ M 2',7'-dichlorofluorescein diacetate (DCFH-DA; Invitrogen) and the plate was incubated in
55 a 37°C/5% CO₂ incubator for 30 minutes. Excess DCFH-DA was removed by washing with HBSS. ROS
56
57
58
59
60
61
62
63
64
65

1 production was then induced by incubation with indicated concentrations of tert-butyl
2 hydroperoxide (TBHP) (Sigma). The increase in DCF fluorescence (indicative of ROS production) was
3 measured (λ_{Ex} 485nm; λ_{Em} 530nm) on a Flexstation 3 fluorescence microplate reader (Molecular
4 Devices, USA) over a 3 hour period. Rates of ROS production were determined across the linear
5 region of the time course (typically 60-90 minutes) and fold-changes in the rate of ROS production
6 were determined compared to cells without TBHP treatment.
7
8
9

10 11 12 13 **Wound scratch assay**

14 Cells (0.5×10^6 in DMEM containing 10% FBS) were seeded in duplicate sterile 6-well plates and
15 incubated overnight in 37°C/5% CO₂. A scratch was then made across the centre of each well with a
16 sterile 200 μ l pipette tip. Cell migration was assessed with time lapse microscopy (LEICA DMIRB
17 Microscope, LEICA Microsystem) at 37°C for up to 65 hours. Images were taken with brightfield 10x
18 phase contrast objective lens, every 30 minutes at three different positions along each scratch. All
19 images were analysed using ImageJ software to determine the wound size relative to the initial
20 scratch.
21
22
23
24
25
26
27
28
29
30
31

32 33 **Transwell migration assay**

34 A transwell migration assay was performed to determine the migration response under the influence
35 of a chemo-attractant (10% FBS). Experiments were performed using the ChemoTx®System kit
36 (Neuro Probe) according to the manufacturer's instructions. Cells (2×10^4) were added to the upper
37 filter and migration towards the chemo-attractant (with HBSS as control) occurred during incubation
38 for 24 hours (37°C/5% CO₂). Following the incubation, the media was aspirated and cells were fixed
39 (4% PFA in PBS, 20 minutes), prior to nuclear DNA staining with Hoechst 33342 for 5 minutes. Filters
40 were washed once with PBS and the microplate was scanned on an ImageXpressMICRO (Molecular
41 Devices, USA) to determine the total number of migrated cells.
42
43
44
45
46
47
48
49
50
51
52

53 **Statistical analysis**

54 All experiments were conducted on at least three independent occasions. GraphPad Prism
55 (GraphPad Software, San Diego California USA) was used for all data fitting and statistical data
56 analysis. Clonogenic assay data were fitted to the general sigmoidal dose response equation. Wound
57
58
59
60
61
62
63
64
65

1 closure data was fitted using exponential decay equations. ROS production rates were determined
2 using linear regression. Where data from multiple experiments are displayed in column format the
3 error bars represent the standard error of the mean. Statistical significance was analysed by one way
4 ANOVA or unpaired t-test (as described in figure legends) to compare between cell lines. A *P* value
5 smaller than 0.05 ($P < 0.05$) was considered significant.
6
7
8
9

10 11 **Results:**

12
13 Understanding of glioma drug resistance and local invasion is critical if we are to develop better
14 therapies for this type of brain tumour. We were interested in observing whether changes in drug
15 resistance and invasion occurred, in a model glioma cell line (C6), following exposure to
16 topoisomerase I and topoisomerase II inhibition. Initially, we explored whether etoposide
17 (topoisomerase II inhibitor) and irinotecan (topoisomerase I inhibitor) were cytotoxic against the C6
18 cell line and used a clonogenic assay to determine $IC_{50,70}$ values (the concentration of drug required
19 to achieve 50 or 70%-maximal inhibition of colony growth; Figure 1). The colony formation of the C6
20 cell line following a single exposure to either drug could be inhibited by low micromolar
21 concentrations of both drugs (IC_{50} for etoposide $8.6 \pm 0.6 \mu\text{M}$; IC_{50} for irinotecan $5.3 \pm 0.4 \mu\text{M}$). Since
22 chemotherapy is normally given to patients in cycles, consisting of a weekly dose for several weeks
23 then a period of rest followed by further cycles, we investigated how C6 cells would respond to
24 chronic cyclical exposure to topoisomerase inhibitors at cytotoxic concentrations. Therefore, we
25 cultured cells for multiple passages in the presence of IC_{70} concentrations of etoposide or irinotecan
26 ($16 \mu\text{M}$ and $8 \mu\text{M}$ respectively), resulting in the production of sub-lines called C6-Et (for a cell line
27 cultured in the presence of etoposide) and C6-Ir (for a cell line cultured in the presence of irinotecan.
28 A further cell line (C6-V) was established after continuous culture in the presence of DMSO (0.05%
29 v/v), the solvent for both drugs.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 A common mechanism that might be expected to show alteration upon drug selection is the up-
49 regulation of ABC transporters of the multidrug resistance family [20]. We saw increased expression
50 of ABCB1/P-glycoprotein in both cell lines compared to a DMSO (vehicle control) selected C6 cell line
51 (Figure 2A). Similarly, a functional assay, based upon extrusion of the fluorescent dye rhodamine 123
52 (R123) by ABCB1-expressing cells, demonstrated that the percentage of cells capable of exporting
53 R123 increased to over 75% in both selected cell lines (Figure 2B). For comparison, the DMSO-
54 treated C6 cell line showed R123 extrusion in 35% of cells, a value identical to that of the parental C6
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

cell line (Figure 2B). Thus, both cell lines displayed elevated ABCB1 expression and function following continuous exposure to topoisomerase inhibitors. In contrast, ABCG2 (breast cancer resistance protein) was not detected either before or after drug selection of the C6 cell line, and ABCC1 (multidrug resistance associated protein 1) was only expressed after exposure of cell lines to topoisomerase inhibitors for greater than 10 passages, culture conditions that were not typical in our experiments (data not shown).

We then examined whether the increase in ABCB1 expression was associated with increased resistance to the topoisomerase inhibitor used in their selection, or with acquired cross-resistance. In clonogenic assays, the C6-Et cell line showed an increase (2.5-fold; $p < 0.05$; Figure 3A) in the IC_{70} for etoposide but showed no cross-resistance to irinotecan, although both drugs are ABCB1 substrates [21]. The modest increase in resistance to etoposide upon continuous exposure without associated cross-resistance leads us to propose that the mechanism of elevated resistance is not solely via a classical multidrug export pump, and that other mechanism(s) of drug resistance, or protection against the damaging effects of etoposide is more active in the C6-Et cell line. Conversely, the C6-Ir cell line maintained the same IC_{70} for irinotecan compared to the C6-V cell line (Figure 3B), i.e. growing C6 cells in the presence of irinotecan did not result in the selection of cells with elevated irinotecan resistance, in agreement with studies indicating that ABCB1, although capable of transporting irinotecan [21], is not able to establish substantial concentration gradients of this drug [22].

Resistance to chemotherapy in cancer cell lines has also been associated with the emergence of a sub-population of cells that displays lower levels of reactive oxygen species, and concomitant elevation of antioxidant enzyme expression [23]. We investigated whether this was true for the drug exposed C6 cell lines by determining their ROS production rates following incubation with tertiary butyl hydroperoxide. ROS production rates were linear for at least 60 minutes following TBHP administration, and were consistent in all three cell lines at multiple concentrations of TBHP (Figure 4A; ANOVA $P > 0.05$), indicating that the cell lines do not differ in their ability to respond to ROS production. In confirmation of this, the expression level of ROS-detoxifying enzymes (e.g. catalase) was equal across the 3 cell lines (Figure 4B, C). This suggests that the prolonged exposure to irinotecan or etoposide does not result in the emergence of a cell population better able to respond to oxidative stress.

1
2 One of the hallmarks of glioma is local invasion [5, 7] , and it was therefore of interest to determine
3 cell migration following drug exposure in a glioma cell line model. Monolayers of C6-V, C6-Et and C6-
4 Ir cells were scratch-wounded with a sterile 0.2ml pipette tip and the rate of migration into the cell
5 free area monitored by time-lapse microscopy. Apparent from the still images taken from time-lapse
6 videos is the reduced wound closure in irinotecan treated cells (Figure 5A, compare for example the
7 16 and 24 hour time points). The size of the wound was measured at 30 minute intervals allowing for
8 a single exponential fit of the data (Figure 5B) to determine the half-time for wound closure. Two of
9 the three cell lines fitted a single exponential equation, whilst the fitting of the C6-Et cell line data
10 required the presence of a short lag phase before an exponential decay. Non-linear regression of the
11 data shows that the C6-Ir cell line closed the wound significantly slower ($t_{1/2} = 720 \pm 35$ mins) than
12 the other two cell lines (C6-Et $t_{1/2} 457 \pm 24$ mins, parental C6 cell line $t_{1/2} 294 \pm 24$ mins). A similar
13 reduction in migratory ability for the C6-Ir cell line was observed with transwell migration assays
14 (Figure 5C). A potential molecular basis of this reduced rate of migration was investigated by
15 determining the expression of collagens Col1A2 and Col3A1, both of which are implicated in tumour
16 invasion and/or epithelia to mesenchymal transition [24]. For Col3A1, the C6-Ir cell line showed a
17 significantly reduced gene expression compared to the C6-Et and C6-V cell line (Figure 5D), and for
18 Col1A2 C6-Ir also showed reduced gene expression compared to the C6-Et (Figure 5E).

32 33 34 35 36 **Discussion**

37
38
39 In this study we compared the effect of two established antineoplastic agents (the topoisomerase II
40 inhibitor etoposide and the topoisomerase I inhibitor irinotecan) on glioma cell line drug resistance
41 and invasion. Instead of monitoring glioma cell response to a single exposure, we focused instead on
42 cellular response after several rounds of treatment (mirroring the clinical regimen). What we
43 observed was that C6 glioma cells do not acquire additional resistance to irinotecan when
44 continuously treated with this type I topoisomerase I inhibitor. In contrast, C6 cells became more
45 resistant to etoposide after several rounds of treatment with this type II topoisomerase inhibitor.
46 Analysis of C6-Et cells revealed that they had increased expression of the multidrug transporter
47 ABCB1. An increase in ABCB1 expression was however observed in cells treated with both etoposide
48 and irinotecan, but only resistance to etoposide was increased, and this was in the absence of cross-
49 resistance to irinotecan. Hence, although ABCB1 is upregulated this does not appear to be a major
50 drug resistance mechanism in these cells. We also assayed the ability of our continuously exposed
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 cells to withstand reactive oxygen species, but saw no change in this ability. The resistance to
2 etoposide therefore appears to be highly specific and the majority of resistance is probably due to a
3 mutation within topoisomerase II itself preventing interaction with etoposide [9] or an increase in
4 the rate of DNA repair [10]. These data indicate that etoposide is a poor choice for cyclical therapy in
5 glioma as it leads to elevated etoposide resistance. Continuous exposure to irinotecan, on the other
6 hand, does not lead to increased resistance to itself or cross-resistance to etoposide.
7
8
9

10
11
12
13
14 Local invasion is a major problem in glioma and makes it difficult to completely surgically remove
15 these tumours. Glioma cells have been shown to be able to produce their own extracellular matrix
16 factors including collagens [25] and high concentrations of collagen in the glioma microenvironment
17 have been shown to correlate with more effective migration [24]. C6 glioma cells produce a diffusely
18 invading tumour in animal models and so may be more useful for modelling invasion *in vitro*
19 compared to U87 cells which produce more circumscribed tumours. We observed a reduced
20 migratory capacity of C6 cells treated with repeated cycles of irinotecan, and a reduction in
21 expression of collagen Col3A1 suggestive that impaired migration may be due to inhibition of cell
22 matrix interactions that are essential for tumour invasion. In contrast, etoposide treated cells which
23 show rapid wound closure also display elevated collagen expression, suggesting that treatment has
24 selected cells with an increased migratory capacity which would be an undesirable effect in therapy.
25 Although there are no clinical reports of increased metastatic potential of etoposide treated
26 tumours, there are growing reports that tumour drug resistance and invasion are interlinked and our
27 data further contributes to this emerging picture [26-29].
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 In conclusion, repeated rounds of treatment of glioma cells with the topoisomerase I inhibitor
44 irinotecan does not result in increased drug resistance or ROS resistance, rather it produces a cell
45 line with impaired migration, supporting further the investigation of irinotecan and derivatives in
46 treatment of glioma.
47
48
49
50
51
52

53 **Acknowledgements**

54
55
56 We deeply appreciate support from the Royal Thai government for a PhD scholarship for WP, and
57 King Abdulaziz University for a PhD scholarship for ABA. This work was supported by funding from
58
59
60
61
62
63
64
65

1 the Brain Tumour Charity (SDBTT 17/3). The authors are grateful to Dr Peter Jones (Life Sciences,
2 University of Nottingham) for the generous donation of transwell migration plates.
3
4
5

6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65

1. Dolecek TA, Propp JM, Stroup NE, Kruchko C (2012) CBTRUS statistical report: primary brain and central nervous system tumors diagnosed in the United States in 2005-2009. *Neuro-oncology* 14 Suppl 5: v1-49 doi:10.1093/neuonc/nos218
2. Stupp R, Mason WP, van den Bent MJ, Weller M, Fisher B, Taphoorn MJ, Belanger K, Brandes AA, Marosi C, Bogdahn U, Curschmann J, Janzer RC, Ludwin SK, Gorlia T, Allgeier A, Lacombe D, Cairncross JG, Eisenhauer E, Mirimanoff RO (2005) Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *The New England journal of medicine* 352: 987-996 doi:10.1056/NEJMoa043330
3. Gilbert MR, Wang M, Aldape KD, Stupp R, Hegi ME, Jaeckle KA, Armstrong TS, Wefel JS, Won M, Blumenthal DT, Mahajan A, Schultz CJ, Erridge S, Baumert B, Hopkins KI, Tzuk-Shina T, Brown PD, Chakravarti A, Curran WJ, Jr., Mehta MP (2013) Dose-dense temozolomide for newly diagnosed glioblastoma: a randomized phase III clinical trial. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 31: 4085-4091 doi:10.1200/JCO.2013.49.6968
4. Haar CP, Hebbar P, Wallace GC, Das A, Vandergrift WA, 3rd, Smith JA, Giglio P, Patel SJ, Ray SK, Banik NL (2012) Drug resistance in glioblastoma: a mini review. *Neurochemical research* 37: 1192-1200 doi:10.1007/s11064-011-0701-1
5. Cuddapah VA, Robel S, Watkins S, Sontheimer H (2014) A neurocentric perspective on glioma invasion. *Nature reviews Neuroscience* 15: 455-465 doi:10.1038/nrn3765
6. Paw I, Carpenter RC, Watabe K, Debinski W, Lo HW (2015) Mechanisms regulating glioma invasion. *Cancer letters* 362: 1-7 doi:10.1016/j.canlet.2015.03.015
7. Sanai N, Berger MS (2008) Glioma extent of resection and its impact on patient outcome. *Neurosurgery* 62: 753-764; discussion 264-756 doi:10.1227/01.neu.0000318159.21731.cf
8. Prabhu S, Harris F, Lea R, Snape TJ (2014) Small-molecule clinical trial candidates for the treatment of glioma. *Drug discovery today* 19: 1298-1308 doi:10.1016/j.drudis.2014.02.007
9. Chikamori K, Grabowski DR, Kinter M, Willard BB, Yadav S, Aebersold RH, Bukowski RM, Hickson ID, Andersen AH, Ganapathi R, Ganapathi MK (2003) Phosphorylation of serine 1106 in the catalytic domain of topoisomerase II alpha regulates enzymatic activity and drug sensitivity. *The Journal of biological chemistry* 278: 12696-12702 doi:10.1074/jbc.M300837200
10. Adachi N, Suzuki H, Iizumi S, Koyama H (2003) Hypersensitivity of nonhomologous DNA end-joining mutants to VP-16 and ICRF-193: implications for the repair of topoisomerase II-mediated DNA damage. *The Journal of biological chemistry* 278: 35897-35902 doi:10.1074/jbc.M306500200
11. Benda P, Lightbody J, Sato G, Levine L, Sweet W (1968) Differentiated rat glial cell strain in tissue culture. *Science* 161: 370-371
12. Schmidek HH, Nielsen SL, Schiller AL, Messer J (1971) Morphological studies of rat brain tumors induced by N-nitrosomethylurea. *Journal of neurosurgery* 34: 335-340 doi:10.3171/jns.1971.34.3.0335
13. Auer RN, Del Maestro RF, Anderson R (1981) A simple and reproducible experimental in vivo glioma model. *The Canadian journal of neurological sciences Le journal canadien des sciences neurologiques* 8: 325-331
14. Barth RF, Kaur B (2009) Rat brain tumor models in experimental neuro-oncology: the C6, 9L, T9, RG2, F98, BT4C, RT-2 and CNS-1 gliomas. *Journal of neuro-oncology* 94: 299-312 doi:10.1007/s11060-009-9875-7
15. Hande KR (1998) Etoposide: four decades of development of a topoisomerase II inhibitor. *Eur J Cancer* 34: 1514-1521

16. Pollack IF, Erff M, Bom D, Burke TG, Strode JT, Curran DP (1999) Potent topoisomerase I inhibition by novel silatecans eliminates glioma proliferation in vitro and in vivo. *Cancer research* 59: 4898-4905
17. Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F (2002) Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome biology* 3: RESEARCH0034
18. Pfaffl MW (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic acids research* 29: e45
19. Franken NA, Rodermond HM, Stap J, Haveman J, van Bree C (2006) Clonogenic assay of cells in vitro. *Nature protocols* 1: 2315-2319 doi:10.1038/nprot.2006.339
20. Kartner N, Riordan JR, Ling V (1983) Cell surface P-glycoprotein associated with multidrug resistance in mammalian cell lines. *Science* 221: 1285-1288
21. Szakacs G, Paterson JK, Ludwig JA, Booth-Genthe C, Gottesman MM (2006) Targeting multidrug resistance in cancer. *Nature reviews* 5: 219-234
22. Goldwirth L, Beccaria K, Carpentier A, Farinotti R, Fernandez C (2014) Irinotecan and temozolomide brain distribution: a focus on ABCB1. *Cancer chemotherapy and pharmacology* 74: 185-193 doi:10.1007/s00280-014-2490-0
23. Achuthan S, Santhoshkumar TR, Prabhakar J, Nair SA, Pillai MR (2011) Drug-induced senescence generates chemoresistant stemlike cells with low reactive oxygen species. *The Journal of biological chemistry* 286: 37813-37829 doi:10.1074/jbc.M110.200675
24. Kaufman LJ, Brangwynne CP, Kasza KE, Filippidi E, Gordon VD, Deisboeck TS, Weitz DA (2005) Glioma expansion in collagen I matrices: analyzing collagen concentration-dependent growth and motility patterns. *Biophysical journal* 89: 635-650 doi:10.1529/biophysj.105.061994
25. Paulus W, Huettner C, Tonn JC (1994) Collagens, integrins and the mesenchymal drift in glioblastomas: a comparison of biopsy specimens, spheroid and early monolayer cultures. *International journal of cancer Journal international du cancer* 58: 841-846
26. Coyle B, Kessler M, Sabnis DH, Kerr ID (2015) ABCB1 in children's brain tumours. *Biochemical Society transactions* 43: 1018-1022 doi:10.1042/BST20150137
27. Othman RT, Kimishi I, Bradshaw TD, Storer LC, Korshunov A, Pfister SM, Grundy RG, Kerr ID, Coyle B (2014) Overcoming multiple drug resistance mechanisms in medulloblastoma. *Acta neuropathologica communications* 2: 57 doi:10.1186/2051-5960-2-57
28. Colone M, Calcabrini A, Toccaceli L, Bozzuto G, Stringaro A, Gentile M, Cianfriglia M, Ciervo A, Caraglia M, Budillon A, Meo G, Arancia G, Molinari A (2008) The multidrug transporter P-glycoprotein: a mediator of melanoma invasion? *The Journal of investigative dermatology* 128: 957-971 doi:10.1038/sj.jid.5701082
29. Miletto-Gonzalez KE, Chen S, Muthukumaran N, Saglimbeni GN, Wu X, Yang J, Apolito K, Shih WJ, Hait WN, Rodriguez-Rodriguez L (2005) The CD44 receptor interacts with P-glycoprotein to promote cell migration and invasion in cancer. *Cancer research* 65: 6660-6667 doi:10.1158/0008-5472.CAN-04-3478

Figure Legends

Figure 1. The glioma cell line C6 is sensitive to DNA topoisomerase inhibitors. A, B clonogenic assays of C6 cells were performed in the presence of increasing concentrations of etoposide (**A**) or irinotecan (**B**) resulting in dose-dependent inhibition of colony growth. The mean IC₇₀ concentrations are 16.2 μ M for etoposide and 7.6 μ M for irinotecan. All data are derived from at least 3 independent experiments and error bars denote the standard error of the mean.

Figure 2. Expression and function of ABCB1 is increased in drug exposed C6 sub-lines. A. Microsomal membrane fractions (40 μ g) of the indicated C6 cell lines were resolved on 8% w/v polyacrylamide gels and, following electro-transfer, membranes were probed with anti-ABCB1 antibody C219 (1:1000). Blots were subsequently re-probed with anti-tubulin antibodies. Results are representative of at least 3 independent experiments. ABCB1 is observed as a two bands indicative of glycosylation. **B.** ABCB1 function was confirmed in a flow cytometry assay in which the percentage of cells exporting rhodamine-123 (R123) was determined. Results are the mean (s.e.m.) of at least 3 independent experiments. Data were compared using ANOVA and Newman-Kuels post-test. Cell lines showing increased % of rhodamine extruding cells compared to vector treated controls are designated by * ($p < 0.05$) and ** ($p < 0.01$).

Figure 3. Continuous exposure of C6 to etoposide, but not to irinotecan, results in selection of a more drug resistant population. The resistance of parental C6 and drug exposed sub-lines lines was assessed by clonogenic assays (**A, B**). The C6-Et line showed increased resistance to etoposide compared to the other cell lines (**A**, $p < 0.05$; ANOVA), whereas the C6-Ir cell line showed no increase in resistance (**B**). The C6-Et cell line did not exhibit cross-resistance to irinotecan.

Figure 4. Prolonged exposure to DNA topoisomerase inhibitors is not associated with increased resistance to ROS. A Cells were cultured in the absence or presence of a reactive oxygen species inducer (TBHP) and a fluorescent indicator of ROS production (DCF). A linear rate of ROS production was measured over 60 minutes following TBHP addition (relative fluorescence units/minute). Relative rates of ROS production compared to untreated cells were calculated over this linear period and displayed as a function of TBHP concentration. **B, C.** Expression of catalase was determined by western blotting using polyclonal anti catalase antibody at 1:500 dilution (Abcam) and relative


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

expression determined by comparison to actin band intensity (AC-74 monoclonal antibody at 1:5000 dilution) (C). All data represents the mean (s.e.m.) of at least 3 independent experiments.

Figure 5. Exposure to irinotecan results in reduced migration and expression of collagens. A Wound scratch healing assays were analysed by time-lapse microscopy, and representative images in the time-lapse are shown. **B** Wound closure is represented in the form of percentage initial scratch size as a function of time. Individual data points and errors are omitted for clarity (as each line results from >130 data points). Non-linear regression of the data shows that the C6-Ir cell line closed the wound significantly slower ($t_{1/2} = 720 \pm 35$ mins) than the C6-Et (457 ± 24 mins) and parental C6 cell line. **C**. Transwell migration assays demonstrate that C6-Ir displays a reduced migration in response to a chemoattractant compared to C6-Et. **D, E** qPCR of collagens 3A1 (**D**) and 1A2 (**E**) demonstrates reduced collagen expression in the C6-irinotecan cell line compared to the C6-etoposide cell line (p-values obtained from post-test comparisons following ANOVA, * $p < 0.05$, ** $P < 0.01$, *** $P < 0.001$). The relative expression of the collagens was determined compared to the geometric mean of two “housekeeping genes” GAPDH and B-actin. All data represents the mean (s.e.m.) of at least 3 independent experiments.

Figure 1

A



B





Figure 2

A



B




Figure 3






Figure 4

A



B



C









Figure 5



Supplementary Table 1: Primer pairs designed for qRT-PCR

Gene	Forward primer	Reverse primer	PCR product size
COL1A2	5'-GCCCACATGCCGTGACTT-3'	5'-GGCCTGGATGCAGGTTTCA-3'	142 bp
COL3A1	5'-TGGTGGACAGATGCTGGTGCTG-3'	5'-GGCCGGCTGGAAAGAAGTCTG-3'	150 bp
GAPDH	5'-CCCCTGGCCAAGGTCATCCA-3'	5'-CGCCATCACGCCACAGCTT-3'	126 bp
β-Actin	5'-ACCGAGGCCCTCTGAACCC-3'	5'-CCAGTGGTACGACCAGAGGCA-3'	134 bp