



The University of  
**Nottingham**

UNITED KINGDOM · CHINA · MALAYSIA

Bonardi, Charlotte and Pardon, Marie-Christine and Armstrong, Paul (2016) Deficits in object-in-place but not relative recency performance in the APPswe/PS1dE9 mouse model of Alzheimer's disease: implications for object recognition. *Behavioural Brain Research*, 313 . pp. 71-81. ISSN 1872-7549

**Access from the University of Nottingham repository:**

<http://eprints.nottingham.ac.uk/34700/1/Bonardi%20Pardon%20%26%20Armstrong%202016.pdf>

**Copyright and reuse:**

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the Creative Commons Attribution Non-commercial No Derivatives licence and may be reused according to the conditions of the licence. For more details see: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

**A note on versions:**

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact [eprints@nottingham.ac.uk](mailto:eprints@nottingham.ac.uk)

Deficits in object-in-place but not relative recency performance in the APPswe/PS1dE9 mouse model  
of Alzheimer's disease: Implications for object recognition.

Charlotte Bonardi\*<sup>1</sup>, Marie-Christine Pardon<sup>2</sup> & Paul Armstrong<sup>1</sup>

<sup>1</sup> School of Psychology, University of Nottingham, University Park, Nottingham, NG7 2RD, United  
Kingdom

<sup>2</sup> School of Biomedical Sciences, University of Nottingham, University Park, Nottingham, NG7 2RD,  
United Kingdom

Paul Armstrong; lpxpa1@nottingham.ac.uk

Marie-Christine Pardon; marie.pardon@nottingham.ac.uk; 0044 115 82 30149

Charlotte Bonardi; charlotte.bonardi@nottingham.ac.uk; 0044 115 8467927

Word Count (Main Text): 8,675

Number of Figures: 5

\*Corresponding author:

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Abstract

Performance was examined on three variants of the spontaneous object recognition (SOR) task, in 5-month old APP<sup>swe</sup>/PS1<sup>dE9</sup> mice and wild-type littermate controls. A deficit was observed in an *object-in-place* (OIP) task, in which mice are preexposed to four different objects in specific locations, and then at test two of the objects swap locations (Experiment 2). Typically more exploration is seen of the objects which have switched location, which is taken as evidence of a *retrieval-generated priming* mechanism. However, no significant transgenic deficit was found in a *relative recency* (RR) task (Experiment 1), in which mice are exposed to two different objects in two separate sample phases, and then tested with both objects. Typically more exploration of the first-presented object is observed, which is taken as evidence of a *self-generated priming* mechanism. Nor was there any impairment in the simplest variant, the *spontaneous object recognition* (SOR) task, in which mice are preexposed to one object and then tested with the familiar and a novel object. This was true regardless of whether the sample-test interval was 5 minutes (Experiment 1) or 24 hours (Experiments 1 and 2). It is argued that SOR performance [depends on retrieval-generated priming as well as self-generated priming, and our preliminary evidence suggests that the retrieval-generated priming process is especially impaired in these young transgenic animals.](#)

## 1. Introduction

1 Alzheimer's disease (AD) is the most common senile dementia, and its prevalence in western  
2 society is a major economic and social challenge. A problem in treatment of AD is the difficulty of  
3 diagnosis; in its early stages it is hard to distinguish from normal aging or Mild Cognitive Impairment  
4 (MCI), which does not always progress into AD. Thus current drug therapies are not optimally  
5 effective because they are administered only once clear clinical symptoms are manifest [1]. But the  
6 neuropathological changes underlying AD begin many years before symptoms emerge [2], meaning  
7 early intervention is possible. Increasing importance is thus being placed on gaining a better  
8 understanding of the biomarkers and cognitive changes that characterise preclinical AD [3], to  
9 facilitate early detection and give a better idea of how and when to administer treatment. This has  
10 been addressed in part through the use of genetically modified mice which over-express one or more  
11 of three genes implicated in familial AD, and display both its neuropathological symptoms as well as  
12 its characteristic cognitive degeneration. Although imperfect analogues of human AD, these models  
13 provide a valuable shortcut for identifying potential early cognitive symptoms, and are regarded by  
14 many as a fundamental tool in understanding AD [4].

15 The aim of our research has been to identify early cognitive signs of AD in one specific  
16 genetic model, the double-transgenic APP<sup>swe</sup>/PS1<sup>dE9</sup> mouse. This may be the best-characterised  
17 transgenic model of AD to date, co-expressing the mutated Swedish APP gene and also the exon-9  
18 deleted variant of the PS1 gene [5]. Elevated levels of oligomeric A $\beta$  in the cortex and hippocampus  
19 have been observed at 3.5 months of age in these mice; these changes are accompanied by synaptic  
20 deficits [6, 7], and are also closely associated with swollen dystrophic cholinergic neurites [8].  
21 Although the amyloid plaques characteristic of AD have been reported at 4 months of age in these  
22 mice, it is only from 6 months that they are consistently observed [9]. A $\beta$  deposition is paralleled by  
23 progressive degeneration of monoaminergic [10, 11] and striatal [12] neurons, and neuroinflammatory  
24 reactions [13] which mirror human AD pathology. These animals also recapitulate the age-related  
25 cognitive decline characteristic of AD [14], which is thought to depend on these neuropathological  
26 changes.

27 The neuropathology that develops in AD in general, and in this mouse model in particular, is  
28 well specified. But precisely *which* aspects of this brain pathology underlie the cognitive deficits that  
29 are such a central feature of AD is still under debate [15]. In this particular mouse line, some findings  
30

1 are consistent with the view that the degree of cognitive impairment is related to the level of plaque  
2 deposition [14]; but other work has cast doubt on this suggestion. First, cognitive deficits have been  
3 observed *before* plaque deposition in these animals [16, 17, 18]. This suggests that A  $\beta$  deposition is  
4 unrelated to cognitive decline, a conclusion supported by the fact that spatial learning deficits in older  
5 mice are correlated not with plaque load but with levels of soluble amyloid [19]. There have also been  
6 reports that manipulations that increase plaque levels *improve* performance on a spatial memory task  
7 [17]. These findings all point to the suggestion that the cognitive markers characterising the early  
8 stages of AD stem from the increased levels of soluble oligomeric A  $\beta$  which precede plaque  
9 formation rather than the plaques themselves. For example, high levels of A  $\beta$  produce local synaptic  
10 abnormalities and breakage of neuronal branches [20], and impair long-term potentiation [21, 22];  
11 soluble A  $\beta$  is also synaptotoxic, producing a reduction in synaptic density that occurs even when  
12 plaques are absent [23]. Evidence like this has led some to argue that these A  $\beta$  -induced changes in  
13 synaptic function underlie the cognitive deterioration [24]. This interpretation is supported by the fact  
14 that cognitive decline correlates with synaptic loss in human AD [25]. Our approach has thus been  
15 based on the assumption that elevated A  $\beta$  is likely to be responsible for the earliest impairments in  
16 cognition seen in AD. Thus we have focussed on examining cognitive ability at about 4-5 months of  
17 age in these mice - by which point levels of oligomeric A  $\beta$  are elevated, but substantial plaque  
18 deposition has not yet occurred.

19 We concentrated on one specific component of recognition memory, the perception of  
20 familiarity [26], as a potential early cognitive symptom. A subset of patients with MCI show selective  
21 impairments in visual recognition memory - a task that relies on familiarity judgements - and distinctive  
22 patterns of grey matter loss similar to those seen in AD [27]. Thus it has been suggested that visual  
23 recognition deficits might be a diagnostic marker of the early stages of AD [28, 29]. In non-human  
24 animals this type of memory is assessed in the spontaneous object recognition task (SOR), which  
25 exploits the observation that rodents will preferentially explore a novel object in preference to one that  
26 is familiar [30, 31]. Animals are exposed to a pair of identical, junk objects, and then after a retention  
27 interval returned to the apparatus, where one of the preexposed objects has been replaced with a  
28 novel item. Selective exploration of the novel object is taken as evidence that the preexposed object  
29 is recognised as familiar. This widely-used task has revealed deficits in a wide range of different

1 transgenic models of AD [32], and impairments are routinely observed in older APP<sup>swe</sup>/PS1<sup>dE9</sup> mice  
2 [33, 34, 35, 36, 37; but see 38]. SOR deficits are also occasionally reported in these mice at 6-7  
3 months of age [34, 39, 40, 41], but never in animals younger than 6 months [40, 42]. As elevated  
4 levels of A $\beta$  are present from around 3.5 months of age in this strain [6, 7], it is difficult to explain the  
5 SOR deficits in terms of this factor. However, a more detailed theoretical analysis suggests that this  
6 assessment might be misleading.  
7

8  
9  
10  
11  
12 Although a number of different theoretical accounts of SOR performance have been proposed  
13 [43 44], as a starting point we focus on one, which is based on *SOP* (Sometimes Opponent Process).  
14 This is an influential theory of associative learning [45] that has proved effective in predicting and  
15 explaining a wide variety of learning phenomena [46, 47, 48]. Because *SOP* is unique in explaining  
16 associative learning through a specific conceptualisation of memory, it has also been successfully  
17 applied to recognition memory [49, 50, 51, 52, 53]. *SOP* asserts that any stimulus may be regarded  
18 as a set of elements. These are normally inactive, but stimulus presentation probabilistically activates  
19 a subset of its elements into a state of primary activation termed *A1*. *A1* is of limited capacity, and  
20 elements in this state decay rapidly into a secondary, *A2* activation state, and thence more slowly to  
21 the inactive state. These activation states differ in critical ways. First, it is typically assumed that an  
22 element in *A1* elicits more vigorous responding than one in *A2*. Second, once an element has  
23 reached the *A2* state it must become inactive again before it can re-enter *A1* - no direct transition from  
24 *A2* to *A1* is possible. This creates a refractory period during which a second presentation of a stimulus  
25 will not create as strong a response as the first, because many of its elements are 'waiting' to decay  
26 into the inactive state - meaning fewer are available for recruitment into *A1*. This transient ability of  
27 stimulus presentation to reduce the impact of subsequent presentations is termed *self-generated*  
28 *priming*. Stimulus elements may also reach *A2* via *retrieval-generated priming*: if two stimuli co-occur  
29 in *A1* an association forms between them, such that presentation of one is able to activate the  
30 representation of the other - *and this activation puts its elements directly into A2*. Thus when the  
31 predicted stimulus actually occurs, fewer of its elements are available to enter *A1* and the response to  
32 the stimulus is reduced.  
33

34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55 According to this analysis both self- and retrieval-generated priming may contribute to  
56 performance on the SOR task [52]. Initial presentation of the preexposed object will prime its  
57 elements into *A1*, from where they decay into *A2*. If the test occurs before the preexposed object's  
58  
59  
60  
61  
62  
63  
64  
65

1 elements have returned to the inactive state, re-presentation of the object will produce less *A1* activity  
2 than a completely novel item, and result in less exploration of the familiar item through self-generated  
3 priming. **But in addition**, during initial preexposure associations may form between the preexposed  
4 item and the surrounding context. At test these contextual cues can prime some of the preexposed  
5 object's elements directly into *A2*, which *also* reduces exploration via retrieval-generated priming.  
6  
7

8  
9  
10 Experimental evidence has been provided in support of this proposal that retrieval-generated  
11 priming can contribute to SOR performance. In a series of studies with rats, Whitt et al. [54] exposed  
12 rats to two objects *P* and *Q*; *P* was presented in context *X* and *Q* in context *Y* (*X* and *Y* were either  
13 other objects, or patterned inserts placed round the perimeter of the experimental arena). Then the  
14 rats experienced a presentation of *X*, and were then immediately tested with *P* and *Q* in the absence  
15 of either *X* or *Y*. The rationale was that, during the sample phase,  $X \rightarrow P$  and  $Y \rightarrow Q$  associations would  
16 form, giving *X* the capacity to prime *P*, and *Y* to prime *Q*. The subsequent presentation of *X* would  
17 thus produce selective retrieval-generated priming of *P*, so that in the test that immediately followed,  
18 the elements of *P* would be placed in the *A2* state, resulting in selective exploration of *Q*. This is what  
19 was observed.  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29

30 The fact that, in terms of this analysis, SOR may be multiply determined raises the possibility  
31 that one of the mechanisms underlying it could be impaired, but SOR performance overall could  
32 appear unaffected if the other mechanism remains intact and can mediate performance to a sufficient  
33 level. The purpose of this work was thus to explore the possibility that one of these two mechanisms  
34 underlying SOR performance might be impaired in the younger animals, even if SOR performance is  
35 not. Thus we tested performance of 5-month-old APP<sup>swe</sup>/PS1<sup>dE9</sup> mice on tasks that independently  
36 assess the self- and retrieval-generated priming processes.  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

## 47 2. Experiment 1

48  
49  
50  
51 Experiment 1 employed a relative recency (RR) task, which establishes if animals can  
52 discriminate objects on the basis of how recently they have been encountered, and provides a  
53 relatively pure measure of self-generated priming (see Figure 1). The animal is first allowed to explore  
54 object, *B* and then a different object, *A*, in the same apparatus. After a delay, the animal receives a  
55 test with both objects [51, 55]. Animals typically show a preference for object *B*, the object  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 encountered earlier in the series. According to SOP this task is primarily a measure of self-generated  
2 priming: as both the more recent *A* and the less recent *B* are presented before test, their elements  
3 should have all been primed into *A1*, and started to decay into *A2*; however, because *B* is  
4 encountered first, by the test *B*'s elements will have had more of a chance to return to the inactive  
5 state, and so be ready to enter *A1* and elicit a strong response, than those of *A*. Critically, as both  
6 items have been encountered in the test apparatus, both have had the same opportunity to become  
7 associated with the surrounding context and hence suffer retrieval-generated priming. Thus  
8 differences in exploration of *A* and *B* are more obviously attributed to the self-generated priming  
9 process. We conducted this RR task in 5 month old APP<sup>swe</sup>/PS1<sup>dE9</sup> and their wild-type littermates,  
10 and then examined performance on the SOR task to confirm no deficit was evident, as the previous  
11 literature suggests. We first conducted the SOR task with a 5-min retention interval between the  
12 sample and test; as no deficit was found, we went on to employ a more difficult version of the task in  
13 which the retention interval was 24-hours in duration.

## 2.1. Materials and Methods

### 2.1.1. Animals

14 All experimental animals were bred in the University of Nottingham's transgenic animal facility  
15 from breeding stock purchased from the Jackson laboratory. Experiment 1 employed 15  
16 experimentally naïve male mice, 8 APP<sup>swe</sup>/PS1<sup>dE9</sup> transgenic mice and 7 wild-type littermates  
17 (Groups APP/PS1 and WT respectively). Testing began when they were approximately 20 weeks old  
18 and lasted about three weeks. All mice were housed in the same room, which was maintained on a  
19 12/12 hour light cycle, with lights on at 07:00 hours; the room temperature, relative humidity and air  
20 exchange were automatically controlled. Animals were group-housed with *ad libitum* access to food  
21 and water, and provided with nesting material and a play tube.

### 2.1.2. Apparatus and Stimuli

22 The experiments were conducted in a rectangular arena with walls and floors of white  
23 translucent plastic (length × width × height: 60 cm × 40 cm × 45 cm), located in a quiet, brightly lit  
24 room. A camera was suspended from a frame 90.0 cm above the centre of the arena, flanked by two



1 LED spotlights 22 cm apart, producing a floor-level illumination of 50 lux. The camera view (~ 45° arc)  
2 included the entire floor and the lower part of the four walls. The trajectory of the animals' heads was  
3 tracked by Any-maze software (Version 4.5; Stoelting, Wood Dale, Illinois). Four copies of each of ten  
4 assorted junk objects (*i*) - (*x*) served as stimuli (see supplementary materials); copies of a specific  
5 object were randomly selected from this 4-object pool for any sample or test phase requiring that  
6 object. The RR task employed *i* & *ii*, and *iiii* & *iv*, the SOR (5-min) task *v* & *vi*, and the SOR (24-hour)  
7 task *vii* & *viii*, and *ix* & *x*. A square zone of length 9.5 cm was defined around each object in the arena,  
8 allowing *exploration time* - the duration of time the mouse's head was within the active zone for a  
9 particular object - to be computed. The zone size was chosen to be large enough to include a  
10 perimeter of between about 2 to 3.5 cm around the various objects, and the objects themselves were  
11 constructed on the basis of pilot work establishing the mice did not show much tendency to climb or  
12 sit on them. Visual inspection confirmed that the time the mice spent with their head in the active zone  
13 normally reflected the mouse orienting toward the object, and so we adopted the automated measure  
14 of time in the active zone as a relatively uncontaminated measure of exploration behaviour.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

### 29 2.1.3. Procedure

30  
31 2.1.3.1. *Preexposure*. Before the start of training each mouse was habituated to the empty arena. In  
32 each of seven sessions the mouse was placed in the centre of the apparatus and allowed to explore  
33 for 5 min. The floor and walls of the apparatus were cleaned with diluted alcohol before each mouse  
34 was placed in the arena.  
35  
36  
37  
38  
39  
40

41 2.1.3.2. *General procedures*. Both tasks involved 5-minute sample phases - two in the RR task and  
42 one in the SOR task - and a 3-minute test phase (see Figure 1). In the sample phases mice were  
43 exposed to two copies of the same object, and in the test to two different objects, *A* and *B*; *A* was the  
44 most recently experienced object in the RR task and the preexposed object in the SOR task; *B* was  
45 the less recently experienced object in the RR task, and the novel object in the SOR task. At the start  
46 of each phase the mouse was placed in the arena centre facing the gap between the two objects; on  
47 its removal the objects, walls and floor were cleaned with diluted alcohol.  
48  
49  
50  
51  
52  
53  
54  
55

56 In order to avoid ceiling effects, in both Experiments 1 and 2 animals received two repetitions  
57 of each task unless discrimination between *B* and *A* was very strong in the first repetition  
58 (power > .85). Thus in the present study all mice received two repetitions of the RR task, followed by  
59  
60  
61  
62  
63  
64  
65

1 one repetition of the SOR 5-min task, and finally two of the more difficult SOR 24-hour task, giving a  
2 total of five repetitions. In each repetition objects could be placed in two of the four zones, which were  
3 situated in opposite corners of the arena. The first repetition employed the bottom left and top right  
4 corners as the two active zones, the second the top left and bottom right, and the position of the  
5 active zones continued to alternate in the remaining three task repetitions. Within each repetition  
6 stimulus identity and position were counterbalanced: thus, for example, in the first RR repetition  
7 (roughly) half the mice in each group had object *i* as *A* and *ii* as *B*, and the remainder the opposite;  
8 then all were tested with *i* at top left and *ii* at bottom right, so object identity and position were  
9 counterbalanced across *A/B* and genotype.  
10

11  
12 2.1.3.3. *RR*: Mice were exposed to two copies of *B* in the first sample phase, and 24 hours later to two  
13 copies of *A* in the second sample phase. The test with *A* and *B* occurred approximately five minutes  
14 after the second sample phase.  
15

16  
17 2.1.3.4. *SOR 5-min*: In the sample phase each mouse was exposed to two copies of *A*, and then after  
18 approximately five minutes was tested with *A* and *B*.  
19

20  
21 2.1.3.5. *SOR 24-hour*. This task was identical to the 5-minute version except that the test phase  
22 occurred approximately 24 hours after the sample phase.  
23

## 24 2.2. Results

### 25 2.2.1. Data Treatment

26 Exploration time was computed in 1-minute bins for each of the objects in each phase for each mouse.  
27 Data from the sample phases were summed across the entire phase and all objects. Data from the  
28 test phase were computed separately for *A* and *B* in three 1-minute bins<sup>1</sup>. Data were analysed using  
29 mixed ANOVAs, and significant two-way interactions explored with simple main effects analysis using  
30 the pooled error term.  $\eta_p^2$  was reported for significant effects and interactions.  
31

### 32 2.2.2. *RR* Results

---

33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58 <sup>1</sup> Raw exploration rates were used as the primary measure, rather than the more usual discrimination  
59 ratio, because they provide a more direct index of behaviour and do not mask potential differences in  
60 baseline exploration.  
61  
62  
63  
64  
65

1 Exploration was calculated separately for the first and second sample phases, to assess potential  
2 differences in exploration of *A* and *B*; for example, if *B* were explored *less* during preexposure it might  
3 be explored more at test simply because it was less familiar than *A*, rather than because it was less  
4 recent. Group mean exploration of *B* was, for Groups APP/PS1 and WT respectively, 25.98 and  
5 22.98s; the corresponding means for *A* were 19.95 and 20.10s. ANOVA with group (APP/PS1/WT),  
6 sample (B/A) and repetition as factors revealed significant interactions between repetition and both  
7 group,  $F(1, 13) = 4.92, p = .045, MSe = 115.05, \eta_p^2 = .27$ , and sample,  $F(1, 13) = 7.35, p = .018, MSe$   
8  $= 71.95, \eta_p^2 = .36$ . Exploration of the Repetition \* Group interaction revealed that Group APP/PS1  
9 explored more in the first repetition (25.98s) than in the second (20.03s),  $F(1, 13) = 4.80, p = .047,$   
10  $MSe = 115.05$ , whereas Group WT did not (with means of 20.13 s and 22.95 s respectively),  $F < 1$ .  
11 Exploration of the Repetition \* Sample interaction revealed more exploration of *B* (27.35s) than of *A*  
12 (18.69s) in repetition 1,  $F(1, 13) = 14.95, p < .001, MSe = 78.95$ , but not in repetition 2 (with means of  
13 21.61s and 21.37s respectively),  $F < 1$ . However, the higher exploration of *B* in repetition 1 would if  
14 anything reduce, not enhance, the size of any relative recency effect.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

29 The results of the test are presented in Figure 2 (upper panel) as *difference scores* -- time  
30 spent exploring *B* minus time spent exploring *A* -- in each minute of test (separate exploration times  
31 for *B* and *A* are presented in Table 1). In the first two minutes the less recent *B* was explored more  
32 than the more recent *A* in both groups, but by the third minute this effect had dissipated in the  
33 transgenic animals. However, ANOVA with group (APP/PS1/WT), object (*A/B*), repetition and minute  
34 as factors revealed only a significant effect of object,  $F(1, 13) = 7.26, p = .018, MSe = 24.39, \eta_p^2 = .36$ ;  
35 there was no effect of group,  $F(1, 13) = 2.77, p = .12, MSe = 24.56$ , and no interaction between these  
36 two factors,  $F(1, 13) = 1.01, p = .33, MSe = 24.39$ ; nothing else was significant, largest  $F(2, 26) = 2.12,$   
37  $p = .14, MSe = 15.92$  for the effect of minute. An additional analysis performed on the data for minute  
38 3 revealed only a main effect of group,  $F(1, 13) = 4.76, p = .048, MSe = 16.12, \eta_p^2 = .27$ ; neither the  
39 Group x Object interaction,  $F(1, 13) = 2.66, p = .127, MSe = 33.15$ , nor anything else was significant,  
40 largest,  $F(1, 13) = 2.67, p = .126, MSe = 22.62$ . Thus there was no statistical evidence for the  
41 apparent attenuation of performance in the transgenic animals in the last minute of the test.  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56

### 57 2.2.3. SOR 5-min Results

58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
The mean exploration time during the sample phase was 16.28 s in Group APP/PS1 and 24.61 s in Group WT - somewhat lower in the transgenic mice. ANOVA with group (APP/PS1/WT) as a factor showed this to be significant,  $F(1, 13) = 5.49$ ,  $p = .036$ ,  $MSe = 94.46$ ,  $\eta_p^2 = .30$ . The lower exploration in the transgenic mice would, if anything, increase the likelihood of seeing a deficit at test - yet this was not observed. The test difference scores are shown in Figure 2 (centre panel), and it is evident that both groups explored the novel *B* more than the familiar *A*; ANOVA with group (APP/PS1/WT), object (*A/B*) and minute as factors revealed a significant effect of object  $F(1, 13) = 11.86$ ,  $p = .004$ ,  $MSe = 16.35$ ,  $\eta_p^2 = .48$  which did not interact with group  $F < 1$ . There was also a significant interaction between object and minute,  $F(2, 26) = 6.55$ ,  $p = .005$ ,  $MSe = 17.52$ ,  $\eta_p^2 = .34$ , and the effect of object was significant in minutes 1 and 2,  $F(1, 39) = 19.78$ ,  $p < .001$ ,  $MSe = 17.13$ , and  $F(1, 39) = 4.17$ ,  $p = .048$ , but not in minute 3,  $F < 1$ . Nothing else was significant, largest  $F(1, 13) = 4.06$ ,  $p = .065$ ,  $MSe = 10.89$ . Thus the tendency to preferentially explore the novel *B* was present only in the first two minutes of the test.

#### 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 2.2.4. SOR 24-hour Results

30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
The mean exploration time during the sample phase was 15.00 s in Group APP/PS1 and 26.29 s in Group WT - again lower in the transgenic mice - and ANOVA with group (APP/PS1/WT) and repetition (1/2) as factors revealed that this was significant,  $F(1, 13) = 11.49$ ,  $p = .005$ ,  $MSe = 165.71$ ,  $\eta_p^2 = .47$ ; there was no effect or interaction involving repetition, largest  $F(1, 13) = 1.16$ ,  $p = .30$ ,  $MSe = 72.87$ . The test difference scores are shown in Figure 2 (lower panel). Although the preference for the novel *B* was modest it was consistent across the test, and again there was no sign of a deficit in the transgenic mice, despite their lower levels of exploration in the sample phase. ANOVA with group APP/PS1, object (*A/B*), repetition (1/2) and minute as factors revealed a significant effect of object  $F(1, 13) = 9.31$ ,  $p = .009$ ,  $MSe = 10.15$ ,  $\eta_p^2 = .42$  which did not interact with group  $F < 1$ . There was also a significant effect of group,  $F(1, 13) = 6.77$ ,  $p = .022$ ,  $MSe = 21.81$ ,  $\eta_p^2 = .34$ , again revealing lower levels of exploration in the APP/PS1 animals; nothing else was significant, largest  $F(1, 13) = 3.21$ ,  $p = .096$ ,  $MSe = 4.65$ .

#### 56 57 58 59 60 61 62 63 64 65 2.3 Discussion

1 The results of this first experiment replicated the results of previous studies, showing no  
2 deficit in an SOR task in 5-month-old APP<sup>swe</sup>/PS1<sup>dE9</sup> mice. This was true regardless of whether the  
3 retention interval was 5 minutes or 24 hours, despite the fact that these tasks varied substantially in  
4 difficulty. The novel result was that there was no significant transgenic impairment in the RR task,  
5 which we have argued is a relatively pure measure of self-generated priming. If self-generated  
6 priming is the primary contributor to performance on the SOR task, then the suggestion is that this  
7 process is intact in the transgenic animals.  
8  
9  
10  
11  
12  
13

14 However, the argument that RR depends solely on self-generated priming assumes that the  
15 degree to which the context is associated with *A* and *B* is equated, because they are both presented  
16 in the context for the same amount of time during the sample phases. However, in the second sample  
17 phase in which *A* is preexposed, the context is presented without *B*, which could result in some  
18 extinction of the context  $\rightarrow B$  association. This could reduce the degree to which the context primes *B*  
19 on test, increasing the tendency of the animals to explore this object: thus a component of RR  
20 performance could be explained in terms of retrieval-generated priming. Moreover, one might expect  
21 that the contribution of retrieval-generated priming would become increasingly evident with time, as  
22 the longer the test continues, the more elements from *A* will have decayed from A2 back to inactive,  
23 making the difference in the self-generated priming of *A* and *B* increasingly small. In this respect it is  
24 intriguing that the apparent deficit in transgenic performance was evident only at the end of the test -  
25 something which might be taken to indicate a deficit in retrieval-generated priming in these animals.  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Experiment 2 directly examined this possibility.

### 3. Experiment 2

Experiment 2 employed an object-in-place (OIP) task [56], which has been argued to provide  
a pure measure of the retrieval-generated priming process [52]. In the sample phase four different  
items were presented in an array; in the subsequent test, two of the items remained in their original  
positions while the remaining two items exchanged locations. Exploration of the items that have  
changed location is typically greater than exploration of those that have not [51, 56]. According to

1 SOP, this is due to retrieval-generated priming: during preexposure, associations may form between  
2 the local features of the arena and the objects that are placed there, so at test those items presented  
3 in the preexposure location will be primed by the contextual cues that still surround them; in contrast  
4 the moved items, being tested in a location with which they have not been associated, will not be  
5 primed in this way. Thus greater exploration of the moved items at test is attributed to the fact that  
6 they suffer less retrieval-generated priming than their static counterparts. Critically all test items are  
7 equally familiar, and have been experienced equally recently - so any differential exploration cannot  
8 be attributed to differences in self-generated priming. In addition, all items are presented at test in  
9 equally familiar locations. After the animals had been tested on this task they were again tested on  
10 the SOR task, but in this study only the more difficult, 24-hour version was employed.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

### 24 3.1. *Materials and Methods*

25 All aspects of the method that are not specified were identical to that of the previous experiment.  
26  
27

#### 28 3.1.1. *Animals*

29 Experiment 2 employed 15 experimentally naïve male mice, 7 APP<sup>swe</sup>/PS1<sup>dE9</sup> transgenic mice and  
30 8 wild-type (WT) littermates. Testing began when they were approximately 21 weeks old and lasted  
31 about two weeks. All mice were housed and maintained exactly as in the previous experiment.  
32  
33  
34  
35  
36  
37  
38  
39

#### 40 3.1.2. *Apparatus and Stimuli*

41 The same apparatus was used as in the last experiment, but with the addition of wall inserts  
42 to increase the distinctiveness of the local features of the context. These inserts were made from  
43 medium density fibreboard lined with linoleum, and each one covered the whole of one of the shorter  
44 walls, and half of both longer walls, of the arena; thus two inserts covered the entire arena wall. These  
45 were 45.0 cm high, and when inside the arena reduced the floor space to 42.0 cm x 32.0 cm. Two  
46 different patterns were used, one on each side of the arena: *Mb*, a mosaic of 2.3 cm<sup>2</sup> blue squares  
47 whose edges were 45° from horizontal, and *Dw*, a mosaic of white 272-cm<sup>2</sup> squares whose edges  
48 were 90° from horizontal, with a black, 16-cm<sup>2</sup> square superimposed at each point where four white  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 squares met (see Figure 2). The same objects were used as in the previous experiment: the OIP task  
2 employed objects *i*, *ii*, *iii* & *iv*, and *vii*, *viii*, *ix* & *x*, and the SOR task objects *v* and *vi*.  
3

### 4 5 3.1.3. Procedure

6  
7 3.1.3.1. *Preexposure*. Identical to that of the previous experiment except that the preexposure  
8 sessions were of 10-min duration, and in the seventh, final session the context inserts were present.  
9

10  
11 3.1.3.2. *OIP*. During the sample phase each mouse was exposed to four different objects, *A*, *A'*, *B* and  
12 *B'*, one in each zones. As this task involved four objects rather than two, the duration of the sample  
13 phase was doubled to 10 minutes. The test phase was identical to the sample phase except that two  
14 of the objects, *B* and *B'*, were transposed (see Table 1); the duration of the test was 4 minutes.  
15  
16

17  
18 Animals received two repetitions of this task. Stimulus identity and position were counterbalanced  
19 across object and genotype (see Table 1). Thus, for example, for 4 transgenic and 4 wild-type mice *B*  
20 and *B'* were objects *i* and *ii* in the top left and bottom right corners during the sample phase, and for  
21 the remaining mice *B* and *B'* were objects *iii* and *iv* in the top right and bottom left corners. The  
22 context inserts were present throughout the sample and test phases.  
23  
24  
25  
26  
27  
28  
29  
30

31 3.1.3.3. *SOR 24-hour*. All mice then received one repetition of the SOR 24-hr task, conducted exactly  
32 as in the previous experiment.  
33  
34  
35

### 36 3.2. Results

37  
38 3.2.1. *OIP Results* The mean exploration time in Group APP/PS1 was 107.16 s, and for Group WT  
39 100.52 s. ANOVA with group and repetition as factors revealed no effect of group,  $F(1, 13) = 1.31$ ,  $p$   
40 = .27,  $MSe = 50.27$ , but a significant effect of repetition,  $F(1, 13) = 87.38$ ,  $p < .001$ ,  $MSe = 282.34$ ,  $\eta_p^2$   
41 = .87, as exploration levels were considerably higher in repetition 1 (123.86s) than in repetition 2  
42 (83.37s); there was no Repetition x Group interaction,  $F < 1$ . The results from the test are shown in  
43 Figure 3 (upper panel). In the wild type mice there was numerically more exploration of the displaced  
44 objects *B* and *B'* than of *A* and *A'* throughout the test, whereas for the transgenic mice the results  
45 were far more variable, with less exploration of the displaced objects in minutes 2 and 4. ANOVA with  
46 group (APP/PS1/WT), object (*A,A'/B,B'*), repetition and minute as factors revealed significant effects  
47 of object,  $F(1, 13) = 6.99$ ,  $p = .02$ ,  $MSe = 30.34$ ,  $\eta_p^2 = .35$ , and also of repetition,  $F(1, 13) = 21.68$ ,  $p$   
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

< .001,  $MSe = 37.87$ ,  $\eta_p^2 = .63$ , and minute,  $F(1, 13) = 6.44$ ,  $p = .001$ ,  $MSe = 10.10$ ,  $\eta_p^2 = .33$ ;  
exploration rates continued to be higher in repetition 1 (11.06s) than in repetition 2 (7.57s). Critically  
there was a significant Group x Object x Minute interaction,  $F(3, 18) = 3.25$ ,  $p = .032$ ,  $MSe = 19.22$ ,  
 $\eta_p^2 = .20$ , suggesting that the groups might differ in their ability to perform on the task over the course  
of the test. To explore this interaction further, two-way ANOVAs were performed on the data from  
each group, with object and minute as factors. In Group APP/PS1 this revealed nothing significant,  
largest  $F(3, 18) = 2.68$ ,  $p = .078$ ,  $MSe = 13.94$ ,  $\eta_p^2 = .31$  for the effect of minute; there was no sign of  
an effect of object,  $F < 1$ . A parallel analysis conducted on the data from Group WT revealed a highly  
significant effect of object,  $F(1, 13) = 49.18$ ,  $p < .001$ ,  $MSe = 5.86$ ,  $\eta_p^2 = .88$ , and also of minute,  $F(3,$   
 $18) = 5.89$ ,  $p = .004$ ,  $MSe = 6.80$ ,  $\eta_p^2 = .46$ ; the interaction was not significant,  $F(3, 18) = 1.34$ ,  $p = .29$ ,  
 $MSe = 15.88$ . As an alternative means of exploring the interaction we conducted an additional  
ANOVA on the difference scores, with group and minute as factors. This revealed a significant  
interaction between minute and group,  $F(3, 39) = 3.25$ ,  $p = .032$ ,  $MSe = 30.34$ ,  $\eta_p^2 = .2$ ; the main  
effects of group and object were not significant,  $F(1, 39) = 2.45$ ,  $p = .140$ ,  $MSe = 30.34$  and  $F < 1$   
respectively. Simple main effects analysis performed on the interaction revealed that Group WT  
showed superior performance on minutes 2 and 4,  $F(1, 52) = 5.97$ ,  $p = .018$ ,  $MSe = 22.00$  and  $F(1, 52)$   
 $= 5.30$ ,  $p = .025$ ,  $MSe = 22.00$ , but not on minutes 1 and 3,  $F_s < 1$ .

3.2.2. *SOR 24-hour Results* The mean exploration time during the preexposure phase was 22.31 s for  
Group APP/PS1 and 19.28 s for Group WT, and these scores did not differ,  $F < 1$ . The results of the  
test are shown in Figure 3 (lower panel); exploration of the novel object was higher in both groups  
throughout the test, and although there was a tendency for accuracy to decline in the transgenic mice  
over the course of the test, this proved not to be significant. ANOVA with group (APP/PS1/WT), object  
(B/A) and minute as factors revealed a significant main effect of object,  $F(1, 13) = 26.76$ ,  $p < .001$ ,  
 $MSe = 6.91$ ,  $\eta_p^2 = .67$ , but there were no significant effects or interactions involving group, largest  $F(1,$   
 $13) = 1.10$ ,  $p = .34$ ,  $MSe = 6.20$ , and nothing else was significant, largest  $F(1, 13) = 1.65$ ,  $p = .21$ ,  
 $MSe = 9.84$ .

### 3.3. $\beta$ -Amyloid Pathology



1 To confirm the presence of  $\beta$ -Amyloid pathology in male APPswe/PS1dE9 mice of this age,  
2 we examined the brains from a different cohort of 4.5-month old male APPswe/PS1dE9. Brains were  
3 post fixed in 4% paraformaldehyde for 6h and kept in 70% ethanol overnight before being embedded  
4 in paraffin wax on a tissue embedding station (Leica TP1020). Immunostaining was carried out using  
5 standard procedures at room temperature on 7 $\mu$ m-thick coronal sections. Briefly, all the solutions  
6 were freshly prepared using PBS + 1% Tween 80, except DAB solution that was prepared in distilled  
7 water. The tissue was re-hydrated in consecutive rinses in xylene, 100% ethanol, 70% ethanol and  
8 distilled water. Antigen retrieval was performed by incubation in 10mM EDTA pH 6.0 for 20 min at  
9 95°C, followed by incubation in formic acid for 1 min. Tissue was then blocked in 5% normal horse  
10 serum, incubated in mouse monoclonal anti-  $\beta$  -amyloid antibody (1:2000, A5213 Sigma Aldrich, St.  
11 Louis, MO, USA) for 1h followed by 1 h incubation with anti-mouse secondary antibody (1:200; Vector  
12 Laboratories Inc. Burlingame, CA). After washing, sections were incubated with Vectastain Elite ABC  
13 kit (Vector Laboratories Inc. Burlingame, CA) and labelled with DAB peroxidase substrate (Vector  
14 Laboratories, Burlingame, CA) according to manufacturer's instructions. To reveal histological  
15 morphology, sections were then counterstained with haematoxylin (purplish-blue nuclear stain) and  
16 eosin (pink cytoplasmic stain) and mounted with DPX-mount media. Digital focused photo-scanning  
17 images were acquired using a Hamamatsu NanoZoomer-XR with TDI camera technology. [Figure 4](#)  
18 shows illustrative examples of amyloid- $\beta$  42 staining generated from brain tissue from 4.5 month old  
19 male mice APPswe/PS1dE9, and it is evident that although amyloid pathology had begun to emerge,  
20 it was slight and largely confined to cortex and hippocampus.  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44

### 45 *3.4 Discussion*

46  
47 In this experiment a deficit in performance on the OIP task was observed in the transgenic  
48 mice: the wild type animals showed a consistent preference for exploring the displaced objects  
49 throughout testing whereas the transgenic animals did not, performing significantly worse than their  
50 wild type counterparts on minutes 2 and 4 of the 4-minute test. Once again these same transgenic  
51 animals showed normal performance in the SOR task. In combination with the results of Experiment 1,  
52 the results are interpretable in terms of SOP if it is assumed that the transgenic animals have a deficit  
53 in retrieval-generated, but not self-generated, priming.  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

But before this conclusion is accepted, alternative interpretations of these results should be considered. First, this task differed from the SOR and RR tasks in that it required the animals to form an association between the contextual inserts and the objects. If the transgenic animals had some difficulty processing distal visual cues which, for example, rendered them unable to see the contexts clearly while they were exploring the objects, this could provide an alternative explanation of these results. However, we are not aware of any evidence suggesting that visual deficits are responsible for impaired performance in these animals. For example, Jardanhazi-Kurutz et al. [34] demonstrated that 4.5-month old APP<sup>swe</sup>/PS1<sup>dE9</sup> mice were impaired in a spatial learning task in the Morris water maze - yet performed as well as wild types locating the platform when it was visible. But this raises a second possibility - that the present results could be interpreted as the spatial learning deficit that we know can be present in transgenic mice of this age. However, the extent to which this should be viewed as an alternative interpretation depends on the specific interpretation of spatial learning that is adopted. Some have argued that performance on spatial tasks can be explained in terms of associative learning processes just like those underlying retrieval-generated priming [57]. If this is the case, then impairments in both spatial learning and the OIP task studied here may be interpreted as a failure of retrieval-generated priming.

33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

On a more procedural note, we attempted to make the three tasks comparable: in all versions the test locations were equally familiar, and the identity and location of novel and familiar items counterbalanced. But there were other differences between them; for example, the SOR task involved preexposure of one item, the RR task of two items, and the OIP task of four; an alternative interpretation is thus that a deficit emerges with an increase in the total number of items to be explored. While this is logically possible, it would imply that some evidence of a deficit should have been observed in the RR task -- but although there was a numerical tendency for transgenic mice to perform worse at the end of the RR test, this was not significant. Moreover, a similar nonsignificant tendency for a decline in performance in the transgenic mice was evident in the SOR task in Experiment 2. In addition, solution of the OIP task does not require the mice to encode all four objects - the reason we employed four rather than two was to ensure that all test locations were equated in familiarity. Finally, although the OIP task requires more objects to be explored in the same sample phase, we doubled the length of preexposure in this experiment to accommodate this. These

1 arguments notwithstanding, further experimental work would be necessary to definitely rule out such  
2 alternative interpretations.  
3  
4  
5  
6

#### 7 **4. General Discussion**

8  
9

10 Two experiments examined the performance of 5-month old APP<sup>swe</sup>/PS1<sup>dE9</sup> mice on three  
11 variants of the SOR task. Although SOR performance is typically unaffected in this model at this age,  
12 it was argued that according to one specific model of recognition memory, SOP [45], performance on  
13 SOR depends on two independent cognitive mechanisms: self-generated and retrieval-generated  
14 priming [52]. Thus even if one of these mechanisms were impaired, SOR performance could appear  
15 unaffected provided the other remained intact enough to support accurate performance. This  
16 appeared to be the case. In Experiment 2 we examined performance on the OIP task, which we argue  
17 provides a pure measure of retrieval-generated priming. The transgenic mice performed significantly  
18 worse than the wild types on two of the four minutes of the test, and overall - in contrast to the wild  
19 type animals - showed no significant preference for the displaced objects. In contrast, in Experiment 1  
20 mice of the same age were not significantly impaired on a relative recency task, supposedly a  
21 measure of self-generated priming [52]. In both experiments the transgenic mice performed normally  
22 on the SOR task, mirroring previous work using animals of this age [40, 42]. These results, although  
23 preliminary, are consistent with the suggestion that the mice suffer a selective deficit in retrieval- but  
24 not self-generated priming at 5 months of age, and that it is only at 6 months of age that the self-  
25 generated priming condition is also affected, and hence a net impairment in SOR observed. In fact the  
26 suggestion that the self- and retrieval generated priming mechanisms may be dissociated is not  
27 without precedent. Recent work on the GluA1 knockout mouse - in which the GluA1 subunit of the  
28 AMPA receptor, an important mediator of synaptic plasticity in the hippocampus, is deleted - has  
29 revealed a pattern complementary to that reported here, a deficit in self- but not retrieval-generated  
30 priming. Specifically, these mice showed an impairment on SOR and RR tasks, but performed  
31 normally on an SOR task variant that relied on retrieval-generated priming [58].  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55

56 This argument implies that young transgenic animals perform accurately on the SOR task  
57 using *only* self-generated priming: because elements of the stimulus presented during the sample  
58 phase have not all decayed from the A2 state by test, they will not all be available to enter A1 and  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

elicit a strong exploratory response. But while assuming decay from A2 is incomplete after 5 minutes is plausible, it is less so when the sample-test interval is 24 hours. With such a long delay it seems natural to attribute SOR performance to retrieval-generated priming, as this depends on the formation of associations which do not decay appreciably with the passage of time. So if retrieval-generated priming were impaired in the transgenic mice, how could they perform accurately on the 24-hour version of the task? One possibility is that there are two types of association that can support this retrieval-generated priming mechanism, only one of which is affected in the transgenic animals. We appealed solely to formation of an association between the preexposed object and the surrounding context, which is precisely what was tapped by the OIP task; but it might also be the case that associations form *among the elements of the object itself* [59]. If elements of each object were inter-associated in this manner, when a mouse begins to explore a familiar object the associations among its elements could result in retrieval-generated priming of the rest of the object. This would ensure that *all* the object's elements can be primed into A2 reducing exploration. This process could also mediate accurate SOR performance (although it would have no effect in the OIP task as all items are equally familiar, and so the possibility of such intra-object associations forming is equated). If the transgenic mice were less able to form associations between the object and the surrounding context than among the object's elements themselves, this could explain why the transgenic animals showed a deficit in OIP but not the 24-hour SOR task. For example, different elements of the same object are more likely to be experienced in close temporal proximity than are the object and the contextual cues that surround it - and imposing a delay between events makes them more difficult to associate [60]. Interestingly, deficits in taste aversion learning - in which ingestion of a flavoured liquid is followed some time later with illness - have been reported in 2-5 month old APP<sup>swe</sup>/PS1<sup>dE9</sup> mice [18, 61]. Thus maybe a difficulty in associating temporally distant events is an early cognitive manifestation of AD in this mouse model. Alternatively, contextual cues may have a spatial component - so that associating the object with the context involves associating elements in different dimensions (visual and spatial), in a way that associating elements of the same object may not. If transgenic mice were poorer at forming associations between visual and spatial cues, this could explain the results. Indeed, something very similar was reported by Swainson et al. [62], in a longitudinal study in which a battery of cognitive tests was performed on participants with either mild AD, questionable dementia or depression, as well as healthy controls. The test that best discriminated those with AD from other

1 participants was a '*visuo-spatial paired associative learning task*' in which participants had to encode  
2 where in a display a particular shape had been presented. Poor performance in this task was also  
3 evident in a subgroup of those with questionable dementia, and correlated with their subsequent  
4 cognitive deterioration - suggesting the task might be a useful diagnostic indicator of incipient AD.  
5  
6  
7

8  
9 These results may also be accommodated by other accounts of recognition memory, which  
10 attempt to make sense of dissociations in recognition performance evident in lesion studies. For  
11 example, Brown and Aggleton [63] proposed that recognition tasks can be divided into tasks such as  
12 SOR and RR, which require that only one item be remembered at a time and depend on perirhinal  
13 cortex, and those like OIP which rely on memory with a spatial or associative component, are  
14 hippocampal-dependent, and often involve the rearrangement or re-pairing of familiar items. A related  
15 approach has been taken by Saksida, Bussey and colleagues [64], who argued that tasks like SOR  
16 and RR are in part supported by memory for stimulus conjunctions (as opposed to individual stimulus  
17 features), and mediated by the perirhinal cortex; in contrast tasks like OIP, which require  
18 representation of the object and the temporal or spatial aspects of the context, are more dependent  
19 on the hippocampus. These dissociations imply that the deficits observed in our mice are likely to  
20 stem from a selective disturbance in hippocampal function. In fact there is good evidence that  
21 neuropathological changes begin emerge in both cortex and hippocampus in the APP<sup>swe</sup>/PS1<sup>dE9</sup>  
22 mouse at the age at which this study was conducted [9, 17]. This is an observation we confirmed in  
23 our own animals, showing that at 4.5 months of age the handful of plaques that are evident are  
24 confined to cortex and hippocampus. However we are not aware of any evidence that hippocampus is  
25 affected *before* the perirhinal cortex, as such models might predict on the basis of the results we have  
26 presented.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44

45  
46 [If our results may be taken to demonstrate a selective deficit in the OIP task in transgenic](#)  
47 [animals in the pre-plaque stage of the disease, this would suggest](#) that performance on this task  
48 might be a diagnostic marker for the early stages of AD. There are other findings from the human  
49 literature that are also consistent with this suggestion. Several authors have reported selective deficits  
50 in AD patients on tasks which require the association of different aspects of a stimulus - termed  
51 *memory binding* [62, 65, 66, 67], and one particular study by Parra et al. [67] described results that  
52 closely mirrored our findings. Participants were presented with two visual stimuli with various colours  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 (C) and shapes (S). In different variants of the task these two stimuli differed either in colour (C<sub>1</sub>S<sub>1</sub> &  
2 C<sub>2</sub>S<sub>1</sub>) or shape (C<sub>1</sub>S<sub>1</sub> and C<sub>1</sub>S<sub>2</sub>) or both (C<sub>1</sub>S<sub>1</sub> & C<sub>2</sub>S<sub>2</sub>) - the *binding* condition. In the former two  
3 conditions the participants with tested with the original items, but with the originally discriminating  
4 feature replaced by a novel one - i.e. C<sub>1</sub>S<sub>1</sub> & **C<sub>3</sub>S<sub>1</sub>** or C<sub>1</sub>S<sub>1</sub> & C<sub>1</sub>**S<sub>3</sub>**, meaning they had to discriminate  
5 items with familiar and novel features - a parallel of the SOR task. In contrast, in the binding condition  
6 they were tested either with the sample items, or with items in which the two features of the sample  
7 items were rearranged (C<sub>1</sub>S<sub>2</sub> & C<sub>2</sub>S<sub>1</sub>), meaning they had to discriminate between equally familiar  
8 features in either familiar or unfamiliar combinations - a parallel of the OIP task. They found a  
9 selective deficit in the binding task in patients with a form of familial AD linked to a specific mutation,  
10 and also in their asymptomatic relatives who were carriers of the mutation, relative to healthy non-  
11 carrier controls. Reports of this type encourage the suggestion that our results are not confined to the  
12 mouse model in which they were found, but may also have parallels in experiments using human  
13 participants, lending them translational value.

14 We are not aware of any other studies conducted on this particular transgenic model of AD  
15 which examine performance on recognition tasks more complex than the standard SOR task.  
16 However a recent series of studies by Davis and colleagues [68, 69] evaluated performance of a  
17 triple-transgenic model of AD, which has tau pathology as well as APP and PS1 mutations, on several  
18 variants of the SOR task. They found that younger mice performed normally on SOR, RR and OIP  
19 tasks, but showed a relatively selective impairment in a '*what-where-which*' task, in which the mice  
20 were preexposed to the same pair of objects in the transposed positions in two distinctive contexts  
21 (Figure 5), before being tested with two replicas of one of the objects in one of the contexts. Correct  
22 performance required the mouse to selectively explore the object that had not been experienced in  
23 that position in that context, and the authors interpreted this in terms of an episodic-like memory  
24 impairment. One interpretation is that these results represent a conflict of data, in that the young  
25 double-transgenic mice were impaired on the simpler task whereas the triple-transgenic mice were  
26 not. However, even within the class of APP<sup>swe</sup>/PS1 transgenic models there can be substantial  
27 variation in the pathology expressed (for example, in the amount of A $\beta$  deposited, and the structure  
28 and appearance of the plaques [70]. In addition one marked difference was that their version of the  
29 OIP task was simpler than the one we employed, involving preexposure of two different objects and  
30 then testing two identical copies of the same object (Figure 5). Any such procedural differences could

1  
2 have been responsible for the difference in results, and further work would be needed to resolve  
3 these issues.  
4

5 We interpreted our findings within the framework of a general model of learning and memory,  
6 SOP [45] which, although primarily used in the context of animal behaviour, can be applied more  
7 generally. In this sense work of this type may provide a novel perspective on cognitive phenomena  
8 from that offered by theories developed solely to explain human cognition. As we have seen, SOP  
9 offers a comprehensive account of the processes underlying recognition memory, interpreting them in  
10 terms of more fundamental principles of associative learning. It also interprets the SOR, RR and OIP  
11 tasks we employed within this same framework. As these tasks may be regarded as respectively  
12 tapping the *what*, *when* and *where* components that characterise episodic memory [71], SOP may  
13 also offer a new perspective on this important memory phenomenon. Interpreting both animal and  
14 human findings in terms of this associative learning model could therefore yield new insights into the  
15 early cognitive deficits of AD, as well as underpinning the translational work that is inevitable for  
16 successful drug development.  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Disclosure

The authors disclose that they have no actual or potential conflicts of interest, financial or otherwise, related to the present work. All animal procedures were carried out in accordance with the UK Animals Scientific Procedures Act and approved by the Home Office under Project Licence 40/3444.



## Author Note

1  
2  
3 Correspondence may be addressed to: C. Bonardi: School of Psychology, University of  
4 Nottingham, Nottingham, NG7 2RD, England, UK.  
5  
6

7  
8 Work reported here was in partial fulfilment of Paul Armstrong's PhD, which was funded by the  
9 School of Psychology, University of Nottingham. We would like to thank Alessandra Agostino for  
10 supplying the amyloid deposition images.  
11  
12  
13

14  
15 Procedures were authorized under UK law, carried put in accordance with the Animals (Scientific  
16 Procedures) Act (1986), and also the EU Directive 2010/63/EU for animal experiments. This work  
17 was funded by the School of Psychology, University of Nottingham, who otherwise had no  
18 involvement in this research.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## References

- 1  
2  
3  
4  
5 [1] M.S., Fiandaca, M.E. Mapstone,, A.K. Cheema & H. Federoff. The critical need for defining  
6 preclinical biomarkers in Alzheimer's disease. *Alzheimer's & Dementia*, 10 (2014) S196-S212.  
7  
8  
9  
10 [2] H.J., Aizenstein, R.D., Nebes, J.A., Saxton, J.C. Price, C.A. Mathis, N.D. Tsopelas et al. Frequent  
11 amyloid deposition without significant cognitive impairment among the elderly. *Archives of Neurology*,  
12 65 (2008) 1509-1517.  
13  
14  
15  
16 [3] R.A. Sperling, P.A. Aisen, L.A. Beckett, D.A. Bennett, S. Craft, A.M. Fagan et al. Toward defining  
17 th preclinical stages of Alzheimer's disease: Recommendations from the National Institute of Aging-  
18 Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's &*  
19 *Dementia*, 7 (2011) 280-292.  
20  
21  
22  
23  
24  
25 [4] C. Balducci, & G. Forloni. AP transgenic mice: Their use and limitations. *Neuromolecular Medicine*,  
26 13 (2011) 117-137.  
27  
28  
29  
30 [5] M.K. Lee, D.R. Borchelt, G. Kim, G. Thinakaran, H.H. Slunt, T. Ratovitski et al. Hyper-  
31 accumulation of FAD-linked presenilin 1 variants in vivo. *Nature Medicine*, 3 (1997) 756–60.  
32  
33  
34  
35 [6] Y.S. Hu, P. Xu, G. Pigino,,S.T. Brady, J. Larson, O. Lazarov. Complex environment experience  
36 rescues impaired neurogenesis, enhances synaptic plasticity, and attenuates neuropathology in  
37 familial Alzheimer's disease-linked APP- swe/PS1DeltaE9 mice. *The Faseb J.: Official Publication of*  
38 *the Federation of American Societies For Experimental Biology*, 24 (2010) 1667–81.  
39  
40  
41  
42  
43  
44 [7] I. Shemer, C. Holmgren, R. Min, L. Fulop, M. Zilberter, K.M. Sousa, et.al. Non-fibrillar beta-amyloid  
45 abates spike-timing-dependent synaptic potentiation at excitatory synapses in layer 2/3 of the  
46 neocortex by targeting postsynaptic AMPA receptors. *The European J. of Neuroscience*, 23 (2006)  
47 2035–47.  
48  
49  
50  
51  
52  
53 [8] S.E. Perez, S. Dar, M.D. Ikonovic, S.T. DeKosky, & E.J. Mufson. Cholinergic forebrain  
54 degeneration in the APPswe/PS1ΔE9 transgenic mouse. *Neurobiology of Disease*, 28 (2007) 3-15.  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- [9] M. Garcia-Alloza, E.M.Robbins, S.X. Zhang-Nune, S.M. Purcell, R.A. Betensky, S. Raju, et al. Characterization of amyloid deposition in the APPswe/PS1DE9 mouse model of Alzheimer's disease. *Neurobiology of Disease*, 24 (2006) 516–24.
- [10] M.E. Szapacs, A.L. Numis, & A.M. Andrews Late onset loss of hippocampal 5-HY and NE is accompanied by increases in BDNF protein expression in mice co-expressing mutant APP and PS1. *Neurobiology of Disease*, 16 (2004) 572-580.
- [11] L. Liu, T. Heikkinen, T. Tapiola, T. van Groen & H. Tanila. The effects of long-term treatment with metrifonate, a cholinesterase inhibitor, on cholinergic activity, amyloid pathology, and cognitive function in APP and PS1 doubly transgenic mice. *Experimental Neurology*, 173 (2002) 196-204.
- [12] M. Richner, G. Bach, & M.J. West. Over expression of amyloid beta-protein reduces the number of neurons in the striatum of the APPswe/PS1DE9. *Brain Research*, 1266 (2009) 87-92.
- [13] L. Holcomb, M.N. Gordon, E. McGowan, X. Yu, S. Benkovic, P. Jantzen, et al. Accelerated Alzheimer-type phenotype in transgenic mice carrying both mutant amyloid precursor protein and presenilin 1 transgenes. *Nature Medicine*, 4 (1998) 97–100.
- [14] G.W. Arendash, D.L. King, M.N. Gordon, D. Morgan, J.M. Hatcher, C.E. Hope. et al. Progressive, age-related behavioral impairments in transgenic mice carrying both mutant amyloid precursor protein and presenilin-1 transgenes. *Brain Research*, 891 (2001) 42–53.
- [15] W.L.Klein, G.A. Krafft, C.E. Finch. Targeting small Abeta oligomers: the solution to an Alzheimer's disease conundrum? *Trends in Neurosciences*, 24 (2001) 219–24.
- [16] L. Holcomb, M.N. Gordon, P. Jantzen, K. Hsiao, K. Duff & D. Morgan. Behavioural changes in transgenic mice expressing both amyloid precursor protein and presenelin-1 mutations: lack of association with amyloid deposits. *Behavior Genetics*, 29 (1999) 177-185.
- [17] J.L. Jankowsky, T. Melnikova, D.J. Fadale, G.M. Xu, H.H. Slunt, V. Gonzales et al. Environmental enrichment mitigates cognitive deficits in a mouse model of Alzheimer's disease. *J. of Neuroscience*, 25 (2005) 5217–24.

- 1  
2 [18] P.J. Pistell, M., Zhu & D.K. Ingram. Acquisition of conditioned taste aversion is impaired in the  
3 amyloid precursor protein/presenilin 1 mouse model of Alzheimer's disease. *Neuroscience*, 152 (2008)  
4 594–600.  
5  
6 [19] W. Zhang, J. Hao, R. Liu, Z. Zhang, G. Lei, C. Su et al. Soluble Ab levels correlate with cognitive  
7 deficits in the 12-month-old APPswe/PS1dE9 mouse model of Alzheimer's disease. *Behavioural Brain*  
8 *Research*, 222 (2011) 342-350.  
9  
10 [20] C-C. Wu, F. Chawla, D. Games, R.E. Ryden, S. Freedman et al. Selective vulnerability of dentate  
11 granule cells prior to amyloid deposition in PDAPP mice: Digital morphometric analyses. *Proceedings*  
12 *of the National Academy of Sciences*, 101 (2004) 7141-7146.  
13  
14 [21] D.B. Freir, C. Holscher, & C.E. Herron. Blockade of long-term potentiation by b-amyloid peptides  
15 in the CA1 region of the rat hippocampus In vivo. *J. of Physiology*, 85 (2001) 708-713.  
16  
17 [22] A. Itoh, T. Akaike, M. Sokabe, A. Nitta, R. Iida et al. Impairments of long-term potentiation in  
18 hippocampal slices of  $\beta$ -amyloid-infused rats. *European J. of Pharmacology*, 382 (1999) 167-175.  
19  
20 [23] L. Mucke, E. Masliah, G-Q. Yu, M. Mallory, E.M. Rockenstein et al. High-level neuronal  
21 expression of  $A\beta_{1-42}$  in wild-type human amyloid protein precursor transgenic mice: Synaptotoxicity  
22 without plaque formation. *J. of Neuroscience*, 20 (2000) 4050-4058.  
23  
24 [24] B. Gong, O.V. Vitolo, F. Trinchese, S. Liu, M. Shelanski, & O. Arancio. Persistent improvement in  
25 synaptic and cognitive functions in an Alzheimer mouse model after rolipram treatment. *The J. of*  
26 *Clinical Investigation*, 114 (2004) 1624-1634.  
27  
28 [25] Terry, R.D., Masliah, E., Salmon, D.P., Butters, N., DeTeresa, R., Hill, R., Hansen, L.A., &  
29 Katzman, R. Physical basis of cognitive alterations in Alzheimer's disease: Synapse loss is the major  
30 correlate of cognitive impairment. *Annals of Neurology*, 30 (1991) 572-580.  
31  
32 [26] G. Mandler. Recognizing: The judgement of previous occurrence. *Psychological Review*, 87  
33 (1980) 252-271.  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- [27] E.J. Barbeau, J.P. Ranjeva, M. Didic, S. Confort-Gouny, O. Felician, E. Soulier, P.J. Cozzone, M. Ceccaldi & M. Poncet. Profile of memory impairment and gray matter loss in amnesic mild cognitive impairment. *Neuropsychologia*, 46 (2008) 1009-1019.
- [28] M. Didic, J.P. Ranjeva, E. Barbeau, S. Confort-Gouny, Y. Le Fur, O. Felician et al. Impaired visual recognition memory in amnesic mild cognitive impairment is associated with mesiotemporal metabolic changes on magnetic resonance spectroscopic imaging. *J. of Alzheimer's Disease*, 22 (2010) 1269-1279.
- [29] S.M. Zola, C.M. Manzanares, P. Clopton, J.J. Lah, & A.I. Levey. A behavioral task predicts conversion to mild cognitive impairment and Alzheimer's disease. *American J. of Alzheimer's Disease and Other Dementias*, 28 (2012) 179-184.
- [30] A. Ennaceur, J. Delacour. (1988). A new one-trial test for neurobiological studies of memory in rats. I. Behavioural data. *Behavioural Brain Research*, 31 47–59.
- [31] P.N. Kivy, R.W. Earl, E.L. Walker. Stimulus context and satiation. *J. of Comparative and Physiological Psychology*, 49 (1956) 90–92.
- [32] B. Grayson, M. Leger, C. Piercy, L. Adamson, M. Harte, & J.C. Neill. Assessment of disease-related cognitive impairments using the novel object recognition (NOR) task in rodents. *Behavioural Brain Research*, 285 (2015) 176-193.
- [33] J.J. Donkin, S. Stukas, V. Hirsch-Reinshagen, D. Namjoshi, A. Wilkinson, S. May et al. ATP-binding cassette transporter A1 mediates the beneficial effects of the liver X receptor agonist GW3965 on object recognition memory and amyloid burden in amyloid precursor protein/presenilin 1 mice. *The J. of Biological Chemistry*, 285 (2011) 34144–34154.
- [34] D. Jardanhazi-Kurutz, M.P. Kummer, D. Terwel, K. Vogel, T. Dyrks, A. Thiele et al. Induced LC degeneration in APP/PS1 transgenic mice accelerates early cerebral amyloidosis and cognitive deficits. *Neurochemistry International*, 57 (2010) 375–82
- [35] W. Li, Y. Liu, X. Huang, N. Abumaria, Y. Zhu, X. Huang. et al. Elevation of brain magnesium prevents synaptic loss and reverses cognitive deficits in Alzheimer's disease mouse model. *Molecular Brain*, 7 (2014) 65-85.

1 [36] J. Yan, J-S. Jung, T-K. Kim, A. Hasan, C-W. Hong, J-S., Nam, & D-K. Song. Protective effects of  
2 ferulic acid in amyloid precursor protein plus presenelin-1 transgenic mouse model of Alzheimer  
3 disease. *Biology Pharmacology Bulletin*, 36 (2013) 140-143.  
4

5  
6 [37] Y. Yoshiike, T. Kimura, S. Yamashita, H. Furudate, T. Mizoroki, M. Murayama. et al. GABA(A)  
7 receptor-mediated acceleration of aging-associated memory decline in APP/PS1 mice and its  
8 pharmacological treatment by picrotoxin. *PLoS One*, 3 (2008) e3029.  
9  
10

11  
12 [38] C.A. Frye, & A.A. Walfe. Effects of progesterone administration and APPswe+PSEN1De9  
13 mutation for cognitive performance of mid-aged mice. *Neurobiology of Learning and Memory*, 89  
14 (2008) 17-26.  
15  
16

17  
18 [39] D. Cheng, J.K. Low, W. Logge, B. Garner, & T. Karl. Chronic cannabidiol treatment improves  
19 social and object recognition in double transgenic APPswe/PS1ΔE9 mice. *Psychopharmacology*, 231  
20 (2014) 3009-3017.  
21  
22

23  
24 [40] I. Pedros, D. Petrov, M. Allgaier, F. Sureda, E. Barroso, C. Beas-Zarate et al. Early alterations in  
25 energy metabolism in the hippocampus of APPswe/PS1dE9 mouse model of Alzheimer's disease.  
26 *Biochimica et Biophysica Acta*, 1842 (2014) 1556-1566.  
27  
28

29  
30 [41] E. Barbero-Camps, A. Fernández, L. Martínez, J.C. Fernández-Checa, A. & Colell. APP/PS1  
31 mice overexpressing SREBP-2 exhibit combined Ab accumulation and tau pathology underlying  
32 Alzheimer's disease. *Human Molecular Genetics* (2013) 1-17.  
33  
34

35  
36 [42] C. Bonardi, F. de Pulford, D. Jennings, & M-C. Pardon. A detailed analysis of the early context  
37 extinction deficits seen in APPswe/PS1dE9 female mice and their relevance to pre-clinical  
38 Alzheimer's disease. *Behavioural Brain Research*, 222 (2011) 89-97.  
39  
40

41  
42 [43] J.P. Aggleton, M.W. Brown. Interleaving brain systems for episodic and recognition memory.  
43 *Trends in Cognitive Science*, 10 (2006) 455–63.  
44  
45

46  
47 [44] R.A. Cowell, T.J. Bussey, & L.M. Saksida. Why does brain damage impair memory? A  
48 connectionist model of object recognition memory in perirhinal cortex. *J. of Neuroscience*, 26 (2006)  
49 12186-12197.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 [45] A.R. Wagner. SOP: A model of automatic memory processing in animal behavior. In: N.E. Spear  
2 & R.R. Miller (Eds.). *Information processing in animals: Memory Mechanisms*. Hillsdale, New Jersey:  
3 Erlbaum. (1981) pp. 5–47.  
4

5  
6 [46] G. Hall, & R.C. Honey. Contextual effects in conditioning, latent inhibition, and habituation:  
7 associative and retrieval functions of contextual cues. *J. of Experimental Psychology: Animal*  
8 *Behavior Processes*, 15 (1989) 232–41.  
9

10  
11 [47] R.C. Honey, G. Hall, & C. Bonardi. Negative priming in associative learning: Evidence from serial  
12 conditioning procedures. *J. of Experimental Psychology: Animal Behavior Processes*, 19 (1993) 90-  
13 97.  
14

15  
16 [48] E.H. Vogel, S.E. Brandon, & A.R. Wagner. Stimulus representation in SOP: II. An application to  
17 inhibition of delay. *Behavioural Processes*, 62 (2003) 27-48.  
18

19  
20 [49] R.C. Honey, M. Good. Associative components of recognition memory. *Current Opinion in*  
21 *Neurobiology*, 10 (2000) 200–204.  
22

23  
24 [50] D.J. Sanderson, D.M. Bannerman. Competitive short-term and long-term memory processes in  
25 spatial habituation. *J. of Experimental Psychology: Animal Behavior Processes*, 37 (2011) 189–99,  
26

27  
28 [51] S.K.E. Tam, J. Robinson, D. J. Jennings, & C. Bonardi. Dissociations in the effect of delay on  
29 object recognition and the effect of dorsal hippocampal damage: Evidence for an associative model of  
30 recognition memory *J. of Experimental Psychology: Animal Behavior Processes*, 40 (2014) 106-115.  
31

32  
33 [52] J. Robinson, & C. Bonardi. An Associative Analysis of Object Memory. *Behavioural Brain*  
34 *Research*, 285 (2015) 1-9.  
35

36  
37 [53] E. Whitt, J. Robinson. Improved spontaneous object recognition following spaced preexposure  
38 trials: evidence for an associative account of recognition memory. *J. of Experimental Psychology:*  
39 *Animal Behavior Processes*, 39 (2013) 174–179.  
40

41  
42 [54] E. Whitt, M. Haselgrove, J. Robinson. (2012). Indirect object recognition: Evidence for  
43 associative processes in recognition memory. *J. of Experimental Psychology: Animal Behavior*  
44 *Processes*, 38 (2012)., 74–83.  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2 [55] J.B. Mitchell, J. Laiacona. The medial frontal cortex and temporal memory: tests using  
3 spontaneous exploratory behaviour in the rat. *Behav Brain Research*, 97 (1998) 107–13.  
4  
5 [56] S. Dix, J.A. Aggleton. Extending the spontaneous preference test of recognition: evidence of  
6 object-location and object-context recognition. *Behav Brain Research*, 99 (1999) 191–200.  
7  
8  
9 [57] J.M. Pearce. An associative analysis of spatial learning. *Quarterly J. of Experimental Psychology*,  
10 62(2009), 1665-1684.  
11  
12  
13 [58] D.J. Sanderson, E. Hindley, E. Smeaton, N. Denny, A. Taylor et al. Deletion of the GluA1 AMPA  
14 receptor subunit impairs recency-dependent object recognition memory. *Learning & Memory*, 18  
15 (2011), 181-190.  
16  
17  
18 [59] I.P.L. McLaren, & N.J. Mackintosh. An elemental model of associative learning: I. Latent  
19 inhibition and perceptual learning. *Animal Learning & Behavior*, 28 (2000) 211-246.  
20  
21  
22 [60] W.J. Mahoney, & J.J.B Ayres. One-trial simultaneous and backward fear conditioning as reflected  
23 in conditioned suppression of licking in rats. *Animal Learning and Behavior*, 4 (1976) 357-362.  
24  
25  
26 [61] L. Ramírez-Lugo, M.S. Jensen, A. Soderman, & M.J. West. Deficits in Aaversive but not in safe  
27 taste memory in the APP<sup>sw</sup>/PS1<sup>dE9</sup> mice. *J. of Alzheimer's Disease*, 18 (2009) 281-293.  
28  
29  
30 [62] R. Swainson, J.R. Hodges, C.J. Galton, J. Semple, A. Michael, B.D. Dunn et al. Early detection  
31 and differential diagnosis of Alzheimer's disease and depression with neuropsychological tasks.  
32 *Dementia and Geriatric Cognitive Disorders*, 12 (2001) 265-280.  
33  
34  
35 [63] M.W. Brown, & J.P. Aggleton. Recognition memory: What are the roles of the perirhinal cortex  
36 and hippocampus? *Nature Reviews Neuroscience*, 2 (2001) 51-61.  
37  
38  
39 [64] R.A. Cowell, T.J. Bussey, & L.M. Saksida. Components of recognition memory: Dissociable  
40 cognitive processes or just differences in representational capacity? *Hippocampus*, 20 (2010) 1245-  
41 1262.  
42  
43  
44 [65] E. Granholm, & N. Butters. Associative encoding and retrieval in Alzheimer's and Huntington's  
45 disease. *Brain & Cognition*, 7 (1998) 335-347.  
46  
47  
48 [66] M.A. Parra, S. Abrahams, K. Fabi, R.H. Logie, S. Luzzi & S. Della Sala. Short-term memory  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



binding deficits in Alzheimer's disease. *Brain*, 132 (2009) 1057–1066.

[67] M.A. Parra, S. Abrahams, R.H. Logie, L.G. Méndez, F. Lopera, & S. Della Sala. Visual short-term memory binding deficits in familial Alzheimer's disease. *Brain*, 133 (2010) 2702-2713.

[68] K.E. Davis, A. Easton, M.J. Eacott, & J. Gigg. Episodic-like memory for what-where-which occasion is selectively impaired in the 3xTgAD mouse model of Alzheimer's disease. *J. of Alzheimer's Disease*, 33 (2013) 681-698.

[69] K.E. Davis, M.J. Eacott, A. Easton, & J. Gigg. Episodic-like memory is sensitive to both Alzheimer's-like pathological accumulation and normal aging processes in mice. *Behavioural Brain Research*, 254 (2013) 73-82.

[70] A. Snellman, F.R. López-Picón, J. Rokka, M. Salmona, G. Forloni et al. Longitudinal amyloid imaging in mouse brain with <sup>11</sup>C-PIB: Comparison of APP23, Tg2576, APP<sub>swe</sub>-PS1<sub>de9</sub> mouse models of Alzheimer's disease. *J. of Nuclear Medicine*, 54 (2013) 1434-1441.

[71] E. Tulving. *Elements of Episodic Memory*. Oxford: Clarendon Press (1983).

## Figure Legends

Figure 1: Schematic of the spontaneous object recognition (SOR), Relative Recency (RR) and Object-in-Place (OIP) tasks. A, B, A' and B' refer to various junk objects. For further details see text.

Figure 2: Experiment 1: Group mean difference scores (exploration of B minus exploration of A) in each minute of test in the Relative Recency (upper panel), Spontaneous Object Recognition 5-min (centre panel) and Spontaneous Object Recognition 24-hr (lower panel) tasks. Error bars show standard error of the mean.

Figure 3: Experiment 2: Group mean difference scores (exploration of B/B' minus exploration of A/A') in each minute of test in the Object-in-Place task (upper panel), and corresponding scores (exploration of B minus exploration of A) for the Spontaneous Object Recognition 24-hr task (lower panel). Error bars show standard error of the mean.

Figure 4: Schematic of the What-Where-Which and Object-in-Place tasks employed by Davis et al. (2013). A, B, A' and B' refer to various junk objects. For further details see text.

Figure 5: Some representative samples of b-amyloid 42 staining generated from brain tissue of 4.5-month old male APP<sup>swe</sup>/PS1<sup>dE9</sup> mice.

Figure 1: Schematic of the spontaneous object recognition (SOR), Relative Recency (RR) and Object-in-Place (OIP) tasks. A, B, A' and B' refer to various junk objects. For further details see text.




Figure 2: Experiment 1: Group mean difference scores (exploration of B minus exploration of A) in each minute of test in the Relative Recency (upper panel), Spontaneous Object Recognition 5-min (centre panel) and Spontaneous Object Recognition 24-hr (lower panel) tasks. Error bars show standard error of the mean.







Figure 3: Experiment 2: Group mean difference scores (exploration of B/B' minus exploration of A/A') in each minute of test in the Object-in-Place task (upper panel), and corresponding scores (exploration of B minus exploration of A) for the Spontaneous Object Recognition 24-hr task (lower panel). Error bars show standard error of the mean.



*Figure 4: Some representative samples of b-amyloid 42 staining generated from brain tissue of 4.5-month old male APPswe/PS1dE9 mice.*






Figure 5: Schematic of the What-Where-Which and Object-in-Place tasks employed by Davis et al. (2013). A, B, A' and B' refer to various junk objects. For further details see text.

:




Table 1: Group mean time (sec) exploring the novel (B) and familiar (A) objects in each replication of the Relative Recency (RR), Spontaneous Object Recognition (SOR) 5-min and 24-hour tasks in each minute of the test of Experiment 1.

Minute		1		2		3	
RR		B	A	B	A	B	A
Rep 1	APP/PS1	5.9	3.6	7.0	5.6	6.1	7.0
	WT	5.3	3.8	6.5	5.5	10.1	3.4
Rep 2	APP/PS1	5.5	3.7	6.6	3.2	3.4	3.8
	WT	6.0	5.0	9.6	5.2	8.7	7.0
SOR 5-min							
	APP/PS1	8.3	2.5	6.4	3.5	4.4	5.6
	WT	11.5	3.8	8.1	4.9	4.9	5.8
SOR 24-hr							
	APP/PS1	2.7	2.5	5.1	3.2	3.7	4.2
	WT	6.2	4.6	6.2	5.7	6.0	4.4
	APP/PS1	2.6	2.0	5.5	2.0	4.8	1.3
	WT	6.3	3.2	4.7	3.5	5.4	4.9



Table 2: Group mean time (sec) exploring the displaced (B) and static (A), or novel (B) and familiar (A), objects in each replication of the Object-in-Place (OIP) and Spontaneous Object Recognition (SOR) 24-hour tasks in each minute of the test of Experiment 2.

Minute		1		2		3		4	
OIP		B	A	B	A	B	A	B	A
Rep 1	APP/PS1	14.2	9.6	12.1	14.1	13.6	8.3	11.2	13.0
	WT	12.8	12.2	12.0	7.2	12.4	9.8	13.1	9.7
Rep 2	APP/PS1	12.1	8.7	8.1	8.3	5.5	6.7	6.3	8.4
	WT	11.7	7.6	10.5	5.4	6.1	6.4	9.1	5.2
SOR 24-hr									
	APP/PS1	7.3	2.9	5.5	3.0	4.0	2.9		
	WT	5.6	2.7	7.1	4.1	5.3	1.9		

**Supplementary Material**

[Click here to download Supplementary Material: Supplementary materials.docx](#)