

Behnke, Jerzy M. and Barnard, Christopher J. and Hurst, J.L. and McGregor, Peter K. and Gilbert, Francis and Lewis, John W. (1993) The prevalence and intensity of infection with helminth parasites in Mus spretus from the Setubal Peninsula of Portugal. Journal of Helminthology, 67 . pp. 115-122. ISSN 0022-149X

Access from the University of Nottingham repository:

http://eprints.nottingham.ac.uk/1183/1/Behnke_et_al_1993._JHelm_67%2C_115._Helms_in _Mus_spretus_in_Portugal.pdf

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see: http://eprints.nottingham.ac.uk/end user agreement.pdf

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

The prevalence and intensity of infection with helminth parasites in *Mus spretus* from the Setubal Peninsula of Portugal

J. M. BEHNKE, C. BARNARD, J. L. HURST, P. K. McGREGOR, F. GILBERT and J. W. LEWIS2

¹Department of Life Science, University of Nottingham, University Park, Nottingham, NG7 2RD, UK; ²Department of Biology, Royal Holloway, University of London, Egham, Surrey, TW20 9TY, UK

ABSTRACT

The results of a 5 year study of helminth parasites of *Mus spretus*, are reported. Six nematode and 5 cestode species were identified but no helminth showed 100% prevalence in *M. spretus*, the most commonly encountered nematode and cestode species being *Syphacia obvelata* (46-6%) and *Taenia taeniaeformis* (22-4%). Among the more unusual helminth species identified was *Eucoleus bacillatus*, a capillariid nematode inhabiting the stomach musculature. This species was identified in 3 of the 5 years of the study. The results are discussed in the broader context of previous studies and the epidemiology of rodent helminth infections in general.

KEY WORDS: Mus spretus, Portugal, Nematoda, Cestoda, prevalence

INTRODUCTION

Although intensive studies of parasitic infections in rodents have been systematically carried out in the UK over many years (SHARPE, 1964; LEWIS, 1968a,b; LEWIS & TWIGG, 1972; LANGLEY & FAIRLEY, 1982; MONTGOMERY & MONTGOMERY, 1988) and some parts of Europe, notably in Poland (FURMAGA, 1957; SOLTYS, 1957; KISIELEWSKA, 1970), Czechoslovakia (BAER & TENORA, 1970) and Hungary (TENORA & MURAI, 1972), reports on rodent parasites from Mediterranean countries usually represent no more than species lists. We are not aware of any detailed quantitative data on parasitic infections in Mus species from Mediterranean countries which consider such effects as host age, sex and seasonal fluctuations in parasite numbers and which contribute to a picture of the ecological relationships between the hosts and parasites concerned. Furthermore, almost nothing is known about the parasites of the little studied rodent Mus spretus, an aboriginal murine species found locally in western Mediterranean regions from France, through Spain, Portugal to Libya. In Portugal M. spretus occupies a niche in shrub-grassland habitats and seldom comes into contact with man in human habitations (SAGE, 1981). This species is thought to be closely related to M. musculus domesticus although it is now considered to merit separate specific status rather than the subspecific status reported earlier (SAGE, 1981).

In a series of two-week investigations carried out annually over 5 years, we accumulated quantitative information on parasitic helminth infections in *M. spretus*. This report describes our study sites and presents data on the identity, prevalence and intensity of infection with the major helminth taxa and species affecting *M. spretus*.

MATERIALS AND METHODS

Study site

The project was carried out during a two-week period in mid April from 1988 until 1992 and formed part of a terrestrial zoology field course. Most mice were caught at the Quinta de Sao Pedro in Sobreda near Lisboa but the Quinta Niza, about 3 km further south east also provided some animals for the study.

The Quinta de Sao Pedro (longitude 9°11′ west, latitude 38°39′ north) comprises 4 ha of land, mainly wooded with overgrown fields last used for arable purposes in 1965 and only occasionally grazed by sheep since then. It is located in the vicinity of Lisboa on the Setubal peninsula, 5 km from the resort of Costa da Caparica on the Atlantic coast. The entire plot is now walled off, although the boundary walls were not completed until 1990 and there are many potential entrance and exit points for small mammals. However, the site has become increasingly isolated from other reservoirs of small mammals in recent years. In 1988, dwellings surrounded about a third of the perimeter of the plot, the remainder being open grazing land, and a 50 m stretch adjoined a local road. Since 1991, however, there has been an extensive building programme in the region and in 1992, almost the entire perimeter, except for the road, had housing in various stages of completion.

Capture and treatment of mice

All mice were caught in standard Longworth traps baited with grain and peanut butter using conventional procedures. New mice, which had not previously been caught in that year, were identified to species, sexed, weighed to the nearest 0.5 g and the stretched body (tip of nose to anus) and tail length were measured. Mice destined to be released or to be kept in captivity for a day or more were marked by fur clippings. Those to be autopsied immediately were killed by cervical dislocation.

Within minutes of death, the body cavity was opened and inspected for overt cysts such as those of Taenia taeniaeformis. The bile duct was located and examined for evidence of the swelling characteristic of Hymenolepis microstoma infection. The entire small intestine was then removed and sectioned into oesophagus, stomach, anterior and posterior halves of the small intestine, caecum and colon. Each section was placed on a small piece of gauze, opened longitudinally and then suspended in a 50 ml plastic beaker containing Hanks's saline at 37°C. The only section to be treated slightly differently was the stomach, the contents of which were first removed by gentle immersion in saline. A different procedure was also adopted when H. microstoma infection was suspected on the basis of a hypertrophied bile duct. In such cases the intestine was removed with the entire liver. All portions of intestine from the posterior half of the small intestine and the stomach were treated as described above. The liver and anterior small intestine were then immersed in warm Hanks's saline in a Petri dish and the small intestine was opened from the posterior moving progressively anteriorly towards the bile duct. All beakers were incubated for a minimum of 2 h before examination. The contents of each beaker were transferred to a plastic Petri dish and examined under a binocular dissecting microscope. Parasites were then transferred with a Pasteur pipette, or using forceps for larger cestodes, to a separate dish for subsequent closer scrutiny. All gauzes were also examined and in no case were any parasites found to have been left attached to the gauze. All the tissues contained within the gauzes were also examined, first by shaking any remaining loose materials into saline and then after the entire mucosa had been scraped off with the edge of a glass slide.

Very few parasites were found after such examination of gut tissues. Only in exceptionally heavy infections with pinworms, were larvae occasionally detected and on rare occasions portions of cut worms were also identified. In general, all the parasites migrated out of the tissues and were easily located in the saline contained in the bottom of the beakers. The exception was *E. bacillatus* which did not detach from host tissues and required careful removal from the intestinal mucosa after the incubation period.

As far as possible, parasites were identified as they were encountered in Petri dishes and counted as they were removed to collecting dishes. Any unusual specimens were reexamined carefully. Tapeworms were measured and their scolices subjected to close

scrutiny. Hooks were counted, photographed and measured. All specimens were fixed: nematodes in 70% ethanol, cestodes in AFA.

Statistical procedures

Careful morphometric and parasitological records were kept of all data collected on each mouse. Where appropriate, data are presented as prevalence of infection (% of mice infected) or as Mean \pm Standard Error of the Mean (S.E.M.) parasite taxa per host species.

RESULTS

Hosts

Table I shows the number of animals of each sex examined in each year. As can be seen, significantly more male mice than females ($\chi^2=5.59$, p<0.02) were examined but, overall, the sample size was sufficient to enable quantitative analysis.

Parasites

Of the 58 mice examined, 75.8% were infected with at least one species of helminth parasite. In addition to the 10 helminth species recorded in Table II, one nematode and one cestode species were not identified. Each was found in only one mouse and sufficient material was recovered to establish that neither belonged to any of the species listed but insufficient to establish firmly the identity of the organisms involved. Data for these two unidentified species were excluded from the analysis. The mean number of

TABLE I. The number of Mus spretus examined during the study by gender and year.

| Year | Male | Female | |
|-------|--------|-------------|--|
| 1988 | 6 | 4 | |
| 1989 | 12 | 4 3 5 | |
| 1990 | 5 | 5 | |
| 1991 | 5 4 | 3 | |
| 1992 | 11 | 5 | |
| Total | 38 | 20 | |

TABLE II. Prevalence (%), total and mean numbers of taxa, and mean worm burden of helminth parasites in *Mus spretus*.

| Taxa | Prevalence | Total no. of taxa | Mean no. of $taxa \pm S.E.M.$ | Mean worm burden ± S.E.M. |
|------------------------------|------------|-------------------|-------------------------------|---------------------------|
| ALL HELMINTHES | 75-8 | 10 | 1.5 ± 0.2 | 15·6 ± 4·8 |
| NEMATODA | 60.3 | 5 | 0.9 ± 0.1 | 14·0 ± 4·8 |
| Nippostrongylus brasiliensis | 17.2 | == | = | 0.4 ± 0.2 |
| Aspiculuris tetraptera | 3.4 | 100 | = | 1.5 ± 1.5 |
| Syphacia obvelata | 46.6 | _ | _ | 11.5 ± 3.7 |
| Heterakis spumosum | 6.9 | - | = | 0.07 ± 0.03 |
| Eucoleus bacillatus | 8.6 | <u></u> | = | 0.4 ± 0.3 |
| PLATYHELMINTHES | 53.5 | 5 | 0.6 ± 0.1 | 1.6 ± 0.3 |
| CESTODA | 44-8 | 4 | 0.5 ± 0.1 | 1.0 ± 0.3 |
| Taenia taeniaeformis | 22-4 | - | _ | 0.3 ± 0.1 |
| Hymenolepis microstoma | 13-8 | = | = | 0.5 ± 0.2 |
| Hymenolepis nana | 6.9 | - | - | 0.1 ± 0.1 |
| Hymenolepis diminuta | 1.7 | _ | - | 0.02 ± 0.02 |
| DIGENEA | 15.5 | 1 | 0.2 ± 0.1 | 0.6 ± 0.3 |
| Brachylaemus spp. | 15.5 | 24 | ATOM | 0.6 ± 0.3 |

taxa recorded per mouse was 1.5 (Table II) although, as can be seen from Fig. 1, this ranged from 0 to 4 with the largest category supporting 2 species (18/58, 31%). The Shannon Index of diversity was calculated as 0.267 ± 0.045 and the effective number of abundant species (Hill's Index 1) as 1.38 ± 0.0768 .

Table II lists the species of helminth parasites recovered and identified. Although identical numbers of nematode and platyhelminth taxa were identified (5 each), nematodes were the most common (60·3% prevalence) and accounted for the majority of worms recovered with a mean worm burden of 14 (Table II). Among the nematodes, the most common species was S. obvelata with a prevalence of (46·6%) and a mean worm burden of 11·5 which was largely dependent on a few heavily infected animals (Fig. 2). All other species were found more rarely. The dominant tapeworms were T. taeniaeformis (22·4%) and H. microstoma which varied from year to year but was found in a total of 8 M. spretus representing a prevalence of 13·8%. However, the mean worm burden for H. microstoma was slightly higher because in 1989 one female mouse was found to harbour 12 specimens (Fig. 3).




FIG. 1. Number of uninfected mice compared with mice from which parasites belonging to 1-4 helminth taxa were recovered.




FIG. 2. The frequency distribution of Syphacia obvelata in Mus spretus. The index of dispersion I and the negative binomial constant k were estimated at 70.029 and 0.167 respectively.

DISCUSSION

This study, despite limited sample size, has provided for the first time detailed quantitative data on infections with parasitic helminths in *M. spretus*, a murine host which is relatively poorly studied and about which little is known. From available publications it appears that, unlike *M. m. domesticus*, this species shuns contact with man, preferring to live in a bush-grassland habitat in Mediterranean regions. Our experience of *M. spretus* suggests that it is relatively timid in comparison to *M. domesticus* and shows different patterns of behaviour (Hurst, personal observations).

Quite clearly, infections with *S. obvelata* accounted for the majority of helminth parasites in the mouse population and these were, in the main, attributable to relatively heavy infections in a few individuals (Fig. 2). Species of *Syphacia* are transmitted through eggs which embryonate within hours of deposition on the perianal skin surface of infected hosts (CHAN, 1952; Lewis & D'Silva, 1980; Kerboeuf & Lewis, 1987). Transmission is therefore likely to be most efficient among hosts which congregate and occupy permanent territories and runs. In fact, transmission of *Syphacia* species is believed to be maximal when hosts congregate for rest and sleep, which in rodents coincides with midday. Lewis & D'Silva (1980) showed that transmission of *S. muris* in caged rats coincided with noon and the early afternoon hours with virtually no deposition of eggs in the early evening when rats were most active in feeding and when most faeces were produced. Similar observations have been made for *S. obvelata* (Lewis & Shava, 1977). In contrast, *A. tetraptera*, whose eggs take about a week to embryonate (Anya, 1966) and are released in host faeces, showed maximum output shortly before dawn when faecal output was at its highest (Phillipson, 1974).

In our study, *M. spretus* exploited widely scattered resources in open grassland. As a consequence, individuals were widely distributed. *M. spretus* share semi-permanent tunnels through dense grass, but exhibit an interesting behavioural trait not shown by either *M. m. domesticus* nor *Apodemus sylvaticus* (Hurst, personal observations): these mice pick up or roll their own freshly-deposited faeces out of tunnels and away from




FIG. 3. The frequency distribution of *Taenia taeniaeformis* and *Hymenolepis microstoma* in *Mus spretus*. The index of dispersion I and the negative binomial constant k for T. taeniaeformis were estimated at 1.99 and 0.348 respectively and for H. microstoma at 6.461 and 0.095 respectively.

pathways. Transmission of parasites dependent on faecal eggs is thus likely to be less efficient in *M. spretus* than among *M. m. domesticus*: in this context it is interesting to note that nematode species transmitted through eggs voided in host faeces and requiring a period of days or even months to embryonate to the infective stage (e.g. *E. bacillatus*, *A. tetraptera*), were rare in *M. spretus* (Table I). The group-territorial social structure of commensal house mice (*M. musculus* and *M. m. domesticus*) allows them to live at very high densities where food resources are abundant (Bronson, 1979; Hurst, 1987a) and they share permanent well-made pathways to move between food and nest sites within buildings (Hurst, 1987b). Under these conditions, parasite transmission through contact with conspecifics and their faeces is likely to be very high.

The patterns of infection with platyhelminth species also deserve comment. Five species of cestodes were recovered although the prevalence of none exceeded 25% among the sampled population. All the tapeworm species identified in this study are known to infect *M. m. domesticus*, which are found locally among farm buildings and human dwellings. Local cats are known to carry *T. taeniaeformis* with a relatively low prevalence of 5% (DA SILVA LEITAO & CRUZ E. SILVA, 1967) and are known to range throughout the study sites. The hymenolepiid infections are all transmitted via insect vectors (beetles and fleas) although *H. nana* has the potential for direct transmission.

All of the other parasites we identified were common helminths of house mice but some deserve special mention. Foremost among these is Eucolleus bacillatus (syn. Trichosoma bacillatum, Eberth, 1863; Capillaria (Throminx) bacillata, Hall 1916) a capillariid nematode species which has not been previously reported from M. spretus in Europe although it has been recorded from this rodent in Tunisia (BERNARD, 1987) and from other rodents in Spain (Feliu et al., 1987a). Oldham (1931) listed E. bacillatus as a parasite of rats and SKRJABIN et al. (1957) gave Norwegian rats (Mus [Epimys] norvegicus) and house mice (Mus musculus) as the primary hosts but it has also been reported from A. sylvaticus (Spain, Feliu et al., 19887b; Tunisia, Bernard 1987; JUSTINE, 1989), Arvicola terrestris (Spain, Feliu et al., 1987c; Belgium, BERNARD, 1969), Arvicola terrestris, Clethrionomys glareolus (Belgium, BERNARD,, 1969) and Microtus majori (in USSR, KURASHVILI, 1989). The adults of this species were located intertwined deep in the stomach walls with only a small section of the female worm protruding through the mucosa to allow eggs to be shed into the gut lumen. As far as we can ascertain, nothing more is known about the biology and life cycle of E. bacillatus, for example whether an intermediate host is required or whether transmission of embryonated eggs can take place directly. In this context we were able to embryonate eggs which were fed to laboratory house mice kept on immunosuppressive agents to maximize the chances of establishment and development, but without success.

Among the rarer species, which we believe to be incidental infections, was Heterakis spumosum, as parasite of rats (WINFIELD, 1933; SMITH, 1953) reported previously from local Rattus norvegicus (ARANDAS REGO, 1966; AFONSO-ROQUE et al., 1984). This species was found at low prevalence and never more than 2 worms/host. Although primarily a parasite of rats, H. spumosum is known to cause transient, patent infections in house mice (ZINTZ & FRANK, 1982). Likewise, the few N. brasiliensis which we identified, also in low intensities, were likely to have been incidental infections from local rats which are known to carry this species with a high prevalence (>80%, see AFONSO-ROQUE et al., 1984). A. tetraptera is primarily a parasite of M. m. domesticus (and laboratory house mice, M. musculus) and was also occasionally encountered in our M. spretus population.

Finally, although we were limited by the small sample sizes examined in some years, and our study was conducted only in a two-week period in April of each of the years, the present work has provided original detailed information on parasitic helminths in a little-studied rodent host species, *M. spretus*.

ACKNOWLEDGEMENTS

We wish to thank Mr. A. Pircher, the Warden of the Quinta de Sao Pedro Research Station for his hospitality, his advice and support throughout our study. We are also grateful to the undergraduate students who participated in this study and without whose assistance the work would not have been possible, and to Mrs. E. Harris, Department of Zoology, The Natural History Museum, London, for assistance with the identification of *E. bacillatus*.

REFERENCES

- AFONSO-ROQUE, M. M., DE MENDONCA, M. M., MARCOS, M. V. & LOPES, F. J. (1984) Endoparasitas encontrados no rato cinzento (Rattus norvegicus Berk.) da zona de Lisboa. Revista Portuguesa de Doencas Infecciosas, 7, 101-109.
- ANYA, A. O. (1966) Studies on the biology of some oxyuroid nematodes. 1. Factors in the development of eggs of Aspiculuris tetraptera Schulz. Journal of Helminthology, 40, 253-260.
- ARANDAS REGO, L. A. (1966) Sobre alguns helmintas parasitas de *Rattus norvegicus* Berk, de Portugal. Anais Escola Superior Medicine Veterinaria, 8, 151-161.
- BAER, J. G. & TENORA, F. (1970) Some species of Hymenolepis (Cestoidea) from rodents and from primates. Acta Scientiarum Naturalium Academiae Scientiarum Bohemoslovacae Brno, 4, 1-32.
- BERNARD, J. (1969) Observations sur les helminthes parasites de Mammiferes et d'Oiseaux de la faune de Belgique. Archives de l'Institut Pasteur de Tunis, 46, 137-193.
- BERNARD, J. (1987) Faune des nematodes parasites des mammiferes de Tunisie et des contrees voisines. Archives de l'Institut Pasteur de Tunis, 64, 265-319.
- BRONSON, F. H. (1979) The reproductive ecology of the house mouse. *Quarterly Review of Biology*, **54**, 265–299.
- CHAN, K. F. (1952) Life cycle studies on the nematode Syphacia obvelata. American Journal of Hygiene, 56, 14-21.
- DA SILVA LEITAO, J. L. & CRUZ E SILVA, J. A. (1967) Contribuicao para o estudo de indice parasitario do gato domestico (Felis catus L.) da cidade de Lisboa. Anais da Escola Superior de Medicina Veterinaria, 9, 45-56.
- EBERTH, C. J. (1863) Untersuchungen uber Nematoden. pp 77. Leipzig.
- FELIU, C., MAS-COMA, S. & TENORA, F. (1987a) The peninsula effect on parasitofaunas: the helminths of Arvicolidae (Rodentia) in Spain. In: Program and Abstracts of the 6th European Multicolloquium of Parasitology, September 4-9th 1988, Budapest. pp. 146. Hungary, European Federation of Parasitologists.
- FELIU, C., GRACENEA, M. & TORREGROSA, M. (1987b) Consideraciones ecologicas sobre la helmintofauna de Apodemus sylvaticus (Linnaeus, 1758) (Rodentia: Muridae) en el Pirineo Oriental Espanol. In: Mamiferos y Helmintos. Volumen Homenaje al Prof. Dr. Herman Kahmannen su 81 Aniversario. Eds. V. Sans-Coma, S. Mas-Coma & J. Gosalbez. pp. 175-181. Ketres Editora, Barcelona.
- FELIU, C., TORRES, J., GOSALBEZ, J., VENTURA, J. & GRACENEA, M. (1987c) Influencias de algunos factores ecologicos sobre las helmintofaunas de Arvicola spp. (Rodentia: Arvicolidae) en el Nordeste Iberico. In: Mamiferos y Helmintos. Volumen Homenaje al Prof. Dr. Herman Kahmann en su 81 Aniversario. Eds. V. Sans-Coma, S. Mas-Coma & J. Gosalbez. pp. 183–193. Ketres Editora, Barcelona.
- FURMAGA, S. (1957) Helmintofauna gryzoni polnych (Rodentia) okolic Lublina. Acta Parasitologica Polonica, 5, 9-50.
- HALL, M. C. (1916) Nematode parasites of mammals of the Orders Rodentia, Lagomorpha and Hyracoidea. Proceedings of the U.S. National Museum, 50, 1–258.
- HURST, J. L. (1987a) Behavioural variation in wild house mice (*Mus domesticus* Rutty): a quantitative assessment of female social organization. *Animal Behaviour*, 35, 1846–1857.
- HURST, J. L. (1987b) The functions of urine marking in a free-living population of house mice, Mus domesticus Rutty. Animal Behaviour, 35, 1433-1442.
- JUSTINE, J. L. (1989) Liste des Capillaria (Nematoda, Capillariinae) parasites de mammiferes africains. Bulletin du Museum National d'Histoire Naturelle, Paris, 11, 755-762.
- KERBOEUF, D. & LEWIS, J. W. (1987) Rhythmic behaviour of intestinal helminths in rodents. *Mammal Review*, 17, 127-134.
- KISIELEWSKA, K. (1970) Ecological organization of intestinal helminth groupings in *Clethrionomys glareolus* (Schreb.) (Rodentia). I. Structure and seasonal dynamics of helminth groupings in a host population in the Bialowieza National Park. *Acta Parasitologica Polonica*, 18, 121–147.
- KURASHVILI, B. E. (1989) Parasitic Worms of the Small Mammal Fauna of the Transcaucasus. Metsniereba, Tbilisi (In Russian).
- LANGLEY, R. & FAIRLEY, J. S. (1982) Seasonal variations in infestations of parasites in a wood mouse Apodemus sylvaticus population in the west of Ireland. Journal of Zoology, 198, 249–261.
- LEWIS, J. W. (1968a) Studies on the helminth parasites of the long-tailed field mouse *Apodemus sylvaticus* sylvaticus from Wales. *Journal of Zoology*, **154**, 287-312.
- LEWIS, J. W. (1968b) Studies on the helminth parasites of voles and shrews from Wales. *Journal of Zoology*, 154, 313-331.

LEWIS, J. W. & D'SILVA, J. (1980) Rhythmic egg deposition by the oxyurid nematode Syphacia muris in the rat. Journal of Zoology, 191, 429–433.

- LEWIS, J. W. & SHAVA, F. H. M. (1977) Rhythmic egg deposition by intestinal nematodes of laboratory mice. Parasitology, 75, iv.
- LEWIS, J. W. & TWIGG, G. I. (1972) A study of the internal parasites of small rodents from woodland areas in Surrey. *Journal of Zoology*, **166**, 61–77.
- MONTGOMERY, S. S. J. & MONTGOMERY, W. I. (1988) Cyclic and non-cyclic dynamics in poulations of helminth parasites of wood mice, Apodemus sylvaticus. Journal of Helminthology, 62, 78–90.
- OLDHAM, J. N. (1931) The helminth parasites of common rats. Journal of Helminthology, 9, 49-90.
- PHILLIPSON, R. F. (1974) Intermittent egg release by Aspiculuris tetraptera in mice. Parasitology, 69, 207-213.
- SAGE, R. D. (1981) Wild mice. In: The Mouse in Biomedical Research. Volume 1, pp. 39-90. Academic Press, London.
- SHARPE, G. I. (1964) The helminth parasites of some small mammal communities. I. The parasites and their hosts. *Parasitology*, 54, 145–154.
- SKRJABIN, K. I., SHIKHOBALOVA, N. P. & ORLOV, I. V. (1957) In: *Trichocephalidae and Capillariidae* of Animals and Man and the Diseases Caused by Them. Gel'mintologicheskaya Laboratoriya Osnovy Nematodologii, Vol. 6. pp. 391. USSR: Akademiya Nauk SSR, Translated from Russian and publiched by Israel Program for Scientific Translations, Jerusalem, 1970.
- SMITH, P. E. (1953) Life history and host-parasite relationship of *Heterakis spumosa*, a nematode parasite in the colon of the rat. *American Journal of Hygiene*, 57, 194–221.
- SOLTYS, A. (1957) Badania nad robakami pasozytniczymi drobnych gryzoni Parku Narodowego w Bialowiezy. Acta Parasitologica Polonica, 12, 487–504.
- TENORA, F. & MURAI, E. (1972) Recent data on five species of the genus *Hymenolepis* (Weinland, 1858) (Cestoidea, Hymenolepididae) parasitizing rodents in Hungary. *Acta Zoologica Academiae Scientiarum Hungaricae*, 18, 129–145.
- WINFIELD, G. A. (1933) Quantitative experimental studies on the rat nematode Heterakis spumosa, Schneider, 1866. American Journal of Hygiene, 17, 168–228.
- ZINTZ, K. & FRANK, W. (1982) Ultrastructural modifications in *Heterakis spumosa* after treatment with febantel or mebendazole. *Veterinary Parasitology*, 10, 47–56.

Accepted 31st March, 1993.