



The University of
Nottingham

UNITED KINGDOM · CHINA · MALAYSIA

Richens, Joanna L. and Vere, Kelly-Ann and Light, Roger A. and Soria, Daniele and Garibaldi, Jonathan and Smith, A. David and Warden, Donald and Wilcock, Gordon and Bajaj, Nin and Morgan, Kevin and O'Shea, Paul (2014) Practical detection of a definitive biomarker panel for Alzheimer's disease: comparisons between matched plasma and cerebrospinal fluid. *International Journal of Molecular Epidemiology and Genetics*, 5 (2). pp. 53-70. ISSN 1948-1756

Access from the University of Nottingham repository:

http://eprints.nottingham.ac.uk/28163/1/IJMEG0000141R1_accepted.pdf

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the Creative Commons Attribution licence and may be reused according to the conditions of the licence. For more details see:
<http://creativecommons.org/licenses/by/2.5/>

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

Practical detection of a definitive biomarker panel for Alzheimer's Disease; comparisons between plasma and cerebrospinal fluid

Joanna L Richens¹, Kelly-Ann Vere¹, Roger A Light², Daniele Soria³, Jonathan Garibaldi³, A David Smith⁴, Donald Warden⁴, Gordon Wilcock⁵, Nin Bajaj⁶, Kevin Morgan⁷, Paul O'Shea¹

¹Cell Biophysics Group, Institute of Biophysics, Imaging and Optical Science, School of Life Sciences, University of Nottingham, University Park, Nottingham, United Kingdom;

²Institute of Biophysics, Imaging and Optical Science, University of Nottingham, University Park, Nottingham, United Kingdom;

³School of Computer Science, University of Nottingham, Jubilee Campus, Nottingham, United Kingdom and Advanced Data Analysis Centre, University of Nottingham, Nottingham, United Kingdom;

⁴Oxford Project to Investigate Memory and Ageing (OPTIMA), University Department of Physiology, Anatomy and Genetics, University of Oxford, Oxford, United Kingdom;

⁵Oxford Project to Investigate Memory and Ageing, Nuffield Department of Clinical Medicine, John Radcliffe Hospital, Oxford, United Kingdom;

⁶Department of Neurology, Nottingham University Hospitals NHS Trust, Queen's Medical Centre, Nottingham, United Kingdom;

⁷School of Life Sciences, Translational Cell Sciences – Human Genetics Group, University of Nottingham, Queen's Medical Centre, Nottingham, United Kingdom;

Address correspondence to: Professor Paul O'Shea, Cell Biophysics Group, Institute of Biophysics, Imaging and Optical Science (IBIOS), School of Life Sciences, University of Nottingham, Nottingham, England, UK, NG7 2RD. Phone: +44(0)1159513209, Fax +44(0)1158466580, E-mail: paul.oshea@nottingham.ac.uk,

Running title: Plasma biomarker panel for Alzheimer's disease

Disclosure of conflict of interest: None.

Abbreviations:

AD: Alzheimer's disease; CSF: Cerebrospinal fluid

Abstract

Previous mass spectrometry analysis of cerebrospinal fluid (CSF) has allowed the identification of a panel of molecular markers that are associated with Alzheimer's disease (AD). The panel comprises Amyloid beta, Apolipoprotein E, Fibrinogen alpha chain precursor, Keratin type I cytoskeletal 9, Serum albumin precursor, SPARC-like 1 protein and Tetranectin. Here we report the development and implementation of immunoassays to measure the abundance and diagnostic capacity of these putative biomarkers in matched lumbar CSF and blood plasma samples taken in life from individuals confirmed at post-mortem as suffering from AD (n=10) and from screened 'cognitively healthy' subjects (n=18). The inflammatory components of Alzheimer's disease were also investigated. Employment of supervised learning techniques permitted examination of the interrelated expression patterns of the putative biomarkers and identified inflammatory components, resulting in biomarker panels with a diagnostic accuracy of 87.5% and 86.7% for the plasma and CSF datasets respectively. This is extremely important as it offers an ideal high-throughput and relatively inexpensive population screening approach. It appears possible to determine the presence or absence of AD based on our biomarker panel and it seems likely that a cheap and rapid blood test for AD is feasible.

Keywords: Alzheimer's disease, Biomarker, Blood Plasma, Cerebrospinal fluid

Introduction

Neurodegenerative disorders, and in particular Alzheimer's disease (AD), are having an increasing impact on our society. By 2040 dementia, of which AD is the most common cause, is predicted to affect 81.1 million individuals worldwide [1]. Although AD therapeutics including Donepezil, Rivastigmine, Galantamine and Memantine are available, AD remains incurable as these drugs only provide some symptomatic relief and do not modify disease progression [2, 3]. As AD is associated with a progressive decline in health, earlier detection may result in enhanced therapeutic success [3]. Current diagnostic procedures are, however, inadequate for early disease detection or the accurate differentiation of AD from other forms of dementia. Provision of enhanced cost-effective diagnostic strategies is vital and a biochemical test that is diagnostic and predictive of AD would help achieve this aim.

Investigations into biomarkers for AD have, to date, failed to identify a single molecule capable of fully depicting the disease state. Genetically AD is one of the best characterised diseases [4-9], however only approximately 24% of heritability has been resolved [10] and possession of genetic indicators (e.g. the E4 allele of the *APOE* gene) does not guarantee disease onset [11]. Diagnosis based solely on a genetic marker would, therefore, result in many inaccurate diagnoses. Studies into individual AD protein biomarkers have tended to focus on the constituents of amyloid beta deposits and neurofibrillary tangles as these are the characteristic hallmarks of the disease [12] but such studies have demonstrated varying levels of success [13-16]. A recent study examining amyloid beta peptide 1-42 (A β 42) has, however, lent support to the belief that the underlying causative factors of AD are initiated many years before the symptoms of the late-onset form of AD (LOAD) manifest [17]. In this study, cerebrospinal fluid (CSF) concentrations of A β 42 were shown to have reached pathological levels 5-10 years prior to conversion from mild cognitive impairment (MCI) to AD, highlighting the huge benefits that biomarker identification could have in the early diagnosis of AD. However, as is the case with single biomarkers in most complex disorders, A β 42 does not appear to have the necessary power to act as a stand-alone AD marker. The findings of this study indicate that 10% of individuals with MCI who had pathological levels of A β 42 did not go on to develop AD. As such these measurements would need to be utilised in conjunction with additional diagnostic procedures and it may be that a panel of biomarkers incorporating A β 42 may be more appropriate.

Vafadar-Isfahani and colleagues recently identified a panel of CSF biomarkers capable of differentiating between healthy individuals and those with AD [18]. Comprising Amyloid beta, Apolipoprotein E, Fibrinogen alpha chain precursor, Keratin type I cytoskeletal 9, Serum albumin precursor, SPARC-like 1 protein and Tetranectin, the diagnostic

performance of this biomarker panel was found to improve as more markers were sequentially added to the model for diagnosis i.e. the effect was additive suggesting that all the markers are necessary for accurate diagnosis. The panel of markers also demonstrated its potential utility in early diagnosis of AD by mapping individuals with Mild Cognitive Impairment (MCI) at an intermediate point between samples from healthy and AD individuals. To enable routine screening of a population, however, it would be preferable for any biochemical test developed to be analysed in blood plasma as this is a far less invasive clinical sample to collect from a patient than CSF. In this study we first aim to identify the components of the recently identified CSF AD biomarker panel [18] in blood plasma. We then proceed to determine their plasma and CSF concentrations and assess their potential utility as diagnostics tools. As Tau is considered one of the foremost AD biomarkers, we have included it in this study along with Clusterin which was recently identified as a potential blood plasma biomarker for AD [19].

Materials and methods

Patient Samples

Sample cohorts used in this study were obtained from the Oxford Project to Investigate Memory and Ageing (OPTIMA; University of Oxford, UK). The OPTIMA study received approval from the Central Oxford Ethics Committee and all individuals gave informed written consent to participate in the study. For 10 patients with a clinical diagnosis of probable AD, 'definite' AD was diagnosed pathologically by the established CERAD criteria. The 18 control subjects were cognitively screened annually for at least 3 years and 6 came to autopsy and were classified as CERAD 'negative'. See a recent report for a brief description of the OPTIMA cohort, CSF sampling procedure and post-mortem analysis [20]. The average interval between CSF sampling and death was 2090 days in the controls and 1806 days in the AD patients.

Protein detection by immunoassay: ELISA

SPARCL1 was analysed by sandwich ELISA using capture and biotinylated detection antibodies from Creative Biomart (CAB-701MH) and R&D Systems (BAF2728) respectively. The wells of ELISA plates were coated overnight with 5 µg/ml capture antibody (CAB-701MH in PBS). Following this and every subsequent incubation wells were washed three times with PBS-0.05% Tween20 (PBST). Antibody-coated wells were blocked with PBST-1% BSA for 1 hr, 50 µl sample (either clinical sample or protein standard) was added for 2 hr and then biotinylated detection antibody (BAF2728) was added at 500 ng/ml for 2 hr. The reaction was developed using streptavidin-HRP and TMB substrate, stopped with 1 M HCl and read at 450 nm. Concentrations of SPARCL1 in the samples were determined from a standard curve generated with recombinant SPARCL1 (R&D Systems; 2728-SL). Keratin 9 was detected according to the manufacturer's protocol using an ELISA kit purchased from antibodies-online GmbH (ABIN417500). ELISA kits against β -Amyloid peptide 1-40 (A β 40) and A β 42 were purchased from Wako Chemicals GmbH (298-64601 and 296-64401) and used according to manufacturer's instructions.

Antibodies and recombinant proteins

Appropriate pairs of monoclonal capture antibody and biotinylated polyclonal detection antibody and a corresponding recombinant protein standard were sourced as displayed in Table 1 and verified by western blot and ELISA.

Antibody coupling to microspheres

COOH-coated fluorescently dyed microspheres (Bio-Plex) were purchased from Bio-Rad (Hercules, CA). Bead regions 011, 020, 027, 033 and 042 were assigned to ApoE,

Tetranectin, Fibrinogen alpha chain, Clusterin and Tau respectively. Monoclonal capture antibodies were coupled to the microspheres using the Bio-Plex Amine Coupling Kit (Bio-Rad, 171-406001) according to manufacturer's instructions. When necessary, pre-processing with Micro Bio-Spin 6 columns (Bio-Rad) was undertaken to ensure antibodies were in PBS buffer containing no additives prior to the attachment procedure. Assays were initially developed and optimised individually for target protein before being combined step-by-step to identify any problems due to reagent cross-reactivity that may have occurred.

Luminex procedure

Designated wells of a filter plate were pre-wetted with wash buffer (PBS-0.05% Tween20 (PBST)). A bead suspension containing 5000 of each antibody-conjugated bead set was added to each well, washed twice with PBST and resuspended in incubation buffer (PBS-1% BSA). To this, a sample of appropriately diluted clinical sample or protein standard was added, and the plate was incubated for 2 hr at 25°C on a rotating plate shaker (600 rpm). Wells were then washed twice with PBST, and incubated with a cocktail of biotinylated detection antibodies (containing each antibody at a pre-determined optimal concentration). The plate was incubated for 1 hr at 25°C on a rotating plate shaker after which wells were again washed twice with PBST and the reporter molecule (Streptavidin-RPE) was added to the appropriate wells. Finally, following a 30 min incubation at 25°C on a rotating plate shaker and two washes with PBST, PBST was added to each well in preparation for analysis. Data were acquired on a Bio-Plex 200 system and analysed with associated software (Bio-Rad). All multiplex assays were performed in duplicate. In each well, a minimum of 100 beads per target molecule were analysed for both bead designation and R-phycoerythrin fluorescence. Clinical sample concentrations of each target protein were determined from standard curves generated using recombinant proteins.

Human cytokine 30-plex Luminex Assay

Cytokine levels in both plasma and CSF were measured using the Luminex human cytokine 30-plex panel (Biosource, Camarillo, CA). This kit is able to simultaneously measure human IL-1 β , IL-1RA, IL-2, IL-2R, IL-4, IL-5, IL-6, IL-7, IL-8, IL-10, IL-12p40/p70, IL-13, IL-15, IL-17, TNF- α , IFN- α , IFN- γ , GM-CSF, MIP-1 α , MIP-1 β , IP-10, MIG, Eotaxin-1, RANTES, MCP-1, VEGF, G-CSF, EGF, FGF-basic and HGF. Samples were analysed according to manufacturer's instructions. Briefly, incubation buffer and 1:2 diluted plasma samples were pipetted into wells and incubated with the beads for 2 hours. Wells were washed, incubated with biotinylated detector antibody for 1 hour, washed again and incubated for 30 mins with streptavidin-RPE. Wells were washed to remove unbound streptavidin-RPE prior to analysis. All samples and standard curves were performed in duplicate. Data was acquired

on a Bio-Plex 200 system and analysed with associated software (Bio-Rad). Standard curves for each cytokine were generated using the pre-mixed lyophilised standards provided in the kits and the cytokine concentrations in samples were determined from the appropriate standard curve.

Individual biomarker analysis

Data was analysed using GraphPad Prism Version 5.02. Concentrations of the markers in the cohorts were compared using the Mann-Whitney test and Spearman correlation coefficients were determined as appropriate. A p-value ≤ 0.05 was considered to be a practical level of clinical significance.

Biomarker panel analysis

To determine the effectiveness of each individual marker, the statistical data was used to calculate the proportion of both the healthy and the diseased population that would be incorrectly diagnosed. It was assumed that both the healthy and the diseased population are normally distributed, but the principle is valid for other distributions.

Considering first the healthy population, take $P_h(c)$ as the probability distribution with respect to marker concentration c . The first step involves calculating the range of marker concentrations where the diseased population overlaps with the healthy population. This is entirely covered by the range of concentrations for the diseased population, so the range can be set to $\mu_d \pm n\sigma_d$, where μ_d and σ_d are the mean and standard deviation of the diseased population respectively, and n is the number of standard deviations from the mean. In this work, n was chosen as 3. We define the proportion of the healthy population that lies within this range to be O_h . An example can be seen in Figure 1, where the healthy population that cannot be correctly diagnosed with the single example marker is shown in grey.

Individuals with marker concentrations in the overlap range are at risk of being misdiagnosed. Determining O_h for a marker gives a measure of its usefulness. O_h can be calculated as the cumulative distribution of the healthy population, P_h , that lies within the overlap range. This can be found by integrating the probability density function for the healthy population, P_h , over the overlap range, as shown in Equation 1.

$$\text{Equation 1} \quad O_h = \int_{\mu_d - n\sigma_d}^{\mu_d + n\sigma_d} P_h(c) dc$$

In some cases the lower integration limit will be negative, implying a negative marker concentration which is of course not possible, so the general case above must be modified.

The probability density function, $P_h(c)$, is multiplied by the step function $H(c)$ as defined in Equation 2.

$$\text{Equation 2} \quad H(c) = \begin{cases} 0, & c < 0 \\ 1, & c \geq 0 \end{cases}$$

This removes the possibility of negative marker concentrations. It also means that the total area of $P_h(c)$ that is under consideration, defined as T_h , is less than 1. T_h must be found so that O_h can be calculated as a proportion. T_h is calculated in Equation 3.

$$\text{Equation 3} \quad T_h = \int_0^{\infty} P_h(c)dc$$

T_h may also be calculated using Equation 4, where the upper integration limit is set based on the healthy population mean and standard deviation to give a more practical limit. The parameter $m = 6$ was chosen in this work.

$$\text{Equation 4} \quad T_h = \int_0^{\mu_h + m\sigma_d} P_h(c)dc$$

The final result for O_h is given in Equation 5 with the same result for the diseased population in Equation 6.

$$\text{Equation 5} \quad O_h = \frac{1}{T_h} \int_{\mu_d - n\sigma_d}^{\mu_d + n\sigma_d} P_h(c)H(c)dc$$

$$\text{Equation 6} \quad O_d = \frac{1}{T_d} \int_{\mu_h - n\sigma_h}^{\mu_h + n\sigma_h} P_d(c)H(c)dc$$

Supervised learning techniques

The parameters O_h and O_d give an indication of how well each individual marker performs as a diagnostic tool, but to get the full power they must be considered together as a panel of markers. To validate the data without preconception of how to classify the subject groupings, three data analysis techniques based on supervised learning techniques were employed; a decision tree classifier (C4.5), a Bayesian classifier (Naïve Bayes) and a Multilayer Perceptron artificial neural network (ANN).

In the C4.5 algorithm each attribute of the data can be used to make a decision that splits the data into smaller subsets. It examines the normalised information gain that results from choosing an attribute for splitting the data. The attribute with the highest normalised information gain is used to make the decision. The algorithm then recurs on the smaller sublists. The system outputs as a decision tree, or a set of if-then rules, which can be used

to classify new cases, with an emphasis on making the model understandable and accurate [21].

The Naive Bayes is a probabilistic classifier based on Bayes' theorem with strong independence assumptions that can deal with large numbers of cases and variables. It aims to predict the class of test instances as accurately as possible and is termed naive because it is based on two simplifying assumptions: (1) That the predictive attributes are conditionally independent given the class and (2) The values of numeric attributes are normally distributed within each class [22].

The Multilayer Perceptron is a feed-forward ANN model that maps sets of input data onto a set of appropriate output. It has three distinctive characteristics: (1) The model of each neuron in the network includes a nonlinear activation function. (2) The network contains one or more layers of hidden neurons (not part of the input or output of the network) which enable the network to learn complex tasks by extracting progressively more meaningful features from the input patterns. (3) The network exhibits a high degree of connectivity, determined by the synapses of the network. A change in the connectivity of the network requires a change in the population of synaptic connections or their weights. The Multilayer Perceptron derives its computing power from these characteristics together with the ability to learn from experience through training [23]. This system is trained to solve problems using the error back-propagation algorithm.

The classification algorithms were applied to the two data sets (CSF/Plasma) to examine whether the classifications into Healthy/AD could be reproduced. Each technique was implemented using a 'leave-one-out' cross-validation, to estimate how accurately the predictive model performs in practice. At each step, the whole data but one point are used for training and the remaining point is used for testing. The process is then repeated n-times, where n is the total number of patients.

Results

The concentrations of eight putative AD protein biomarkers were determined in plasma and CSF samples collected simultaneously from individual 'donors'. The 'matched' nature of the samples makes it feasible for comparisons to be made between the protein constituents of each sample, but this also led to restrictions of the overall cohort sizes. All of the proteins examined (A β 40, A β 42, ApoE, Clusterin, Fibrinogen alpha chain, Keratin 9, SPARCL1, Tau, Tetranectin) could be detected in CSF and plasma using immunoassays (Figure 2). The mean concentrations of the proteins measured in both sample sets are shown in Table 1. SPARCL1 ($p = 0.035$) and A β 42 ($p = 0.049$) were found to be significantly decreased in CSF from AD patients in comparison to healthy controls. Keratin 9 was only detectable in CSF samples from three AD patients (Mean concentration \pm SD; 3.75 ± 5.9 pg/ml) and not any healthy individuals. In plasma from AD patients, the concentrations of A β 42 ($p = 0.0008$) and Tau ($p = 0.047$) were significantly decreased in comparison to those observed in healthy controls. Significant correlations between the plasma and CSF concentrations of ApoE, SPARCL1, Tau and Tetranectin were determined in healthy individuals but were found to be weaker in the AD patient cohort (Table 2).

To investigate the inflammatory component of AD, samples were examined using Luminex technology, allowing simultaneous assessment of 30 cytokines. In the plasma and CSF samples, 11 and 12 respectively of the cytokines examined were not present at concentrations that could be detected using this technology and are omitted from Table 3. Furthermore, some samples contained levels of RANTES that exceeded the maximum range of the standard curve so these values have also been omitted from Table 3. Significant differences (p -value ≤ 0.05) in concentration between AD and healthy samples were observed for FGF-basic ($p = 0.05$), IL-1RA ($p = 0.02$), MCP-1 ($p = 0.01$) and MIP-1 β ($p = 0.04$) in plasma samples as illustrated in Figure 3. These differences were not replicated in the matched CSF samples in which only IL-12 ($p = 0.04$) demonstrated a significant change.

In order to gain more insight into the mechanisms of disease and potential pathways implicated, we sought to identify any correlations that exist between components of the targeted biomarker panel and the inflammatory proteins shown to be significantly altered in AD (FGF-basic, IL-12, IL-1RA, MCP-1 and MIP-1 β). The correlation of Keratin 9 with the other proteins could only be established in the disease cohort as it was not detectable in samples from the healthy cohort. Whilst all potential combinations of proteins were analysed, only those shown to be significantly correlated are displayed in Table 4 along with the p -value of the correlation and the effect of disease on the correlation. All of the targeted

biomarker panel and inflammatory proteins were found to be significantly correlated ($p \leq 0.05$) with at least one other protein (Table 4). Identification of a correlation in CSF samples did not automatically infer a correlation between that protein pair in the corresponding plasma samples as only the A β 40/A β 42 and MCP-1/Tetranectin pairings demonstrated the same patterns of correlations in CSF and plasma samples. In these cases a correlation that existed between the proteins in healthy samples was disrupted in the disease cohorts. Interestingly, some of the protein targets that were not significantly altered in the disease state (Figure 2 and Table 2) were found to correlate with some of those that exhibited significant changes. For example, clusterin, which had p-values of 0.90 and 0.55 in plasma and CSF respectively, was found to correlate with A β 40, A β 42, FGF-basic, MCP-1 and Tau demonstrating its potential importance in the disease network.

Despite having identified significant differences in the concentrations of several proteins in AD samples, it would still not be possible to accurately differentiate an AD sample from a healthy sample using purely individual protein biomarkers as there is still significant overlap present between the ranges of concentrations of the two separate cohorts (Figure 2). Examination of this overlapping region allowed numerical assessment of the diagnostic power of each individual marker in the healthy and disease states for both CSF and plasma samples (Table 5). Keratin 9 values were determined using the detection limits of the ELISA kit used during sample analysis. When analysed in this manner, the individual markers that provided the most power in identifying healthy CSF, AD CSF, healthy plasma and AD plasma were IL-1RA (66.43%), Keratin 9 (98.76%), FGF-basic (82.13%) and MIP-1 β (28.05%) respectively. It is important, however, that these single protein markers are investigated as a combined panel of biomarkers to determine whether this can provide improved diagnostic ability and permit identification of a subject's disease state. Data analysis techniques based on supervised learning techniques were, therefore, employed in order to validate our data without preconception of how to classify the subject groupings. The biomarker panel examined comprised the original biomarker panel plus all detectable inflammatory markers. Due to the nature of the analysis, i.e. analysis of a complete biomarker panel, only samples where measurements had been obtained for all of these markers could be utilised.

In the CSF samples, the C4.5 algorithm did not perform well, resulting in only seven (46.7%) patients being correctly classified, whilst the remaining eight were misclassified. The naïve Bayes classifier performed slightly better, returning 60% (i.e. nine patients) accuracy. The remaining six patients were not correctly assigned to their category. The most accurate results were obtained with the neural network, with 13 patients been correctly classified

(86.7%). When the whole data set was used for both training and testing, all three classification techniques reached an accuracy of 100% (all patients correctly classified). For the measurements made in plasma samples, the C4.5 algorithm classified 14 out of 16 (87.5%) correctly. The C4.5 algorithm shows that, based upon this dataset, it is enough to check the values of the M7 variable ($A\beta_{42}$) to establish whether a patient should be classified as Healthy or as AD. However, when the whole data were used for both training and testing, not all patients were correctly classified (15/16). In plasma samples the naïve Bayes algorithm reached an accuracy of 75% (i.e. 12 patients correctly classified). This method was also able to classify correctly all patients when both training and testing were performed using the whole data. Finally, the ANN was less accurate than for the CSF samples, assigning only 13 out of 16 patients (81.25%) to the correct group, but it reached 100% accuracy when the whole data was used for training and testing. It is evident that the classification techniques performed better on the plasma samples with more data available. The naïve Bayes classifier was unable to outperform the other two techniques; this is likely to be due to the cohort size as this method is known to perform well with big datasets and if data are normally distributed. For the CSF samples the ANN was the best classifier, but unfortunately it is difficult to access the 'rules' used for classification by this method as they are not 'visible'.

Discussion

Confirmation of the presence of a disease to acceptable levels of clinical confidence from a single molecular indicator (i.e. a biomarker) is not a trivial task. Practical examples of such markers do, however, exist. Prostate-specific antigen (PSA) and Human Epidermal Growth Factor Receptor 2 (HER2), for example, are employed diagnostically for prostate and breast cancers respectively. Whilst even these exemplars display limitations (HER2 is not definitive of all breast cancer subtypes [24] and the diagnostic assay for PSA produces large numbers of false positives [25]), they provide useful diagnostic tools and, particularly in the case of HER2, have led to improved therapeutic intervention [26]. Investigations into biomarkers for AD have, to date, failed to identify a single molecule capable of fully depicting the disease state due in part to the complex nature of the pathways underlying the disease [27]. An alternative approach whereby panels of biomarkers are investigated (as opposed to a singular marker) has, in comparison, heralded some success an example of which being the recent identification of a panel of cerebrospinal fluid (CSF) biomarkers capable of differentiating between AD and healthy individuals. This panel comprised some novel protein biomarkers not previously associated with AD and, encouragingly, was applied in the identification of a cohort of individuals with mild cognitive impairment (MCI) [18].

The initial aim of this study was to demonstrate that the components of this biomarker panel could be detected and quantified in blood plasma. Further to this our aim was to determine the concentrations of these proteins in CSF and to examine their utility as individual biomarkers in their own rights in both types of clinical sample (CSF and plasma). It is well documented that albumin is the most abundant protein component of both blood plasma [28] and CSF [29]. It also has limited utility as a biomarker due to its physical properties, the inter-individual variation in ranges that it exhibits and the vastly higher concentrations that it exists at compared to the other biomarker components of the panel under consideration [28, 30]. Instead of providing an accurate measure of disease onset and/or progression, we now consider that it is more likely that albumin is acting as a marker of damage to the blood-CSF barrier [31] and therefore it was omitted from our current study. As an association between inflammatory genes/proteins and AD has been previously demonstrated [32-34], the inflammatory profiles of the samples were also assessed to investigate whether they would positively augment the diagnosis obtained using the original biomarker panel.

As illustrated in Figure 2, all of the components of the biomarker panel that were under consideration (A β 40, A β 42, ApoE, Clusterin, Fibrinogen alpha chain, Keratin 9, SPARCL1, Tau and Tetranectin) could be measured in blood plasma samples from both healthy and diseased individuals using immunoassay. All proteins could also be detected in CSF

samples from both healthy and diseased individuals with the exception of Keratin 9 which was not detectable in CSF from healthy individuals. To our knowledge, this is only the second demonstration of the presence of Keratin 9, a protein normally considered to be associated with skin, in blood. Previous data suggests that Keratin 9 may act as a serum marker of metastasis of hepatocellular carcinoma [35]. It is also the first immunoassay validation of the presence of Keratin 9 in CSF; the only previous association between CSF Keratin 9 and AD was reported in a study by Vafadar-Isfahani and colleagues [18].

The cohort sizes under investigation in this study were modest, but are in keeping with other similar studies [36]. Perhaps unexpectedly given these relatively small cohorts, significant differences in the mean concentrations of some of the biomarker panel components and inflammatory proteins were found in both CSF and plasma samples (Figures 2 and 3; Tables 2 and 3) suggesting that these proteins may have potential utility as individual biomarkers in AD. The most notable potential AD biomarker identified in this study is CSF Keratin 9 as this was found to only be present in CSF samples collected from AD patients, not healthy individuals. Another promising protein marker is SPARCL1, levels of which were significantly altered in CSF samples of AD patients when compared to healthy controls ($p = 0.035$) and approaching significance (0.089) in plasma samples. SPARCL1 has been linked to several diseases including various forms of cancer [37-43], uterine leiomyomas [44] and multiple sclerosis [45]. An association between SPARCL1 and AD has previously been demonstrated by Yin *et al* using 1D electrophoresis followed by LC-MS/MS [46]. The study also identified SPARCL1 as a potential target for Parkinson's disease [46]. The other proteins found to be significantly altered in the disease state during this study, A β 42 and Tau have extensively been linked to AD and thus this study provides further validation of their potential utility in AD diagnostics.

When looking at the significance of the association of the proteins with AD in CSF, the most powerful (excluding Keratin 9 for which a p value cannot be determined) are SPARCL1, A β 42 and Fibrinogen. These three proteins were also identified as the three most important components of the AD biomarker panel derived in the mass spectrometry study that preceded this study [18]. The levels of correlation between plasma and CSF protein concentrations were found to be variable (Table 2). The blood brain barrier restricts molecular diffusion (including capillary walls among other structures) and associated biochemical and physiological processes to maintain a dynamical blood/CSF concentration gradient. The observed differences that we found seem most likely due to compromises in this barrier function which have been demonstrated to be compromised in neurological conditions including AD [47].

Evidence of an inflammatory component to AD has been emerging in the literature over the last few years, however the nature and extent of the inflammatory response has yet to be fully elucidated. A variety of cytokines and chemokines have been implicated in the disease process [36, 48, 49]. Indeed, the first suggestion of a biomolecular fingerprint for AD included a large number of inflammatory markers [34]. The majority of the cytokines examined in this study were detectable in both plasma and CSF which is in contrast to a previous study undertaken using Luminex in which only one cytokine out of the 22 examined was detectable [50]. These differences could be due to the sensitivity of the detection kit being utilised or the disease stage of the patients being studied. Correlation between cytokine levels and disease progression have been previously identified [51]. Motta and colleagues demonstrated an initial large increase in immune responsiveness for some cytokines in mild-stage AD with respect to comparative healthy individuals. As the disease progressed from the early stages a gradual decline in cytokine levels were observed until concentrations in individuals with late-stage disease had dropped below those observed in the healthy cohort [51]. When taking the findings of this previous study into consideration, the decrease in levels of many of the inflammatory markers observed within this study may have been expected as the samples investigated represent late-stage disease.

Of the detectable cytokines, we observed significant decreases in CSF levels of IL-12 ($p = 0.04$) and plasma levels of FGF-basic ($p = 0.05$), IL-1RA ($p = 0.02$), MCP-1 ($p = 0.01$) and a significant increase in plasma MIP-1 β ($p = 0.04$) (Figure 3 and Table 3). All five of these proteins have previously been linked to AD but the extent and strength of these associations is varied. Whilst FGF-basic has been linked to AD processes [52] and MIP-1 β (CCL4) has been identified as a component of some A β 1-42 pathways [53, 54], to our knowledge this is one of the first instances of them being highlighted as potential AD biomarkers. In contrast, several studies have examined expression patterns of MCP-1 (CCL2) in AD with slightly conflicting findings, but generally it appears that it increases in AD CSF and decreases in AD plasma [55-58] which is in accordance with the findings of this study. The IL-1 cytokine family has been widely implicated in AD pathology [59-61] and decreased expression of IL-1RA in CSF has previously been linked to AD [62]. Levels of IL-12 have previously been shown to decrease in CSF from AD patients [63] corroborating the findings of this study. Expression of IL-12 in plasma samples has also been tracked throughout AD progression with an increased concentration being observed during mild stage disease which gradually decreases as the health of the individual declines [51]. However these results were not replicated in this study as we observed comparable levels of IL-12 in plasma from AD patients and healthy individuals.

In addition to the foregoing points, Figures 2 and 3 highlight the difficulties encountered whilst trying to define a singular biomarker indicative of the presence of a complex disease condition such as AD. Even if, as demonstrated by some of the data presented here, a prospective biomarker is shown to be significantly different in the disease condition compared to that in the healthy, there still remains uncertainty as to the accuracy of the clinical decision based on the marker measurement from a patient. This uncertainty has its origins in the overlap between the marker distributions (Figure 1). Only those concentration values falling outside of the overlapping ranges can be considered as an affirmative diagnosis, thus extensive levels of overlap may render the practical application of a respective marker untenable.

One solution to the foregoing marker screening problem involves combination of the measurements of separate molecular markers into a singular analytical protocol to yield a more reliable clinical decision. During this study this approach was investigated through application of established supervised learning techniques to the datasets. Using the C4.5 algorithm, a diagnostic accuracy (in correctly classifying either an AD or Healthy sample) of 87.5% was obtained for the plasma data sets whilst application of the neural network algorithm to the CSF dataset yielded a diagnostic accuracy of 86.7%. It is not appropriate to assign sensitivity or specificity values to the results obtained within this study due to the nature of the analytical methods used which were implemented using a 'leave-one-out' cross validation of the entire healthy and AD cohorts. The values of diagnostic accuracy obtained are promising, particularly for the plasma samples, but need to be validated further using larger sample cohorts. We envisage that this type of multiparametric biomarker testing would be used as a screening tool to inform decisions about further diagnostic requirements e.g. whether a patient should be referred for an MRI scan. When used in conjunction with clinical and neuropsychometric evaluation, the ability to define a healthy individual from a blood test would enable a subset of the population to avoid the need for referral for imaging tests such as MRI [64] which are expensive and tend to be of limited availability outside specialist centres.

Acknowledgements

This work was funded by a Medical Research Council Developmental Pathways Grant, grant number 1001500 (<http://www.mrc.ac.uk/index.htm>). The work of OPTIMA has been supported over 20 years by major grants from Bristol-Myers Squibb, Merck, USA Inc, MRC, Charles Wolfson Charitable Trust. GW was partly funded by the NIHR Biomedical Research

Centre Programme, Oxford. No funding bodies had any role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Disclosure Statement

There are no actual or potential conflicts of interest to disclose.

References

- [1] Ballard C, Gauthier S, Corbett A, Brayne C, Aarsland D and Jones E. Alzheimer's disease. *The Lancet* 2011; 377: 1019-1031.
- [2] Herrmann N, Chau SA, Kircanski I and Lanctot KL. Current and emerging drug treatment options for Alzheimer's disease: a systematic review. *Drugs* 2011; 71: 2031-2065.
- [3] Salomone S, Caraci F, Leggio GM, Fedotova J and Drago F. New pharmacological strategies for treatment of Alzheimer's disease: focus on disease modifying drugs. *Br J Clin Pharmacol* 2012; 73: 504-517.
- [4] Bertram L, Lill CM and Tanzi RE. The genetics of Alzheimer disease: back to the future. *Neuron* 2010; 68: 270-281.
- [5] Harold D, Abraham R, Hollingworth P, Sims R, Gerrish A, Hamshere ML, Pahwa JS, Moskvina V, Dowzell K, Williams A, Jones N, Thomas C, Stretton A, Morgan AR, Lovestone S, Powell J, Proitsi P, Lupton MK, Brayne C, Rubinsztein DC, Gill M, Lawlor B, Lynch A, Morgan K, Brown KS, Passmore PA, Craig D, McGuinness B, Todd S, Holmes C, Mann D, Smith AD, Love S, Kehoe PG, Hardy J, Mead S, Fox N, Rossor M, Collinge J, Maier W, Jessen F, Schurmann B, van den Bussche H, Heuser I, Kornhuber J, Wiltfang J, Dichgans M, Frolich L, Hampel H, Hull M, Rujescu D, Goate AM, Kauwe JS, Cruchaga C, Nowotny P, Morris JC, Mayo K, Sleegers K, Bettens K, Engelborghs S, De Deyn PP, Van Broeckhoven C, Livingston G, Bass NJ, Gurling H, McQuillin A, Gwilliam R, Deloukas P, Al-Chalabi A, Shaw CE, Tsolaki M, Singleton AB, Guerreiro R, Muhleisen TW, Nothen MM, Moebus S, Jockel KH, Klopp N, Wichmann HE, Carrasquillo MM, Pankratz VS, Younkin SG, Holmans PA, O'Donovan M, Owen MJ and Williams J. Genome-wide association study identifies variants at *CLU* and *PICALM* associated with Alzheimer's disease. *Nat Genet* 2009; 41: 1088-1093.
- [6] Hollingworth P, Harold D, Sims R, Gerrish A, Lambert JC, Carrasquillo MM, Abraham R, Hamshere ML, Pahwa JS, Moskvina V, Dowzell K, Jones N, Stretton A, Thomas C, Richards A, Ivanov D, Widdowson C, Chapman J, Lovestone S, Powell J, Proitsi P, Lupton MK, Brayne C, Rubinsztein DC, Gill M, Lawlor B, Lynch A, Brown KS, Passmore PA, Craig D, McGuinness B, Todd S, Holmes C, Mann D, Smith AD, Beaumont H, Warden D, Wilcock G, Love S, Kehoe PG, Hooper NM, Vardy ER, Hardy J, Mead S, Fox NC, Rossor M, Collinge J, Maier W, Jessen F, Ruther E, Schurmann B, Heun R, Kolsch H, van den Bussche H, Heuser I, Kornhuber J, Wiltfang J, Dichgans M, Frolich L, Hampel H, Gallacher J, Hull M, Rujescu D, Giegling I, Goate AM, Kauwe JS, Cruchaga C, Nowotny P, Morris JC, Mayo K, Sleegers K, Bettens K, Engelborghs S, De Deyn PP, Van Broeckhoven C, Livingston

- G, Bass NJ, Gurling H, McQuillin A, Gwilliam R, Deloukas P, Al-Chalabi A, Shaw CE, Tsolaki M, Singleton AB, Guerreiro R, Muhleisen TW, Nothen MM, Moebus S, Jockel KH, Klopp N, Wichmann HE, Pankratz VS, Sando SB, Aasly JO, Barcikowska M, Wszolek ZK, Dickson DW, Graff-Radford NR, Petersen RC, van Duijn CM, Breteler MM, Ikram MA, Destefano AL, Fitzpatrick AL, Lopez O, Launer LJ, Seshadri S, Berr C, Champion D, Epelbaum J, Dartigues JF, Tzourio C, Alperovitch A, Lathrop M, Feulner TM, Friedrich P, Riehle C, Krawczak M, Schreiber S, Mayhaus M, Nicolhaus S, Wagenpfeil S, Steinberg S, Stefansson H, Stefansson K, Snaedal J, Bjornsson S, Jonsson PV, Chouraki V, Genier-Boley B, Hiltunen M, Soininen H, Combarros O, Zelenika D, Delepine M, Bullido MJ, Pasquier F, Mateo I, Frank-Garcia A, Porcellini E, Hanon O, Coto E, Alvarez V, Bosco P, Siciliano G, Mancuso M, Panza F, Solfrizzi V, Nacmias B, Sorbi S, Bossu P, Piccardi P, Arosio B, Annoni G, Seripa D, Pilotto A, Scarpini E, Galimberti D, Brice A, Hannequin D, Licastro F, Jones L, Holmans PA, Jonsson T, Riemenschneider M, Morgan K, Younkin SG, Owen MJ, O'Donovan M, Amouyel P and Williams J. Common variants at ABCA7, MS4A6A/MS4A4E, EPHA1, CD33 and CD2AP are associated with Alzheimer's disease. *Nat Genet* 2011; 43: 429-435.
- [7] Lambert JC, Heath S, Even G, Champion D, Sleegers K, Hiltunen M, Combarros O, Zelenika D, Bullido MJ, Tavernier B, Letenneur L, Bettens K, Berr C, Pasquier F, Fievet N, Barberger-Gateau P, Engelborghs S, De Deyn P, Mateo I, Franck A, Helisalmi S, Porcellini E, Hanon O, de Pancorbo MM, Lendon C, Dufouil C, Jaillard C, Leveillard T, Alvarez V, Bosco P, Mancuso M, Panza F, Nacmias B, Bossu P, Piccardi P, Annoni G, Seripa D, Galimberti D, Hannequin D, Licastro F, Soininen H, Ritchie K, Blanche H, Dartigues JF, Tzourio C, Gut I, Van Broeckhoven C, Alperovitch A, Lathrop M and Amouyel P. Genome-wide association study identifies variants at CLU and CR1 associated with Alzheimer's disease. *Nat Genet* 2009; 41: 1094-1099.
- [8] Naj AC, Jun G, Beecham GW, Wang LS, Vardarajan BN, Buross J, Gallins PJ, Buxbaum JD, Jarvik GP, Crane PK, Larson EB, Bird TD, Boeve BF, Graff-Radford NR, De Jager PL, Evans D, Schneider JA, Carrasquillo MM, Ertekin-Taner N, Younkin SG, Cruchaga C, Kauwe JS, Nowotny P, Kramer P, Hardy J, Huentelman MJ, Myers AJ, Barmada MM, Demirci FY, Baldwin CT, Green RC, Rogava E, George-Hyslop PS, Arnold SE, Barber R, Beach T, Bigio EH, Bowen JD, Boxer A, Burke JR, Cairns NJ, Carlson CS, Carney RM, Carroll SL, Chui HC, Clark DG, Corneveaux J, Cotman CW, Cummings JL, Decarli C, Dekosky ST, Diaz-Arrastia R, Dick M, Dickson DW, Ellis WG, Faber KM, Fallon KB, Farlow MR, Ferris S, Frosch MP, Galasko DR, Ganguli M, Gearing M, Geschwind DH, Ghetti B, Gilbert JR,

- Gilman S, Giordani B, Glass JD, Growdon JH, Hamilton RL, Harrell LE, Head E, Honig LS, Hulette CM, Hyman BT, Jicha GA, Jin LW, Johnson N, Karlawish J, Karydas A, Kaye JA, Kim R, Koo EH, Kowall NW, Lah JJ, Levey AI, Lieberman AP, Lopez OL, Mack WJ, Marson DC, Martiniuk F, Mash DC, Masliah E, McCormick WC, McCurry SM, McDavid AN, McKee AC, Mesulam M, Miller BL, Miller CA, Miller JW, Parisi JE, Perl DP, Peskind E, Petersen RC, Poon WW, Quinn JF, Rajbhandary RA, Raskind M, Reisberg B, Ringman JM, Roberson ED, Rosenberg RN, Sano M, Schneider LS, Seeley W, Shelanski ML, Slifer MA, Smith CD, Sonnen JA, Spina S, Stern RA, Tanzi RE, Trojanowski JQ, Troncoso JC, Van Deerlin VM, Vinters HV, Vonsattel JP, Weintraub S, Welsh-Bohmer KA, Williamson J, Woltjer RL, Cantwell LB, Dombroski BA, Beekly D, Lunetta KL, Martin ER, Kamboh MI, Saykin AJ, Reiman EM, Bennett DA, Morris JC, Montine TJ, Goate AM, Blacker D, Tsuang DW, Hakonarson H, Kukull WA, Foroud TM, Haines JL, Mayeux R, Pericak-Vance MA, Farrer LA and Schellenberg GD. Common variants at MS4A4/MS4A6E, CD2AP, CD33 and EPHA1 are associated with late-onset Alzheimer's disease. *Nat Genet* 2011; 43: 436-441.
- [9] Seshadri S, Fitzpatrick AL, Ikram MA, DeStefano AL, Gudnason V, Boada M, Bis JC, Smith AV, Carassquillo MM, Lambert JC, Harold D, Schrijvers EM, Ramirez-Lorca R, Debette S, Longstreth WT, Jr., Janssens AC, Pankratz VS, Dartigues JF, Hollingworth P, Aspelund T, Hernandez I, Beiser A, Kuller LH, Koudstaal PJ, Dickson DW, Tzourio C, Abraham R, Antunez C, Du Y, Rotter JI, Aulchenko YS, Harris TB, Petersen RC, Berr C, Owen MJ, Lopez-Arrieta J, Varadarajan BN, Becker JT, Rivadeneira F, Nalls MA, Graff-Radford NR, Campion D, Auerbach S, Rice K, Hofman A, Jonsson PV, Schmidt H, Lathrop M, Mosley TH, Au R, Psaty BM, Uitterlinden AG, Farrer LA, Lumley T, Ruiz A, Williams J, Amouyel P, Younkin SG, Wolf PA, Launer LJ, Lopez OL, van Duijn CM and Breteler MM. Genome-wide analysis of genetic loci associated with Alzheimer disease. *JAMA* 2010; 303: 1832-1840.
- [10] Lee SH, Harold D, Nyholt DR, Goddard ME, Zondervan KT, Williams J, Montgomery GW, Wray NR and Visscher PM. Estimation and partitioning of polygenic variation captured by common SNPs for Alzheimer's disease, multiple sclerosis and endometriosis. *Hum Mol Genet* 2013; 22: 832-841.
- [11] Ward A, Crean S, Mercaldi CJ, Collins JM, Boyd D, Cook MN and Arrighi HM. Prevalence of apolipoprotein E4 genotype and homozygotes (APOE e4/e4) among patients diagnosed with Alzheimer's disease: a systematic review and meta-analysis. *Neuroepidemiology* 2012; 38: 1-17.

- [12] Prvulovic D and Hampel H. Amyloid beta (Abeta) and phospho-tau (p-tau) as diagnostic biomarkers in Alzheimer's disease. *Clin Chem Lab Med* 2011; 49: 367-374.
- [13] Ewers M, Mielke MM and Hampel H. Blood-based biomarkers of microvascular pathology in Alzheimer's disease. *Exp Gerontol* 2010; 45: 75-79.
- [14] Hampel H, Burger K, Teipel SJ, Bokde AL, Zetterberg H and Blennow K. Core candidate neurochemical and imaging biomarkers of Alzheimer's disease. *Alzheimers Dement* 2008; 4: 38-48.
- [15] Mattsson N, Zetterberg H, Hansson O, Andreasen N, Parnetti L, Jonsson M, Herukka SK, van der Flier WM, Blankenstein MA, Ewers M, Rich K, Kaiser E, Verbeek M, Tsolaki M, Mulugeta E, Rosen E, Aarsland D, Visser PJ, Schroder J, Marcusson J, de Leon M, Hampel H, Scheltens P, Pirtila T, Wallin A, Jonhagen ME, Minthon L, Winblad B and Blennow K. CSF biomarkers and incipient Alzheimer disease in patients with mild cognitive impairment. *JAMA* 2009; 302: 385-393.
- [16] Schneider P, Hampel H and Buerger K. Biological marker candidates of Alzheimer's disease in blood, plasma, and serum. *CNS Neurosci Ther* 2009; 15: 358-374.
- [17] Buchhave P, Minthon L, Zetterberg H, Wallin AK, Blennow K and Hansson O. Cerebrospinal fluid levels of beta-amyloid 1-42, but not of tau, are fully changed already 5 to 10 years before the onset of Alzheimer dementia. *Arch Gen Psychiatry* 2012; 69: 98-106.
- [18] Vafadar-Isfahani B, Ball G, Coveney C, Lemetre C, Boocock D, Minthon L, Hansson O, Miles AK, Janciauskiene SM, Warden D, Smith AD, Wilcock G, Kalsheker N, Rees R, Matharoo-Ball B and Morgan K. Identification of SPARC-like 1 Protein as Part of a Biomarker Panel for Alzheimer's Disease in Cerebrospinal Fluid. *J Alzheimers Dis* 2012; 28: 625-636.
- [19] Thambisetty M, Simmons A, Velayudhan L, Hye A, Campbell J, Zhang Y, Wahlund LO, Westman E, Kinsey A, Guntert A, Proitsi P, Powell J, Causevic M, Killick R, Lunnon K, Lynham S, Broadstock M, Choudhry F, Howlett DR, Williams RJ, Sharp SI, Mitchelmore C, Tunnard C, Leung R, Foy C, O'Brien D, Breen G, Furney SJ, Ward M, Kloszewska I, Mecocci P, Soininen H, Tsolaki M, Vellas B, Hodges A, Murphy DG, Parkins S, Richardson JC, Resnick SM, Ferrucci L, Wong DF, Zhou Y, Muehlboeck S, Evans A, Francis PT, Spenger C and Lovestone S. Association of plasma clusterin concentration with severity, pathology, and progression in Alzheimer disease. *Arch Gen Psychiatry* 2010; 67: 739-748.
- [20] Williams JH, Wilcock GK, Seeburger J, Dallob A, Laterza O, Potter W and Smith AD. Non-linear relationships of cerebrospinal fluid biomarker levels with cognitive function: an observational study. *Alzheimers Res Ther* 2011; 3: 5.

- [21] Quinlan JR. C4.5: programs for machine learning. Morgan Kaufmann Publishers Inc., 1993.
- [22] John GH and Langley P. Estimating continuous distributions in Bayesian classifiers. Proceedings of the Eleventh conference on Uncertainty in artificial intelligence 1995
- [23] Haykin S. Neural Networks: A Comprehensive Foundation. Prentice Hall PTR, 1998.
- [24] Gutierrez C and Schiff R. HER2: biology, detection, and clinical implications. Arch Pathol Lab Med 2011; 135: 55-62.
- [25] Lim LS and Sherin K. Screening for prostate cancer in U.S. men ACPM position statement on preventive practice. Am J Prev Med 2008; 34: 164-170.
- [26] de Bono JS and Ashworth A. Translating cancer research into targeted therapeutics. Nature 2010; 467: 543-549.
- [27] Richens JL, Morgan K and O'Shea P. Reverse engineering of Alzheimer's disease based on biomarker pathways analysis. Neurobiol Aging In press;
- [28] Anderson NL and Anderson NG. The human plasma proteome: history, character, and diagnostic prospects. Mol Cell Proteomics 2002; 1: 845-867.
- [29] Yuan X, Russell T, Wood G and Desiderio DM. Analysis of the human lumbar cerebrospinal fluid proteome. Electrophoresis 2002; 23: 1185-1196.
- [30] Richens JL, Lunt EA, Sanger D, McKenzie G and O'Shea P. Avoiding nonspecific interactions in studies of the plasma proteome: practical solutions to prevention of nonspecific interactions for label-free detection of low-abundance plasma proteins. J Proteome Res 2009; 8: 5103-5110.
- [31] Chalbot S, Zetterberg H, Blennow K, Fladby T, Andreassen N, Grundke-Iqbal I and Iqbal K. Blood-cerebrospinal fluid barrier permeability in Alzheimer's disease. J Alzheimers Dis 2011; 25: 505-515.
- [32] Morgan K. The three new pathways leading to Alzheimer's disease. Neuropathol Appl Neurobiol 2011; 37: 353-357.
- [33] Jones L, Holmans PA, Hamshere ML, Harold D, Moskvina V, Ivanov D, Pocklington A, Abraham R, Hollingworth P, Sims R, Gerrish A, Pahwa JS, Jones N, Stretton A, Morgan AR, Lovestone S, Powell J, Proitsi P, Lupton MK, Brayne C, Rubinsztein DC, Gill M, Lawlor B, Lynch A, Morgan K, Brown KS, Passmore PA, Craig D, McGuinness B, Todd S, Holmes C, Mann D, Smith AD, Love S, Kehoe PG, Mead S, Fox N, Rossor M, Collinge J, Maier W, Jessen F, Schurmann B, Heun R, Kolsch H, van den Bussche H, Heuser I, Peters O, Kornhuber J, Wiltfang J, Dichgans M, Frolich L, Hampel H, Hull M, Rujescu D, Goate AM, Kauwe JS, Cruchaga C, Nowotny P, Morris JC, Mayo K, Livingston G, Bass NJ, Gurling H, McQuillin A, Gwilliam R, Deloukas P, Al-Chalabi A, Shaw CE, Singleton AB, Guerreiro R, Muhleisen TW, Nothen MM, Moebus S, Jockel KH, Klopp N, Wichmann HE, Ruther

- E, Carrasquillo MM, Pankratz VS, Younkin SG, Hardy J, O'Donovan MC, Owen MJ and Williams J. Genetic evidence implicates the immune system and cholesterol metabolism in the aetiology of Alzheimer's disease. *PLoS One* 2010; 5: e13950.
- [34] Ray S, Britschgi M, Herbert C, Takeda-Uchimura Y, Boxer A, Blennow K, Friedman LF, Galasko DR, Jutel M, Karydas A, Kaye JA, Leszek J, Miller BL, Minthon L, Quinn JF, Rabinovici GD, Robinson WH, Sabbagh MN, So YT, Sparks DL, Tabaton M, Tinklenberg J, Yesavage JA, Tibshirani R and Wyss-Coray T. Classification and prediction of clinical Alzheimer's diagnosis based on plasma signaling proteins. *Nat Med* 2007; 13: 1359-1362.
- [35] Fu BS, Liu W, Zhang JW, Zhang T, Li H and Chen GH. [Serum proteomic analysis on metastasis-associated proteins of hepatocellular carcinoma]. *Nan Fang Yi Ke Da Xue Xue Bao* 2009; 29: 1775-1778.
- [36] Lee KS, Chung JH, Lee KH, Shin MJ, Oh BH and Hong CH. Bioplex analysis of plasma cytokines in Alzheimer's disease and mild cognitive impairment. *Immunol Lett* 2008; 121: 105-109.
- [37] Hu H, Zhang H, Ge W, Liu X, Loera S, Chu P, Chen H, Peng J, Zhou L, Yu S, Yuan Y, Zhang S, Lai L, Yen Y and Zheng S. Secreted Protein Acidic and Rich in Cysteines-like 1 Suppresses Aggressiveness and Predicts Better Survival in Colorectal Cancers. *Clin Cancer Res* 2012; 18: 5438-5448.
- [38] Zhang H, Widegren E, Wang DW and Sun XF. SPARCL1: a potential molecule associated with tumor diagnosis, progression and prognosis of colorectal cancer. *Tumour Biol* 2011; 32: 1225-1231.
- [39] Yu SJ, Yu JK, Ge WT, Hu HG, Yuan Y and Zheng S. SPARCL1, Shp2, MSH2, E-cadherin, p53, ADCY-2 and MAPK are prognosis-related in colorectal cancer. *World J Gastroenterol* 2011; 17: 2028-2036.
- [40] Hurley PJ, Marchionni L, Simons BW, Ross AE, Peskoe SB, Miller RM, Erho N, Vergara IA, Ghadessi M, Huang Z, Gurel B, Park BH, Davicioni E, Jenkins RB, Platz EA, Berman DM and Schaeffer EM. Secreted protein, acidic and rich in cysteine-like 1 (SPARCL1) is down regulated in aggressive prostate cancers and is prognostic for poor clinical outcome. *Proc Natl Acad Sci U S A* 2012; 109: 14977-14982.
- [41] Esposito I, Kayed H, Keleg S, Giese T, Sage EH, Schirmacher P, Friess H and Kleeff J. Tumor-suppressor function of SPARC-like protein 1/Hevin in pancreatic cancer. *Neoplasia* 2007; 9: 8-17.
- [42] Li P, Qian J, Yu G, Chen Y, Liu K, Li J and Wang J. Down-regulated SPARCL1 is associated with clinical significance in human gastric cancer. *J Surg Oncol* 2012; 105: 31-37.

- [43] Lau CP, Poon RT, Cheung ST, Yu WC and Fan ST. SPARC and Hevin expression correlate with tumour angiogenesis in hepatocellular carcinoma. *J Pathol* 2006; 210: 459-468.
- [44] Mencalha AL, Levinsphul A, Deterling LC, Pizzatti L and Abdelhay E. SPARC-like1 mRNA is overexpressed in human uterine leiomyoma. *Mol Med Report* 2008; 1: 571-574.
- [45] Hammack BN, Fung KY, Hunsucker SW, Duncan MW, Burgoon MP, Owens GP and Gilden DH. Proteomic analysis of multiple sclerosis cerebrospinal fluid. *Mult Scler* 2004; 10: 245-260.
- [46] Yin GN, Lee HW, Cho JY and Suk K. Neuronal pentraxin receptor in cerebrospinal fluid as a potential biomarker for neurodegenerative diseases. *Brain Res* 2009; 1265: 158-170.
- [47] Carmeliet P and De Strooper B. Alzheimer's disease: A breach in the blood-brain barrier. *Nature* 2012; 485: 451-452.
- [48] Lee KS, Chung JH, Choi TK, Suh SY, Oh BH and Hong CH. Peripheral cytokines and chemokines in Alzheimer's disease. *Dement Geriatr Cogn Disord* 2009; 28: 281-287.
- [49] Hu WT, Chen-Plotkin A, Arnold SE, Grossman M, Clark CM, Shaw LM, Pickering E, Kuhn M, Chen Y, McCluskey L, Elman L, Karlawish J, Hurtig HI, Siderowf A, Lee VM, Soares H and Trojanowski JQ. Novel CSF biomarkers for Alzheimer's disease and mild cognitive impairment. *Acta Neuropathol* 2010; 119: 669-678.
- [50] Choi C, Jeong JH, Jang JS, Choi K, Lee J, Kwon J, Choi KG, Lee JS and Kang SW. Multiplex analysis of cytokines in the serum and cerebrospinal fluid of patients with Alzheimer's disease by color-coded bead technology. *J Clin Neurol* 2008; 4: 84-88.
- [51] Motta M, Imbesi R, Di Rosa M, Stivala F and Malaguarnera L. Altered plasma cytokine levels in Alzheimer's disease: correlation with the disease progression. *Immunol Lett* 2007; 114: 46-51.
- [52] Stopa EG, Gonzalez AM, Chorsky R, Corona RJ, Alvarez J, Bird ED and Baird A. Basic fibroblast growth factor in Alzheimer's disease. *Biochem Biophys Res Commun* 1990; 171: 690-696.
- [53] Ito S, Sawada M, Haneda M, Ishida Y and Isobe K. Amyloid-beta peptides induce several chemokine mRNA expressions in the primary microglia and Ra2 cell line via the PI3K/Akt and/or ERK pathway. *Neurosci Res* 2006; 56: 294-299.
- [54] Smits HA, Rijmsmus A, van Loon JH, Wat JW, Verhoef J, Boven LA and Nottet HS. Amyloid-beta-induced chemokine production in primary human macrophages and astrocytes. *J Neuroimmunol* 2002; 127: 160-168.

- [55] Conduetier Gg, Blondeau N, Guyon A, Nahon J-L and RovÃˆre C. The role of monocyte chemoattractant protein MCP1/CCL2 in neuroinflammatory diseases. *Journal of Neuroimmunology* 2010; 224: 93-100.
- [56] Galimberti D, Schoonenboom N, Scheltens P, Fenoglio C, Bouwman F, Venturelli E, Guidi I, Blankenstein MA, Bresolin N and Scarpini E. Intrathecal chemokine synthesis in mild cognitive impairment and Alzheimer disease. *Arch Neurol* 2006; 63: 538-543.
- [57] Galimberti D, Fenoglio C, Lovati C, Venturelli E, Guidi I, Corra B, Scalabrini D, Clerici F, Mariani C, Bresolin N and Scarpini E. Serum MCP-1 levels are increased in mild cognitive impairment and mild Alzheimer's disease. *Neurobiol Aging* 2006; 27: 1763-1768.
- [58] Sun YX, Minthon L, Wallmark A, Warkentin S, Blennow K and Janciauskiene S. Inflammatory markers in matched plasma and cerebrospinal fluid from patients with Alzheimer's disease. *Dement Geriatr Cogn Disord* 2003; 16: 136-144.
- [59] Payao SL, Goncalves GM, de Labio RW, Horiguchi L, Mizumoto I, Rasmussen LT, de Souza Pinhel MA, Silva Souza DR, Bechara MD, Chen E, Mazzotti DR, Ferreira Bertolucci PH and Cardoso Smith Mde A. Association of interleukin 1beta polymorphisms and haplotypes with Alzheimer's disease. *J Neuroimmunol* 2012; 247: 59-62.
- [60] Kitazawa M, Cheng D, Tsukamoto MR, Koike MA, Wes PD, Vasilevko V, Cribbs DH and LaFerla FM. Blocking IL-1 signaling rescues cognition, attenuates tau pathology, and restores neuronal beta-catenin pathway function in an Alzheimer's disease model. *J Immunol* 2011; 187: 6539-6549.
- [61] Forlenza OV, Diniz BS, Talib LL, Mendonca VA, Ojopi EB, Gattaz WF and Teixeira AL. Increased serum IL-1beta level in Alzheimer's disease and mild cognitive impairment. *Dement Geriatr Cogn Disord* 2009; 28: 507-512.
- [62] Tarkowski E, Liljeroth AM, Nilsson A, Minthon L and Blennow K. Decreased levels of intrathecal interleukin 1 receptor antagonist in Alzheimer's disease. *Dement Geriatr Cogn Disord* 2001; 12: 314-317.
- [63] Rentzos M, Paraskevas GP, Kapaki E, Nikolaou C, Zoga M, Rombos A, Tsoutsou A and Vassilopoulos DD. Interleukin-12 is reduced in cerebrospinal fluid of patients with Alzheimer's disease and frontotemporal dementia. *J Neurol Sci* 2006; 249: 110-114.
- [64] Hampel H, Frank R, Broich K, Teipel SJ, Katz RG, Hardy J, Herholz K, Bokde AL, Jessen F, Hoessler YC, Sanhai WR, Zetterberg H, Woodcock J and Blennow K. Biomarkers for Alzheimer's disease: academic, industry and regulatory perspectives. *Nat Rev Drug Discov* 2010; 9: 560-574.

Figure Legends

Figure 1 Determining Population Distribution Overlaps The overlap (grey) in population distributions of a healthy (black line) and disease (dashed line) cohort for a single disease biomarker.

Figure 2 Immunoassays of Biomarker Panel Components. Quantification of components of the biomarker panel in AD patients and healthy individuals in (A) CSF and (B) plasma samples. Samples were interrogated using either Luminex or ELISA immunoassays as described in the Methods section.

Figure 3 Inflammatory Proteins in AD. Inflammatory proteins were examined in blood plasma samples from healthy and AD cohorts. Samples were interrogated using the Luminex human cytokine 30-plex panel (Biosource). Data was collected using a Bio-Plex 200 system and analysed with associated software (Bio-Rad). Only those proteins demonstrated to differ significantly between the two patient cohorts are displayed: A) FGF-basic; B) IL-1RA; C) MCP-1 and D) MIP-1 β .

Table 1 Immunoassay Reagents. Details of the antibody and protein reagents used in the development of Luminex assays for this study.

Table 2 Protein Biomarkers in CSF and Plasma. Expression levels of the biomarker panel components were quantified in CSF and plasma samples using immunoassays. Variations between the cohorts were evaluated using the Mann-Whitney test. Spearman correlation coefficients were determined for protein values between the two sample types. A p-value 0.05 was considered statistically significant

Table 3 Inflammatory Proteins in CSF and Plasma. Expression levels of inflammatory proteins were quantified in CSF and plasma samples using the Luminex human cytokine 30-plex panel (Biosource). Variations between the cohorts were evaluated using the Mann-Whitney test. A p-value 0.05 was considered statistically significant.

Table 4 Correlations Between Proteins Concentrations in CSF and Plasma. Spearman correlation coefficients were determined for all protein pairings in CSF and plasma. A p-value 0.05 was considered statistically significant. Target markers showing significant correlations either in the healthy or disease state are shown.

Table 5 The Power of Individual Proteins Within the Biomarkers Panels. Contribution of each individual marker to the overall power of the biomarker panels shown in Figure 4. Values are presented as the percentage of a cohort that can be accurately identified using that marker. Each marker is ranked according to its contribution to the strength of the biomarker panel.

Figure 1





Figure 2

(A) CSF



(B) Plasma




Figure 3




Table 1

Target	Capture Antibody		Recombinant Protein		Detection Antibody	
	Supplier	Product	Supplier	Product	Supplier	Product
ApoE	Abcam	ab1907	Merck	178468	Abcam	ab24274
Clusterin	AbD Serotec	MCA2612	R&D Systems	2937-HS	R&D Systems	BAF2937
Fibrinogen α -chain	Abcam	ab19079	Thermo Scientific	RP-43142	Abcam	ab34546
Tau	Abcam	ab80579	Abcam	ab72489	Thermo Scientific	MN1000 B
Tetranectin	Abcam	ab51883	R&D Systems	5170-CL	R&D Systems	BAF5170

Table 2

Target Marker	Mean concentrations (\pm SD)				P-value (change due to AD)		Plasma/CSF correlation	
	Plasma		CSF		Plasma	CSF	Healthy	AD
	Healthy	AD	Healthy	AD				
A β 40 (pmol/L)	9.36 \pm 1.65	9.00 \pm 1.10	161 \pm 76.6	175 \pm 53.4	0.8968	0.5787	0.93	0.66
A β 42 (pmol/L)	4.91 \pm 0.90	3.54 \pm 0.41	43.8 \pm 21.6	25.5 \pm 10.1	0.0008*	0.0493*	0.61	0.14
ApoE (μ g/ml)	5.86 \pm 1.28	4.98 \pm 0.99	10.08 \pm 2.88	10.05 \pm 1.62	0.1718	0.9809	0.02	0.54
Clusterin (μ g/ml)	6.12 \pm 1.31	5.93 \pm 1.31	7.14 \pm 1.02	7.38 \pm 1.19	0.9046	0.5490	0.14	0.18
Fibrinogen α -chain (μ g/ml)	348 \pm 401	323 \pm 354	8.77 \pm 13.08	12.2 \pm 8.36	0.7191	0.0887	0.30	0.87
Keratin 9 (pg/ml)	422 \pm 219	625 \pm 379	0.0 \pm 0.0	3.75 \pm 5.9	0.2721	-	-	0.24
SPARCL1 (ng/ml)	1957 \pm 1136	1271 \pm 557	1286 \pm 620	815 \pm 245	0.0887	0.0349*	0.05	0.89
Tau (ng/ml)	75.1 \pm 14.3	63.4 \pm 11.9	61.5 \pm 17.6	63.9 \pm 11.7	0.0466*	0.5490	\leq 0.0001	0.47
Tetranectin (μ g/ml)	29.9 \pm 14.3	21.4 \pm 5.4	6.59 \pm 3.09	6.08 \pm 2.18	0.3498	0.9046	\leq 0.0001	0.58

Table 3

Marker	Mean concentrations (\pm SD)				P-value (change due to AD)	
	Plasma		CSF		Plasma	CSF
	Healthy	AD	Healthy	AD		
EGF	8.08 \pm 5.60	9.54 \pm 8.04	5.24 \pm 6.91	3.93 \pm 8.68	0.82	0.07
Eotaxin	54.7 \pm 44.1	39.8 \pm 19.0	25.2 \pm 26.2	10.1 \pm 21.1	0.47	0.26
FGF-basic	2.30 \pm 3.44	0.124 \pm 0.392	3.80 \pm 3.00	2.62 \pm 1.11	0.05*	0.29
G-CSF	10.6 \pm 10.4	17.8 \pm 14.1	11.8 \pm 24.8	2.97 \pm 4.28	0.42	0.32
HGF	111 \pm 38.2	84.9 \pm 14.5	89.6 \pm 30.2	77.7 \pm 40.9	0.10	0.18
IFN- α	256 \pm 163	165 \pm 116	93.3 \pm 143	59.4 \pm 111	0.27	0.73
IL-1RA	41.0 \pm 53.5	2.98 \pm 9.43	7.82 \pm 22.5	1.65 \pm 3.53	0.02*	0.95
IL-2R	143 \pm 132	64.9 \pm 80.2	40.6 \pm 50.7	38.1 \pm 78.4	0.14	0.57
IL-6	0.293 \pm 0.927	0.647 \pm 2.05	0.833 \pm 1.52	0.266 \pm 0.841	1.00	0.11
IL-8	3.22 \pm 5.82	0.576 \pm 1.82	11.6 \pm 11.6	6.99 \pm 8.45	0.23	0.58
IL-10	34.8 \pm 110	3.68 \pm 7.76	0.0 \pm 0.0	0.084 \pm 0.266	0.67	-
IL-12	116 \pm 99.2	109 \pm 52.7	52.1 \pm 48.8	25.6 \pm 52.0	0.97	0.04*
IP-10	13.8 \pm 9.23	14.0 \pm 6.83	8.32 \pm 5.22	6.46 \pm 6.00	0.91	0.36
MCP-1	235 \pm 102	150 \pm 62.9	237 \pm 54.4	240 \pm 96.7	0.01*	0.72
MIG	45.5 \pm 34.5	39.3 \pm 24.1	25.3 \pm 35.0	11.4 \pm 19.5	0.45	0.45
MIP-1 α	13.4 \pm 12.2	6.23 \pm 5.71	9.70 \pm 11.9	1.81 \pm 3.94	0.15	0.15
MIP-1 β	62.2 \pm 48.0	152 \pm 88.1	47.7 \pm 77.1	49.7 \pm 131	0.04*	0.36
VEGF	0.477 \pm 0.485	0.205 \pm 0.261	0.730 \pm 0.513	0.533 \pm 0.276	0.21	0.21

Table 4

CSF Correlations					Plasma Correlations				
Target 1	Target 2	Healthy	AD	Effect due to AD	Target 1	Target 2	Healthy	AD	Effect due to AD
A β 40	A β 42	0.0368*	0.5364	Disrupted correlation	A β 40	A β 42	0.0167*	0.9768	Disrupted correlation
A β 40	ApoE	0.4279	0.0154*	Correlation	A β 40	Clusterin	0.0333*	0.7033	Disrupted correlation
A β 40	SPARCL1	0.0279*	0.7033	Disrupted correlation	A β 40	Fibrinogen	0.3556	0.0368*	Correlation
A β 40	Tau	0.1710	0.0458*	Correlation	A β 40	IL-12	0.0167*	0.0962	Disrupted correlation
A β 40	Tetranectin	0.0831	0.0072**	Correlation	A β 40	Tau	0.0333*	0.2675	Disrupted correlation
A β 42	MCP-1	0.4976	0.0072**	Correlation	A β 42	Clusterin	0.0072**	0.0458*	Weaker correlation
ApoE	MCP-1	0.0067**	0.0831	Disrupted correlation	A β 42	Tau	0.0154*	0.1323	Disrupted correlation
ApoE	Tau	0.0368*	0.0458*	No change in correlation	ApoE	Fibrinogen	0.7033	0.0368*	Correlation
ApoE	Tetranectin	0.0458*	0.0368*	No change in correlation	ApoE	IL-12	0.0154*	0.7930	Disrupted correlation
Clusterin	FGF-basic	0.5560	0.0107*	Correlation	ApoE	Tau	0.1710	0.0046**	Correlation
Clusterin	MCP-1	0.0341*	0.9349	Disrupted correlation	ApoE	Tetranectin	0.4618	0.0368*	Correlation
FGF-basic	MIP-1 beta	0.0480*	0.3268	Disrupted correlation	Clusterin	Tau	0.0046**	0.2162	Disrupted correlation
MCP-1	Tau	0.0123*	0.3599	Disrupted correlation	Fibrinogen	Tau	0.9768	0.0154*	Correlation
MCP-1	Tetranectin	0.0480*	0.2992	Disrupted correlation	Fibrinogen	Tetranectin	0.0831	0.0107*	Correlation
Tau	Tetranectin	0.0022**	0.0107*	Weaker correlation	IL-12	Tau	0.0107*	0.7520	Disrupted correlation
					MCP-1	Tetranectin	0.0458*	0.3268	Disrupted correlation

	Tau	Tetranectin	0.5364	0.0072**	Correlation
--	-----	-------------	--------	----------	-------------

Table 5

Marker	Healthy CSF		AD CSF		Healthy Plasma		AD Plasma	
	Rank	Power(%)	Rank	Power(%)	Rank	Power(%)	Rank	Power(%)
A β 40	4	40.58	21	0	14	5.13	12	0.002
A β 42	5	29.68	17	0	3	56.67	10	0.07
ApoE	11	10.48	6	4.50	20	0.002	4	4.52
Clusterin	17	0.57	8	2.71	18	0.03	8	1.97
EGF	21	0.02	9	0.82	22	0.0003	5	3.21
Eotaxin	14	4.00	14	0.0006	7	19.08	18	0
FGF-basic	6	26.37	19	0	1	82.13	15	0
Fibrinogen	7	18.41	18	0	21	0.002	7	2.41
G-CSF	2	52.30	22	0	23	0.0001	3	5.02
HGF	22	0.013	10	0.62	5	35.51	19	0
IFN-a	16	2.45	13	0.002	13	6.19	20	0
IL-12	19	0.46	12	0.06	12	7.33	23	0
IL-1RA	1	66.43	20	0	2	73.54	16	0
IL-2R	23	0.0003	7	3.54	10	12.74	21	0
IL-6	8	13.92	23	0	-	-	-	-
IL-8	13	4.34	15	0.0002	4	44.19	22	0
IP-10	20	0.10	11	0.21	17	1.37	11	0.003
Keratin 9	24	0	1	98.76	24	0	2	7.96
MCP-1	24	0	4	8.57	9	15.56	17	0
MIG	9	13.14	24	0	16	3.06	13	0.0003
MIP-1a	3	46.84	24	0	6	23.96	24	0
MIP-1B	24	0	5	6.23	24	0	1	28.05
SPARCL1	12	6.66	16	0	11	7.68	14	0
Tau	15	3.92	2	15.68	19	0.02	6	2.48
Tetranectin	18	0.56	3	14.95	15	3.86	9	1.49
VEGF	10	11.85	24	0	8	17.48	24	0