



The University of
Nottingham

UNITED KINGDOM • CHINA • MALAYSIA

Fiset, Matthew and Kasprzyk, Alexander M. (2008) A note on palindromic δ -vectors for certain rational polytopes. *Electronic Journal of Combinatorics*, 15 . N18/1-N18/4. ISSN 1077-8926

Access from the University of Nottingham repository:

<http://eprints.nottingham.ac.uk/30719/1/893-953-1-PB.pdf>

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see:
http://eprints.nottingham.ac.uk/end_user_agreement.pdf

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

A note on palindromic δ -vectors for certain rational polytopes

Matthew H. J. Fiset and Alexander M. Kasprzyk*

Department of Mathematics and Statistics
University of New Brunswick
Fredericton, NB, Canada
u0a35@unb.ca, kasprzyk@unb.ca

Submitted: May 19, 2008; Accepted: Jun 1, 2008; Published: Jun 6, 2008
Mathematics Subject Classifications: 05A15, 11H06

Abstract

Let P be a convex polytope containing the origin, whose dual is a lattice polytope. Hibi's Palindromic Theorem tells us that if P is also a lattice polytope then the Ehrhart δ -vector of P is palindromic. Perhaps less well-known is that a similar result holds when P is rational. We present an elementary lattice-point proof of this fact.

1 Introduction

A *rational polytope* $P \subset \mathbb{R}^n$ is the convex hull of finitely many points in \mathbb{Q}^n . We shall assume that P is of maximum dimension, so that $\dim P = n$. Throughout let k denote the smallest positive integer for which the dilation kP of P is a *lattice polytope* (i.e. the vertices of kP lie in \mathbb{Z}^n).

A *quasi-polynomial* is a function defined on \mathbb{Z} of the form:

$$q(m) = c_n(m)m^n + c_{n-1}(m)m^{n-1} + \dots + c_0(m),$$

where the c_i are periodic coefficient functions in m . It is known ([Ehr62]) that for a rational polytope P , the number of lattice points in mP , where $m \in \mathbb{Z}_{\geq 0}$, is given by a quasi-polynomial of degree $n = \dim P$ called the *Ehrhart quasi-polynomial*; we denote this by $L_P(m) := |mP \cap \mathbb{Z}^n|$. The minimum period common to the cyclic coefficients c_i of L_P divides k (for further details see [BSW08]).

*The first author was funded by an NSERC USRA grant. The second author is funded by an ACEnet research fellowship.

Stanley proved in [Sta80] that the generating function for L_P can be written as a rational function:

$$\text{Ehr}_P(t) := \sum_{m \geq 0} L_P(m)t^m = \frac{\delta_0 + \delta_1 t + \dots + \delta_{k(n+1)-1} t^{k(n+1)-1}}{(1-t)^{n+1}},$$

whose coefficients δ_i are non-negative. For an elementary proof of this and other relevant results, see [BS07] and [BR07]. We call $(\delta_0, \delta_1, \dots, \delta_{k(n+1)-1})$ the (*Ehrhart*) δ -vector of P .

The *dual polyhedron* of P is given by $P^\vee := \{u \in \mathbb{R}^n \mid \langle u, v \rangle \leq 1 \text{ for all } v \in P\}$. If the origin lies in the interior of P then P^\vee is a rational polytope containing the origin, and $P = (P^\vee)^\vee$. We restrict our attention to those P containing the origin for which P^\vee is a lattice polytope.

We give an elementary lattice-point proof that, with the above restriction, the δ -vector is palindromic (i.e. $\delta_i = \delta_{k(n+1)-1-i}$). When P is *reflexive*, meaning that P is also a lattice polytope (equivalently, $k = 1$), this result is known as *Hibi's Palindromic Theorem* [Hib91]. It can be regarded as a consequence of a theorem of Stanley's concerning the more general theory of Gorenstein rings; see [Sta78].

2 The main result

Let P be a rational polytope and consider the Ehrhart quasi-polynomial L_P . There exist k polynomials $L_{P,r}$ of degree n in l such that when $m = lk + r$ (where $l, r \in \mathbb{Z}_{\geq 0}$ and $0 \leq r < k$) we have that $L_P(m) = L_{P,r}(l)$. The generating function for each $L_{P,r}$ is given by:

$$\text{Ehr}_{P,r}(t) := \sum_{l \geq 0} L_{P,r}(l)t^l = \frac{\delta_{0,r} + \delta_{1,r}t + \dots + \delta_{n,r}t^n}{(1-t)^{n+1}}, \quad (2.1)$$

for some $\delta_{i,r} \in \mathbb{Z}$.

Theorem 2.1. *Let P be a rational n -tope containing the origin, whose dual P^\vee is a lattice polytope. Let k be the smallest positive integer such that kP is a lattice polytope. Then:*

$$\delta_{i,r} = \delta_{n-i, k-r-1}.$$

Proof. By Ehrhart–Macdonald reciprocity ([Ehr67, Mac71]) we have that:

$$L_P(-lk - r) = (-1)^n L_{P^\circ}(lk + r),$$

where L_{P° enumerates lattice points in the strict interior of dilations of P . The left-hand side equals $L_P(-(l+1)k + (k-r)) = L_{P, k-r}(-(l+1))$. We shall show that the right-hand side is equal to $(-1)^n L_P(lk + r - 1) = (-1)^n L_{P, r-1}(l)$.

Let $H_u := \{v \in \mathbb{R}^n \mid \langle u, v \rangle = 1\}$ be a bounding hyperplane of P , where $u \in \text{vert } P^\vee$. By assumption, $u \in \mathbb{Z}^n$ and so the lattice points in \mathbb{Z}^n lie at integer heights relative to

H_u ; i.e. given $u' \in \mathbb{Z}^n$ there exists some $c \in \mathbb{Z}$ such that $u' \in \{v \in \mathbb{R}^n \mid \langle u, v \rangle = c\}$. In particular, there do not exist lattice points at non-integral heights. Since:

$$P = \bigcap_{u \in \text{vert } P^\vee} H_u^-,$$

where H_u^- is the half-space defined by H_u and the origin, we see that $(mP^\circ) \cap \mathbb{Z}^n = ((m-1)P) \cap \mathbb{Z}^n$. This gives us the desired equality.

We have that $L_{P, k-r}(-l+1) = (-1)^n L_{P, r-1}(l)$. By considering the expansion of (2.1) we obtain:

$$\begin{aligned} \sum_{i=0}^n \delta_{i, k-r} \binom{-(l+1) + n - i}{n} &= L_{P, k-r}(-l+1) \\ &= (-1)^n L_{P, r-1}(l) = (-1)^n \sum_{i=0}^n \delta_{i, r-1} \binom{l + n - i}{n}. \end{aligned}$$

But $\binom{-(l+1)+n-i}{n} = (-1)^n \binom{l+n-i}{n}$, and since $\binom{l}{n}, \binom{l+1}{n}, \dots, \binom{l+n}{n}$ form a basis for the vector space of polynomials in l of degree at most n , we have that $\delta_{i, k-r} = \delta_{n-i, r-1}$. \square

Corollary 2.2. *The δ -vector of P is palindromic.*

Proof. This is immediate once we observe that:

$$\text{Ehr}_P(t) = \text{Ehr}_{P,0}(t^k) + t \text{Ehr}_{P,1}(t^k) + \dots + t^{k-1} \text{Ehr}_{P, k-1}(t^k).$$

\square

3 Concluding remarks

The crucial observation in the proof of Theorem 2.1 is that $(mP^\circ) \cap \mathbb{Z}^n = ((m-1)P) \cap \mathbb{Z}^n$. In fact, a consequence of Ehrhart–Macdonald reciprocity and a result of Hibi [Hib92] tells us that this property holds if and only if P^\vee is a lattice polytope. Hence rational convex polytopes whose duals are lattice polytopes are characterised by having palindromic δ -vectors. This can also be derived from Stanley’s work [Sta78] on Gorenstein rings.

References

- [BR07] Matthias Beck and Sinai Robins, *Computing the continuous discretely*, Undergraduate Texts in Mathematics, Springer, New York, 2007, Integer-point enumeration in polyhedra.
- [BS07] Matthias Beck and Frank Sottile, *Irrational proofs for three theorems of Stanley*, European J. Combin. **28** (2007), no. 1, 403–409.

- [BSW08] Matthias Beck, Steven V. Sam, and Kevin M. Woods, *Maximal periods of (Ehrhart) quasi-polynomials*, J. Combin. Theory Ser. A **115** (2008), no. 3, 517–525.
- [Ehr62] Eugène Ehrhart, *Sur les polyèdres homothétiques bordés à n dimensions*, C. R. Acad. Sci. Paris **254** (1962), 988–990.
- [Ehr67] ———, *Sur un problème de géométrie diophantienne linéaire. II. Systèmes diophantiens linéaires*, J. Reine Angew. Math. **227** (1967), 25–49.
- [Hib91] Takayuki Hibi, *Ehrhart polynomials of convex polytopes, h -vectors of simplicial complexes, and nonsingular projective toric varieties*, Discrete and computational geometry (New Brunswick, NJ, 1989/1990), DIMACS Ser. Discrete Math. Theoret. Comput. Sci., vol. 6, Amer. Math. Soc., Providence, RI, 1991, pp. 165–177.
- [Hib92] ———, *Dual polytopes of rational convex polytopes*, Combinatorica **12** (1992), no. 2, 237–240.
- [Mac71] I. G. Macdonald, *Polynomials associated with finite cell-complexes*, J. London Math. Soc. (2) **4** (1971), 181–192.
- [Sta78] Richard P. Stanley, *Hilbert functions of graded algebras*, Advances in Math. **28** (1978), no. 1, 57–83.
- [Sta80] ———, *Decompositions of rational convex polytopes*, Ann. Discrete Math. **6** (1980), 333–342, Combinatorial mathematics, optimal designs and their applications (Proc. Sympos. Combin. Math. and Optimal Design, Colorado State Univ., Fort Collins, Colo., 1978).