

Reducing the impact of physical inactivity: evidence to support the case for targeting people with chronic mental and physical conditions

Dr Everson-Hock, E.S., Research Fellow, School of Health and Related Research, University of Sheffield^{1,*}

Dr Green, M.A., Research Associate, School of Health and Related Research, University of Sheffield¹

Prof Goyder, E.C., Professor of Public Health, School of Health and Related Research, University of Sheffield¹

Dr Copeland, R.J., Reader in Physical Activity and Health, Centre for Sport and Exercise Science, Sheffield Hallam University and National Centre for Sport and Exercise Medicine^{2,6}

Dr Till, S.H., Consultant Rheumatologist and Reader in Sport and Exercise Medicine, Sheffield Teaching Hospital NHS Trust and Sheffield Hallam University^{3,4}

Dr Heller, B., Principal Research Fellow, Centre for Sports Engineering Research, Sheffield Hallam University⁴

Dr Hart, O., General Practice Partner, Sloan Medical Centre⁵

¹School of Health and Related Research, University of Sheffield, Regent Court, 30 Regent Street, Sheffield, S1 4DA, UK

²Centre for Sport and Exercise Science, Sheffield Hallam University, Collegiate Hall, Collegiate Campus, Sheffield, S10 2BP

³Sheffield Teaching Hospital NHS Trust, Royal Hallamshire Hospital, Glossop Road, Sheffield, S10 2JF

⁴Centre for Sports Engineering Research, Sheffield Hallam University, Collegiate Hall, Collegiate Campus, Sheffield, S10 2BP

⁵Sloan Medical Centre, 2 Little London Road, Sheffield, S8 0YH

⁶National Centre for Sport and Exercise Medicine, Sheffield

*Corresponding author. E-mail: e.everson-hock@sheffield.ac.uk

Running head: Reducing the impact of physical inactivity

Keywords: physical activity, physical inactivity, chronic disease

Word count (excluding references): 2974

Date of submission: December 2014

Abstract

Background

Recent evidence suggests that small increases in the physical activity of those considered least active can have a bigger health impact than raising levels of those already achieving or close to achieving recommendations. Profiling the characteristics of those who are least active allows for appropriate targeting of interventions. This study therefore examined the characteristics of people in the lowest physical activity bracket.

Methods

Data were taken from the Collaboration for Leadership in Applied Health Research and Care (CLAHRC) funded 'South Yorkshire Cohort', a longitudinal observational dataset of residents of South Yorkshire, England. Five separate outcomes based on a shortened version of the GPPAQ were used to represent the lowest levels of physical activity. Potential predictors examined were; age, sex, Body Mass Index (BMI), ethnicity, chronic conditions, current employment and deprivation. Descriptive statistics and logistic regression were conducted.

Results

Individuals with chronic mental and physical conditions (fatigue, insomnia, anxiety, depression, diabetes, breathing problems, high blood pressure, heart disease, stroke and cancer) were more likely to report the lowest levels of physical activity across all five outcomes. Demographic variations were also observed.

Conclusions

Targeting people with chronic mental and physical conditions has the potential reduce the impact of physical inactivity.

Introduction

According to a recent economic analysis, physical inactivity related ill health cost the UK National Health Service £0.9 billion in 2006-2007.¹ Physical inactivity has been attributed to 6% of coronary heart disease, 7% type 2 diabetes, 10% breast cancer, 10% colon cancer and 9% premature mortality incidence globally.² Increasing physical activity has been associated with improvements in physical and mental health and wellbeing.³⁻⁶ A recent review and network meta-analysis has demonstrated comparative effectiveness of physical activity with drug treatment on mortality for coronary heart disease, stroke, heart failure and prediabetes.⁷

Recent evidence suggests that smaller increases in the physical activity of those who are least active can have a bigger health and cost-effectiveness impact than raising levels of those already slightly active to guideline levels, at a population level.^{8,9} With rates of physical inactivity increasing (for example walking and cycling for transport rates have declined over the past 18 years¹⁰), it is important to understand the factors associated with low physical activity so that interventions can target segments of the population whose health stands to benefit the most.

Research has concentrated on the predictors of levels of physical activity. Physical activity participation is related to various demographic and health factors such as age, sex, socioeconomic status and health status.¹¹⁻¹⁸ However, there has been less consideration of how these factors may differ as predictors of physical inactivity. Individuals who are inactive constitute a distinct subgroup that have not been profiled. They differ from individuals who are active since physical activity can represent a range of activity levels and hence incorporate different subgroups of individuals. Given that it is the physically inactive that can reap the largest health benefits, they form an important group to focus on compared to individuals of all levels of physical activity who may not gain as much.

The few studies that have focused on physical inactivity have provided an insight into important predictors of the population subgroup. Being female, older age and higher level of education were significantly associated with physical inactivity in a large-scale survey in five Asian countries.¹⁹ Being an ethnic minority, of older age, less education and providing care to others were found to be correlated with physical inactivity in a large-scale survey of United States women aged 40 and over.²⁰ Physical inactivity among employees of a major academic institution in the United States was associated with higher prevalence of cardiovascular diseases, fair or poor health status and absenteeism from work.²¹

However, there has been little understanding of how chronic mental and physical health conditions may act as barriers to physical activity. If they are associated with physical inactivity, then they offer

a potential point of intervention given that individuals could be targeted through their use of health care. The demographic and health characteristics of those with the lowest levels of activity in a more general population in the United Kingdom (UK), however, have not yet been examined in detail. Profiling the characteristics of the least active will allow for appropriate targeting of interventions to increase physical activity among this group, thus providing the potential for greater improvements in health not just in those individuals but also at population level.

The current study aimed to expand on such findings by examining predictors of the lowest levels of physical activity, to identify segments of the population whose health could be most improved the most by increasing physical activity.

Methods

CLAHRC South Yorkshire Cohort

Data were obtained from the first wave of the Collaboration for Leadership in Applied Health Research and Care (CLAHRC) South Yorkshire Cohort (2010-2012), a longitudinal observational dataset set up to collect health data on the residents of South Yorkshire, England.^{22,23} The cohort contains information on physical activity, current and long-term health, health care usage and demographic characteristics.

The current study examined data from the 10,628 participants recruited via 14 Sheffield GP practices within the CLAHRC South Yorkshire Cohort (response rate of 17.1%). The restriction to Sheffield only data was to help inform 'Move More', a recently launched citywide campaign to increase physical activity across the population of Sheffield (<http://www.movemoresheffield.com>).²⁴ Therefore, a focus on examining the predictors of the least active among Sheffield residents contributes to the effort to understand where targeted community and individual interventions could have the greatest impact in terms of population health and reducing inequalities. In addition, analyses here intend to provide a baseline of physical activity for Sheffield, providing direction for interventions linked to the 'Move More' programme.

Physical activity measure

Frequency and intensity of self-reported physical activity were measured by mailed and online questionnaire using the question, "During the last WEEK, how many hours did you spend on each of the following activities?" The different types of activities were categorised into:

- Physical exercise such as swimming, jogging, aerobics, football, tennis, gym workout etc.
- Cycling, including cycling to work and during leisure time
- Walking, including walking to work, shopping, for pleasure etc.

Response options were: 'none', 'some but less than 1 hour', 'at least 1 hour but less than 3 hours' and '3 hours or more'.^{22,23} This item is a shortened version of the General Practice Physical Activity Questionnaire (GPPAQ), a validated physical activity screening tool for primary care.^{25,26}

Outcomes

Five separate outcomes were used to represent the lowest levels of physical activity:

- Combined outcome – lowest level of physical activity across all three categories, i.e. those people who reported 'none' for physical exercise and cycling and 'none' or 'less than 1 hour' for walking
- Lowest level of physical activity for physical exercise and cycling categories combined (i.e. those reporting 'none' to both)
- Lowest level of physical exercise (i.e. those reporting 'none')
- Lowest level of cycling (i.e. those reporting 'none')
- Lowest level of walking (i.e. those reporting 'none' or 'less than 1 hour')

The combined outcome aimed to examine the predictors of those who have reported doing none of the activities listed in the three categories. We also included minimal walking (less than one hour a week) since these low levels of movement still represent an inactive lifestyle that may be missed by using 'none' alone (e.g. walking around the house, or from work to the car). The analyses were repeated using only 'none' for the walking measure, however they produced similar results. The second outcome, combining the physical exercise and cycling categories but excluding walking is a more sensitive outcome, since walking was excluded in the GPPAQ due to the difficulty in determining the contribution of the self-reported walking to overall physical activity (e.g., speed, duration of bouts, intensity).²⁶ Finally, each category (physical exercise, cycling and walking) was examined separately as they were found to be measuring distinct behaviours.

Potential predictors measured in the CLAHRC cohort

The following predictors were examined²², from self-report measures:

- Age
- Sex
- Body mass index (BMI)

- Ethnicity (split as 'White' (93%) and 'Non-White' (overall 7%; consists of 1% 'Mixed', 3.7% 'Asian', 1.7% 'Black' and 0.6% 'Other'))
- Employed or not
- Deprivation (measured using postcode and Indices of Deprivation 2010)¹⁶
- Chronic conditions (tiredness/fatigue, insomnia, anxiety/nerves, depression, diabetes, breathing problems, high blood pressure, heart disease, stroke and cancer)

These predictor variables were selected as they encompass all the demographic and health characteristics of the cohort sample that were in the dataset and are potential predictors of ability to benefit from physical activity, which research has suggested are related to physical activity.¹¹⁻¹⁸ The mental and physical conditions were all included because there is relatively robust evidence that physical activity is associated with a decreased risk of development, progression or symptoms of the condition.^{5,6,27-56}

Analysis

The percentages of individuals identified as physically inactive according to the five outcomes were examined across the explanatory variables (Table I). Patterns of low activity/inactivity were also examined by age and gender (Figures I and II). A logistic regression analysis was undertaken to test the predictive value of each explanatory variable on each of the five outcomes (Table II). Finally, utilising the results and applying the sample weights, it was possible to extrapolate the results to explore the numbers of people that could be targeted through interventions (Table III).

Results

Table I presents summary descriptive statistics for our outcome variables. The highest proportions of those reporting the lowest levels of physical activity across all five outcomes were found in those with chronic mental and physical conditions. In addition, the proportions of people reporting the lowest levels of activity increased incrementally across most outcomes with increasing age, weight (although higher proportions of underweight than normal weight individuals reported the lowest levels of activity across all outcomes) and deprivation quintile. Higher proportions of unemployed (versus employed) people reported the lowest levels of activity on all outcomes. For sex, the relationship was less clear-cut, with a higher proportion of males reporting lowest levels in terms of the combined outcome and walking and females reporting lowest levels in terms of the combined outcome excluding walking, physical exercise and cycling. Similarly, for ethnicity, non-white

participants reported lowest levels on the combined outcome and walking, with white participants reporting lowest levels on the combined outcome excluding walking, physical exercise and cycling.

The incremental increase in the proportion of people reporting the lowest levels of activity with increasing age held for males and females across the five activity outcomes (Figures I and II), with a particularly marked increase in the proportion reporting the lowest levels of physical exercise and cycling in males with age. There were also slight increases in the proportions of males and females reporting the lowest levels of activity around the time of state retirement age (i.e. from the 65-74 to the ≥ 75 age group).

The results of the logistic regression analysis suggest that certain chronic conditions were significant independent predictors of the lowest levels of physical activity on four of the five outcomes (Table II). Fatigue significantly predicted the lowest levels of physical activity on the combined outcome, the combined outcome excluding walking, physical exercise and walking. Anxiety significantly predicted the lowest levels of physical activity on the combined outcome excluding walking and physical exercise. Depression significantly predicted the lowest levels of physical activity on combined outcome, and breathing problems significantly predicted the lowest levels of physical activity on the combined outcome and walking. Variation in cycling was significantly accounted for overall by the presence of long term conditions. Whilst the predictor variables were correlated in our data, these correlations were not strong suggesting that their multiple inclusion in the model is not problematic.

Age is consistently positively associated with physical inactivity across each measure, as are BMI and deprivation. Being employed was positively associated with cycling, possibly reflecting individuals who cycle to work. Ethnic minorities were over twice as likely to be inactive overall as compared to the individuals who were White and this is due to inactivity in relation to walking. Whilst males were significantly more likely to be physically inactive, this was due to differences in walking behaviour after accounting for the other variables (but the opposite was seen for physical exercise and cycling).

Robust data from the cohort enables extrapolation to the Sheffield population in terms of numbers of people in Sheffield who will benefit from interventions specifically targeted at those with certain chronic conditions (Table III). Depending on the physical activity variable targeted, between 72,245 and 225,853 people would stand to benefit from increased physical activity if those with all chronic conditions were targeted.

Discussion

Main finding of this study

Our study has presented a detailed profiling of the characteristics of inactive individuals across a range of demographic factors and health conditions. These findings could help to inform interventions aimed at improving physical activity levels amongst inactive individuals.

What is already known on this topic

Research has demonstrated that raising physical activity levels in the inactive can have the greatest improvement in health than compared to individuals of low physical activity levels.^{8,9} However, there has been little research into the factors associated with physical inactivity, with most research concentrated in understanding individuals who are active.

What this study adds

Together, the Cohort figures for those with chronic mental and physical conditions can be extrapolated to between 72,245 and 225,853 inactive people in Sheffield (depending on the measure), who may potentially benefit from a targeted intervention. Since small increases in the physical activity of those who are least active can result in greater population health benefit than increasing the levels of more active people to the recommended levels,^{8,9} targeting people with chronic mental and physical conditions might optimally reduce the impact of physical inactivity on population health.

The issue of cause and effect may be pertinent. The data examined in the current study is cross-sectional, so it could be that a low level of physical activity was an initiating factor in the chronic condition, or that the chronic condition resulted in low levels of activity, or a combination of the two. The implication of the current findings, then, is that if interventions can successfully increase physical activity in those with chronic mental and physical conditions then health improvements may be noticed that in turn might make it easier to further increase participation in physical activity.

Since individuals with chronic conditions often access health services, these sub-populations of people could be targeted for increasing activity through health service settings. For example, brief counselling could be added to clinic appointments or relevant, tailored written information could be available in waiting rooms. The healthcare environment could be changed to make it easier for physical activity to be part of NHS care pathways for people with chronic disease, for example NHS clinics being co-located with swimming pools and gym facilities.

There is a need for more innovative evidence-based approaches that can overcome the significant barriers to behaviour change, and particularly increasing physical activity, for individuals with health problems and physical limitations. Significant cultural and attitudinal changes may still be required to ensure the benefits of physical activity are seen to outweigh any condition-associated risks, and ensure it is seen as important and worthwhile to tackle the barriers to increased activity. Improving physical activity in these groups of people may also have the added benefit of improvement in the prognosis or management of their conditions, as evidence suggests a beneficial effect of physical activity on fatigue,^{43,49} insomnia,^{47,50} anxiety,^{27,30,37} depression,^{33,39,48} diabetes,^{6,28,29,41,51} breathing problems,^{38,42,45} high blood pressure,^{44,52} heart disease,^{5,34,35,53} stroke^{31,46,54} and cancer,^{3,32,36,55}. Since April 2013, the Quality and Outcomes Framework (QOF) for UK general practices has included screening (using the GPPAQ) and brief intervention to reduce inactivity for patients with hypertension. Similar incentives, if shown to be effective, could be extended to other chronic conditions.

The proportions of people with the lowest physical activity in the current analysis are similar to those in previous studies examining general practice patient populations using similar measures. For example, 45% of general practice patients in London were found to be inactive (no physical exercise or cycling) as measured on the GPPAQ in an evaluation of the Let's Get Moving physical activity intervention (compared with our estimate of 42.8%).⁵⁶ In another study examining GPPAQ use, 43% of GP patients in low SES areas were categorised as inactive (no physical exercise or cycling) on the GPPAQ.⁵⁷ Although our sample is similar to other patient-based populations, non-patient populations suggest differences. For example, a report for the British Heart Foundation estimated physical inactivity to be 30% for males and 38% for females.⁵⁸ This may suggest that our estimates for individuals who could benefit from intervention may be upwardly biased.

Limitations of this study

The CLAHRC South Yorkshire Cohort consists of a self-selected sample, and it is known that volunteer samples can differ from the overall population on a number of important characteristics.^{28,29} The South Yorkshire Cohort is broadly representative, only containing a slightly older, more affluent and female population resulting in a small bias in our results.²² Although the physical activity measure used in the South Yorkshire Cohort data was based on a validated measure, the measure makes no reference to household or occupational physical activity, levels of which could potentially be high even among those with low levels of physical exercise, cycling and walking. The measure is not validated to discriminate between minimal levels of walking, however the guidance given to participants is not totally clear and so this type of minimal activity may have been reported.

Likewise, the measure does not capture sedentary behaviour, which still confers health risk even among those who meet Department of Health (2011) guidelines for physical activity.⁵⁹

The data were also entirely self-reported, which can potentially be a source of recall bias and presentation bias. For example, correlations between self-reported physical activity and more objective measures such as accelerometry and pedometry tend to be weak.^{16,60,61} This may be less problematic for our study, however, as self-reported physical activity is typically overestimated compared with objective measures and we were concerned with examining the lowest levels of physical activity.

The data were cross-sectional and this restricts the potential to use our analysis to make any causal associations. This is particularly relevant in the context of chronic conditions and physical activity, because of the bi-directional influences of some modelled relationships (a chronic condition may influence physical inactivity, but physical inactivity could also influence the onset and prognosis of a chronic condition). Finally, the Cohort dataset did not include data on commuting patterns or car ownership. However, these factors have been shown to be related to physical inactivity, with increased levels of car ownership and usage associated with lower levels of walking and cycling.^{17,62,63} Further research should incorporate such information into analyses to help better design interventions.

Conclusion

Targeting people with chronic mental and physical conditions may potentially be a very effective and cost-effective strategy to reduce the impact of physical inactivity. The impact would be twofold, since people with chronic conditions represent an important group to target as a large and relatively inactive population and physical activity is known to improve such conditions. Profiling the wider characteristics of individuals who are physically inactive can also usefully inform the design and targeting of interventions.

Funding

This publication presents independent research as part of the Obesity Theme in CLAHRC SY 2008-2013 which is supported by the National Institute for Health Research, Collaboration for Leadership in Applied Health Research and Care, Yorkshire and Humber (NIHR CLAHRC YH) and The University of Sheffield. The views and opinions expressed are those of the authors, and not necessarily those of the NHS, the NIHR or the Department of Health.

Acknowledgements

We are grateful to all the individuals who have enrolled in the cohort. We also acknowledge the GP practice staff for their contribution in the recruiting process. This publication is the work of the authors and does not necessarily reflect the views of the South Yorkshire Cohort Management Team or Steering Committee.

This publication presents independent research by the National Institute for Health Research Collaboration for Leadership in Applied Health Research and Care for South Yorkshire (NIHR CLAHRC SY) a pilot which ended in 2013. Further details about the new NIHR CLAHRC Yorkshire and Humber can be found at www.clahrc-hy.nihr.ac.uk. The views and opinions expressed are those of the authors, and not necessarily those of the NHS, the NIHR or the Department of Health.

We are also grateful to South Yorkshire CLAHRC, the University of Sheffield, Sheffield Hallam University and the National Centre for Sport and Exercise Medicine in Sheffield. Thanks to Hugh Balmer for his helpful advice.

References

1. Scarborough P, Bhatnagar P, Wickramasinghe KK, Allender S, Foster C, Rayner M. The economic burden of ill health due to diet, physical inactivity, smoking, alcohol and obesity in the UK: an update to 2006-07 NHS costs. *J Public Health (Oxf)*. 2011;33(4):527–35. doi:10.1093/pubmed/fdr033.
2. Lee I-M, Shiroma EJ, Lobelo F, Puska P, Blair SN, Katzmarzyk PT. Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *Lancet*. 2012;380(9838):219–29. doi:10.1016/S0140-6736(12)61031-9.
3. Brown JC, Winters-Stone K, Lee A, Schmitz KH. Cancer, physical activity, and exercise. *Compr Physiol*. 2012;2(4):2775–809. doi:10.1002/cphy.c120005.
4. Cramp F, Byron-Daniel J. Exercise for the management of cancer-related fatigue in adults. *Cochrane database Syst Rev*. 2012;11:CD006145. doi:10.1002/14651858.CD006145.pub3.
5. Murtagh EM, Murphy MH, Boone-Heinonen J. Walking: the first steps in cardiovascular disease prevention. *Curr Opin Cardiol*. 2010;25(5):490–6. doi:10.1097/HCO.0b013e32833ce972.
6. De Lemos ET, Oliveira J, Pinheiro JP, Reis F. Regular physical exercise as a strategy to improve antioxidant and anti-inflammatory status: benefits in type 2 diabetes mellitus. *Oxid Med Cell Longev*. 2012;2012:741545. doi:10.1155/2012/741545.
7. Naci H, Ioannidis JPA. Comparative effectiveness of exercise and drug interventions on mortality outcomes: metaepidemiological study. *BMJ*. 2013;347(oct01 1):f5577–f5577. doi:10.1136/bmj.f5577.
8. Minton J, Dimairo M, Everson-Hock E, Scott E, Goyder E. Exploring the relationship between baseline physical activity levels and mortality reduction associated with increases in physical activity: a modelling study. *BMJ Open*. 2013;3(10):e003509. doi:10.1136/bmjopen-2013-003509.
9. Woodcock J, Franco OH, Orsini N, Roberts I. Non-vigorous physical activity and all-cause mortality: systematic review and meta-analysis of cohort studies. *Int J Epidemiol*. 2011;40(1):121–38. doi:10.1093/ije/dyq104.
10. Department for Transport. *National Travel Survey: England 2013*. London; 2014. Available at: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/337473/nt_s2013-01.pdf. Accessed August 5, 2014.
11. Bauman AE, Reis RS, Sallis JF, Wells JC, Loos RJF, Martin BW. Correlates of physical activity: why are some people physically active and others not? *Lancet*. 2012;380(9838):258–71. doi:10.1016/S0140-6736(12)60735-1.
12. Beenackers MA, Kamphuis CBM, Giskes K, et al. Socioeconomic inequalities in occupational, leisure-time, and transport related physical activity among European adults: a systematic review. *Int J Behav Nutr Phys Act*. 2012;9(1):116. doi:10.1186/1479-5868-9-116.

13. Boone-Heinonen J, Diez Roux A V, Kiefe CI, Lewis CE, Guilkey DK, Gordon-Larsen P. Neighborhood socioeconomic status predictors of physical activity through young to middle adulthood: the CARDIA study. *Soc Sci Med*. 2011;72(5):641–9. doi:10.1016/j.socscimed.2010.12.013.
14. Stalsberg R, Pedersen A V. Effects of socioeconomic status on the physical activity in adolescents: a systematic review of the evidence. *Scand J Med Sci Sports*. 2010;20(3):368–83. doi:10.1111/j.1600-0838.2009.01047.x.
15. Barnett I, Guell C, Ogilvie D. The experience of physical activity and the transition to retirement: a systematic review and integrative synthesis of qualitative and quantitative evidence. *Int J Behav Nutr Phys Act*. 2012;9(1):97. doi:10.1186/1479-5868-9-97.
16. Health and Social Care Information Centre. *Health Survey for England - 2008: Physical activity and fitness*. Leeds: Health and Social Care Information Centre; 2009. Available at: <http://www.hscic.gov.uk/pubs/hse08physicalactivity>. Accessed January 24, 2014.
17. Adams J. Prevalence and socio-demographic correlates of “active transport” in the UK: analysis of the UK time use survey 2005. *Prev Med (Baltim)*. 2010;50(4):199–203. doi:10.1016/j.ypmed.2010.01.006.
18. Fox KR, Hillsdon M, Sharp D, et al. Neighbourhood deprivation and physical activity in UK older adults. *Health Place*. 2011;17(2):633–40. doi:10.1016/j.healthplace.2011.01.002.
19. Ng N, Hakimi M, Van Minh H, et al. Prevalence of physical inactivity in nine rural INDEPTH Health and Demographic Surveillance Systems in five Asian countries. *Glob Health Action*. 2009;2. doi:10.3402/gha.v2i0.1985.
20. King AC, Castro C, Wilcox S, Eyler AA, Sallis JF, Brownson RC. Personal and environmental factors associated with physical inactivity among different racial–ethnic groups of U.S. middle-aged and older-aged women. *Heal Psychol*. 2000;19(4):354–364.
21. Birdee GS, Byrne DW, McGown PW, et al. Relationship between physical inactivity and health characteristics among participants in an employee-wellness program. *J Occup Environ Med*. 2013;55(5):514–9. doi:10.1097/JOM.0b013e31827f37d7.
22. Green MA, Li J, Relton C, et al. Cohort Profile: The Yorkshire Health Study. *Int J Epidemiol*. 2014;doi: 10.1093/ije/dyu121. doi:10.1093/ije/dyu121.
23. Relton C, Bissell P, Smith C, et al. South Yorkshire Cohort: a “cohort trials facility” study of health and weight - protocol for the recruitment phase. *BMC Public Health*. 2011;11(1):640. doi:10.1186/1471-2458-11-640.
24. Tew GA, Copeland RJ, Till SH. Sport and exercise medicine and the Olympic health legacy. *BMC Med*. 2012;10(1):74. doi:10.1186/1741-7015-10-74.
25. Department of Health. General Practice Physical Activity Questionnaire (GPPAQ) - Publications - GOV.UK. 2013. Available at: <https://www.gov.uk/government/publications/general-practice-physical-activity-questionnaire-gppaq>. Accessed January 14, 2014.

26. National Institute for Health and Clinical Excellence (NICE). *The General Practice Physical Activity Questionnaire (GPPAQ)*. London; 2006. Available at: <http://www.nice.org.uk/nicemedia/live/11927/40195/40195.pdf>.
27. Asmundson GJG, Fetzner MG, Deboer LB, Powers MB, Otto MW, Smits JAJ. Let's get physical: a contemporary review of the anxiolytic effects of exercise for anxiety and its disorders. *Depress Anxiety*. 2013;30(4):362–73. doi:10.1002/da.22043.
28. Avery L, Flynn D, van Wersch A, Sniehotta FF, Trenell MI. Changing physical activity behavior in type 2 diabetes: a systematic review and meta-analysis of behavioral interventions. *Diabetes Care*. 2012;35(12):2681–9. doi:10.2337/dc11-2452.
29. Chimen M, Kennedy A, Nirantharakumar K, Pang TT, Andrews R, Narendran P. What are the health benefits of physical activity in type 1 diabetes mellitus? A literature review. *Diabetologia*. 2012;55(3):542–51. doi:10.1007/s00125-011-2403-2.
30. Conn VS. Anxiety outcomes after physical activity interventions: meta-analysis findings. *Nurs Res*. 59(3):224–31. doi:10.1097/NNR.0b013e3181dbb2f8.
31. Cumming TB, Tyedin K, Churilov L, Morris ME, Bernhardt J. The effect of physical activity on cognitive function after stroke: a systematic review. *Int Psychogeriatr*. 2012;24(4):557–67. doi:10.1017/S1041610211001980.
32. Davies NJ, Batehup L, Thomas R. The role of diet and physical activity in breast, colorectal, and prostate cancer survivorship: a review of the literature. *Br J Cancer*. 2011;105 Suppl :S52–73. doi:10.1038/bjc.2011.423.
33. Dinas PC, Koutedakis Y, Flouris AD. Effects of exercise and physical activity on depression. *Ir J Med Sci*. 2011;180(2):319–25. doi:10.1007/s11845-010-0633-9.
34. Doering L V, Eastwood J-A. A literature review of depression, anxiety, and cardiovascular disease in women. *J Obstet Gynecol Neonatal Nurs*. 2011;40(3):348–61. doi:10.1111/j.1552-6909.2011.01236.x.
35. Ferrier S, Blanchard CM, Vallis M, Giacomantonio N. Behavioural interventions to increase the physical activity of cardiac patients: a review. *Eur J Cardiovasc Prev Rehabil*. 2011;18(1):15–32. doi:10.1097/HJR.0b013e32833ace0e.
36. Fong DYT, Ho JWC, Hui BPH, et al. Physical activity for cancer survivors: meta-analysis of randomised controlled trials. *BMJ*. 2012;344:e70. doi:10.1136/bmj.e70.
37. Moylan S, Eyre HA, Maes M, Baune BT, Jacka FN, Berk M. Exercising the worry away: how inflammation, oxidative and nitrogen stress mediates the beneficial effect of physical activity on anxiety disorder symptoms and behaviours. *Neurosci Biobehav Rev*. 2013;37(4):573–84. doi:10.1016/j.neubiorev.2013.02.003.
38. Nellesen A, Hernandez NA, Pitta F. Physiotherapy and rehabilitative interventions in patients with chronic respiratory diseases: exercise and non-exercise treatment. *Panminerva Med*. 2013;55(2):197–209. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23676960>. Accessed January 24, 2014.

39. Robertson R, Robertson A, Jepson R, Maxwell M. Walking for depression or depressive symptoms: A systematic review and meta-analysis. *Ment Health Phys Act.* 2012;5(1):66–75. doi:10.1016/j.mhpa.2012.03.002.
40. Soares FHR, de Sousa MBC. Different types of physical activity on inflammatory biomarkers in women with or without metabolic disorders: a systematic review. *Women Health.* 2013;53(3):298–316. doi:10.1080/03630242.2013.782940.
41. Umpierre D, Ribeiro PAB, Kramer CK, et al. Physical activity advice only or structured exercise training and association with HbA1c levels in type 2 diabetes: a systematic review and meta-analysis. *JAMA.* 2011;305(17):1790–9. doi:10.1001/jama.2011.576.
42. Butts JF, Belfer MH, Gebke KB. Exercise for patients with COPD: an integral yet underutilized intervention. *Phys Sportsmed.* 2013;41(1):49–57. doi:10.3810/psm.2013.02.1999.
43. Chambers D, Bagnall A-M, Hempel S, Forbes C. Interventions for the treatment, management and rehabilitation of patients with chronic fatigue syndrome/myalgic encephalomyelitis: an updated systematic review. *J R Soc Med.* 2006;99(10):506–20. doi:10.1258/jrsm.99.10.506.
44. Cornelissen VA, Smart NA. Exercise training for blood pressure: a systematic review and meta-analysis. *J Am Heart Assoc.* 2013;2(1):e004473. doi:10.1161/JAHA.112.004473.
45. Gloeckl R, Marinov B, Pitta F. Practical recommendations for exercise training in patients with COPD. *Eur Respir Rev.* 2013;22(128):178–86. doi:10.1183/09059180.00000513.
46. Pang MYC, Charlesworth SA, Lau RWK, Chung RCK. Using aerobic exercise to improve health outcomes and quality of life in stroke: evidence-based exercise prescription recommendations. *Cerebrovasc Dis.* 2013;35(1):7–22. doi:10.1159/000346075.
47. Passos GS, Poyares DLR, Santana MG, Tufik S, Mello MT de. Is exercise an alternative treatment for chronic insomnia? *Clinics (Sao Paulo).* 2012;67(6):653–60. Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3370319&tool=pmcentrez&rendertype=abstract>. Accessed January 24, 2014.
48. Rethorst CD, Trivedi MH. Evidence-based recommendations for the prescription of exercise for major depressive disorder. *J Psychiatr Pract.* 2013;19(3):204–12. doi:10.1097/01.pra.0000430504.16952.3e.
49. Yancey JR, Thomas SM. Chronic fatigue syndrome: diagnosis and treatment. *Am Fam Physician.* 2012;86(8):741–6. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23062157>. Accessed January 24, 2014.
50. Yang P-Y, Ho K-H, Chen H-C, Chien M-Y. Exercise training improves sleep quality in middle-aged and older adults with sleep problems: a systematic review. *J Physiother.* 2012;58(3):157–63. doi:10.1016/S1836-9553(12)70106-6.
51. Colberg SR, Sigal RJ, Fernhall B, et al. Exercise and type 2 diabetes: the American College of Sports Medicine and the American Diabetes Association: joint position statement. *Diabetes Care.* 2010;33(12):e147–67. doi:10.2337/dc10-9990.

52. Millar PJ, Goodman JM. Exercise as medicine: Role in the management of primary hypertension. *Appl Physiol Nutr Metab*. 2014;39(7):856–8. doi:10.1139/apnm-2014-0006.
53. Charansonney OL, Vanhees L, Cohen-Solal A. Physical activity: from epidemiological evidence to individualized patient management. *Int J Cardiol*. 2014;170(3):350–7. doi:10.1016/j.ijcard.2013.11.012.
54. Gallanagh, Siobhan; Quinn, Terry J; Alexander, Jen; Walters MR. Physical activity in the prevention and treatment of stroke. *ISRN Neurol*. 2011;2011:953818. doi:10.5402/2011/953818.
55. Singh F, Newton RU, Galvão DA, Spry N, Baker MK. A systematic review of pre-surgical exercise intervention studies with cancer patients. *Surg Oncol*. 2013;22(2):92–104. doi:10.1016/j.suronc.2013.01.004.
56. Bull FC, Milton KE. A process evaluation of a “physical activity pathway” in the primary care setting. *BMC Public Health*. 2010;10(1):463. doi:10.1186/1471-2458-10-463.
57. Heron N, Tully MA, McKinley MC, Cupples ME. Physical activity assessment in practice: a mixed methods study of GPPAQ use in primary care. *BMC Fam Pract*. 2014;15(1):11. doi:10.1186/1471-2296-15-11.
58. Townsend N, Bhatnagar P WK, Scarborough P, Foster C RM. *Physical activity statistics 2012*. London; 2012. Available at: file:///C:/Users/User/Downloads/M130 BHF_Physical Activity Supplement_2012.pdf. Accessed August 5, 2014.
59. Department of Health. *Start Active, Stay Active: A report on physical activity for health from the four home countries’ Chief Medical Officers*. London; 2011. Available at: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/216370/dh_128210.pdf. Accessed March 7, 2014.
60. Igelström H, Emtner M, Lindberg E, Asenlöf P. Level of agreement between methods for measuring moderate to vigorous physical activity and sedentary time in people with obstructive sleep apnea and obesity. *Phys Ther*. 2013;93(1):50–9. doi:10.2522/ptj.20120123.
61. Celis-Morales CA, Perez-Bravo F, Ibañez L, Salas C, Bailey MES, Gill JMR. Objective vs. self-reported physical activity and sedentary time: effects of measurement method on relationships with risk biomarkers. Dasgupta K, ed. *PLoS One*. 2012;7(5):e36345. doi:10.1371/journal.pone.0036345.
62. Cerin E, Leslie E, Owen N. Explaining socio-economic status differences in walking for transport: an ecological analysis of individual, social and environmental factors. *Soc Sci Med*. 2009;68(6):1013–20. doi:10.1016/j.socscimed.2009.01.008.
63. Saelens BE, Sallis JF, Frank LD. Environmental correlates of walking and cycling: Findings from the transportation, urban design, and planning literatures. *Ann Behav Med*. 2003;25(2):80–91. doi:10.1207/S15324796ABM2502_03.