

which has been published in final form at <https://doi.org/10.1111/anu.12389>. This article may be used for non-commercial purposes in accordance With Wiley Terms and Conditions for self-archiving.

Effect of dietary saturated and monounsaturated fatty acids in juvenile barramundi *Lates calcarifer*.

Michael J. Salini <sup>1,2\*</sup>, Giovanni M. Turchini <sup>1</sup>, Brett D. Glencross <sup>3,4</sup>

<sup>1</sup> School of Life and Environmental Sciences, Deakin University, Warrnambool, Princess Hwy, Sherwood Park, Victoria, Australia

<sup>2</sup> CSIRO Agricultural Productivity Flagship, 144 North Street, Woorim, Queensland, Australia.

<sup>3</sup> CSIRO Agricultural Productivity Flagship, 41 Boggo Road, Dutton Park, Queensland, Australia

<sup>4</sup> Present address: Ridley Aquafeed, 12 Neon St, Narangba, Queensland, Australia

\*Corresponding author: Email [msalini@deakin.edu.au](mailto:msalini@deakin.edu.au)

Phone +61 (0)457 499 258

Running title: Dietary SFA and MUFA in juvenile barramundi

Keywords: Barramundi; SFA; MUFA; Fatty acid; Metabolism;  $\beta$ -oxidation;

## Abstract

Barramundi (*Lates calcarifer*), a catadromous teleost of commercial interest, perform well when fed a wide range of different dietary oils in varying combinations. However, as critical limits of fish oil (FO) are approached, a range of alternative oils are being explored that are typically richer in shorter chain and more saturated fatty acids. In this study, the response of juvenile barramundi (47.0 g/fish initial weight) fed isolipidic and isoenergetic diets with 8.2% added oil was tested. The experimental test diets were either a 2:1 or 1:2 ratio of SFA to MUFA (SFA-D and MUFA-D respectively) compared to a control diet (CTRL-D) fed for a period of eight weeks. The replacement of FO with diets containing mostly olive oil (diet MUFA-D) and mostly refined palm oil (diet SFA-D) did not impact the growth performance or feed utilisation parameters of fish in this study. However, the composition of the tissues generally reflected the diets and mass-dependent retention of energy was observed in the SFA-D and MUFA-D fed fish. The *in vivo* beta-oxidation activity was consistent with the diet fatty acid composition with the most dominant FA being heavily oxidised. *In vivo* elongation and desaturation activity indicated both a demand for initial synthesis of fatty acids and energy production. However, despite the similarities, some vagaries in the results were observed and discussed. This study provides evidence that additional dietary MUFA and SFA are suitable lipid classes for juvenile barramundi and they are both equally efficient at sparing LC-PUFA from an oxidative fate.

## Introduction

In the context of feeding fish alternative oils, many studies have focused on the essential fatty acids and their bioactive long-chain derivatives, eicosapentaenoic and docosahexaenoic acid (20:5n-3 and 22:6n-3). However, it is suggested that saturated and monounsaturated fatty acids (SFA and MUFA) are preferred substrates for  $\beta$ -oxidation in fish, sparing the more essential fatty acids for their functional and biological roles (review by Henderson 1996). As alternative oils are being increasingly used in aquafeeds it is important to understand the effect of increasing dietary levels of SFA and MUFA on lipid metabolism in fish.

Carnivorous fish such as barramundi (*Lates calcarifer*), also known as the Asian seabass, must rely heavily on oxidation of lipid for their energetic requirements. Past studies have consistently shown positive responses to the replacement of fish oil (FO) with a range of alternative oils such as soybean, canola, rapeseed, Echium, usually included as blends (Williams et al. 2006, Alhazzaa et al. 2011, Raso & Anderson 2003). Moreover, it is also clear that higher dietary levels of long-chain polyunsaturated (LC-PUFA), originating from FO, are oxidised or potentially retro-converted, rather than being efficiently deposited in the flesh (Salini et al. 2015a).

Atlantic salmon (*Salmo salar*) also performed well when fed diets with either low n-3 and high monounsaturated fatty acids (MUFA) and in addition MUFA were found to be good substrates for  $\beta$ -oxidation (Menoyo et al. 2003). In the same species,  $\beta$ -oxidation was found to be dependent on energy demand and that when fatty acids were provided in excess to synthetic demands they were increasingly  $\beta$ -oxidised for energy production (Stubhaug et al. 2007, Torstensen et al. 2004). Denstadli et al. (2011) found that the medium chain length decanoic acid (10:0) was rapidly oxidised for energy production whereas oleic acid (18:1n-9) was primarily deposited in the intramuscular fat in Atlantic salmon. In other species such as

Atlantic cod (*Gadus morhua*) reduced lipid deposition in the liver was likely caused by selective  $\beta$ -oxidation of 16:0 and 18:0 saturated fatty acids (SFA) present in palm oil (PO) and rapeseed oil (Jobling et al. 2008). In agreement, polka dot grouper (*Cromileptes altivelis*) more readily oxidised lipid in diets containing SFA (coconut fat) for energy rather than MUFA (olive oil); however this was at the expense of efficient growth (Smith et al. 2005). Further to this, the authors of a recent study on European sea bass (*Dicentrarchus labrax*) speculated that a single fatty acid (eg. 18:1n-9) might be more efficient at sparing LC-PUFA from  $\beta$ -oxidation than oil blends (Eroldogan et al. 2013).

Using an *in-vivo* whole-body fatty acid balance approach for determining fatty acid metabolism in rainbow trout, Turchini & Francis (2009) found that the most heavily oxidised fatty acids were also the most dominant. However, in the fish oil fed fish the most predominant dietary fatty acid (16:0) was not readily oxidised for energy production, but rather was preferentially elongated to 18:0 or desaturated to 16:1n-7. The same research group also found that the rate of  $\beta$ -oxidation of LC-PUFA was reduced when Murray cod (*Maccullochella peelii peelii*) were fed diets containing abundant MUFA and SFA (Turchini et al. 2011). Moreover, a recent study on hybrid striped bass (White Bass *Morone chrysops*  $\times$  Striped Bass *M. saxatilis*) concluded that diets containing SFA-rich lipid and to a lesser extent MUFA-rich lipid were able to conserve tissue fatty acid profiles possibly due to preferential oxidation (Trushenski et al. 2015).

Consistent responses of a range of fish species indicate that SFA and MUFA provided in surplus are heavily oxidised as substrates for energy, effectively sparing LC-PUFA for biological needs and deposition. Likewise, it is unfavourable to provide excess n-3 LC-PUFA, typical of FO, as these FA can also be readily  $\beta$ -oxidised (Turchini et al. 2013, Torstensen et al. 2000). Recent studies have shown that barramundi are capable of utilising

poultry oil, characterised by high MUFA content, however to what extent SFA are preferentially utilised or able to 'spare' LC-PUFA remains unclear. The proposed experiment examined the effect of diets containing a 2:1 or 1:2 ratio of SFA to MUFA in the lipid as it was hypothesised that these lipid classes would be metabolised differently. The digestibility of the diets was also measured and mass-balance computations used to estimate the fatty acid metabolism *in vivo*.

## **Materials and methods**

### *Ingredient and diet preparation*

A single basal diet was formulated and prepared without the addition of dietary lipids. The dry ingredients were passed separately through a hammermill (Mikro Pulverizer, type 1 SH, New Jersey, USA) such that the maximum particle size was less than 750 µm. All dry ingredients were then thoroughly mixed using an upright commercial mixer (Bakermix, Model 60 A-G, NSW, Australia). Fish meal was defatted prior to use by manually mixing hexane and fish meal (2:1) in a large drum. The mix was left to soak for 3h before draining the excess hexane and repeating the process a second time. The fish meal was oven dried overnight at 60 °C to a constant dry matter. The chemical composition of the main dietary ingredients is presented in Table 1. The lipid blends were first heated in an oven to 70 °C then mixed gently. The basal diet was then separated into smaller batches and aliquots of lipid (8.2% of the diet) were added to form the three treatment diets. Fresh water was added at approximately 30% of dry mash weight and mixed to form consistent dough then the dough was subsequently screw pressed through a 4 mm die. The pellets were dried overnight at 60 °C to a constant dry matter. The dietary treatments provided protein at 60%, lipid at 13% with an energetic value of 22 MJ/kg. The three dietary treatments consisted of a control diet with added FO (designated as CTRL-D), a diet containing a blend of FO and olive oil (designated

as MUFA-D) and a diet containing a blend of FO and refined palm products (designated as SFA-D). The present study used a blend of two common palm fractions, refined, bleached and deodorised palm oil (RBDPO) and palm flake, a highly refined stearin fraction. The RDBPO fraction used in the present study contained approximately 35% MUFA whereas the palm flake product was mostly SFA with a fatty acid profile dominated by roughly equal percentages of 16:0 and 18:0 FA. This blend of oils was used to achieve the desired SFA to MUFA ratio of the diets. The diets were stored at -20 °C until required. The formulation and chemical composition of the three diets are presented in Table 2.

#### *Barramundi husbandry and growth*

Juvenile barramundi (*Lates calcarifer*) were sourced from the Betta Barra fish hatchery (Atherton, QLD, Australia), on-grown in a 10,000L tank and fed a commercial diet (Marine Float; Ridley Aquafeed, Narangba, QLD, Australia). Prior to commencement of the experiment the fish were transferred to a series of experimental tanks (300L) with flow-through seawater (salinity =35 g/kg; dissolved oxygen  $4.6 \pm 0.15$  mg /L) maintained at  $30.0 \pm 0.01$  °C (mean  $\pm$  SD) with a supply rate of about 3 L/min to each of the tanks. At the beginning of the experiment, the tanks held 26 fish of  $47.0 \pm 0.3$  g (mean  $\pm$  SD, n =234 individually weighed fish). The three experimental diets were randomly distributed amongst the nine tanks with each treatment having three replicate tanks. The diets were fed once daily to apparent satiety as determined over three separate feeding events, for eight weeks. Any uneaten feed was collected shortly after and a correction factor was applied (Helland et al. 1996). Briefly, the correction factor was calculated as the proportion of soluble losses after immersion in water for 1 h.

#### *Sample collection, preparation and digestibility analysis*

Ethical clearance was approved for the experimental procedures by the CSIRO animal ethics committee A12/2013. Six fish of similar size from the original stock were euthanized by an overdose of AQUI-S™ (Lower Hutt, New Zealand) at the beginning of the experiment and stored at -20 °C until biochemical analysis. A further six fish were dissected and a sample of liver tissue was then removed and placed into 1.5 mL screw-top vial and kept on dry ice before being transferred to a -80 °C freezer until biochemical analysis.

Prior to the termination of the growth assay, faeces were collected using established settlement protocols for the digestibility assessment (Blyth et al. 2014). Briefly, a collection chamber was filled with water and frozen then attached to the evacuation line of a swirl separator and left overnight. The following morning, the collection chamber was removed and the chilled faeces were captured in a plastic sample container and stored at -20 °C until chemical analysis.

### *Chemical analyses*

Prior to analysis the diets were each ground to a fine powder using a bench grinder (KnifeTec™ 1095, FOSS, Denmark). The whole fish were passed through a commercial meat mincer (MGT – 012, Taiwan) twice to obtain a homogeneous mixture. A sample was taken for dry matter analysis and another sample was freeze-dried along with the faecal samples until no further loss of moisture was observed (Alpha 1-4, Martin Christ, Germany). Dry matter was calculated gravimetrically following oven drying at 105°C for 24 h. Total yttrium concentrations were determined in the diet and faecal samples after nitric acid digestion in a laboratory microwave digester (Ethos One, Milestone, Italy) using inductively coupled plasma-mass spectrophotometry (ICP-MS) (ELAN DRC II, Perkin Elmer, USA) following McQuaker et al. (1979). Crude protein was calculated after the determination of total nitrogen by organic elemental analysis (CHNS-O, Flash 2000, Thermo Scientific, USA), based on N x

6.25. Total lipid content was determined following extraction of the lipids using chloroform:methanol (2:1) following Folch et al. (1957). Gross ash content was determined gravimetrically following loss of mass after combustion of a sample in a muffle furnace at 550°C for 12 h. Gross energy was determined by adiabatic bomb calorimetry (Parr 6200 Calorimeter, USA).

Fatty acid composition was determined following the methods of Christie (2003). Lipids were esterified by an acid-catalysed methylation and 0.3 mg of an internal standard was added to each sample (21:0 Supelco, PA, USA). The fatty acids were identified relative to the internal standard following separation by gas chromatography (GC). An Agilent Technologies 6890N GC system (Agilent Technologies, California, USA) fitted with a DB-23 (60m x 0.25mm x 0.15 µm, cat 122-2361 Agilent Technologies, California) capillary column and flame ionisation detection was used. The temperature program was 50–175 °C at 25 °C /min then 175–230 °C at 2.5 °C /min. The injector and detector temperatures were set at 250 °C and 320 °C, respectively. The column head pressure was set to constant pressure mode at 170 kPa using hydrogen as the carrier gas. The peaks were identified by comparing retention times to the internal standard and further referenced against known standards (37 Comp. FAME mix, Supelco, PA, USA). The resulting peaks were then corrected by the theoretical relative FID response factors (Ackman 2002) and quantified relative to the internal standard.

#### *Calculations and statistical analysis*

Differences in the ratio of dry matter, protein, lipid and energy to yttrium in the diet and faeces were calculated to determine the apparent digestibility using the formula: (Maynard & Loosli 1979):



$$AD_{diet} = \left( 1 - \left( \frac{Y_{diet} \times Parameter_{faeces}}{Y_{faeces} \times Parameter_{diet}} \right) \right) \times 100$$

Where  $Y_{diet}$  and  $Y_{faeces}$  represent the yttrium content in the diet and faeces respectively and the  $Parameter_{diet}$  and  $Parameter_{faeces}$  represent the nutritional parameter (dry matter, protein, lipid or energy) and specific fatty acids in the diet and faeces respectively. Nutrient retention efficiencies were calculated as the ratio of the nutrient or specific fatty acid gained relative to their respective consumption during the study period using the formula (Maynard & Loosli 1979):

$$Retention \% = \left( \frac{Nutrient_f - Nutrient_i}{Nutrient_c} \right) \times 100$$

Where  $Nutrient_f$  is the nutrient/energy content of the fish from each replicate upon termination of the experiment and the  $Nutrient_i$  is the mean nutrient/energy content of the initial fish (n=6 fish).  $Nutrient_c$  is the amount of nutrient/energy consumed during the experiment. The computation of apparent *in vivo* fatty acid elongation, desaturation and  $\beta$ -oxidation was performed using the whole-body fatty acid balance method (WBFABM) following Turchini et al. (2007). Briefly, this involved determination of the appearance/disappearance of specific fatty acids by mass balance. The resulting values of net appearance/disappearance were then transformed to a molecular weight basis per gram of body weight per day (nmol/g fish/d). Subsequent back calculations along the known fatty acid bioconversion pathways were used to determine the fate of specific fatty acids.

All data are expressed as mean  $\pm$  SEM (n=3) unless otherwise specified. Data with values < 0.05 are reported as 0.1. All data were checked for normal distribution and homogeneity of variance by qualitative assessment of residual and normal Q-Q plots. All data were analysed by one-way ANOVA or t-test using the RStudio package v.0.98.501 (R Core Team 2012).

Any percentage data were arcsine transformed prior to analysis. Levels of significance were compared using Tukey's HSD *a posteriori* test with significance among treatments defined as  $P < 0.05$ .

## **Results**

### *Growth and feed utilisation*

During the 56 d growth assay, the fish in all treatments responded readily to the experimental diets and growth in the CTRL-D group was consistent with the predicted model growth, achieving 106% of the modelled potential (Glencross & Bermudes 2012). Live-weight measurements were conducted after four weeks of feeding and then again after eight weeks and although there was a tendency towards lower growth in the SFA-D and MUFA-D fish at both time points, there were no significant differences (Table 3). The same trend was observed in terms of daily feed intake and growth rate with no significant differences and there was no difference in FCR values (Table 3). There were no differences in terms of survival with only one fish that died and was removed from the system over the term of the experiment (Table 3).

The apparent digestibility of macro nutrients and specific fatty acids was affected by the diets (Table 4). Both dry matter (DM) and lipid were significantly less digestible in the SFA-D fed group of fish. There were no significant differences in either protein or gross energy (GE) digestibility among the treatments (Table 4). There were no significant differences in the digestibility of individual and total LC-PUFA among the diets, each showing almost complete digestion in most cases. Similarly, there were no significant differences in the digestibility of individual and total PUFA and 18:1. However, the saturated fatty acids were

significantly affected with 18:0 and 16:0 being less digestible in the SFA-D fed fish compared to the other treatments. The total SFA digestibility was also significantly reduced in the SFA-D fed fish (Table 4).

### *Biochemical analysis*

The whole-body DM, protein and lipid composition on a wet weight basis was not significantly affected by the diets; however, the GE was significantly lower in the SFA-D fed fish (Table 5). Generally, the fatty acid composition of the diets was reflected in the whole fish and also the liver tissue. Whole-body total fatty acids were not significantly different however there were some significant differences between the specific fatty acids. Among the LC-PUFA the whole-body 22:6n-3 and 22:5n-3 was significantly higher in the CTRL-D fish compared to the MUFA-D fish. The whole-body 18:2n-6 was significantly highest in the MUFA-D and lowest in the CTRL-D (Table 5). The whole-body 18:1 was significantly highest in the MUFA-D fed fish whereas 18:0 and 16:0 were lowest in the MUFA-D fed (Table 5). The total LC-PUFA and the total n-3 fatty acids were significantly lowest in the MUFA-D fed fish (Table 5).

The liver fatty acid composition was affected in a similar fashion to that of the whole-body where 22:6n-3, 22:5n-3, 20:5n-3 and 20:4n-6 had significantly higher composition in the CTRL-D fed fish. However, there were no significant differences among the MUFA-D or SFA-d diets (Table 5). There was no significant difference in PUFA composition of the liver. The total LC-PUFA and n-3 composition were significantly higher in the CTRL-D fish and there was no significant difference in n-6 composition of the liver (Table 5).

### *Mass-balance computations*

There were no significant differences in protein or lipid retention however there was significantly higher energy retention in the CTRL-D fish (Table 3). There was significantly higher  $\beta$ -oxidation of specific n-3 series fatty acids (22:6n-3, 20:5n-3 and 18:4n-3) and n-6 series fatty acids (18:3n-6 and 20:4n-6) in the CTRL-D diet compared to the other treatment diets (Table 6). There was no recorded  $\beta$ -oxidation for any of the dominant saturates (including 12:0, 14:0, 16:0 and 18:0) however there was a significant increase in  $\beta$ -oxidation of n-7 series FA (16:1n-7 and 18:1n-7) in the CTRL-D diet compared to the other treatment diets. There was a significant increase in the  $\beta$ -oxidation of 18:1 fatty acids in the MUFA-D diet compared to the other treatment diets (Table 6).

There was significantly higher fatty acid *de novo* production of 12:0 (neogenesis) in the MUFA-D fed fish compared to the other treatment diets (Table 6). The recorded elongation activity of 14:0 to 14:1n-5 was significantly higher in the CTRL-D and MUFA-D fed fish while the elongation of 16:0 to 16:1n-7 was highest in the CTRL-D and SFA-D fed fish. Elongation of 20:5n-3 to 22:5n-3 was only recorded in the fish fed the SFA-D and MUFA-D diets and there was no significant difference (Table 6). The  $\Delta$ -9 desaturation activity was significantly highest in the SFA-D fed fish followed by the control fed fish and there was none recorded in the MUFA-D fed fish (Table 6). There was no  $\Delta$ -5 or  $\Delta$ -6 desaturation activity detected in any of the diet treatments (data not presented).

Relative to the total intake of specific fatty acids, there was a significantly greater proportion of total LC-PUFA deposited in the SFA-D and MUFA-D fed fish; however, there was no difference between these two groups (Table 7). Likewise, there was proportionally less  $\beta$ -oxidation in the SFA-D and MUFA-D fish; however there was no difference between these two groups (Table 7). There was only a minor proportion of the consumed LC-PUFA converted or excreted in the groups of fish.

## Discussion

The present study demonstrated that the inclusion of a 2:1 or 1:2 ratio of either SFA or MUFA, did not affect the growth performance of juvenile barramundi. The simultaneous reduction of dietary LC-PUFA (~11% LC-PUFA in SFA- and MUFA-D vs 21.5% LC-PUFA in CTRL-D), provided that they were still above reported requirements of 1.2% LC-PUFA, also had no effect on performance (Williams et al. 2006). Moreover, the feed intake and FCR values were unaffected. This is consistent with a range of species showing that substitution of FO with lipid rich in either SFA or MUFA does not affect fish growth performance (Turchini et al. 2009). The fish in the present study more than tripled in size suggesting that trial duration or nutrient turnover was not a confounding issue. The fish in the SFA-D and MUFA-D treatments did show a numerical reduction in growth however this was not confirmed statistically. It is uncertain whether longer trial duration would have resulted in significant differences as recent studies with barramundi have demonstrated that changes to the lipid profile of the diets can have rapid metabolic effects (Salini et al. 2015b). Lipid and in particular the saturated fatty acids are generally less digestible at lower environmental temperature and as a result less energy availability for growth (Ng et al. 2004, Olsen & Ringø 1998). The barramundi is a tropical species adapted to high water temperature and potentially better able to cope with dietary SFA and MUFA rich lipid.

The replacement of FO with MUFA rich lipid such as that from poultry oil had no effect on growth or FCR in a range of species including rainbow trout (*Oncorhynchus mykiss*) juveniles (Fonseca-Madrigal et al. 2005), post-smolt Atlantic salmon (*Salmo salar*) (Bell et al. 2002) or large 1.5 kg Atlantic salmon (Torstensen et al. 2000), juvenile red hybrid tilapia (*Oreochromis* sp.) (Bahurmiz & Ng 2007) humpback grouper (*Cromileptes altivelis*) (Shapawi et al. 2008) and African catfish (*Clarias gariepinus*) (Ng et al. 2003). However,

consistent among these studies when FO was completely replaced by MUFA rich lipid was the modified tissue FA profile resulting in reduced concentration of the beneficial n-3 LC-PUFA. The same effect was clearly noted in the present study. However, proportional to the LC-PUFA intake, the results of most studies and those of the present study conclude that MUFA rich lipid is an ideal energy source capable of ‘sparing’ the more valuable LC-PUFA from  $\beta$ -oxidation.

An important consideration when formulating with SFA rich palm oil (PO) is the fraction used (Ng & Gibon 2011). Past studies demonstrated that growth performance (Shapawi et al. 2008, Ng et al. 2003) and digestibility of a range of species was not affected by different PO fractions (Bahurmiz & Ng 2007) while the digestibility of palm products was significantly reduced compared to FO. Recent studies have also concluded that other sources of lipid rich in SFA such as beef tallow did not affect the growth of Atlantic salmon (Emery et al. 2014) or rainbow trout (Trushenski et al. 2011). Consistent with these reports, the present study demonstrated that the blend of palm products used did not compromise growth performance in barramundi while there were notable reductions to the lipid and specific fatty acid digestibility.

In the present study, the digestibility of total lipid was significantly reduced in the SFA fed fish however this did not lead to a reduction in energy availability or any changes in whole-body lipid composition or retention. The reduction in lipid digestibility was evidently a result of the greatly reduced digestibility of the saturates, including both 16:0 and 18:0 FA. This is in agreement with other studies showing that the digestibility of the saturated fatty acids was reduced when a range of species were fed SFA rich oil (Torstensen et al. 2000, Ng et al. 2004, Turchini et al. 2013, Caballero et al. 2002). Moreover, in the present study the digestibility of SFA's decreased with increasing chain length, consistent with other studies

(Caballero et al. 2002, Ng et al. 2004, Johnsen et al. 2000). The reduced rate of lipolysis and absorption of longer chain SFA, such as 18:0 is caused by the lower ability of these FA to form lipid emulsions prior to digestion (Olsen et al. 1998).

In contrast to the present study, negative effects of the high level inclusion of SFA (e.g. as occurs with high level use of PO) have been reported and that these effects are arguably more pronounced in carnivorous and/or marine fish with higher n-3 LC-PUFA requirements.

Fountoulaki et al. (2009) found that gilthead sea bream growth was negatively impacted by reduced digestibility of SFA rich lipid over an extended growth trial of six months. Japanese seabass (*Lateolabrax japonicas*) growth performance was significantly reduced with a high inclusion of SFA rich lipid after only 50 days of growth and similarly reduced digestibility was inferred to be the cause (Gao et al. 2012). In agreement, Turchini et al. (2013) found that over a long duration (27 weeks) rainbow trout fed SFA rich lipid (at 75% replacement level) showed depressed performance compared to that of a control diet.

An *in vivo* whole-body fatty acid balance method (WBFABM) was used in the present study in order to understand the apparent fate of specific fatty acids (Turchini et al. 2007).

Theoretically,  $\beta$ -oxidation of specific fatty acids should be recorded if they are provided in excess (Turchini et al. 2013, Eroldogan et al. 2013, Stubhaug et al. 2007). However, in the present study there was no recorded  $\beta$ -oxidation of any SFA despite the relatively high proportion of SFA in all three diets. This effect is likely to be caused by the low lipid levels of all three diets, a strategy that was intended to highlight the potential effects of the lipid classes. Based on the  $\beta$ -oxidation results, it appears that MUFA is marginally better at sparing LC-PUFA from oxidation (Table 6) which is in agreement with past studies (Turchini et al. 2011, Codabaccus et al. 2012). However, in the present study the final composition of the fish suggests that the SFA-D fed fish were significantly more efficient at depositing or

'sparing' LC-PUFA in the whole body (Table 5). To resolve this discrepancy, it is necessary to look at the net intake and total intake budgets to clarify the situation (Table 7). When expressed proportional to FA intake, the SFA-D and MUFA-D fed fish consequently deposited almost exactly the same LC-PUFA and the differences between the two diets were insignificant.

Previous studies have demonstrated that limited *de novo* FA production (neogenesis) occurs when diets with adequate SFA were fed to Atlantic salmon (Emery et al. 2014) or rainbow trout (Turchini et al. 2013). However, neogenesis was evident in MUFA rich (canola oil) fed trout (Turchini et al. 2013). This is broadly similar to the results obtained in the present study in that neogenesis in barramundi was significantly higher in the MUFA fed fish. This may be partly explained by the up-regulation of genes related to lipogenic activity in response to vegetable oil observed in other species (Tocher et al. 2002, Bell et al. 2001, Bell et al. 2002). In contrast, dietary PUFA clearly demonstrated a suppression of the lipogenic enzyme, fatty acid synthase (FAS) in the rat (Blake & Clarke 1990). Moreover, in rainbow trout hepatocytes, FAS expression was strongly inhibited by PUFA (18:3n-3 and 20:5n-3). These results and those of the present study confirm that the energetically expensive process of initial synthesis of palmitic acid from acetyl- and malonyl-CoA is avoided by the presence of dietary SFA thus allowing energy to be utilised more efficiently for growth and other cellular processes.

There was also elongation and desaturation activity occurring in the CTRL-D fed fish. This may indicate that the total lipid content of the diet was limiting as intended based on the recommended 14 to 16 % specification for growing barramundi (Glencross 2006). However, there was clearly adequate n-3 LC-PUFA in the diets based on the known requirement data for barramundi of around 1.2% (Glencross & Rutherford 2011, Williams et al. 2006).



Consistent with Glencross & Rutherford (2011), there was no elongation of 20:5n-3 to 22:5n-3 in the control fed fish. However, in the SFA-D and MUFA-D fed fish, there was a slight increase in the elongation of available 20:5n-3 to 22:5n-3 possibly in an attempt to achieve 22:6n-3 synthesis. However, barramundi, like most marine fish are not equipped with the complete set of enzymes required to endogenously synthesise sufficient LC-PUFA from precursor FA (Mohd-Yusof et al. 2010). Moreover, a recent study also demonstrated a similar increase in 22:5n-3 from 20:5n-3, lending further support to the elongation results obtained in the present study (Salini et al. 2015a).

## **Conclusions**

The results of the present study demonstrate that the inclusion of 2:1 or 1:2 ratios of SFA and MUFA did not lead to a reduction in growth performance of juvenile barramundi. However, a range of other metabolic modifications were observed. Notable was the LC-PUFA sparing effect of both MUFA and SFA. Additionally, SFA and MUFA were preferentially metabolised and deposited in the whole body and liver tissue proportional to their respective intake. The low digestibility of specific fatty acids (18:0 and 16:0) is consistent with other studies and may have an impact in the long term utilisation of the SFA rich diet. These results clearly indicate that consideration must be given to the proportion of either SFA or MUFA during diet formulation, as these two classes of fatty acids can influence the *in vivo* metabolism of fatty acids and the final fatty acid composition of the whole fish.

## **Acknowledgements**

The authors wish to acknowledge the CSIRO Agricultural Productivity Flagship for financial support. The authors would also like to thank the technical staff at the Bribie Island Research Centre (BIRC), QLD, Australia, for their valued contributions.

## References

- Ackman, R.G. (2002) The gas chromatograph in practical analysis of common and uncommon fatty acids for the 21st century. *Anal. Chim. Acta*, 465, 175-192.
- Alhazzaa, R., Bridle, A.R., Nichols, P.D. & Carter, C.G. (2011) Replacing dietary fish oil with Echium oil enriched barramundi with C18 PUFA rather than long-chain PUFA. *Aquaculture*, 312, 162-171.
- Bahurmiz, O.M. & Ng, W.-K. (2007) Effects of dietary palm oil source on growth, tissue fatty acid composition and nutrient digestibility of red hybrid tilapia, *Oreochromis* sp., raised from stocking to marketable size. *Aquaculture*, 262, 382-392.
- Bell, J.G., Henderson, R.J., Douglas, R.T., Fiona, M., James, R.D., Allan, P., Richard, P.S. & John, R.S. (2002) Substituting fish oil with crude palm oil in the diet of Atlantic salmon (*Salmo salar*) affects muscle fatty acid composition and hepatic fatty acid metabolism. *J. Nutr.*, 132, 222-30.
- Bell, J.G., McEvoy, J., Tocher, D.R., McGhee, F., Campbell, P.J. & Sargent, J.R. (2001) Replacement of fish oil with rapeseed oil in diets of Atlantic salmon (*Salmo salar*) affects tissue lipid compositions and hepatocyte fatty acid metabolism. *J. Nutr.*, 131, 1535-1543.
- Blake, W.L. & Clarke, S.D. (1990) Suppression of rat hepatic fatty-acid synthase and s-14 gene-transcription by dietary polyunsaturated fat. *J. Nutr.*, 120, 1727-1729.
- Blyth, D., Tabrett, S., Bourne, N. & Glencross, B.D. (2014) Comparison of faecal collection methods and diet acclimation times for the measurement of digestibility coefficients in barramundi (*Lates calcarifer*). *Aquac. Nutr.*, 21, 248-255.
- Caballero, M.J., Obach, A., Rosenlund, G., Montero, D., Gisvold, M. & Izquierdo, M.S. (2002) Impact of different dietary lipid sources on growth, lipid digestibility, tissue fatty acid composition and histology of rainbow trout, *Oncorhynchus mykiss*. *Aquaculture*, 214, 253-271.
- Christie, W.W. (2003) Lipid analysis, isolation, separation, identification and structural analysis of lipids., 3rd edn, PJ Barnes and Associates, Bridgewater, UK.
- Codabaccus, B.M., Carter, C.G., Bridle, A.R. & Nichols, P.D. (2012) The “n-3 LC-PUFA sparing effect” of modified dietary n-3 LC-PUFA content and DHA to EPA ratio in Atlantic salmon smolt. *Aquaculture*, 356-357, 135-140.
- Denstadli, V., Bakke, A.M., Berge, G.M., Krogdahl, A., Hillestad, M., Holm, H. & Ruyter, B. (2011) Medium-Chain and Long-Chain Fatty Acids Have Different Postabsorptive Fates in Atlantic Salmon. *J. Nutr.*, 141, 1618-1625.
- Emery, J.A., Smullen, R.P. & Turchini, G.M. (2014) Tallow in Atlantic salmon feed. *Aquaculture*, 422-423, 98-108.
- Eroldogan, T.O., Yilmaz, A.H., Turchini, G.M., Arslan, M., Sirkecioglu, N.A., Engin, K., Ozsahinoglu, I. & Mumogullarinda, P. (2013) Fatty acid metabolism in European sea bass (*Dicentrarchus labrax*): effects of n-6 PUFA and MUFA in fish oil replaced diets. *Fish Physiol. Biochem.*, 39, 941-55.
- Folch, J., Lees, M. & Sloane-Stanley, G. (1957) A simple method for the isolation and purification of total lipids from animal tissues. *J. Biol. Chem.*, 226, 497-509.
- Fonseca-Madrigal, J., Karalazos, V., Campbell, P.J., Bell, J.G. & Tocher, D.R. (2005) Influence of dietary palm oil on growth, tissue fatty acid compositions, and fatty acid metabolism in liver and intestine in rainbow trout (*Oncorhynchus mykiss*). *Aquac. Nutr.*, 11, 241-250.
- Fountoulaki, E., Vasilaki, A., Hurtado, R., Grigorakis, K., Karacostas, I., Nengas, I., Rigos, G., Kotzamanis, Y., Venou, B. & Alexis, M.N. (2009) Fish oil substitution by vegetable oils in commercial diets for gilthead sea bream (*Sparus aurata* L.); effects on growth performance, flesh quality and fillet fatty acid profile: Recovery of fatty acid profiles by a fish oil finishing diet under fluctuating water temperatures. *Aquaculture*, 289, 317-326.

- Gao, J., Koshio, S., Ishikawa, M., Yokoyama, S., Ren, T., Komilus, C.F. & Han, Y. (2012) Effects of dietary palm oil supplements with oxidized and non-oxidized fish oil on growth performances and fatty acid compositions of juvenile Japanese sea bass, *Lateolabrax japonicus*. *Aquaculture*, 324–325, 97-103.
- Glencross, B.D. (2006) The nutritional management of barramundi, *Lates calcarifer* - a review. *Aquac. Nutr.*, 12, 291-309.
- Glencross, B.D. & Bermudes, M. (2012) Adapting bioenergetic factorial modelling to understand the implications of heat stress on barramundi (*Lates calcarifer*) growth, feed utilisation and optimal protein and energy requirements – potential strategies for dealing with climate change? *Aquac. Nutr.*, 18, 411-422.
- Glencross, B.D. & Rutherford, N. (2011) A determination of the quantitative requirements for docosahexaenoic acid for juvenile barramundi (*Lates calcarifer*). *Aquac. Nutr.*, 17, e536-e548.
- Helland, S.J., Grisdale-Helland, B. & Nerland, S. (1996) A simple method for the measurement of daily feed intake of groups of fish in tanks. *Aquaculture*, 139, 157-163.
- Henderson, R.J. (1996) Fatty acid metabolism in freshwater fish with particular reference to polyunsaturated fatty acids. *Archiv für Tierernaehrung*, 49, 5-22.
- Jobling, M., Leknes, O., Sæther, B.-J. & Bendiksen, E.Å. (2008) Lipid and fatty acid dynamics in Atlantic cod, *Gadus morhua*, tissues: Influence of dietary lipid concentrations and feed oil sources. *Aquaculture*, 281, 87-94.
- Johnsen, R.I., Grahl-Nielsen, O. & Roem, A. (2000) Relative absorption of fatty acids by Atlantic salmon *Salmo salar* from different diets, as evaluated by multivariate statistics. *Aquac. Nutr.*, 6, 255-261.
- Maynard, L.A. & Loosli, J.K. (1979) *Animal Nutrition*, 6th edn., McGraw-Hill Book Co., New York, NY.
- McQuaker, N.R., Brown, D.F. & Kluckner, P.D. (1979) Digestion of Environmental Materials for Analysis by Inductively Coupled Plasma - Atomic Emission Spectrometry. *Anal. Chem.*, 51, 1082-1084.
- Menoyo, D., Lopez-Bote, C.J., Bautista, J.M. & Obach, A. (2003) Growth, digestibility and fatty acid utilization in large Atlantic salmon (*Salmo salar*) fed varying levels of n-3 and saturated fatty acids. *Aquaculture*, 225, 295-307.
- Mohd-Yusof, N.Y., Monroig, O., Mohd-Adnan, A., Wan, K.L. & Tocher, D.R. (2010) Investigation of highly unsaturated fatty acid metabolism in the Asian sea bass, *Lates calcarifer*. *Fish Physiol. Biochem.*, 36, 827-843.
- Ng, W.-K. & Gibon, V. (2011) Palm oil and saturated fatty acid-rich vegetable oils. In: *Fish oil replacement and alternative lipid sources in aquaculture feeds* (Turchini, G. M., Ng, W.-K. & Tocher, D. eds) 1st edn, CRC Press, Taylor and Francis group, FL, USA.
- Ng, W.-K., Lim, P.-K. & Boey, P.-L. (2003) Dietary lipid and palm oil source affects growth, fatty acid composition and muscle  $\alpha$ -tocopherol concentration of African catfish, *Clarias gariepinus*. *Aquaculture*, 215, 229-243.
- Ng, W.-K., Sigholt, T. & Bell, J.G. (2004) The influence of environmental temperature on the apparent nutrient and fatty acid digestibility in Atlantic salmon (*Salmo salar* L.) fed finishing diets containing different blends of fish oil, rapeseed oil and palm oil. *Aquac. Res.*, 35, 1228-1237.
- Olsen, R.E., Henderson, R.J. & Ringø, E. (1998) The digestion and selective absorption of dietary fatty acids in Arctic charr, *Salvelinus alpinus*. *Aquac. Nutr.*, 4, 13-21.
- Olsen, R.E. & Ringø, E. (1998) The influence of temperature on the apparent nutrient and fatty acid digestibility of Arctic charr, *Salvelinus alpinus* L. *Aquac. Res.*, 29, 695-701.
- R Core Team (2012) *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Raso, S. & Anderson, T.A. (2003) Effects of dietary fish oil replacement on growth and carcass proximate composition of juvenile barramundi (*Lates calcarifer*). *Aquac. Res.*, 34, 813-819.

- Salini, M.J., Irvin, S.J., Bourne, N., Blyth, D., Cheers, S., Habilay, N. & Glencross, B.D. (2015a) Marginal efficiencies of long chain-polyunsaturated fatty acid use by barramundi (*Lates calcarifer*) when fed diets with varying blends of fish oil and poultry fat. *Aquaculture*, In press.
- Salini, M.J., Turchini, G.M., Wade, N. & Glencross, B.G. (2015b) Defining essential fatty acid deficiency in juvenile barramundi *Lates calcarifer*. *Br. J. Nutr.*, Submitted.
- Shapawi, R., Mustafa, S. & Ng, W.-K. (2008) Effects of dietary fish oil replacement with vegetable oils on growth and tissue fatty acid composition of humpback grouper, *Cromileptes altivelis* (Valenciennes). *Aquac. Res.*, 39, 315-323.
- Smith, D.M., Williams, I.H., Williams, K.C., Barclay, M.C. & Venables, W.N. (2005) Oxidation of medium-chain and long-chain fatty acids by polka dot grouper *Cromileptes altivelis*. *Aquac. Nutr.*, 11, 41-48.
- Stubhaug, I., Lie, Ø. & Torstensen, B.E. (2007) Fatty acid productive value and  $\beta$ -oxidation capacity in Atlantic salmon (*Salmo salar* L.) fed on different lipid sources along the whole growth period. *Aquac. Nutr.*, 13, 145-155.
- Tocher, D., Fonseca-Madrigal, J., Bell, J.G., Dick, J., Henderson, R.J. & Sargent, J. (2002) Effects of diets containing linseed oil on fatty acid desaturation and oxidation in hepatocytes and intestinal enterocytes in Atlantic salmon (*Salmo salar*). *Fish Physiol. Biochem.*, 26, 157-170.
- Torstensen, B.E., Froyland, L. & Lie, O. (2004) Replacing dietary fish oil with increasing levels of rapeseed oil and olive oil - effects on Atlantic salmon (*Salmo salar* L.) tissue and lipoprotein lipid composition and lipogenic enzyme activities. *Aquac. Nutr.*, 10, 175-192.
- Torstensen, B.E., Lie, O. & Froyland, L. (2000) Lipid metabolism and tissue composition in Atlantic salmon (*Salmo salar* L.) - Effects of capelin oil, palm oil, and oleic acid-enriched sunflower oil as dietary lipid sources. *Lipids*, 35, 653-664.
- Trushenski, J., Rosenquist, J. & Gause, B. (2011) Growth performance, tissue fatty acid composition, and consumer appeal of rainbow trout reared on feeds containing terrestrially derived rendered fats. *N. Am. J. Aquac.*, 73, 468-478.
- Trushenski, J.T., Crouse, C.C. & Rombenso, A.N. (2015) Effects of fish oil sparing on fillet fatty acid composition in hybrid striped bass are influenced by dietary levels of saturated and unsaturated fatty acids. *N. Am. J. Aquac.*, 77, 160-169.
- Turchini, G.M. & Francis, D.S. (2009) Fatty acid metabolism (desaturation, elongation and  $\beta$ -oxidation) in rainbow trout fed fish oil- or linseed oil-based diets. *Br. J. Nutr.*, 102, 69-81.
- Turchini, G.M., Francis, D.S. & De Silva, S.S. (2007) A whole body, in vivo, fatty acid balance method to quantify PUFA metabolism (desaturation, elongation and beta-oxidation). *Lipids*, 42, 1065-1071.
- Turchini, G.M., Francis, D.S., Senadheera, S.P.S.D., Thanuthong, T. & De Silva, S.S. (2011) Fish oil replacement with different vegetable oils in Murray cod: Evidence of an "omega-3 sparing effect" by other dietary fatty acids. *Aquaculture*, 315, 250-259.
- Turchini, G.M., Hermon, K., Cleveland, B.J., Emery, J.A., Rankin, T. & Francis, D.S. (2013) Seven fish oil substitutes over a rainbow trout grow-out cycle: I) Effects on performance and fatty acid metabolism. *Aquac. Nutr.*, 19, 82-94.
- Turchini, G.M., Torstensen, B.E. & Ng, W.-K. (2009) Fish oil replacement in finfish nutrition. *Reviews in Aquaculture*, 1, 10-57.
- Williams, K.C., Barlow, C.G., Rodgers, L.J. & Agcopra, C. (2006) Dietary composition manipulation to enhance the performance of juvenile barramundi (*Lates calcarifer* Bloch) reared in cool water. *Aquac. Res.*, 37, 914-927.

**Table 1** Chemical composition of ingredients used in experimental diets, all values are presented as g/kg DM unless otherwise stated.

|                          | DF   |       |       |       |       |       |       |      |       |      |      |      |
|--------------------------|------|-------|-------|-------|-------|-------|-------|------|-------|------|------|------|
|                          | Fis  | Poult | Soy   | Whe   | Whe   | Casei | Whe   | Fis  | Oliv  | Pal  | Pal  |      |
|                          | h    | ry    | isola | at    | at    | n     | at    | h    | e oil | m    | m    |      |
|                          | me   | meal  | te    | glute | flour |       | starc | oil  |       | Oil  | Flak |      |
|                          | al   |       |       | n     |       |       | h     |      |       |      | e    |      |
|                          |      |       |       |       |       |       |       |      |       |      |      |      |
| <i>Composition</i>       |      |       |       |       |       |       |       |      |       |      |      |      |
| Dry matter (g/kg)        | 984  | 958   | 958   | 927   | 839   | 924   | 836   | 99   | 2     | 987  | 100  | 99   |
| Protein                  | 789  | 641   | 895   | 823   | 112   | 870   | 5     | 4    | 4     | 3    | 4    | 4    |
| Ash                      | 163  | 138   | 46    | 1     | 6     | 11    | 3     | 1    | ND    | ND   | ND   | ND   |
| Lipid                    | 46   | 151   | 57    | 121   | 22    | 5     | 0     | 6    | 973   | 963  | 986  | 986  |
| Carbohydrate #           | 1    | 70    | 2     | 55    | 860   | 113   | 992   | 39   | 23    | 34   | 10   | 10   |
| Gross energy (MJ/kg)     | 18.9 | 20.4  | 21.8  | 21.2  | 15.3  | 21.9  | 14.5  | 39.3 | 39.5  | 39.5 | 5    | 39.3 |
| <i>Fatty acids (%) ^</i> |      |       |       |       |       |       |       |      |       |      |      |      |
| 16:0                     | 25.8 | 25.4  | 17.8  | -     | -     | -     | -     | 22.9 | 9.9   | 9    | 46.2 | 46.2 |
| 18:0                     | 8.7  | 8.6   | 4.5   | -     | -     | -     | -     | 5.1  | 3.0   | 4.6  | 51.3 | 51.3 |
| 18:1                     | 15.5 | 43.8  | 25.1  | -     | -     | -     | -     | 18.6 | 73.8  | 6    | 0.4  | 0.4  |
| 18:2n-6                  | 1.7  | 11.0  | 46.5  | -     | -     | -     | -     | 2.0  | 11.0  | 6.7  | ND   | ND   |
| 18:3n-3                  | 0.8  | 1.0   | 5.1   | -     | -     | -     | -     | 1.0  | 1.0   | ND   | ND   | ND   |
| 20:4n-6                  | 2.5  | 0.6   | ND    | -     | -     | -     | -     | 1.5  | ND    | ND   | ND   | ND   |
| 20:5n-3                  | 9.0  | 0.5   | ND    | -     | -     | -     | -     | 11.3 | ND    | ND   | ND   | ND   |
| 22:5n-3                  | 2.2  | ND    | ND    | -     | -     | -     | -     | 2.1  | ND    | ND   | ND   | ND   |
| 22:6n-3                  | 19.9 | ND    | ND    | -     | -     | -     | -     | 14.2 | ND    | ND   | ND   | ND   |
| SFA                      | 39.8 | 36.1  | 22.6  | -     | -     | -     | -     | 36.4 | 13.4  | 6    | 99.6 | 99.6 |
| MUFA                     | 22.2 | 50.8  | 25.8  | -     | -     | -     | -     | 29.1 | 74.6  | 7    | 0.4  | 0.4  |
| PUFA                     | 3.4  | 12.0  | 51.6  | -     | -     | -     | -     | 4.9  | 12.0  | 6.7  | ND   | ND   |
| LC-PUFA                  | 34.5 | 1.2   | ND    | -     | -     | -     | -     | 29.7 | ND    | ND   | ND   | ND   |
| n-3                      | 32.7 | 1.6   | 5.1   | -     | -     | -     | -     | 30.5 | 1.0   | ND   | ND   | ND   |
| n-6                      | 5.2  | 11.6  | 46.5  | -     | -     | -     | -     | 4.1  | 11.0  | 6.7  | ND   | ND   |

ND, not detected, N/A, not analysed, values <0.1 are reported as 0.1.

# Carbohydrate calculated by difference (eg. carbohydrate = 1000 – (protein + lipid + ash)

^ All fatty acids are presented as a percentage of the total fatty acids. Quantitative data can be obtained by multiplying the total FA (mg/g lipid) by specific fatty acids (%). 18:1, sum of 18:1n-7, 18:1n-9 cis, 18:1n-9 trans; saturated fatty acids (SFA), sum of 12:0, 14:0, 16:0, 18:0, 20:, 22:0, 24:0; monounsaturated fatty acids (MUFA), sum of 14:1n-5, 16:1n-7, 18:1n-7, 18:1n-9 (cis and trans), 20:1n-7, 20:1n-9, 22:1n-9, 24:1n-9; polyunsaturated fatty acids (PUFA), sum 18:2n-6 (cis and trans), 18:3n-6, 18:3n-3, 18:4n-3; long chain polyunsaturated fatty acids (LC-PUFA), sum 20:2n-6, 20:3n-6, 20:4n-6, 22:4n-6, 2-:3n-3, 20:5n-3, 22:5n-3, 22:6n-3; n-3 and n-6, sum of omega 3 and 6 PUFA and LC-PUFA.

**Table 2** Formulation and composition of experimental diets. All values are g/kg DM unless otherwise stated.

|  | CTRL-D<br>(Fish oil) | SFA-D<br>(Palm oil) | MUFA-D<br>(Olive oil) |
|--|----------------------|---------------------|-----------------------|
| <i>Formulation</i>                         |                      |                     |                       |
| DF Fish meal <sup>a</sup>                  | 150                  | 150                 | 150                   |
| Poultry meal <sup>a</sup>                  | 150                  | 150                 | 150                   |
| Soy protein isolate <sup>b</sup>           | 150                  | 150                 | 150                   |
| Wheat gluten <sup>b</sup>                  | 150                  | 150                 | 150                   |
| Wheat flour <sup>b</sup>                   | 109                  | 109                 | 109                   |
| Casein <sup>c</sup>                        | 100                  | 100                 | 100                   |
| Pregelged wheat starch <sup>b</sup>        | 80                   | 80                  | 80                    |
| DL-Methionine <sup>c</sup>                 | 10                   | 10                  | 10                    |
| Di-calcium phosphate                       | 10                   | 10                  | 10                    |
| Pre-mix vitamins and minerals <sup>d</sup> | 8                    | 8                   | 8                     |
| Yttrium oxide <sup>e</sup>                 | 1                    | 1                   | 1                     |
| Fish oil <sup>a</sup>                      | 82                   | 41                  | 41                    |
| Olive oil <sup>f</sup>                     | 0                    | 0                   | 41                    |
| Palm Oil <sup>f</sup>                      | 0                    | 15                  | 0                     |
| Palm Flake <sup>f</sup>                    | 0                    | 26                  | 0                     |
| <i>Composition as analysed</i>             |                      |                     |                       |
| Dry matter (g/kg)                          | 940                  | 952                 | 979                   |
| Protein                                    | 598                  | 601                 | 585                   |
| Ash  | 64                   | 63                  | 60                    |
| Lipid                                      | 126                  | 137                 | 135                   |
| Carbohydrate <sup>#</sup>                  | 204                  | 192                 | 217                   |
| Gross energy (MJ/kg DM)                    | 21.5                 | 21.6                | 22.2                  |
| <i>Fatty acids (%) <sup>^</sup></i>        |                      |                     |                       |
| Total FA (mg/g lipid)                      | 801.7                | 734.7               | 717.4                 |
| 16:0                                       | 22.9                 | 32.0                | 18.4                  |
| 18:0                                       | 5.5                  | 16.1                | 4.8                   |
| 18:1                                       | 22.7                 | 20.6                | 42.6                  |
| 18:2n-6                                    | 10.0                 | 10.5                | 13.3                  |
| 18:3n-3                                    | 1.3                  | 1.0                 | 1.3                   |
| 20:4n-6                                    | 1.2                  | 0.7                 | 0.7                   |
| 20:5n-3                                    | 8.1                  | 4.1                 | 4.0                   |
| 22:5n-3                                    | 1.6                  | 0.9                 | 0.9                   |
| 22:6n-3                                    | 10.6                 | 5.4                 | 5.4                   |
| SFA  | 34.5                 | 51.8                | 26.4                  |
| MUFA                                       | 31.3                 | 25.0                | 47.3                  |
| PUFA                                       | 12.7                 | 12.1                | 15.3                  |
| LC-PUFA                                    | 21.5                 | 11.0                | 10.9                  |
| n-3  | 22.9                 | 12.0                | 12.2                  |
| n-6  | 11.3                 | 11.1                | 14.0                  |

ND, not detected, N/A, not analysed, values <0.1 are reported as 0.1.

<sup>#</sup> Carbohydrate calculated by difference (eg. carbohydrate = 1000 – (protein + lipid + ash))

<sup>a</sup> Ridley aquafeeds, Narangba, QLD, Australia

<sup>b</sup> Manildra Group, Rocklea, QLD, Australia

<sup>c</sup> Bulk Powders, [www.bulkpowders.com.au](http://www.bulkpowders.com.au)

d Vitamin and mineral premix (IU kg<sup>-1</sup> or g/kg of premix): vitamin A, 2.5MIU; vitamin D3, 0.25 MIU; vitamin E, 16.7 g; vitamin K3, 1.7 g; vitamin B1, 2.5 g; vitamin B2, 4.2 g; vitamin B3, 25 g; vitamin B5, 8.3; vitamin B6, 2.0 g; vitamin B9, 0.8; vitamin B12, 0.005 g; biotin, 0.17 g; vitamin C, 75 g; choline, 166.7 g; inositol, 58.3 g; ethoxyquin, 20.8 g; copper, 2.5 g; ferrous iron, 10.0 g; magnesium, 16.6 g; manganese, 15.0 g; zinc, 25.0 g

e Yttrium oxide; Stanford Materials, Aliso Viejo, California, United States

f Sydney Essential Oil Co. (Sydney, NSW, Australia)

^ Refer to Table 1 for details.

**Table 3** Growth performance and feed utilisation of juvenile barramundi fed experimental diets for eight weeks.

|                        | CTRL-D                  | SFA-D                   | MUFA-D                  | TEST ^        |
|------------------------|-------------------------|-------------------------|-------------------------|---------------|
| Week 0 weight (g)      | 46.9 ± 0.1              | 47.3 ± 0.2              | 47.1 ± 0.1              | F=2.2, P=0.20 |
| Week 4 weight (g)      | 139.2 ± 1.5             | 135.3 ± 7.8             | 136.0 ± 3.0             | F=0.9, P=0.45 |
| Week 8 weight (g)      | 238.3 ± 1.2             | 231.2 ± 9.0             | 230.9 ± 6.2             | F=0.4, P=0.66 |
| Feed intake (g/fish)   | 209.6 ± 2.7             | 202.4 ± 5.0             | 200.7 ± 4.9             | F=0.6, P=0.58 |
| Growth rate (g/fish/d) | 3.4 ± 0.1               | 3.3 ± 0.2               | 3.3 ± 0.1               | F=0.5, P=0.65 |
| FCR                    | 1.10 ± 0.1              | 1.12 ± 0.1              | 1.09 ± 0.1              | F=2.0, P=0.21 |
| Survival (%)           | 98.0 ± 0.1              | 100.0 ± 0.0             | 100.0 ± 0.0             | F=1.0, P=0.42 |
| Protein retention (%)  | 34.1 ± 1.1              | 29.8 ± 1.9              | 32.5 ± 0.9              | F=2.5, P=0.16 |
| Lipid retention (%)    | 48.6 ± 4.5              | 40.2 ± 1.7              | 45.9 ± 0.7              | F=2.4, P=0.17 |
| Energy retention (%)   | 37.8 ± 0.6 <sup>a</sup> | 33.5 ± 1.0 <sup>b</sup> | 34.4 ± 0.3 <sup>b</sup> | F=10.1*       |

ND, not detected, N/A, not analysed, values <0.1 are reported as 0.1.

^ One-way ANOVA, *DF* 2,6, post-hoc Tukey's HSD. Superscript letters indicate different levels of significance between treatment diets, percentage data were arcsine transformed prior to analysis.



**Table 4** Apparent digestibility coefficients of macro nutrients and specific fatty acids of experimental diets fed to juvenile barramundi. All data (n=3 per treatment) are reported as apparent digestibility percentage (%)

|   | CTRL-D                   | SFA-D                   | MUFA-D                  | TEST ^        |
|---|--------------------------|-------------------------|-------------------------|---------------|
| <i>Macro nutrient digestibility (%)</i>         |                          |                         |                         |               |
| Dry matter                                      | 64.3 ± 1.3 <sup>ab</sup> | 60.3 ± 1.5 <sup>b</sup> | 69.4 ± 0.2 <sup>a</sup> | F=9.6*        |
| Protein   | 91.1 ± 0.3               | 92.0 ± 1.2              | 92.1 ± 1.4              | F=0.2, P=0.85 |
| Lipid   | 91.0 ± 0.7 <sup>a</sup>  | 73.7 ± 2.6 <sup>b</sup> | 92.9 ± 1.2 <sup>a</sup> | F=21.8**      |
| Gross energy                                    | 87.3 ± 0.3               | 82.4 ± 3.3              | 89.2 ± 1.6              | F=1.6, P=0.31 |
| <i>Fatty acid digestibility (%)<sup>#</sup></i> |                          |                         |                         |               |
| 16:0  | 82.1 ± 1.4 <sup>a</sup>  | 59.2 ± 3.8 <sup>b</sup> | 86.6 ± 1.7 <sup>a</sup> | F=17.7*       |
| 18:0  | 77.5 ± 3.3 <sup>a</sup>  | 42.4 ± 5.4 <sup>b</sup> | 82.1 ± 2.6 <sup>a</sup> | F=15.5*       |
| 18:1  | 92.4 ± 0.3               | 89.2 ± 1.5              | 94.3 ± 1.0              | F=3.6, P=0.12 |
| 18:2n-6   | 96.0 ± 0.1               | 94.3 ± 0.8              | 96.3 ± 0.9              | F=1.5, P=0.33 |
| 18:3n-3   | 98.6 ± 1.1               | 100.0 ± 0.0             | 97.7 ± 0.6              | T=0.6, P=0.60 |
| 20:4n-6   | 100.0 ± 0.0              | 100.0 ± 0.0             | 100.0 ± 0.0             | N/A           |
| 20:5n-3   | 99.1 ± 0.1               | 100.0 ± 0.0             | 98.6 ± 0.4              | T=0.9, P=0.45 |
| 22:5n-3   | 100.0 ± 0.0              | 100.0 ± 0.0             | 100.0 ± 0.0             | N/A           |
| 22:6n-3   | 98.5 ± 0.2               | 100.0 ± 0.0             | 97.3 ± 0.8              | T=1.1, P=0.35 |
| SFA   | 82.9 ± 1.6 <sup>a</sup>  | 55.6 ± 4.2 <sup>b</sup> | 86.4 ± 1.8 <sup>a</sup> | F=18.3**      |
| MUFA  | 92.9 ± 0.4               | 90.0 ± 1.3              | 94.3 ± 1.0              | F=2.9, P=0.17 |
| PUFA  | 96.7 ± 0.1               | 95.0 ± 0.7              | 96.6 ± 0.9              | F=1.2, P=0.38 |
| LC-PUFA   | 98.9 ± 0.1               | 100.0 ± 0.0             | 98.1 ± 0.6              | T=1.1, P=0.40 |
| n-3   | 98.9 ± 0.1               | 100.0 ± 0.0             | 98.1 ± 0.6              | T=1.2, P=0.36 |
| n-6   | 96.4 ± 0.1               | 94.6 ± 0.8              | 96.5 ± 0.9              | F=1.5, P=0.32 |

ND, not detected, N/A, not analysed, values <0.1 are reported as 0.1.

^ P<0.05\*, P<0.01\*\*, P<0.001\*\*\*; One-way ANOVA, *DF* 2,6, post-hoc Tukey's HSD; T-test, *DF* 4, was used to test two variables; Superscript letters indicate different levels of significance between treatment diets, percentage data were arcsine transformed prior to analysis.

# Refer to Table 1 for details.

**Table 5** Whole body composition data (n=3 per treatment, g/kg live basis). Whole body and liver fatty acid data (n=3 per treatment, % total).

|   | Initial      | CTRL-D                   | SFA-D                   | MUFA-D                  | TEST ^                  |
|---|--------------|--------------------------|-------------------------|-------------------------|-------------------------|
| <i>Whole body composition</i>                 |              |                          |                         |                         |                         |
| Dry matter                                    | 265 ± 0.4    | 329 ± 0.5                | 318 ± 0.3               | 324 ± 0.3               | F=2.1,<br>P=0.21        |
| Protein                                       | 188 ± 0.5    | 206 ± 0.7                | 191 ± 0.8               | 200 ± 0.6               | F=1.2,<br>P=0.36        |
| Lipid   | 51 ± 0.4     | 95 ± 0.6                 | 91 ± 0.2                | 94 ± 0.2                | F=0.3,<br>P=0.73        |
| Gross energy (MJ/kg)                          | 61 ± 0.2     | 84 ± 0.2 <sup>a</sup>    | 77 ± 0.1 <sup>b</sup>   | 79 ± 0.1 <sup>ab</sup>  | F=8.0*                  |
| <i>Whole body fatty acids (%)<sup>#</sup></i> |              |                          |                         |                         |                         |
| Total FA (mg/g lipid)                         | 681.0 ± 35.1 | 616.6 ± 19.2             | 665.7 ± 68.1            | 647.7 ± 44.7            | F=0.4,<br>P=0.69        |
| 16:0  | 26.1 ± 0.2   | 0.5 <sup>a</sup>         | 0.2 <sup>a</sup>        | 0.1 <sup>b</sup>        | F=74.3***<br>F=148.8*** |
| 18:0  | 7.8 ± 0.2    | 7.4 ± 0.1 <sup>b</sup>   | 9.9 ± 0.2 <sup>c</sup>  | 6.4 ± 0.0 <sup>a</sup>  | *                       |
| 18:1  | 37.9 ± 0.8   | 0.5 <sup>a</sup>         | 0.2 <sup>a</sup>        | 0.1 <sup>b</sup>        | F=371.0***              |
| 18:2n-6                                       | 9.7 ± 0.3    | 8.8 ± 0.2 <sup>a</sup>   | 10.2 ± 0.2 <sup>b</sup> | 11.0 ± 0.1 <sup>c</sup> | F=37.4***               |
| 18:3n-3                                       | 1.0 ± 0.1    | 1.0 ± 0.0                | 1.0 ± 0.0               | 1.1 ± 0.0               | F=3.3,<br>P=0.11        |
| 20:4n-6                                       | 0.4 ± 0.0    | 0.7 ± 0.0 <sup>a</sup>   | 0.5 ± 0.0 <sup>b</sup>  | 0.5 ± 0.0 <sup>b</sup>  | F=14.7**                |
| 20:5n-3                                       | 1.4 ± 0.1    | 3.7 ± 0.3 <sup>a</sup>   | 2.6 ± 0.1 <sup>b</sup>  | 2.2 ± 0.1 <sup>b</sup>  | F=13.1**                |
| 22:5n-3                                       | 0.8 ± 0.0    | 1.6 ± 0.1 <sup>a</sup>   | 1.3 ± 0.1 <sup>ab</sup> | 1.2 ± 0.0 <sup>b</sup>  | F=5.3*                  |
| 22:6n-3                                       | 2.1 ± 0.2    | 5.5 ± 0.6 <sup>a</sup>   | 4.3 ± 0.2 <sup>ab</sup> | 3.8 ± 0.0 <sup>b</sup>  | F=6.1*                  |
| SFA   | 38.5 ± 0.2   | 39.3 ± 0.7 <sup>a</sup>  | 41.4 ± 0.4 <sup>a</sup> | 31.9 ± 0.0 <sup>b</sup> | F=102.4***              |
| MUFA  | 45.7 ± 0.5   | 37.7 ± 0.7 <sup>a</sup>  | 36.6 ± 0.1 <sup>a</sup> | 46.6 ± 0.1 <sup>b</sup> | F198.1***               |
| PUFA  | 11.0 ± 0.1   | 11.7 ± 0.3 <sup>a</sup>  | 13.2 ± 0.3 <sup>b</sup> | 13.9 ± 0.1 <sup>b</sup> | F=26.6**                |
| LC-PUFA                                       | 4.7 ± 0.2    | 11.4 ± 1.1 <sup>a</sup>  | 8.8 ± 0.4 <sup>a</sup>  | 7.6 ± 0.1 <sup>b</sup>  | F=7.7*                  |
| n-3   | 4.3 ± 0.0    | 11.9 ± 1.2 <sup>a</sup>  | 9.3 ± 0.4 <sup>a</sup>  | 8.2 ± 0.1 <sup>b</sup>  | F=7.0*                  |
| n-6   | 11.4 ± 0.1   | 11.2 ± 0.2 <sup>a</sup>  | 12.7 ± 0.2 <sup>b</sup> | 13.4 ± 0.1 <sup>b</sup> | F=38.1***               |
| <i>Liver fatty acids (%)<sup>#</sup></i>      |              |                          |                         |                         |                         |
| 16:0  | 27.8 ± 0.4   | 30.8 ± 0.4 <sup>ab</sup> | 33.7 ± 0.7 <sup>b</sup> | 27.6 ± 1.3 <sup>a</sup> | F=12.9**                |
| 18:0  | 11.5 ± 0.2   | 10.6 ± 0.2 <sup>a</sup>  | 13.9 ± 0.4 <sup>b</sup> | 9.8 ± 0.8 <sup>a</sup>  | F=17.5**                |

|         |            |                          |                         |                         |                  |
|---------|------------|--------------------------|-------------------------|-------------------------|------------------|
| 18:1    | 34.8 ± 0.3 | 31.1 ± 0.1 <sup>a</sup>  | 33.3 ± 0.8 <sup>a</sup> | 39.8 ± 0.6 <sup>b</sup> | F=59.3***        |
| 18:2n-6 | 6.4 ± 0.2  | 4.8 ± 0.2                | 4.3 ± 0.6               | 6.9 ± 1.2               | F=3.0,<br>P=0.12 |
| 18:3n-3 | 0.6 ± 0.0  | 0.5 ± 0.0                | 0.4 ± 0.1               | 0.6 ± 0.2               | F=1.5,<br>P=0.29 |
| 20:4n-6 | 0.7 ± 0.0  | 0.8 ± 0.0 <sup>a</sup>   | 0.5 ± 0.0 <sup>b</sup>  | 0.5 ± 0.0 <sup>b</sup>  | F=37.4***        |
| 20:5n-3 | 1.7 ± 0.1  | 2.7 ± 0.1 <sup>a</sup>   | 1.3 ± 0.1 <sup>b</sup>  | 1.5 ± 0.3 <sup>b</sup>  | F=18.0**         |
| 22:5n-3 | 1.1 ± 0.1  | 1.4 ± 0.1 <sup>a</sup>   | 0.7 ± 0.1 <sup>b</sup>  | 0.9 ± 0.1 <sup>b</sup>  | F=14.4**         |
| 22:6n-3 | 4.4 ± 0.1  | 4.8 ± 0.1 <sup>a</sup>   | 2.7 ± 0.2 <sup>b</sup>  | 2.8 ± 0.3 <sup>b</sup>  | F=15.0**         |
| SFA     | 42.9 ± 0.3 | 45.5 ± 0.5 <sup>ab</sup> | 50.5 ± 0.9 <sup>b</sup> | 40.5 ± 1.8 <sup>a</sup> | F=17.4**         |
| MUFA    | 40.0 ± 0.3 | 37.9 ± 0.3 <sup>a</sup>  | 38.2 ± 0.7 <sup>a</sup> | 44.8 ± 0.6 <sup>b</sup> | F=48.9***        |
| PUFA    | 8.5 ± 0.1  | 10.4 ± 0.2               | 5.5 ± 0.7               | 8.4 ± 1.2               | F=3.4,<br>P=0.10 |
| LC-PUFA | 8.6 ± 0.1  | 0.1 <sup>a</sup>         | 5.7 ± 0.4 <sup>b</sup>  | 6.3 ± 0.7 <sup>b</sup>  | F=29.6***        |
| n-3     | 8.6 ± 0.1  | 9.7 ± 0.1 <sup>a</sup>   | 5.6 ± 0.5 <sup>b</sup>  | 6.1 ± 0.9 <sup>b</sup>  | F=17.6**         |
| n-6     | 8.5 ± 0.2  | 6.8 ± 0.2                | 5.7 ± 0.7               | 8.6 ± 1.0               | F=3.9,<br>P=0.08 |

ND, not detected, N/A, not analysed, values <0.1 are reported as 0.1.

<sup>^</sup> P<0.05\*, P<0.01\*\*, P<0.001\*\*\*; One-way ANOVA, *DF* 2,6, post-hoc Tukey's HSD; Superscript letters indicate different levels of significance between treatment diets, percentage data were arcsine transformed prior to analysis..

<sup>#</sup> Refer to Table 1 for details.

**Table 6** The apparent *in vivo* fatty acid  $\beta$ -oxidation, de novo fatty acid synthesis and fatty acid bioconversion (elongation and desaturation) activity in juvenile barramundi fed experimental diets for eight weeks, deduced by the whole body fatty acid balance method. All data (n=3 per treatment) are reported as nmol/g fish/d basis.

|   | CTRL-D                          | SFA-D                         | MUFA-D                        | TEST ^        |
|---|---------------------------------|-------------------------------|-------------------------------|---------------|
| <i><math>\beta</math>-oxidation</i> #       |                                 |                               |                               |               |
| 16:0  | ND                              | ND                            | ND                            | N/A           |
| 18:0  | ND                              | ND                            | ND                            | N/A           |
| 18:1  | 54.0 $\pm$ 2.8 <sup>a</sup>     | 7.7 $\pm$ 3.9 <sup>a</sup>    | 580.4 $\pm$ 48.6 <sup>b</sup> | F=127.6***    |
| 18:2n-6                                     | 298.3 $\pm$ 16.8                | 312.5 $\pm$ 27.4              | 240.5 $\pm$ 14.5              | F=3.5, P=0.09 |
| 18:3n-3                                     | ND                              | ND                            | ND                            | N/A           |
| 20:4n-6                                     | 64.3 $\pm$ 3.3 <sup>a</sup>     | 25.8 $\pm$ 1.3 <sup>b</sup>   | 27.6 $\pm$ 0.7 <sup>b</sup>   | F=108.0***    |
| 20:5n-3                                     | 482.9 $\pm$ 23.9 <sup>a</sup>   | 183.7 $\pm$ 12.8 <sup>b</sup> | 167.7 $\pm$ 4.8 <sup>b</sup>  | F=125.0***    |
| 22:5n-3                                     | 25.7 $\pm$ 9.9                  | ND                            | ND                            | N/A           |
| 22:6n-3                                     | 631.3 $\pm$ 50.9 <sup>a</sup>   | 220.5 $\pm$ 17.3 <sup>b</sup> | 187.9 $\pm$ 3.7 <sup>b</sup>  | F=63.0***     |
| SFA   | ND                              | ND                            | ND                            | N/A           |
| MUFA  | 348.4 $\pm$ 12.9 <sup>b</sup>   | 38.5 $\pm$ 13.8 <sup>c</sup>  | 639.9 $\pm$ 60.9 <sup>a</sup> | F=69.8***     |
| PUFA  | 558.1 $\pm$ 32.0 <sup>a</sup>   | 382.9 $\pm$ 44.0 <sup>b</sup> | 492.7 $\pm$ 19.8 <sup>a</sup> | F=7.3*        |
| LC-PUFA                                     | 1623.6 $\pm$ 115.5 <sup>a</sup> | 574.1 $\pm$ 37.9 <sup>b</sup> | 508.6 $\pm$ 15.4 <sup>b</sup> | F=78.4***     |
| n-3   | 1616.3 $\pm$ 116.4 <sup>a</sup> | 560.1 $\pm$ 38.3 <sup>b</sup> | 494.7 $\pm$ 15.1 <sup>b</sup> | F=78.2***     |
| n-6   | 565.4 $\pm$ 31.1 <sup>a</sup>   | 396.9 $\pm$ 43.6 <sup>b</sup> | 506.6 $\pm$ 20.1 <sup>a</sup> | F=7.1*        |
| <i>Neogenesis</i> #                         |                                 |                               |                               |               |
| 12:0  | 188.3 $\pm$ 31.6 <sup>a</sup>   | 166.0 $\pm$ 29.8 <sup>a</sup> | 312.3 $\pm$ 22.6 <sup>b</sup> | F=7.7*        |
| <i>Elongation</i> #                         |                                 |                               |                               |               |
| 12:0  | 202.6 $\pm$ 28.8                | 184.9 $\pm$ 29.9              | 285.5 $\pm$ 22.4              | F=3.1, P=0.12 |
| 14:0  | 350.8 $\pm$ 30.0 <sup>b</sup>   | 241.4 $\pm$ 30.2 <sup>a</sup> | 363.7 $\pm$ 20.0 <sup>b</sup> | F=6.1*        |
| 16:0  | 216.8 $\pm$ 17.7 <sup>b</sup>   | 235.9 $\pm$ 15.6 <sup>b</sup> | 107.2 $\pm$ 3.6 <sup>a</sup>  | F=25.4**      |
| 20:5n-3                                     | ND                              | 13.9 $\pm$ 4.8                | 12.9 $\pm$ 1.1                | T=0.5, P=0.55 |
| <i><math>\Delta</math>-9 desaturation</i> # |                                 |                               |                               |               |
| 18:0  | 98.0 $\pm$ 22.7 <sup>b</sup>    | 288.3 $\pm$ 28.6 <sup>a</sup> | ND                            | T=5.2**       |

ND, not detected, N/A, not analysed, values <0.1 are reported as 0.1.

# Computations following Turchini et al (2007). Refer to Table 1 for details.

^ P<0.05\*, P<0.01\*\*, P<0.001\*\*\*; One-way ANOVA, *DF* 2,6, post-hoc Tukey's HSD; T-test, *DF* 4, was used to test two variables. Superscript letters indicate different levels of significance between treatment diets, percentage data were arcsine transformed prior to analysis.

**Table 7** Calculated summary of LC-PUFA flux in juvenile barramundi. All data are (n=3) reported as a percentage based on the total intake of each fatty acid.

|                      |             | CTRL-D     | SFA-D      | MUFA-D     | TEST <sup>^</sup> |
|----------------------|-------------|------------|------------|------------|-------------------|
| 22:6n-3 <sup>#</sup> | % Converted | ND         | ND         | ND         | N/A               |
|                      | % Oxidised  | 58.1 ± 5.1 | 38.6 ± 3.5 | 36.7 ± 0.4 | F=10.9*           |
|                      | % Excreted  | 1.5 ± 0.1  | ND         | 2.7 ± 0.6  | T=2.0, P=0.11     |
|                      | % Deposited | 40.4 ± 5.0 | 61.4 ± 3.5 | 60.6 ± 0.3 | F=11.4**          |
| 22:5n-3              | % Converted | ND         | ND         | ND         | N/A               |
|                      | % Oxidised  | 20.5 ± 8.0 | ND         | ND         | N/A               |
|                      | % Excreted  | ND         | ND         | ND         | N/A               |
|                      | % Deposited | 79.5 ± 8.0 | 100 ± 0.0  | 100 ± 0.0  | N/A               |
| 20:5n-3              | % Converted | ND         | 3.5 ± 1.2  | 3.6 ± 0.3  | T=0.1, P=0.96     |
|                      | % Oxidised  | 63.4 ± 3.7 | 47.3 ± 3.9 | 47.3 ± 1.1 | F=8.7*            |
|                      | % Excreted  | 1.0 ± 0.1  | ND         | 1.4 ±      | T=1.5, P=0.21     |
|                      | % Deposited | 35.6 ± 3.6 | 49.2 ± 2.7 | 47.6 ± 0.7 | F=7.8*            |
| n-3 LC-PUFA          | % Converted | ND         | 1.2 ± 0.4  | 1.2 ± 0.1  | T=0.1, P=0.96     |
|                      | % Oxidised  | 47.3 ± 5.6 | 28.6 ± 2.5 | 28.0 ± 0.5 | F=9.7*            |
|                      | % Excreted  | 0.8 ± 0.1  | ND         | 1.4 ± 0.3  | T=1.8, P=0.14     |
|                      | % Deposited | 51.8 ± 5.5 | 70.2 ± 2.1 | 69.4 ± 0.2 | F=9.2*            |

ND, not detected, N/A, not analysed, values <0.1 are reported as 0.1.

<sup>^</sup> P<0.001\*\*\* P<0.01\*\* P<0.05\*; One-way ANOVA, *DF* 2,6, post-hoc Tukey's HSD; T-test, *DF* 4, was used to test two variables; Superscript letters indicate different levels of significance between treatment diets, percentage data were arcsine transformed prior to analysis.

<sup>#</sup>Assumed that no further conversion occurs (Turchini et al 2007).