

POSTER PRESENTATION

Open Access

Center-surround interactions in a network model of layer 4C α of primary visual cortex

Christoph Metzner^{1,2*}, Achim Schweikard¹, Bartosz Zurowski³*From* Twenty Second Annual Computational Neuroscience Meeting: CNS*2013
Paris, France. 13-18 July 2013

Context integration is an ubiquitous principle in cortical processing underlying many perceptual and cognitive functions. Several neuropsychiatric disorders have been associated with an impairment of integration of context information, particularly schizophrenia [5]. One way to investigate the mechanisms underlying context processing and its impairments, is to look at context integration in the well-understood visual system. Center-surround interactions (CSI), i.e. the mutual influencing of stimuli presented in the center and in the surround of the visual or receptive field, respectively, are well established, both in animal neurophysiology and human psychophysical and neuroimaging studies.

We investigated orientation-selective center-surround interactions in a network model of the cortical layer 4C α , the input layer of V1 and the first cortical stage where CSI effects occur. The model is based on a previous one [6] and consists of 1024 excitatory and 256 inhibitory multi-compartment cells coupled via AMPA, NMDA and GABA synapses with a virtual 44 × 44 layer of lateral geniculate nucleus (LGN) processing units. First, we used moving sinusoidal gratings of varying orientation (0°-330°, step size: 30°) to determine orientation selectivity of the neurons [4]. Second, we measured non-specific center-surround suppression by increasing the size of the grating at the preferred orientation and calculating the suppression index (SI) according to [1]. Finally, we measured orientation-specific surround suppression by introducing a second grating surrounding the optimally sized center grating. We varied the orientations of the surround grating and measured the SI for each orientation.

We found orientation and direction selectivity comparable to findings from neurophysiological studies [2,4]

and those from another modeling study [4]. A high proportion (78%) of the model neurons show non-specific suppression with a mean SI of 47.21, which is comparable to animal studies in monkeys [1,3]. The orientation-selective suppression is strongest when the surround has the same orientation as the center grating and reduces with increasing angle deviation (mean SI parallel: 47.21; mean SI orthogonal: 36.72). This is in agreement with animal experiments [1] but also with human psychophysical studies [5,7].

We have presented a neuronal network model that showed non-specific and orientation-specific center-surround suppression in accordance with experimental and psychophysical data. Our model allows for investigating in detail the mechanisms supporting center-surround suppression and context-integration. This comprises changes of the perceptual input, characteristics of neurons and synapses as well as modulatory influences of GABA, for instance. Finally, our model is suited to systematically model the neural substrates of context integration impairments in neuropsychiatric disorders.

Acknowledgements

This work was partially supported by the Graduate School for Computing in Medicine and Life Sciences funded by Germany's Excellence Initiative [DFG GSC 235/1].

Author details

¹Institute for Robotics and Cognitive Systems, University of Luebeck, 23538 Luebeck, Germany. ²Graduate School for Computing in Medicine and Life Sciences, University of Luebeck, 23538 Luebeck, Germany. ³Department of Psychiatry, University Clinics Schleswig-Holstein, 23538 Luebeck, Germany.

Published: 8 July 2013

References

1. Jones H, et al: Surround suppression in primate V1. *J Neurophysiol* 2001, **86**(4):2011-2028.
2. Ringach D, Shapley R, Hawken M: Orientation selectivity in macaque v1: diversity and laminar dependence. *J Neurosci* 2002, **22**(13):5639-5651.

* Correspondence: metzner@rob.uni-luebeck.de

¹Institute for Robotics and Cognitive Systems, University of Luebeck, 23538 Luebeck, Germany

Full list of author information is available at the end of the article

3. Sillito A, et al: **Visual cortical mechanisms detecting focal orientation discontinuities.** *Nature* 1995, **378**:492-496.
4. Zhu W, Xing D, Shelley M, Shapley R: **Correlation between spatial frequency and orientation selectivity in v1 cortex: Implications of a network model.** *Vision research* 2010, **50**(22):2261-2273.
5. Yoon J, et al: **GABA concentration is reduced in visual cortex in schizophrenia and correlates with orientation-specific surround suppression.** *J Neurosci* 2010, **30**(10):3777-3781.
6. Metzner C, Schweikard A, Zurowski B: **Towards Realistic Receptive Field Properties in a Biologically Inspired Spiking Network Model of the Mammalian Primary Visual Cortex.** *Front Comput Neurosci* 2011.
7. Xing J, Heeger D: **Measurement and modeling of center-surround suppression and enhancement.** *Vision Research* 2001, **4**(5):571-583.

doi:10.1186/1471-2202-14-S1-P435

Cite this article as: Metzner et al.: Center-surround interactions in a network model of layer 4C α of primary visual cortex. *BMC Neuroscience* 2013 **14**(Suppl 1):P435.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

