

University of Huddersfield Repository

Freire, Ana C., Basit, Abdul W., Choudhary, Rahul, Piong, Chee W. and Merchant, Hamid A.

Does sex matter? The influence of gender on gastrointestinal physiology and drug delivery.

Original Citation

Freire, Ana C., Basit, Abdul W., Choudhary, Rahul, Piong, Chee W. and Merchant, Hamid A. (2011) Does sex matter? The influence of gender on gastrointestinal physiology and drug delivery. International Journal of Pharmaceutics, 415 (1-2). pp. 15-28. ISSN 03785173

This version is available at http://eprints.hud.ac.uk/29048/

The University Repository is a digital collection of the research output of the University, available on Open Access. Copyright and Moral Rights for the items on this site are retained by the individual author and/or other copyright owners. Users may access full items free of charge; copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational or not-for-profit purposes without prior permission or charge, provided:

- The authors, title and full bibliographic details is credited in any copy;
- A hyperlink and/or URL is included for the original metadata page; and
- The content is not changed in any way.

For more information, including our policy and submission procedure, please contact the Repository Team at: E.mailbox@hud.ac.uk.

http://eprints.hud.ac.uk/

Does sex matter? The influence of gender on gastrointestinal physiology and drug delivery.

Freire AC1, Basit AW, Choudhary R, Piong CW, Merchant HA.

¹Department of Pharmaceutics, The School of Pharmacy, University of London, 29-39 Brunswick Square, London, WC1N 1AX, United Kingdom.

Abstract

We all respond differently to drugs. Personalised medicine aims to improve efficacy and reduce side effects, and efforts are being made to understand the physiological differences that underlie responses to drugs. Genetics, diet and disease state can be key; however, gender also plays an important role in pharmacokinetics, pharmacodynamics and drug toxicity. Differences in metabolism and clearance of drugs as a consequence of distinct hepatic and renal processes in males and females are now much better understood but little is known about gender differences in the gastrointestinal tract. As the recipient of all orally administered medications, differences at this level can have a major impact on drug delivery and bioavailability; yet these continue to be ignored and insufficiently studied in the context of drug disposition. The aim of this review is to highlight the known gender differences in gut physiology. Clinical case studies are presented, where possible, to illustrate the influence of these differences on drug disposition and gaps in current knowledge are highlighted to encourage further research in this area.

* Correspondence at:

Email: a.basit@ucl.ac.uk hamid.merchant@hud.ac.uk

Telephone: +44 207 753 5865 +44 1484 472387 Fax: +44 207 753 5865 +44 1484 472182

FIGURES




Figure 1. Key gender differences at the level of gastrointestinal tract affecting oral drug delivery and bioavailability, refer to Table 1 for details.




Figure 2. Gender differences in iron absorption (58 Fe erythrocyte incorporation) in preadolescent subjects (n=15 each) presented as geometric mean \pm SD of the group. Figure drawn using data from Woodhead et al., 1991.




Figure 3. Percentage change in ranitidine bioavailability (excreted in urine over 24 h) in male and female volunteers (n=6 each). Figure drawn using data from Ashiru et al., 2008.




Figure 4. Median hourly intragastric acidity in 35 healthy females and 36 healthy male subjects. In the figure, **B**- Breakfast, **C**-coffee, **L**-lunch, **T**-tea, **D**-dinner, **N**-nightcap. Reproduced from Prewett *et al.*, 1991. Bar chart on the right represents Mean (± SE) basal pH in healthy subjects (252 men, 113 women). Figure drawn using data from Feldman and Barnett 1991.




Figure 5. Individual and mean (\pm SD) gastric residence time of Heidelberg capsule of 12 healthy male and 12 aged and race-matched female counterparts. Figure reproduced from Mojaverian *et al.*, 1988.




Figure 6. Proportion of total colonic transit time spent in different segments of the colon in 73 normal subjects (M = Male, F = Female). Figure redrawn from Metcalf *et al.*, 1987.




Figure 7. Mean (±SE) peak blood alcohol levels for males and females tested at different times in the menstrual cycle. Figure redrawn from Jones and Jones 1976.




Figure 8. Mean plasma R-verapamil, S-verapamil, R-norverapamil, S-norverapamil concentrationtime profiles at steady state in young and elderly subjects (men and women) following
osmotically-controlled extended release verapamil (180 mg) formulation.

△ young male, ○ elderly male, ▲ young female, ● elderly female.

Figure reproduced from Gupta et al 1995.




Figure 9. Midazolam pharmacokinetics parameters (mean \pm SD) after oral administration in male and female healthy volunteers (n=8 each) after clarithromycin administration (500 mg twice a day for 7 days). Figure drawn using data from Gorski et al 1998.

This article is published in full at:

Int J Pharm. 2011 Aug 30;415(1-2):15-28. doi: 10.1016/j.ijpharm.2011.04.069. Epub 2011 May 26.