



## Strathprints Institutional Repository

**Bradley, J. A. and Anesio, A. M. and Singarayer, J. S. and Heath, M. R. and Arndt, S. (2015) SHIMMER (1.0) : a novel mathematical model for microbial and biogeochemical dynamics in glacier forefield ecosystems. Geoscientific Model Development Discussions, 8 (8). pp. 6143-6216. , <http://dx.doi.org/10.5194/gmdd-8-6143-2015>**

This version is available at <http://strathprints.strath.ac.uk/54015/>

**Strathprints** is designed to allow users to access the research output of the University of Strathclyde. Unless otherwise explicitly stated on the manuscript, Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Please check the manuscript for details of any other licences that may have been applied. You may not engage in further distribution of the material for any profitmaking activities or any commercial gain. You may freely distribute both the url (<http://strathprints.strath.ac.uk/>) and the content of this paper for research or private study, educational, or not-for-profit purposes without prior permission or charge.

Any correspondence concerning this service should be sent to Strathprints administrator: [strathprints@strath.ac.uk](mailto:strathprints@strath.ac.uk)



This discussion paper is/has been under review for the journal Geoscientific Model Development (GMD). Please refer to the corresponding final paper in GMD if available.

# SHIMMER (1.0): a novel mathematical model for microbial and biogeochemical dynamics in glacier forefield ecosystems

J. A. Bradley<sup>1,2</sup>, A. M. Anesio<sup>1</sup>, J. S. Singarayer<sup>3</sup>, M. R. Heath<sup>4</sup>, and S. Arndt<sup>2</sup>

<sup>1</sup>Bristol Glaciology Centre, School of Geographical Sciences, University of Bristol, Bristol, BS8 1SS, UK

<sup>2</sup>BRIDGE, School of Geographical Sciences, University of Bristol, Bristol, BS8 1SS, UK

<sup>3</sup>Department of Meteorology, University of Reading, Reading, RG6 6BB, UK

<sup>4</sup>Department of Mathematics and Statistics, University of Strathclyde, Glasgow, G1 1XH, UK

Received: 9 July 2015 – Accepted: 15 July 2015 – Published:

Correspondence to: J. A. Bradley (j.bradley@bristol.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

GMDD

8, 1–74, 2015

**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

SHIMMER (Soil biogeochemical Model for Microbial Ecosystem Response) is a new numerical modelling framework which is developed as part of an interdisciplinary, iterative, model-data based approach fully integrating fieldwork and laboratory experiments with model development, testing, and application. SHIMMER is designed to simulate the establishment of microbial biomass and associated biogeochemical cycling during the initial stages of ecosystem development in glacier forefield soils. However, it is also transferable to other extreme ecosystem types (such as desert soils or the surface of glaciers). The model mechanistically describes and predicts transformations in carbon, nitrogen and phosphorus through aggregated components of the microbial community as a set of coupled ordinary differential equations. The rationale for development of the model arises from decades of empirical observation on the initial stages of soil development in glacier forefields. SHIMMER enables a quantitative and process focused approach to synthesising the existing empirical data and advancing understanding of microbial and biogeochemical dynamics. Here, we provide a detailed description of SHIMMER. The performance of SHIMMER is then tested in two case studies using published data from the Damma Glacier forefield in Switzerland and the Athabasca Glacier in Canada. In addition, a sensitivity analysis helps identify the most sensitive and unconstrained model parameters. Results show that the accumulation of microbial biomass is highly dependent on variation in microbial growth and death rate constants,  $Q_{10}$  values, the active fraction of microbial biomass, and the reactivity of organic matter. The model correctly predicts the rapid accumulation of microbial biomass observed during the initial stages of succession in the forefields of both the case study systems. Simulation results indicate that primary production is responsible for the initial build-up of substrate that subsequently supports heterotrophic growth. However, allochthonous contributions of organic matter are identified as important in sustaining this productivity. Microbial production in young soils is supported by labile organic matter, whereas carbon stocks in older soils are more refractory. Nitrogen fixing bacteria are respon-

GMDD

8, 1–74, 2015

### SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sible for the initial accumulation of available nitrates in the soil. Biogeochemical rates are highly seasonal, as observed in experimental data. The development and application of SHIMMER not only provides important new insights into forefield dynamics, but also highlights aspects of these systems that require further field and laboratory research. The most pressing advances need to come in quantifying nutrient budgets and biogeochemical rates, in exploring seasonality, the fate of allochthonous deposition in relation to autochthonous production, and empirical studies of microbial growth and cell death, to increase understanding of how glacier forefield development contributes to the global biogeochemical cycling and climate in the future.

## 1 Introduction

Ice fronts in polar and alpine regions are retreating as a result of climate warming, and as a consequence, glacier forefield areas in high latitude and high altitude regions are rapidly expanding (Graversen et al., 2008; ACIA, 2005). Glacier coverage in upland Alpine regions in Europe have declined by up to 30 % from the 1970s to 2003, exposing roughly 860 km<sup>2</sup> of previously ice-covered land area (Paul et al., 2011). Similarly, rapid glacial retreat has been observed in Iceland (Staines et al., 2014), North America (Insam and Haselwandter, 1989; Hahn and Quideau, 2013; Ohtonen et al., 1999; Sattin et al., 2009), Asia (Liu et al., 2012) and Svalbard (Moreau et al., 2008). These vast expanses of newly uncovered land have been locked under ice for tens of thousands of years, are typically highly oligotrophic, with low nutrient budgets and are subject to harsh and rapidly fluctuating environmental conditions. They have potentially significant yet largely unexplored roles on large scale biogeochemical cycling and climate (Dessert et al., 2003; Anderson et al., 2000; Smittenberg et al., 2012; Berner et al., 1983), global methane budgets (Kirschke et al., 2013), the global phosphorus cycle (Filippelli, 2002; Follmi et al., 2009) and the productivity of downstream and coastal ecosystems (Anesio et al., 2009; Mindl et al., 2007; Fountain et al., 2008; Anderson et al., 2000). Furthermore, the initial stages of microbial community development in

GMDD

8, 1–74, 2015

## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



soils is fundamental to understanding life in extreme environments, which may serve as an analogue for survival and habitability of environments currently assumed devoid of life on this planet and others.

Microbial communities are the primary colonisers of recently exposed soils, and are thought to be fundamental in soil stabilisation, the build-up of carbon and nutrient pools, creating a favourable habitat and facilitating the establishment of higher order plants (Schulz et al., 2013; Bradley et al., 2014). A conceptual overview of forefield nutrient cycling is presented in Fig. 1. Recently de-glaciated soils vary in their mineralogical and microbial compositions. Total Organic Carbon (TOC), Total Nitrogen (TN), and Total Phosphorus (TP) content of newly exposed glacial forefield soils is low, in the range of 0.1–40.0 mg C g<sup>-1</sup>, 0.1–2.0 mg N g<sup>-1</sup>, and 2–8 µg P g<sup>-1</sup> (Bradley et al., 2014). However, these typically increase by 1 to 2 orders of magnitude with ageing from newly exposed to well-developed (decades old) soil. Three distinct sources contribute carbon to recently exposed soils: autochthonous primary production by autotrophic microorganisms, allochthonous material deposited on the soil surface (from wind, hydrology, biology and precipitation) and ancient organic pools derived from under the glacier. Organic matter accumulation from all three sources supports the development of heterotrophic communities, yet their relative significance remains unknown. The continual autotrophic production, heterotrophic re-working and allochthonous deposition lead to the accumulation of organic material, which supports higher abundances and diversity of microorganisms (Bradley et al., 2014; Schulz et al., 2013). Nitrogen is derived from active nitrogen-fixing organisms, allochthonous deposition, and degradation of organic substrate. Bioavailable phosphorus is usually abundant in the topsoil or bedrock of glaciated regions from weathering of the mineral surface, and can also be liberated due to the degradation of organic molecules.

Hypotheses relating accumulation of carbon and nutrients to increasing species richness and diversification have tended to be descriptive and qualitative, rather than quantitative. Little is known about the main drivers of microbial succession and controls on abundance, diversity, and activity, and this limits our understanding of glacier forefield

---

**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



contribution to global nutrient pathways and our ability to predict how these rapidly expanding ecosystems may respond in the future, including their potential impact on atmospheric CO<sub>2</sub>, global climate, and downstream productivity.

This lack of understanding can be partly attributed to the difficulty of quantifying the different external organic matter and nutrient fluxes, as well as disentangling the complexity of biogeochemical processes underlying the microbial dynamics and soil carbon and nutrient build-up along the chronosequence by observations and/or laboratory experiments alone. Numerical models are useful tools in this context as they can not only help to disentangle the complex process interplay, diagnose the fluxes between ecosystem components and, ultimately, predict the sensitivity and response of an environment to changing environmental and climatic conditions, but can also help identify important data and knowledge gaps and hence guide the design of efficient field campaigns and laboratory studies directly targeted at closing these gaps. Nevertheless, a modelling framework that could be used to explore microbial dynamics and associated nutrient cycling in glacier forefields currently does not exist.

The development of soil models has been common in the past and important in informing soil management, policy and prediction (McGill, 2007, 1996), for example in understanding the contribution of Soil Organic Matter (SOM) to the formation of stable aggregate soils, the ease of soil cultivation, water holding characteristics, and the decreased risk of physical damage and compaction. The explicit inclusion of soil microbial dynamics has been shown to drastically improve the performance of these models (Wieder et al., 2013). There are many different types of soil models in use today across a range of scales and purposes, such as informing agricultural policy, understanding biogeochemical cycling and soil food webs, and the feedbacks between soil processes, hydrology and the atmosphere (Stapleton et al., 2005; Blagodatsky and Richter, 1998; Knapp et al., 1983; Grant et al., 1993; German et al., 2012; Ingwersen et al., 2008; Leffelaar and Wessel, 1988; Kuijper et al., 2005; Kravchenko et al., 2004; Parton et al., 1988; Garnier et al., 2001; Darrah, 1991; Foereid and Yearsley, 2004; Vandewerf and Verstraete, 1987b; Long and Or, 2005; Maggi and Porporato, 2007;

---

**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Moorhead and Sinsabaugh, 2006; Panikov and Sizova, 1996; Toal et al., 2000; Zelenev et al., 2000; Scott et al., 1995). However, although these models include an explicit microbial component, SOM models are tailored towards research questions that are focused on geochemistry and specifically organic matter dynamics rather than biology.

5 The models presented in the literature quickly become complex and thus require comprehensive data sets for set-up, calibration and evaluation. Many of their parameters cannot be constrained on the basis of information available for glacier forefield ecosystems, thus resulting in large uncertainties, making subsequent analysis difficult and results sometimes unclear (Hellweger and Bucci, 2009; Beven and Binley, 10 1992). Accordingly, their model structure is not suitable for the study of glacial forefields. Unique nutrient cycles, extreme and highly variable environmental conditions and rapidly changing compositions of microbial communities interact to form chronosequence dynamics often seen only in forefield ecosystems (Bradley et al., 2014). There is not a single model that incorporates all of the necessary conditions to model fore- 15 field development without an unacceptable level of abstraction and simplification of the system.

Therefore, we developed the new model framework SHIMMER (Soil biogeochemical Model for Microbial Ecosystem Response) to quantitatively simulate the initial stages of ecosystem development and assess biogeochemical processes in the forefield of 20 glaciers. The code is written and executed in the free open source computing environment and programming language R, which is available to download on the web (<http://www.r-project.org/>). SHIMMER is not only a new model framework, but is also part of an interdisciplinary, iterative, open-source, model-data based approach fully integrating fieldwork and laboratory experiments with model development, testing, and 25 application. More specifically, the model scope and complexity of the first version of SHIMMER is informed by: (1) a comprehensive review of glacier forefield research (Bradley et al., 2014) to determine a suitable level of complexity and identify the most important processes to resolve and (2) the data set collected during a first field campaign to investigate the coupled microbial and geochemical dynamics in the forefield

## GMDD

8, 1–74, 2015

### SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of a retreating Svalbard glacier. The model framework is aimed at identifying the dominant processes and biogeochemical controls on de-glaciated forefield ecosystem development. The model is kept as general as possible, thus is easily transferable to other microbial ecosystems such as desert soils or glacier surfaces. Simplifications are made to account for the quality and availability of observational data and processes will be resolved at greater detail once suitable data becomes available. However, the model is dynamically sufficient, i.e. the minimum processes that are needed to resolve the system and provide useful output are included. Some additional processes are included such as the ability of soil microbes to fix atmospheric nitrogen or assimilate nitrate depending on soil Dissolved Inorganic Nitrogen (DIN) levels, allochthonous substrate and nutrient inputs, varying levels of substrate bioavailability, and the microbial response to daily and seasonally varying environmental fluctuations. An important feature of the first version of the SHIMMER model is the inclusion of multiple pools of microbial biomass, each with certain functions and characteristics, to represent the diversity of species commonly found in a glacier forefield (Bradley et al., 2014). The current version of the model is designed to represent the microbial community prior to the establishment of plants, and therefore only the initial stages of chronosequence development will be assessed. Microbial communities may be heavily structured by establishing vegetation (Brown and Jumpponen, 2014), and the physical properties of vegetated soils are considerably different in terms of water retention, ultra-violet exposure, temperature fluctuations (Ensign et al., 2006; King et al., 2008), and nutrient status (Kastovska et al., 2005; Schutte et al., 2009).

This paper provides a comprehensive description of the modelling framework. The newly developed model presented here is then used to conduct an extensive parameter sensitivity study. In addition, a first model performance test is conducted on the basis of existing published data from the Damma Glacier forefield in Switzerland (Bernasconi et al., 2011; Brankatschk et al., 2011; Guelland et al., 2013b) and from the Athabasca Glacier in Canada (Insam and Haselwandter, 1989). In combination with a review of parameter values published in the literature, the results of this sensitivity study will in-

## GMDD

8, 1–74, 2015

### SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





form a series of future laboratory experiments that will help constrain the most sensitive and/or unconstrained model parameters for our study site. Furthermore, the sensitivity study and the test case studies will inform the design of additional fieldwork that will help further quantify important processes and model forcings. It is expected that the model scope and complexity will increase as more field data becomes available. In addition, it is intended that new model developments will guide and inform future field and laboratory studies so that subsequent versions of the model will run with narrower plausible ranges of parameters, and explicitly resolve processes that currently cannot be constrained on the basis of available data. SHIMMER can thus be considered as the first step of an interdisciplinary, iterative approach (illustrated in Fig. 2 and Table 1) where new data informs new model developments that will result in new insights, which in turn will inform new laboratory and field experiments and so forth. It thus not only contributes to more accurate quantitative predictions that enable a deeper understanding of the processes that control microbial communities, their role on global biogeochemical cycles and their response to climate variations, but also provides an ideal platform for the synthesis and exchange of knowledge and information across different disciplines.

The final developed model presented here is:

- Structurally (i.e. spatial resolution, number of species, processes included) and mechanistically (i.e. process formulation) complex enough to describe the required properties for carbon, nitrogen and phosphorus turnover necessary to address the questions identified in the main aims (Table 1).
- Structurally and mechanistically simple enough to constrain and validate parameters and simulation results by available data and literature.
- Able to operate with numerical efficiency on various timescales from days to decades, in order to represent an entire chronosequence development in sufficient detail.
- Applicable to a range of environments.

## GMDD

8, 1–74, 2015

### SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



– Structurally stable, conserves mass and provides robust numerical output.

## 2 Model description

The following sections provide a detailed description of SHIMMER and its implementation. A conceptual model diagram is presented in Fig. 3. The state-variables are listed in Table 2. Derived variables are included in the model to quantify mass budgets and transfers between pools. These are listed in Table 3, along with their formulation. The mathematical formulation of the model is presented in Table 4. Parameters are summarized in Table 5.

### 2.1 Physical support

The model support of SHIMMER (1.0) is 0-D; i.e. there is no specific spatial discretization (e.g. depth). This choice is informed by the quality of observational data. The model can be easily expanded into higher dimensions by including transport terms. The model represents the top centimetre of the soil surface as a homogeneous mix, and light, temperature, nutrients, organic compounds and microbial biomass are thus assumed to be evenly distributed. External nutrient inputs and leaching are additions to and subtractions from an external environment.

### 2.2 Model components

The model contains pools of microbial biomass, organic matter and both dissolved inorganic and organic nitrogen and phosphorus (Table 2). A system of coupled ordinary differential equations describes the transfers and transformations of these pools (Table 4).

## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



## 2.2.1 Microbial dynamics

In its current form, SHIMMER distinguishes between autotrophic ( $A$ ) and heterotrophic ( $H$ ) microbial communities, which are further subdivided into three categories ( $A_{1-3}$ ,  $H_{1-3}$ ) (Table 2) to account for the different metabolic needs and physiological pathways of microbial organisms commonly found in glacier forefields, but remain manageable in terms of validating the model behaviour with existing datasets.  $A_1$  and  $H_1$  represent microbes derived residually from the subglacial environment, and will be referred to as “subglacial microbes”.  $A_1$  represents chemolithoautotrophic microbes such as the genus *Thiobacillus*, and  $H_1$  represents heterotrophic subglacial microbes such as the family Comamonadaceae.  $A_2$  and  $H_2$  represent microbes commonly found in glacier forefield soils with no “special” characteristics, such as *Leptolyngbya* and *Sphingopyxis* respectively, and will be referred to as “soil microbes”. Whereas the soil-derived microbes are intended to maintain a state of high metabolic versatility, subglacial species are represented in SHIMMER as more energy conserving, and adapted to the extreme environmental conditions. In addition,  $A_3$  and  $H_3$  are able to fix atmospheric  $N_2$  gas as a source of nitrogen when DIN stocks become limiting, giving them a competitive advantage. Examples of  $A_3$  and  $H_3$  are *Nostoc* and Rhizobiales respectively, and will be referred to as “nitrogen fixing microbes” or “nitrogen fixers”. The growth of all other microbial groups depends on the availability of dissolved inorganic nitrogen and phosphorus in the soil.

The overall rate of change of autotrophic and heterotrophic communities is described by:

$$\frac{dA_i}{dt} = U_{A_i} - G_{A_i} - X_{A_i} \quad (1)$$

$$\frac{dH_i}{dt} = U_{H_i} - G_{H_i} - X_{H_i} \quad (2)$$

Microbial dynamics are thus governed by three terms: growth ( $U$ ), losses ( $G$ ), and the rate of exudate and EPS production ( $X$ ).

GMDD

8, 1–74, 2015

**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Growth $U$

Microbial biomass is created by chemolithoautotrophic ( $U_{A_1}$ ), photo-autotrophic ( $U_{A_2}$  and  $U_{A_3}$ ), and heterotrophic ( $U_{H_{1-3}}$ ) growth. Subglacial chemolithoautotrophs ( $A_1$ ) acquire their energy through chemical synthesis of mineral compounds. Autotrophic soil species ( $A_2$  and  $A_3$ ) acquire energy through photosynthesis, and accordingly require light in order to grow. Chemolithoautotrophic growth (Eq. 3) and photoautotrophic growth (Eq. 4) are described as:

$$U_{A_1} = A_1 \times T_f \times d \times \rho_{\text{sub}} \times I_{\text{max}A} \times \left( \frac{\text{DIN}}{\text{DIN} + (K_n \times k_{\text{sub}})} \right) \times \left( \frac{\text{DIP}}{\text{DIP} + (K_p \times k_{\text{sub}})} \right) \quad (3)$$

$$U_{A_2} = A_2 \times T_f \times d \times I_{\text{max}A} \times \left( \frac{L}{L + K_L} \right) \times \left( \frac{\text{DIN}}{\text{DIN} + K_n} \right) \times \left( \frac{\text{DIP}}{\text{DIP} + K_p} \right) \quad (4)$$

where autotrophic growth is calculated according to a maximum growth rate ( $I_{\text{max}}$ ) (at temperature ( $T$ ) =  $T_{\text{ref}}$ ) which is further affected by substrate availability, temperature, photosynthetically available radiation (PAR), and nutrient concentrations (Mur et al., 1999; Van Liere and Walsby, 1982). Temperature dependency is described by a temperature response factor ( $T_f$ ) with a  $Q_{10}$  formulation (Table 4) effectively slowing down or speeding up all life processes (Soetaert and Herman, 2009; Yoshitake et al., 2010; Schipper et al., 2014).  $T$  represents the temperature of the soil. The reference temperature ( $T_{\text{ref}}$ ) is the temperature at which rates ( $I_{\text{max}}$  and  $\alpha$ ) are described. The response of growth rates to temperature allows microorganisms to contend with harsh environmental conditions and promotes their overall longevity in a transient natural setting. The typical response of phototrophic growth ( $A_2$  and  $A_3$ ) to PAR is a saturation curve. Both photosynthesis and phototrophic growth are limited at low irradiance. Phototrophic activity may be inhibited at a very high irradiance, however this is not represented in the SHIMMER framework for sake of maintaining a manageable level of complexity (an inhibition term in light limitation functions typically uses two parameters (Soetaert

## GMDD

8, 1–74, 2015

### SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and Herman, 2009)). PAR sensitivity of photo-autotrophs is described by Monod kinetics, and a half-saturation constant  $K_L$  is fitted to adjust growth rate with PAR. Nutrient (nitrogen and phosphorus) limitation is also expressed as Monod kinetics, with half-saturation ( $K_N$  and  $K_P$ ) expressions.

Subglacial species ( $A_1$  and  $H_1$ ) are represented in SHIMMER as more energy conserving and adapted to harsh environmental conditions. Accordingly, their maximum growth rate ( $I_{\max}$ ) is reduced (by a factor  $\rho_{\text{Sub}}$ ), and their nutrient limitation response (expressed by the half-saturation constants  $K_S$ ,  $K_N$  and  $K_P$ ) is reduced (by a factor  $k_{\text{Sub}}$ ).

The nitrogen fixing groups ( $A_3$  and  $H_3$ ) have the ability to fix atmospheric nitrogen gas under nitrogen-limiting conditions, or assimilate DIN. Bacteria in environmental samples do not usually fix nitrogen in the presence of available DIN sources (Bottomley and Myrold, 2007; Holl and Montoya, 2005) because nitrogen fixation is an energetically expensive process (Liu et al., 2011; Cannell and Thornley, 2000; Phillips, 1980). However, even at very high concentrations of nitrate, there may not be total inhibition of nitrogen fixation, and simultaneous nitrogen fixation and nitrate assimilation may occur (Holl and Montoya, 2005). The inhibition of nitrogen fixation with DIN in SHIMMER thus follows Monod-kinetics (Table 4) (Holl and Montoya, 2005; Rabouille et al., 2006; Goebel et al., 2008). To account for the additional energy expenditure of nitrogen fixation, the growth efficiency ( $Y_A$  and  $Y_H$ ) and maximum growth rates ( $I_{\max}$ ) of nitrogen fixers whilst fixing nitrogen ( $U_{A_3-N_2}$  and  $U_{H_3-N_2}$ ) is multiplied by an adjustable reduction factor ( $n_f$ ) (Breitbarth et al., 2008; LaRoche and Breitbarth, 2005). Whilst nitrogen fixers are actively fixing atmospheric nitrogen, soil DIN concentration is not a limiting factor on their growth rate and accordingly, the growth-limiting half-saturation expression for soil DIN ( $K_n$ ) is discounted from  $U_{A_3-N_2}$  and  $U_{H_3-N_2}$ . Similarly, whilst nitrogen fixers are using  $N_2$  gas as their source of nitrogen, there is no uptake of DIN from the soil DIN pool (Table 4). Nitrogen fixation in the SHIMMER model is sensitive to many of the environmental factors often cited in the literature, including surrounding DIN concentrations, temperature, and carbon and phosphorus limitation (Liu et al., 2011).

---

## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Heterotrophic microbes ( $H_{1-3}$ ) acquire their energy through the degradation of organic substrate ( $S_1$  and  $S_2$ ) rather than by photosynthetic or chemolithoautotrophic processes. Their growth from labile ( $U_{H_2L}$ ) and refractory ( $U_{H_2R}$ ) organic matter is formulated in a similar way to Eqs. (3) and (4) but depends on the bioavailability of organic matter rather than light:

$$U_{H_2L} = H_2 \times T_f \times d \times I_{\max H} \times \left( J_{S_1} \times \frac{S_1}{S_1 + K_s} \right) \times \left( \frac{\text{DIN}}{\text{DIN} + K_n} \right) \times \left( \frac{\text{DIP}}{\text{DIP} + K_p} \right) \quad (5)$$

$$U_{H_2R} = H_2 \times T_f \times d \times I_{\max H} \times \left( J_{S_2} \times \frac{S_2}{S_2 + K_s} \right) \times \left( \frac{\text{DIN}}{\text{DIN} + K_n} \right) \times \left( \frac{\text{DIP}}{\text{DIP} + K_p} \right) \quad (6)$$

The model distinguishes between two pools of organic matter: a highly reactive pool ( $S_1$ ) and less reactive pool ( $S_2$ ). The highly reactive pool comprises highly available and fresh organic compounds that are preferentially degraded by microorganisms and therefore may respond quickly to changing external conditions and inputs. The less reactive pool represents the bulk of substrate present in the non-living organic component of soil. This organic matter degrades over longer timescales and therefore accumulates and may respond more slowly to changes in the environment. In order to express a preference of labile substrate, the parameters  $J_{S_1}$  and  $J_{S_2}$  (with  $J_{S_1} > J_{S_2}$ ) represent factors that scale the maximum rate at which labile carbon substrate ( $S_1$ ) and refractory substrate ( $S_2$ ) are utilised, respectively.

In natural environments, most microorganisms live under fluctuating conditions, and as such their growth is inhibited in response to suboptimal conditions (Cowan et al., 2004). Organisms commonly reduce their metabolic activity and lower their energy expenditure in order to endure adverse environmental conditions. Accordingly, a large fraction of microorganisms in any environmental sample are in a metabolically inactive state (Lennon and Jones, 2011). The active fraction of microbial biomass is represented by parameter  $d$ . This parameter is affixed to both the growth expression and the loss expression.

## Loss $G$

The loss terms ( $G_{A_i}$  and  $G_{H_i}$ ) represent the net removal of biomass from the living biomass pools ( $A_{1-3}$  and  $H_{1-3}$ ). These are integral measures of natural death, viral lysis and grazing, which are lumped together in order to reduce complexity, to keep the number of parameters in a feasible and manageable range, and appropriate to the current availability of data and understanding of the system. The loss terms ( $G_{A_i}$  and  $G_{H_i}$ ) are density dependant, and are also sensitive to variations in soil temperature ( $T$ ), the active fraction  $d$  and adjustable rate parameters  $\alpha_A$  (autotrophs) and  $\alpha_H$  (heterotrophs):

$$G_{A_i} = T_f \times d \times \alpha_A \times A_i^2 \quad (7)$$

$$G_{H_i} = T_f \times d \times \alpha_H \times H_i^2 \quad (8)$$

It is assumed that losses ( $G_{A_i}$  and  $G_{H_i}$ ) form insoluble microbial necromass (organic matter), comprising of organic carbon substrate ( $S$ ), organic nitrogen (ON) and organic phosphorus (OP), which enters the surrounding soil and is immediately available as substrate for heterotrophic growth.

## Exudate production $X$

Exudate production by autotrophs ( $X_{A_i}$ ) and heterotrophs ( $X_{H_i}$ ) (Eqs. 9 and 10) is proportional to growth rates (Allison, 2005), and is modulated by the exudation constants  $ex_A$  (autotrophs) and  $ex_H$  (heterotrophs):

$$X_{A_i} = ex_A \times U_{A_i} \quad (9)$$

$$X_{H_i} = ex_H \times U_{H_i} \quad (10)$$

Exudate is highly reactive; therefore these stocks directly contribute to the labile substrate pools ( $S_1$ ,  $ON_1$ ,  $OP_1$ ).

## GMDD

8, 1–74, 2015

### SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2.2.2 Organic matter dynamics

Organic matter dynamics are described by equations:

$$\frac{dS_1}{dt} = v_{\text{Sub}} \times I_{S_1} + q \times G_{A_i} + q \times G_{H_i} + X_A + X_H - U_{S_1 H_i} - W_{S_1} \quad (11)$$

$$\frac{dS_2}{dt} = v_{\text{Sub}} \times I_{S_2} + (1 - q) \times G_{A_i} + (1 - q) \times G_{H_i} - U_{S_2 H_i} - W_{S_2} \quad (12)$$

Organic matter accumulates in the soil due to additions from microbial loss ( $G_{A_i}$  and  $G_{H_i}$ ), exudate and EPS production ( $X_{A_i}$  and  $X_{H_i}$ ) and allochthonous external carbon substrate inputs ( $I_{S_1}$  and  $I_{S_2}$ ), and is depleted due to heterotrophic growth ( $U_{S_1 H_i}$  and  $U_{S_2 H_i}$ ) and leaching ( $W_{S_i}$ ). Substrate leaching ( $W_{S_i}$ ) is proportional to the mass of substrate and an adjustable parameter  $g_{S_i}$ . The coupling of substrate degradation to biomass growth ( $U_{S_1 H_i}$  and  $U_{S_2 H_i}$ ) is governed by the yield  $Y_H$ , describing bacterial growth efficiency (biomass increase per unit of carbon substrate consumed). The remainder is respired as Dissolved Inorganic Carbon (DIC). The adjustable parameter  $q$  represents the partitioning of substrate into the labile fraction ( $q$ ) and refractory fraction ( $1 - q$ ).

## 2.2.3 Nitrogen and phosphorus

The model accounts for DIN and DIP, as well as ON and OP species. The nitrogen and phosphorus components of living and dead microbial biomass are stoichiometrically coupled to microbial carbon ( $A_{1-3}$  and  $H_{1-3}$ ) and organic matter ( $S_1$  and  $S_2$ ) pools, respectively, by the N:C and P:C ratios of soil bacteria (Bernasconi et al., 2011). DIN dynamics are described as:

$$\frac{dDIN}{dt} = v_{\text{DIN}} \times I_{\text{DIN}} - \text{DIN}_{\text{Consumed}} + \text{DIN}_{\text{Released}} - W_{\text{DIN}} \quad (13)$$

DIN is consumed by microbial growth and recycled through the heterotrophic degradation of organic matter. Atmospheric DIN deposition ( $I_{\text{DIN}}$ ) and leaching ( $W_{\text{DIN}}$ ) are

## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





potentially important processes in the nitrogen cycle in forefield soils (Williams et al., 2009; Brooks and Williams, 1999; Schimel et al., 2004). Phosphorus dynamics are represented in an almost identical way (however, note the absence of atmospheric fixation).

## 3 Implementation

### 3.1 Numerical solution

The model is written and solved in the R statistical environment, an open source computing environment and programming language. Due to the non-linear and complex nature of the equations which comprise the model, they must be solved numerically rather than analytically. SHIMMER uses the adaptive time-step solver “Isoda” from the deSolve package (Soetaert et al., 2010) to calculate the numerical solution. Results are provided at daily time steps. On a standard desktop computer running R, the model usually takes less than 1 min to simulate 10 years of succession.

### 3.2 Forcing and initial conditions

The following external forcings drive and regulate the system’s dynamics:

- PAR (wavelength of approximately 400–700 nm) ( $W m^{-2}$ ).
- Snow depth (m).
- Soil temperature ( $^{\circ}C$ ).
- Allochthonous nutrient inputs ( $\mu g^{-1} d^{-1}$ ).
- Nutrient leaching rate.

**GMDD**

8, 1–74, 2015

## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The presence of snow attenuates sunlight and inhibits PAR from reaching the soil surface. This is accounted for in pre-processing of forcing data according to the equation:

$$n = n_0 e^{-mx} \quad (14)$$

whereby  $n$  is the irradiance ( $\text{Wm}^{-2}$ ),  $x$  is the snow depth (m) and  $m$  is the extinction coefficient for snow. The extinction coefficients for various types of snow can be measured and an estimate of 6 is used in this instance to represent snow in glacier forefields (Greenfell and Maykut, 1977).

External forcing and initial conditions for the Damma Glacier and Athabasca Glacier are presented in Fig. 4 and Table 2, respectively. They are directly informed from field studies (Sect. 4.3.1 and 4.3.2). The sensitivity and uncertainty analysis is forced with data from the Damma Glacier, since results can be interpreted alongside the more detailed contextual observations.

## 4 Model runs

### 4.1 Nominal

The model is run with nominal parameters for a period representing 75 years of succession (the approximate length of the test datasets), starting on the 1 January, in order to provide a baseline model output from which parameters are varied to determine sensitivity. Model output is available in a number of forms: daily output from the entire model run, the “sample day” of each year (7 September/day 244), cumulative totals, rates of production and final year values. Leap years are ignored. The model is forced with meteorological data collected from the Damma Glacier, Switzerland (Case study 1 – see Sect. 4.3.1 for details).

## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 4.2 Sensitivity and uncertainty analysis

A sensitivity study involving 24 model parameters is carried out to assess the stability of model output given variation in all key parameters. This is important when accounting for uncertainty, since high sensitivity of key parameters that have a relatively wide plausible range of values would lead to large uncertainties in predictions. Sensitivity analysis is considered across all state variables to assess the extent to which parameter variation influences whole model behaviour or only single variables. The model is run for 75 years under each sensitivity scenario, starting on 1 January, with daily output. The following model output ( $X$ ) is explored:

- Total autotrophic biomass (average over the entire simulation run).
- Total heterotrophic biomass (average over the entire simulation run).
- Total C substrate (average over the entire simulation run).
- DIN (average over the entire simulation run).
- DIP (average over the entire simulation run).
- Total ON (average over the entire simulation run).
- Total OP (average over the entire simulation run).
- Total nitrogen fixed (cumulative).
- Seasonal variation in microbial biomass (final year of simulation).

Plausible ranges for model parameter values are constrained from values in published literature (Table 5). Many of the parameters show considerable variation, but the most confident values (their applicability to the glacier forefield system, the method by which the value was determined, and their frequency in the literature) are used as nominal values. To explore sensitivity, uncertainty and linearity, plausible ranges are

GMDD

8, 1–74, 2015

**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



split into tenths, and simulations are run sequentially through all eleven possible values for each parameter.

Sensitivity around nominal values is quantified using a variation on the method presented in Xenakis et al. (2008). The relative sensitivity ( $\lambda$ ) of a certain model output ( $X$ ) to a parameter ( $p$ ) is estimated according to:

$$\lambda(X, p) = \frac{p}{X} \times \frac{\delta X}{\delta p} \quad (15)$$

where  $p$  is the nominal value for the parameter,  $X$  is the model output from nominal parameter values,  $\delta p$  is the difference in parameter value either side of the nominal value, and  $\delta X$  is the change in model output (simulated over the range of parameters identified in  $\delta p$ ). The resulting estimate for sensitivity ( $\lambda$ ) quantifies the relation between the model output and variation in a single parameter as a first derivative of their relationship either side of the nominal value, and is normalised based on the magnitude of parameter and model output values. Thus,  $\lambda$  indicates the sensitivity of model output to parameter variation and also the direction (sign) of the change. A positive  $\lambda$  value indicates that an increase in the parameter value yields an increased value in the model output, whereas a negative  $\lambda$  value indicates that an increase in the parameter causes a decrease in the value of the model output. Values of  $\lambda$  further from zero indicate that the model output is highly sensitive to variation in the parameter.

The calculation of  $\lambda$  may yield a “false-negative” result (i.e. a value close to zero) when the variation in model output either side of the nominal value has an opposite sign (i.e. a parabolic relationship between the parameter value and model output). In this instance, a  $\lambda$  value close to zero would indicate little sensitivity, however the model may actually be highly sensitive. To test for this “false-negative”, model output with variation in a parameter is assessed graphically. Either a linear or non-linear relationship is attributed to the change in model output in response to the change in parameter value, based on visual assessment of each plot (e.g. Fig. 5). Non-linear results are assessed to determine if the highest sensitivity is around the nominal parameter value, since this

## GMDD

8, 1–74, 2015

### SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



has implications in interpreting the model output. If parameters are most sensitive near to the nominal values, there is a higher potential variation in model output and therefore potentially greater uncertainty in interpreting results.

To explore uncertainty ( $\emptyset$ ) associated with each parameter, the percentage variation in model output is calculated according to:

$$\emptyset = \frac{X_{\max} - X_{\min}}{X} \times 100 \quad (16)$$

where  $X_{\max}$  and  $X_{\min}$  are the highest and lowest values for model output over the entire plausible range in parameter variation, and  $X$  is the model output given nominal parameters.

### 4.3 Optimization

Model parameters implicitly account for all processes that are not explicitly accounted for in the model and, therefore, may vary across different environments. Based on the outcome of the sensitivity and uncertainty analysis, parameters are adjusted in an optimisation exercises to improve model fit to the validation datasets. Parameters are varied incrementally to determine the effect on the accumulation of microbial biomass and averaged squared error is calculated with each model run. Parameters  $J_{S_1}$  and  $J_{S_2}$  (the relative bioavailability of labile and refractory substrate) are artefacts of the SHIMMER modelling structure. Therefore, these two parameters are the primary free parameters, which are adjusted to reduce mean squared error. Once a known optimum range for these parameters has been determined, the parameters that yield the highest sensitivity, uncertainty and non-linearity are adjusted. Parameters for which we have a high degree of confidence (narrow plausible ranges, lower sensitivity, linear behaviour and low uncertainty) are not adjusted in the optimisation exercise, since even relatively large changes in their value would cause only a small change in model output.

Given the wealth of physical, biological, genomic and chemical data available for the Damma Glacier, the focus of the analysis of model dynamics is on this dataset. The

## GMDD

8, 1–74, 2015

### SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Damma Glacier dataset is unique in this sense. However, data from the Athabasca Glacier forefield provides additional support that the model can respond dynamically to predict the development of soils from a range of environments and study sites. This dataset is more representative of the type of data that is typically available for deglaciated forefields.

### 4.3.1 Case study 1: Damma Glacier, Switzerland

Published data sets of the biogeochemical development of the Damma Glacier forefield in Canton Uri, Switzerland (46.6° N, 8.5° E) are used to test and validate the model, and explore detailed model behaviour (Bernasconi et al., 2011). Over the last two decades, plant and microbial succession at this site has been extensively studied, because of its gradual retreat and the resulting ideal study conditions. Comprehensive data sets have been collected as part of the BigLink project (Bernasconi et al., 2011), with further detailed studies on nutrient cycling (Brankatschk et al., 2011; Bernasconi et al., 2011; Guelland et al., 2013a; Göransson et al., 2014; Smittenberg et al., 2012; Tamburini et al., 2012), microbial community composition (Duc et al., 2009; Frey et al., 2013; Lazzaro et al., 2012; Meola et al., 2014; Sigler and Zeyer, 2002; Zumsteg et al., 2012), soil depths (Rime et al., 2014), and soil activity (Göransson et al., 2011; Guelland et al., 2013b; Zumsteg et al., 2011) across the forefield. Therefore, the site is highly appropriate to gain insight into model behaviour and various biogeochemical processes.

The forefield chronosequence is roughly 650 m long and represents a range of soil ages from zero years old to around 120 years old (Brankatschk et al., 2011). The underlying bedrock is mainly granitic gneiss (Frey et al., 2010) with a silty, sandy soil texture (Lazzaro et al., 2009). The site has a northeast exposition (Bernasconi et al., 2011) and an inclination of 25% (Sigler and Zeyer, 2002). Soil pH decreases from pH 5.1 in initial soils (10 years ice free) to 4.1 in developed soils (ice free for 2000 years) and water holding capacity increases from 26 to 33% (Brankatschk et al., 2011). Recently exposed sites at the Damma Glacier (ice-free for 6 to 13 years) are characterised by mostly unvegetated, sandy-silty sediment, gravel and large rocks. Intermediate soils

# GMDD

8, 1–74, 2015

## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(ice-free for 60 to 80 years) are characterised by increasing vegetation cover and soil structure resembling a typical soil profile. The old sites (ice-free for roughly 120 years) are fully vegetated, with clearly visible soil horizons (Bernasconi et al., 2011). Molecular characterisation suggests that both specialised heterotrophs ( $\alpha$ -,  $\beta$ -,  $\gamma$ -*Proteobacteria*), autotrophs (*Cyanobacteria*) and other nitrogen fixing species are found in all samples from all ages (Duc et al., 2009). There is a clear increase in TOC with soil age along the Damma Glacier forefield (Bernasconi et al., 2011) from around  $700 \mu\text{gCg}^{-1}$  in recently exposed soils to around  $30\,000 \mu\text{gCg}^{-1}$  in developed soils. Similarly, microbial biomass, TN and phosphorus increase by roughly an order of magnitude from recently exposed soils to developed soils (Bernasconi et al., 2011; Göransson et al., 2011).

The model is evaluated using least-squares error against 4 chemical analyses presented in the BigLink dataset (Bernasconi et al., 2011):

- Total microbial biomass ( $A_1 + A_2 + A_3 + H_1 + H_2 + H_3$ ): presented as  $C_{\text{mic}}$ .
- Carbon substrate ( $S_1 + S_2$ ): calculated as  $\text{TOC} - C_{\text{mic}}$ .
- ON ( $\text{ON}_1 + \text{ON}_2$ ): calculated as  $\text{TN} - N_{\text{mic}}$ .
- Available DIP: presented as  $P_{\text{resin}}$ .

Observational data was collected on 7 September (day 244 of the year), and is therefore compared to model output from day 244 of each year. The omission of sites older than 77 years (due to vegetation influence) leaves 16 samples ranging from 5 to 77 years since ice retreat. The 5 year data is used as initial conditions, leaving three data-points in the “early soils” category and 12 from later stages of succession where there is relatively high plat abundance. A secondary dataset of DIP accumulation complements the BigLink dataset (Göransson et al., 2011). Least-square error calculation and minimisation of errors are done only on those data points. The remaining data points from the later stages of succession are used as a test to see if microbial abundance in the late stages of chronosequence development falls within a plausible range, given the scatter in the observational data.

**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion







## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



is prescribed evenly over 10 days when there is significant snowmelt: days 158–167 (7–16 May). DIP is typically liberated by rock weathering, however Frey et al. (2010) analysed the minerals liberated from the weathering of the granitic Damma Glacier bedrock material and did not find any traceable amounts of phosphorus. However, different mineralogy is likely to considerably alter the importance of rock weathering as a source of phosphorus between locations, increasing the uncertainty for the amount of DIP generated by weathering processes. The annual input of DIP is prescribed as  $80 \mu\text{g P cm}^{-2} \text{ yr}^{-1}$  (equal to DIN input), but this release is spread evenly over the first snow-free months of each year, from day 167–206 (16 June–25 July). Prescribed allochthonous inputs are presented in Table 6. The proportion of the allochthonous nutrient input that is available to the soil represented by the model is adjusted by parameters  $v_{\text{Sub}}$  (for all substrate pools),  $v_{\text{DIN}}$  (for DIN) and  $v_{\text{DIP}}$  (for DIP).

### 4.3.2 Case study 2: Athabasca Glacier, Canada

Published data from the Athabasca Glacier forefield, Canada (52.2° N, 117.2° W) are used as a second case study in the validation exercise (Insam and Haselwandter, 1989). The Athabasca Glacier forefield is a high altitude (2740 m) site with soil ages from 5 to 225 years. The mineralogy is medium textured, mostly calcareous and neutral to slightly alkaline pH, with relatively slow plant colonisation (Insam and Haselwandter, 1989). The Athabasca glacier forefield is less intensively studied, and accordingly there is less contextual information on the biogeochemical development of soils than the Damma Glacier. However, the soils in the earlier stages of development (< 100 years) provide a robust test of model behaviour and underlying system dynamics due to the sparseness of vegetation and lack of interference in the microbial signal from vascular plants.

The model is evaluated using least-squares error against two observed bulk biogeochemical variables (Insam and Haselwandter, 1989):

- Total microbial biomass ( $A_1 + A_2 + A_3 + H_1 + H_2 + H_3$ ).

– Carbon substrate ( $S_1 + S_2$ ): calculated as  $C_{\text{org}} - \text{Microbial Biomass}$ .

Observational data was collected in July and is compared to model output from day 196 of each year. Sites older than 50 years should be interpreted cautiously due to the influence of establishing vegetation. The 5 year data is used as initial conditions. Initial microbial biomass is assumed to be evenly distributed between all microbial groups of autotrophs and heterotrophs, and initial substrate bioavailability is assumed to be 30% labile and 70% refractory. Since there are no quantitative estimates of DIN, DIP, ON and OP, initial inorganic nutrient concentrations are assumed to follow the same ratio as the Damma Glacier case study, and organic material follows a stoichiometric ratio (Bernasconi et al., 2011). Annual profiles of monthly average soil temperature (at 5 cm depth) and snow depth are obtained from published literature (Achuff and Coen, 1980) and linearly interpolated to provide daily forcing data (Fig. 4). Daily solar irradiance data from 2014 is obtained from the Alberta Agriculture and Rural Development Agroclimatic Information Service for a nearby meteorological station (Stavely AAFC, 1360 m, 50.2° N, 113.9° NW). These are repeated year-on-year for the duration of the model run. There is no observational, experimental or modelled data of sufficient quality to provide forcings of allochthonous inputs to the Athabasca Glacier forefield. Therefore estimations from the Damma Glacier are used and parameters  $v_{\text{Sub}}$ ,  $v_{\text{DIN}}$  and  $v_{\text{DIP}}$  are adjusted.

As with Case study 1, optimisation is carried out based on the results of the sensitivity study, and a minimisation of least squared error and visual fit to data is carried out based on numerous model runs varying parameters that were identified in the sensitivity test.

## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



## 5 Results and discussion

### 5.1 Nominal

The model behaviour under nominal parameters, and forced with meteorological data and initial conditions from the Damma Glacier is presented in Fig. 6. Total microbial biomass is initially stable at roughly  $3.4 \mu\text{gCg}^{-1}$  (year 1), followed by an exponential growth phase to  $46.8 \mu\text{gCg}^{-1}$  (year 15), and then a decline to near-steady-state around  $31.0 \mu\text{gCg}^{-1}$ , varying seasonally by roughly  $12.0 \mu\text{gCg}^{-1}$ . By the final year of simulation (year 75), the community has evolved (from an even split 16.7% per pool) such that the most dominant pool is soil autotrophs ( $A_2 = 35.8\%$ ), followed by nitrogen fixing autotrophs ( $A_3 = 27.4\%$ ) and subglacial chemoautotrophs ( $A_1 = 12.3\%$ ), with all heterotrophic biomass ( $H_{1-3}$ ) making up the remaining 24.5%. Total bacterial production rises steadily from  $0.3 \mu\text{gCg}^{-1}\text{yr}^{-1}$  (year 1) to  $114.2 \mu\text{gCg}^{-1}\text{yr}^{-1}$  (year 15), after which (year 31 onwards) bacterial production declines by roughly a half. Autotrophs are consistently the highest producers, responsible for between 72.6 to 89.2% of the total bacterial production, and heterotrophs are responsible for 10.8 to 27.4% of the total bacterial production.

There is a steady accumulation of carbon substrate throughout the entire simulation, from  $735.4 \mu\text{gCg}^{-1}$  (year 1) to  $4129.2 \mu\text{gCg}^{-1}$  (year 75). Labile carbon substrate accumulates during the first 7 years of succession from  $290.0 \mu\text{gCg}^{-1}$  to  $337.5 \mu\text{gCg}^{-1}$ , after which (years 7 to 20) there is a sharp decline to a minimum of  $129.0 \mu\text{gCg}^{-1}$  (year 20), as substrate stocks become more refractory (39.4% labile in year 1 to 3.7% in year 75). ON and OP follow similar dynamics to carbon substrate ( $S_1$  and  $S_2$ ). The accumulation of substrate is derived from autotrophic activity, the build-up of microbial necromass and allochthonous deposition. DIN and DIP accumulate during the first 14 years of the simulation. Following the initial increase, DIN stocks continue to accumulate and DIP experiences a decline.

## GMDD

8, 1–74, 2015

### SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 5.2 Sensitivity and linearity

### 5.2.1 Sensitivity

Sensitivity analysis is presented in Fig. 7. The accumulation of autotrophic and heterotrophic biomass is most sensitive to variation in the  $Q_{10}$  value ( $\lambda \geq 0.70$ ). Biomass accumulation is also highly sensitive to adjustments in the active fraction ( $d$ ) ( $\lambda = -0.55$  and  $\lambda = -0.52$  respectively), the bioavailability of refractory substrate ( $J_{S_2}$ ) ( $\lambda = 0.44$  and  $\lambda = 0.64$  respectively), the partitioning of microbial necromass into labile and refractory pools ( $q$ ) ( $\lambda = 0.31$  and  $\lambda = 0.45$  respectively) and microbial growth rates ( $I_{\max}$ ) ( $-0.55 \leq \lambda \leq 0.67$ ). Biomass accumulation is moderately sensitive to death rates ( $\alpha_A$  and  $\alpha_H$ ), the efficiency of heterotrophic growth ( $Y_H$ ), and the allochthonous substrate input ( $v_{\text{Sub}}$ ) ( $\pm 0.15 \leq \lambda \leq \pm 0.41$ ). Biomass accumulation has relatively low sensitivity ( $\lambda \leq \pm 0.15$ ) to variation in half saturation constants ( $K_L$ ,  $K_S$ ,  $K_N$  and  $K_P$ ), parameters affecting only the dynamics of subglacial microbes ( $A_1$  and  $H_1$ ) ( $p_{\text{Sub}}$  and  $k_{\text{Sub}}$ ) and nitrogen fixers ( $A_3$  and  $H_3$ ) ( $n_f$ ,  $K_{N2}$ ), the bioavailability of labile substrate ( $J_{S_1}$ ), exudate rates ( $ex_A$  and  $ex_H$ ), the input of allochthonous nutrients ( $v_{\text{DIN}}$  and  $v_{\text{DIP}}$ ) and the efficiency of autotrophic growth ( $Y_A$ ). Variation in the half saturation for nitrogen ( $K_N$ ) and phosphorus ( $K_P$ ) causes little change to the accumulation of biomass ( $-0.02 \leq \lambda \leq 0.00$  and  $-0.05 \leq \lambda \leq 0.03$  respectively), but has a proportionally large effect on the accumulation of DIN, DIP and total nitrogen fixation ( $0.22 \leq \lambda \leq 0.95$ ). Similarly, the reduction of efficiency and growth rates ( $n_f$ ) for nitrogen fixers whilst fixing nitrogen (rather than assimilating DIN) has a relatively minor effect on the accumulation of biomass ( $\lambda \leq \pm 0.09$ ) but strongly affects DIN, DIP and total nitrogen fixed ( $\lambda = 0.96$ ,  $\lambda = -0.96$  and  $\lambda = 0.60$  respectively).

Microbial communities alter their metabolic state (through the  $Q_{10}$  formulation) to persist during long periods of cold. At cold temperatures typical of glacier forefield environments (soil temperatures  $< T_{\text{ref}}$ ),  $T_f$  is reduced by increasing  $Q_{10}$ , which simultaneously reduces growth rates and death rates. As a result, total bacterial production decreases. Death rates also decline with reduced production, leading to an increase in

## GMDD

8, 1–74, 2015

### SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



microbial biomass over the entire duration of the simulation ( $\lambda > 0$ ). This is supported by experimental evidence that microbial communities from cold environments typically have a higher  $Q_{10}$  response to temperature variation (Schipper et al., 2014) to promote the survival of biomass under prolonged periods of harsh environmental conditions.

Heterotrophic production is critical in supporting the overall establishment of biomass and activity of the entire microbial community. Increasing the maximum growth rate of heterotrophs ( $I_{\max H}$ ) leads to a substantial increase in both heterotrophic and autotrophic biomass ( $\lambda = 0.47$  and  $\lambda = 0.67$  respectively). However, strikingly, an increase in the maximum growth rate of autotrophs ( $I_{\max A}$ ) has the net effect of lowering autotrophic and heterotrophic biomass ( $\lambda = -0.36$  and  $\lambda = -0.55$  respectively). Autotrophic growth is responsible for the build-up of substrate during the initial stages of soil development, thus supporting heterotrophic production. However, the dominance of autotrophic communities rapidly consumes available nutrients, at the expense of heterotrophs. If autotrophs are given a competitive advantage (i.e. increasing  $I_{\max A}$ ), their rapid growth increases the consumption of the essential nutrients DIN ( $\lambda = -0.57$ ) and DIP ( $\lambda = -1.58$ ), and heterotrophic growth becomes nutrient-limited. However, if heterotrophs are given a competitive advantage (i.e. increasing  $I_{\max H}$ ), a higher rate of heterotrophic production is supported and substrate is degraded more rapidly, recycling DIN ( $\lambda = 0.76$ ) and DIP ( $\lambda = 0.38$ ). This is an effective positive feedback effect, whereby the additional nutrients recycled from rapid heterotrophic degradation are able to support the growth of all microbial populations (including autotrophs ( $\lambda = 0.47$ )) in this oligotrophic and relatively nutrient poor environment. Similarly, increasing autotrophic death rates ( $\alpha_A$ ) reduces competition for nutrient resources whilst also increasing the availability of degradable organic matter (autotrophic necromass), thereby increasing heterotrophic biomass ( $\lambda = 0.24$ ).

An increase in the heterotrophic growth efficiency ( $Y_H$ ) leads to an overall more rapid accumulation of heterotrophic biomass ( $\lambda = 0.14$ ). Increasing the heterotrophic growth efficiency suggests that for each mole of organic carbon degraded, the uptake of nutrients by heterotrophic biomass is higher. This decreases nutrient availability, and ac-

---

**SHIMMER (1.0):  
a novel mathematical  
model**J. A. Bradley et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

cordingly increasing  $Y_H$  significantly lowers the accumulation of autotrophic biomass ( $\lambda = -0.41$ ). A decrease in heterotrophic growth efficiency effectively means that for heterotrophs to grow by the same amount, they must degrade more organic matter per mole of carbon incorporated in biomass, thereby increasing the recycling of nutrients, which in turn supports microbial growth in all communities. Previous modelling studies have consistently found microbial growth efficiency ( $Y$ ) to be the most sensitive parameter in determining overall biomass accumulation and the compartmentalisation of carbon between biotic and abiotic pools (Blagodatsky and Richter, 1998; Ingwersen et al., 2008; Toal et al., 2000).

## 5.2.2 Linearity

Linearity of parameter sensitivity is explored qualitatively over the range of parameter values by plotting the change in model output and by visually fitting a linear or polynomial relationship to the trend. Sensitivity and linearity evaluation have important implications in model optimisation. The highest degrees of freedom can be given to those parameters with fairly low sensitivity and a high degree of linearity ( $Q_{10}$ ,  $ex_A$ ,  $ex_H$ ,  $K_L$ ,  $K_S$ ,  $K_N$  and  $K_P$ ), since the parameters affect model output minimally and in a reasonably predictable way (illustrated in Fig. 5a). The parameters  $v_{Sub}$ ,  $p_{Sub}$ ,  $q$  and  $Y_H$  also behave fairly linearly. Changes in model output respond non-linearly to changes in  $I_{maxA}$ ,  $I_{maxH}$ ,  $J_{S_2}$ ,  $d$ ,  $\alpha_A$ ,  $\alpha_H$ ,  $v_{DIN}$  and  $v_{DIP}$ . Therefore, the sensitivity value  $\lambda$  is likely to be subject to change when looking at values away from the nominal parameter value, as illustrated for  $I_{maxA}$  in Fig. 5b. For non-linear parameters  $\alpha_A$ ,  $\alpha_H$ ,  $v_{DIN}$  and  $v_{DIP}$ , the maximum sensitivity is found in the region near the nominal value. This means that slight changes in parameter values will greatly affect the model output. The reverse is true for parameters  $I_{maxA}$ ,  $I_{maxH}$ ,  $J_{S_2}$ , and  $d$ . These parameters behave fairly linearly with comparatively little sensitivity around the nominal value, however their sensitivity increases with distance from the nominal value, leading to substantial changes in model output. This may give the impression of stability around the nominal values however a tipping point may be reached when parameters deviate too much from the nominal value. As

## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



such, non-linear parameters must be given due caution in optimisation exercises since changes in these parameters may yield unexpected model behaviour.

### 5.3 Uncertainty

Uncertainty is evaluated over the entire plausible parameter range and results are presented in Fig. 9. The parameters that yield the highest uncertainty in the accumulation of autotrophic and heterotrophic biomass are  $I_{\max H}$  (397 and 685 % respectively) and  $J_{S_2}$  (333 and 606 % respectively). A high degree of uncertainty ( $\geq 60\%$ ) also results from variation in  $\alpha_A$ ,  $\alpha_H$ ,  $I_{\max A}$ ,  $q$ ,  $Y_H$ ,  $d$ ,  $v_{\text{Sub}}$ ,  $v_{\text{DIP}}$ . This is due to a combination of high sensitivity and wide plausible ranges of parameter values. There is minimal uncertainty ( $\leq 30\%$ ) resulting from variation in parameters  $ex_A$ ,  $ex_H$ ,  $\rho_{\text{Sub}}$ ,  $k_{\text{Sub}}$ ,  $K_L$ ,  $K_N$ ,  $K_P$ ,  $K_{N_2}$ ,  $J_{S_1}$ ,  $Y_A$ , and  $v_{\text{DIN}}$ . Parameters that yield high uncertainty in the accumulation of biomass also tend to cause high degrees of uncertainty in other model outputs, most notably the accumulation of DIN and DIP, the total nitrogen fixed and the seasonality.

Measurements of bacterial growth are fundamental to most aspects of microbial ecology. Consequently, there are many estimates for  $I_{\max A}$  and  $I_{\max H}$  from literature and related modelling studies, however they span over an order of magnitude ( $0.24$  to  $4.80 \text{ d}^{-1}$ ) (Mur et al., 1999; Van Liere and Walsby, 1982; Frey et al., 2010; Ingwersen et al., 2008; Knapp et al., 1983; Zelenev et al., 2000; Stapleton et al., 2005; Darrah, 1991; Blagodatsky and Richter, 1998; Vandewerf and Verstraete, 1987a; Foereid and Yearsley, 2004; Toal et al., 2000; Scott et al., 1995), greatly increasing the uncertainty associated with  $I_{\max A}$  and  $I_{\max H}$  (average uncertainty = 265 and 693 % respectively). Maximum growth rates have been experimentally measured for soils from the Damma Glacier, with a value roughly in the middle of the plausible range (Frey et al., 2010), however there is inherent abstraction when incorporating laboratory measurements into models because of the assumptions and simplifications in model design.

Microbial death rates ( $\alpha_A$  and  $\alpha_H$ ) are difficult and problematic to define experimentally (Toal et al., 2000), and there is a great deal of variation in how losses from microbial biomass are modelled. Death rates yield moderate sensitivity in model outputs

**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





---

**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



( $\lambda \leq -0.13$  and  $\lambda \geq 0.24$ ), and very high uncertainty (77 to 178 %), as a consequence of the large plausible parameter range, which spans almost two orders of magnitude (0.006 to  $0.48 \text{ d}^{-1}$ ). Furthermore, the transferability of microbial death rate constants is compromised by the different mathematical formulations used to describe death rates (e.g. constant fixed rate: German et al., 2012; Grant et al., 1993; Scott et al., 1995; Toal et al., 2000; logistic: Boudreau, 1999; Kravchenko et al., 2004; and variable depending on model conditions: Knapp et al., 1983; Blagodatsky and Richter, 1998; Ingwersen et al., 2008; Zelenev et al., 2000; Lancelot et al., 2005).

The allochthonous deposition of organic matter and nutrients is a source of considerable uncertainty, as plausible parameter ranges for  $v_{\text{Sub}}$ ,  $v_{\text{DIN}}$  and  $v_{\text{DIP}}$  are very wide and the model output is highly sensitive to variation in these parameters. Analysis of the BACWAVE model (Zelenev et al., 2000) found high sensitivity of the spatial and temporal response of bacterial populations to changes in allochthonous carbon sources in soil. Published literature provides estimates of snow nutrient concentrations at the Damma Glacier (Brankatschk et al., 2011), however the fate of these nutrients once deposited on the soil surface is largely unknown. Accordingly, the plausible parameter ranges for  $v_{\text{Sub}}$ ,  $v_{\text{DIN}}$  and  $v_{\text{DIP}}$  are wide, and the resulting uncertainty is very high (75 to 2503 %). Furthermore, variation in parameters  $v_{\text{DIN}}$  and  $v_{\text{DIP}}$  results in highly non-linear behaviour, with comparatively large changes resulting from small changes in the parameter value. However, the effect of  $v_{\text{DIN}}$  on the accumulation of autotrophic and heterotrophic biomass is minimal (5 and 9 %), due to the substantial inputs of nitrogen from nitrogen fixers throughout the development of the forefield.

The relative bioavailability of labile and refractory substrate, as well as the partitioning of necromass and allochthonous substrate between these two pools is an important feature of the SHIMMER model. Adjustment in  $q$ ,  $J_{S_1}$  and  $J_{S_2}$  causes considerable variation in the accumulation of autotrophic and heterotrophic biomass ( $0.07 \leq \lambda \leq 0.64$ ) and relatively large uncertainty (4 to 606 %). The partitioning of the reactive continuum of natural organic matter compounds into a limited number of substrate pools is, although common practice, artificial, and a meaningful value cannot be determined ex-



perimentally or from literature (e.g. Arndt et al., 2013). Therefore, these parameters are ideal for initial tuning and optimisation exercises.

Estimation of the inhibition of nitrogen fixation with DIN ( $K_{N_2}$ ) is based on published literature and a relatively large range is considered in the uncertainty analysis ( $56.4 \leq K_{N_2} \leq 1128 \mu\text{g N g}^{-1}$ ) (LaRoche and Breitbarth, 2005; Rabouille et al., 2006; Holl and Montoya, 2005). The total nitrogen fixed and total accumulation of DIN is fairly sensitive to variation in  $K_{N_2}$  ( $\lambda = 0.08$  and  $\lambda = 0.11$  respectively), however the resulting uncertainty in the overall accumulation of autotrophic and heterotrophic biomass is low (3 and 1 % respectively). Therefore, although this parameter is not well defined, its importance is outweighed by other much more sensitive parameters such as the active fraction ( $d$ ) and microbial death rates ( $\alpha_A$  and  $\alpha_H$ ).

Many of the fairly well constrained parameters result in low uncertainty values. This can be explained by relatively tight parameter bounds explored in the uncertainty analysis, or relatively low sensitivity of model output to variation in this parameter, or a combination of both of these factors. For example, exudate production rates ( $ex_A$  and  $ex_H$ ) (uncertainty  $\leq 2\%$ ) have relatively tight parameter bounds (Allison, 2005), as well as low overall sensitivity ( $-0.01 \leq \lambda \leq 0.02$ ). Half-saturation constants for carbon substrate ( $K_S$ ) have been estimated experimentally (Vandewerf and Verstraete, 1987a, b; Blagodatsky et al., 1998; Anderson and Domsch, 1985) and fitted using models (Darrah, 1991; Ingwersen et al., 2008; Stapleton et al., 2005), with values between 50 and  $1000 \mu\text{g C g}^{-1}$ . This variation can be attributed to differences in experimental technique and medium of substrate used, and differences in model structure and optimisation, however overall model sensitivity is low ( $\lambda = -0.07$ ), as is reflected in other models (Ingwersen et al., 2008; Blagodatsky and Richter, 1998).

---

**SHIMMER (1.0):  
a novel mathematical  
model**J. A. Bradley et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## 5.4 Optimization

### 5.4.1 Case study 1: Damma Glacier

Model parameters are optimised to obtain the best possible fit to observational data from the Damma Glacier, Switzerland. Figure 10 shows the comparison of optimized simulation results with observational data. Using the optimal parameter set (Table 5), total microbial biomass increases rapidly during the initial stages of the soil development from  $3.5 \mu\text{g C g}^{-1}$  (year 1) to a peak of  $61.6 \mu\text{g C g}^{-1}$  (year 12), followed by a decline over the following 10 years to roughly  $37.0 \mu\text{g C g}^{-1}$  where it remains fairly steady for the later stages of succession (years 30 to 75). The microbial community evolves from an even split between all six pools of microbial biomass (16.7%) to a community dominated by autotrophs, with maximum autotrophic biomass ( $A_{1-3} = 78.5\%$ ) in year 13 (Fig. 11). Nitrogen fixing autotrophs ( $A_3$ ) are the dominant functional group during the first 10 years of the simulation (accounting for up to 56.1% of biomass). In the later stages of the forefield development, soil autotrophs ( $A_2$ ) becomes the most dominant community (up to 35.6% of biomass), as DIN concentrations increase. Subglacial microbes ( $A_1$  and  $H_1$ ) are consistently outcompeted throughout the simulation by soil microbes ( $A_2$  and  $H_2$ ) (Fig. 11).

Primary production ( $A_{1-3}$ ) dominates the system, accounting for between 68.7 and 88.9% of total bacterial production, whereas heterotrophic production ( $H_{1-3}$ ) accounts for the remaining 11.1 to 31.3% (Fig. 12b). This trend is also reflected in independent field studies, whereby autotrophic production has been identified as a major source of carbon in young soils at the Damma Glacier (Zumsteg et al., 2013b; Esperschutz et al., 2011; Frey et al., 2013) and elsewhere including the Puca Glacier in Peru (Schmidt et al., 2008). Heterotrophic production is closely associated with the abundance and availability of carbon substrate. A high proportion of labile substrate (39.4% in year 1) supports high rates of heterotrophic production and rapid accumulation of heterotrophic biomass. Labile substrate is rapidly depleted (Fig. 12a) followed by a sharp decline in biomass (Fig. 10). Following the exhaustion of labile organic carbon substrate, het-

GMDD

8, 1–74, 2015

**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





stage successional changes by providing a source of nitrogen (Chapin et al., 1994). The presence of a high diversity of diazotrophs has been attributed to an accumulation of nitrogen in forefield soils from the Damma Glacier (Duc et al., 2009), the Puca Glacier (Peru) (Schmidt et al., 2008; Nemergut et al., 2007), Mendenhall Glacier (Alaska) (Satin et al., 2009; Knelman et al., 2012) and Anvers Island (Antarctica) (Strauss et al., 2012). Initial phosphorus limitation is rapidly alleviated by the accumulation of DIP at an average rate of  $2.0 \mu\text{g P g}^{-1} \text{yr}^{-1}$  (years 1 to 10). Predicted DIP closely resembles field data (Göransson et al., 2011). The main supply of phosphorus to natural terrestrial ecosystems is the underlying parent rock, especially in glaciated settings whereby relatively high erosion rates and crushed rock flour give rise to increased mineral dissolution rates. However, phosphate is a relatively immobile macronutrient due to sorption and interaction with other soil constituents, making it a potential growth-limiting nutrient in terrestrial ecosystems (Hinsinger, 2001). Isotopic analysis (Tamburini et al., 2012) and modelling suggests that biological activity is the main driver of phosphorus cycling in developing soils at the Damma Glacier.

Microbial abundance and production calculated over winter (January, February, March) and summer (July, August, September) varies seasonally (Fig. 10a). Snow-cover attenuates PAR in the winter period causing large seasonal fluctuations in the biomass and overall production of photosynthetic organisms ( $A_2$  and  $A_3$ ). Whilst heterotrophic production is also higher in the summer ( $0.13 \mu\text{g C g}^{-1} \text{d}^{-1}$ ) than winter ( $0.05 \mu\text{g C g}^{-1} \text{d}^{-1}$ ), populations remain more stable. During cold periods, all life processes are slowed down, effectively ensuring the persistence of bacterial populations under sub-optimal conditions. The seasonal evolution of the Damma Glacier forefield is not well understood, however transplantation studies have indicated that microbial communities respond dynamically to changing environmental conditions (Zumsteg et al., 2013a) and soil bacteria photosynthesise, degrade organic material and fix nitrogen at varying rates over spring, summer and autumn (Lazzaro et al., 2012). SHIMMER estimates that 69.6 to 74.5 % of total net  $\text{CO}_2$  efflux occurs during the four month snow-

## GMDD

8, 1–74, 2015

### SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



free summer period between June and October, which agrees well with estimations from field studies at the Damma Glacier (62 to 70 %) (Guelland et al., 2013b).

#### 5.4.2 Case study 2: Athabasca Glacier, Canada

The model is calibrated and validated against a second test dataset from the Athabasca Glacier forefield in Canada. Parameters (Table 5) are varied sequentially to provide a best fit to observations of microbial biomass and carbon substrate (Fig. 13). Microbial biomass accumulates from  $2.6 \mu\text{gCg}^{-1}$  in year 1 to  $11.5 \mu\text{gCg}^{-1}$  in year 40, and then continues to accumulate slowly (roughly  $0.02 \mu\text{gCg}^{-1}\text{yr}^{-1}$ ) in the later stages of the chronosequence (years 50 to 75). These results agree with the observed accumulation of microbial biomass in the validation dataset. During the early stages of succession (< 15 years), nitrogen fixing autotrophs dominate the microbial community, accounting for up to 62.5% of the total biomass. Heterotrophs ( $H_{1-3}$ ) make up 17.2% of the biomass in year 15. Soil heterotrophs ( $H_2$ ) account for only 4.7% of biomass in year 15 ( $0.33 \mu\text{gCg}^{-1}$ ), and are severely limited by DIN ( $11.3 \mu\text{gNg}^{-1}$ ), however exceed to 8.3% of total biomass ( $1.36 \mu\text{gCg}^{-1}$ ) in year 75, in a similar abundance to nitrogen fixing heterotrophs ( $A_3 = 9.0\%$ ), when DIN stocks are more plentiful ( $43.2 \mu\text{gCg}^{-1}$ ).

The model predicts the accumulation of carbon substrate from  $298.2 \mu\text{gCg}^{-1}$  in year 1 to  $995.9 \mu\text{gCg}^{-1}$  in year 75. This is in line with observations. It is interesting to note that best fit is achieved with a substantially lower allochthonous input of substrate ( $v_{\text{Sub}} = 0.05$ ) for the Athabasca Glacier compared to the Damma Glacier ( $v_{\text{Sub}} = 0.60$ ), suggesting that allochthonous inputs may be lower or less dominant in the Athabasca Glacier forefield. Autotrophic production accounts for > 87.5% of total bacterial production. The majority of the DIN assimilation in young soils (< 15 years) is by nitrogen fixation (up to 96.6%), compared to less than half of total DIN assimilated in the later stages of development (years 70 to 75). Our ability to put these model results into context with field data is limited since there the Athabasca Glacier forefield is considerably less intensively studied than the Damma Glacier. Nevertheless, the Athabasca Glacier

GMDD

8, 1–74, 2015

### SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



site acts as a secondary test and validation of model behaviour against published observational and experimental data suggesting an increase in microbial biomass and accumulation of organic carbon.

## 6 Model development, application and recommendations

5 The SHIMMER framework greatly improves our ability to quantify dominant biogeochemical processes and make scenario-based predictions of soil development. Sensitivity studies help identify the most important parameters, and thus allow making recommendations for future laboratory experiments or field campaigns that would help reduce uncertainty. The model design is such that SHIMMER can be transferred to  
10 study other microbial-driven systems such as the establishment of biomass and nutrient cycling in desert soils, ice surfaces (e.g. cryoconite), microbial mats, and the built environment (e.g. fuel and chemical storage). SHIMMER does not have a vegetation component, and can be effectively implemented in locations where environmental conditions are harsh and/or do not support the establishment of permanent vascular  
15 vegetation. An additional result of developing the SHIMMER framework has been an appreciation of the types of data that are necessary to build a mechanistic and fully constrained numerical representation of these systems, and this is presented here as a recommendation to future field and laboratory efforts.

### 6.1 Data availability

20 Model development and confidence in model evaluations would be improved by higher temporal and spatial sampling resolution along the chronosequence and interdisciplinary (microbial and geochemical) data sets (concentrations and rates). At the very least, a proportional split between primary producers (autotrophs) and consumers (heterotrophs) will help in determining net productivity and validating model predictions of  
25 community composition. SHIMMER does not explicitly account for vegetation and thus

# GMDD

8, 1–74, 2015

## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



cannot reproduce the high organic carbon accumulation in vegetated sites (Fig. 10). Therefore, there is a need for test sites where the influence of vegetation is not so great. A full mass budget of inputs (e.g. aerial deposition) and outputs (e.g. leaching) of substrate and nutrients through the soil surface would greatly improve the predictive power of SHIMMER, as demonstrated in sensitivity analysis (Sect. 5.2.1).

## 6.2 Death and dormancy

Many biological processes including microbial growth and death are simplified in order to describe them mathematically. Whilst microbial growth rates are fairly well constrained from laboratory measurements (Frey et al., 2010), cell death is not well constrained, with a lack of empirical measurements, and fundamental differences in how these rates are defined in models (German et al., 2012; Grant et al., 1993; Scott et al., 1995; Toal et al., 2000; Boudreau, 1999; Kravchenko et al., 2004; Knapp et al., 1983; Blagodatsky and Richter, 1998; Ingwersen et al., 2008; Zelenev et al., 2000; Lancelot et al., 2005). Additionally, microbial biomass is able to persist under limiting nutrient conditions, and exist in active or dormant states (Mason et al., 1986). However, in situ measurements of the proportion of active microbial biomass in field studies are rare, thus making validation difficult. Those that are provided in the literature (Toal et al., 2000; Blagodatsky and Richter, 1998; Vandewerf and Verstraete, 1987b) are mostly derived from models and are a snap-shot in time. As such, they do not represent the seasonality of an environment. Appropriate representation of these rates is important since they strongly influence both the microbial populations' stability in size, and the available substrate and necromass. Therefore, more focussed empirical observations are needed to support and inform model processes, increase confidence in predictions and support future development.

**GMDD**

8, 1–74, 2015

### SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





### 6.3 Bacterial growth

Parameters defined as constants in mathematical models are often known to vary in time depending on prevailing environmental and biogeochemical conditions, for example sensitivity to temperature, soil moisture, oxygen availability, C : N ratio and quality of soil organic matter (Manzoni et al., 2012; Erhagen et al., 2015). Growth yield and background maintenance may also be defined separately (Manzoni and Porporato, 2009; Manzoni et al., 2012; Blagodatsky and Richter, 1998; Ingwersen et al., 2008). The addition of variable growth rates and efficiencies for example, would require additional parameters that currently cannot be defined with sufficient confidence. The result of this would be further abstraction from the system, and an increase in overall uncertainty and possible error, rather than improving the model. Laboratory incubations are useful to estimate parameter values experimentally (Blagodatsky et al., 1998). It is our intention with this study that future laboratory analysis can at least in part inform parameters, accounting for simplifications, abstractions and assumptions that may skew the value over a non-physically realistic range, such that confidence in model output is increased.

### 6.4 Discrete pools vs. continuum representation

The SHIMMER model framework divides aspects of the system into discrete categories contained within state variables. In a real soil system, the composition of the microbial community, along with the quality of substrate are continua. Classifying these into multiple categories ultimately provides a simplistic view of the system. This is not a problem as long as the model still contains sufficient detail to accurately represent the overall dynamics and provide useful information to improve understanding. Categorising variables into two or more pools (e.g. labile and refractory) also provides tuning parameters, flexibility and a high degree of generality, making it easier to translate the model framework to the particular system of interest. Other modelling approaches such as individual-based modelling (IBM) would account for population heterogeneity, and the

GMDD

8, 1–74, 2015

**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





ability to link mechanisms to population dynamics at the individual level (Hellweger and Bucci, 2009). However, this falls outside the scope of this exercise.

## 6.5 Deterministic vs. stochastic

The biogeochemical development of a de-glaciated forefield is extremely heterogeneous across a range of spatial scales, which affects the signal of biogeochemical changes in observations (Bernasconi et al., 2011; Bradley et al., 2014). Stochastic variability resulting from disturbances, heterogeneity in biotic and abiotic processes, lateral and vertical particle movement and diffusion, inter-annual variability, and changing environmental conditions affect the biogeochemical development of a chronosequence in such a way that a deterministic model cannot predict. Furthermore, the SHIMMER model does not predict or account for the effect of plant biomass, which is highly abundant in the forefield of the Damma Glacier (Bernasconi et al., 2011). A combination of these factors result in a large amount of scatter in the observational record that is absent from the model prediction. However, a first attempt to gain quantitative insight into process interplay in glacier forefields favours a deterministic framework.

## 6.6 0-D vs. multi-D

Currently, nearly all soil system models assume that at the finest level of detail, soil is a well-mixed homogeneous particle with respect to dominant processes and its composition, whereas microbial populations and metabolic rates are known to be heterogeneous across a number of different spatial scales. This heterogeneity is driven by abiotic factors including climate, hydrology, topography and mineralogy (especially with regard to phosphorus release from rock weathering), and biotic factors that result in response to each of these. Spatial heterogeneity of nutrient distribution in response to abiotic factors is explored in the Damma Glacier forefield (Göransson et al., 2014). A large amount of additional parameters and equations would need to be incorporated in order to include 1-D or 2-D processes. It is currently unfeasible to incorporate this

# GMDD

8, 1–74, 2015

## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





from the Damma Glacier, in the range of  $130.0 \mu\text{g C g}^{-1} \text{ yr}^{-1}$  (Schulz et al., 2013; Guelland et al., 2013a). Although seasonal evolution of glacier forefields is not well understood (Bradley et al., 2014), modelling work is likely to provide insight into the dynamics of the “non-growing season” with modelled summer production accounting for roughly 70 % of total annual respiration, in line with field observations from the Damma Glacier (Guelland et al., 2013b).

The accumulation of microbial biomass is highly sensitive to variation in the  $Q_{10}$  values, the active fraction of biomass, the bioavailability of refractory organic matter, the partitioning of organic matter between labile and refractory pools, bacterial growth efficiency, and rates of microbial growth and death. These parameters also have high uncertainty due to a relatively large range of plausible values. Many of the well constrained parameters (e.g. half saturation constants and exudation rates) have low sensitivity and uncertainty and show mostly linear behaviour. One of the striking outcomes of the sensitivity study is the apparent strong dependence between the heterotrophic and autotrophic microbial communities. Heterotrophic production degrades organic matter and recycles nutrients, in turn supporting autotrophic and heterotrophic growth. Increasing heterotrophic growth rates therefore increases the accumulation in all pools of biomass.

SHIMMER is the first step towards an iterative and interdisciplinary framework (presented in Fig. 2), integrating fieldwork and laboratory experiments with model development, testing and application. The development of SHIMMER (1.0) is informed by previous experimental and field campaigns (Bradley et al., 2014). It is expected that further quantitative analysis of forefield dynamics will guide and inform future studies that will provide new data and insights, which will inform further model development and so forth. SHIMMER thus contributes to more accurate quantitative predictions that enable a deeper understanding of the processes which control microbial communities, their role on global biogeochemical cycles and their response to climate variations in the future.

---

**SHIMMER (1.0):  
a novel mathematical  
model**J. A. Bradley et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Code availability

The SHIMMER source code is freely available by directly contacting the corresponding author. The code is written and executed in the free open source computing environment and programming language R, which is available for download on the web (http://www.r-project.org/).

*Acknowledgements.* We would like to thank Stefano Bernasconi and Anna Lazzaro (ETH Zurich, Switzerland), Beat Frey, Thomas Rime and Tobias Jonas (WSL, Switzerland), and Hans Göransson (BOKU Vienna, Austria) for helpful and insightful discussions, and for support in acquiring meteorological data. This research was supported by NERC grant no. NE/J02399X/1 to A. Anesio. S. Arndt acknowledges support from NERC grant no. NE/IO21322/1.

## References

- Achuff, P. L. and Coen, G. M.: Subalpine cryosolic soils in Banff and Jasper National-Parks, *Can. J. Soil Sci.*, 60, 579–581, 1980.
- ACIA: Arctic Climate Impacts Assessment, Cambridge University Press, Cambridge, 2005.
- Allison, S. D.: Cheaters, diffusion and nutrients constrain decomposition by microbial enzymes in spatially structured environments, *Ecol. Lett.*, 8, 626–635, 2005.
- Anderson, S. P., Drever, J. I., Frost, C. D., and Holden, P.: Chemical weathering in the foreland of a retreating glacier, *Geochim. Cosmochim. Ac.*, 64, 1173–1189, 2000.
- Anderson, T. H. and Domsch, K. H.: Maintenance carbon requirements of actively-metabolizing microbial-populations under insitu conditions, *Soil Biol. Biochem.*, 17, 197–203, 1985.
- Anesio, A. M., Hodson, A. J., Fritz, A., Psenner, R., and Sattler, B.: High microbial activity on glaciers: importance to the global carbon cycle, *Global Change Biol.*, 15, 955–960, 2009.
- Arndt, S., Jorgensen, B. B., Larowe, D. E., Middelburg, J. J., Pancost, R. D., and Regnier, P.: Quantifying the degradation of organic matter in marine sediments: a review and synthesis, *Earth-Sci. Rev.*, 123, 53–86, 2013.
- Bernasconi, S. M., Bauder, A., Bourdon, B., Brunner, I., Bunemann, E., Christl, I., Derungs, N., Edwards, P., Farinotti, D., Frey, B., Frossard, E., Furrer, G., Gierga, M., Goransson, H.,

GMDD

8, 1–74, 2015

## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Gulland, K., Hagedorn, F., Hajdas, I., Hindshaw, R., Ivy-Ochs, S., Jansa, J., Jonas, T., Kiczka, M., Kretzschmar, R., Lemarchand, E., Luster, J., Magnusson, J., Mitchell, E. A. D., Venterink, H. O., Plotze, M., Reynolds, B., Smittenberg, R. H., Stahl, M., Tamburini, F., Tipper, E. T., Wacker, L., Welc, M., Wiederhold, J. G., Zeyer, J., Zimmermann, S., and Zumsteg, A.: Chemical and biological gradients along the Damma Glacier soil chronosequence, Switzerland, *Vadose Zone J.*, 10, 867–883, 2011.

Berner, R. A., Lasaga, A. C., and Garrels, R. M.: The carbonate-silicate geochemical cycle and its effect on atmospheric carbon-dioxide over the past 100 million years, *Am. J. Sci.*, 283, 641–683, 1983.

Beven, K. and Binley, A.: The future of distributed models – model calibration and uncertainty prediction, *Hydrol. Process.*, 6, 279–298, 1992.

Blagodatsky, S. A. and Richter, O.: Microbial growth in soil and nitrogen turnover: a theoretical model considering the activity state of microorganisms, *Soil Biol. Biochem.*, 30, 1743–1755, 1998.

Blagodatsky, S. A., Yevdokimov, I. V., Larionova, A. A., and Richter, J.: Microbial growth in soil and nitrogen turnover: model calibration with laboratory data, *Soil Biol. Biochem.*, 30, 1757–1764, 1998.

Bottomley, P. and Myrold, D.: Biological N inputs, in: *Soil Microbiology, Ecology and Biochemistry*, 3rd Edn., edited by: Paul, E., Elsevier, USA, 365–388, 2007.

Boudreau, B. P.: A theoretical investigation of the organic carbon-microbial biomass relation in muddy sediments, *Aquat. Microb. Ecol.*, 17, 181–189, 1999.

Bradley, J. A., Singarayer, J. S., and Anesio, A. M.: Microbial community dynamics in the forefield of glaciers, *Proceedings Biological Sciences/The Royal Society*, 281, 2793–2802, doi:10.1098/rspb.2014.0882, 2014.

Brankatschk, R., Towe, S., Kleineidam, K., Schloter, M., and Zeyer, J.: Abundances and potential activities of nitrogen cycling microbial communities along a chronosequence of a glacier forefield, *ISME J.*, 5, 1025–1037, 2011.

Breitbarth, E., Wohlers, J., Klas, J., Laroche, J., and Peeken, I.: Nitrogen fixation and growth rates of *Trichodesmium* IMS-101 as a function of light intensity, *Mar. Ecol.-Prog. Ser.*, 359, 25–36, 2008.

Brooks, P. D. and Williams, M. W.: Snowpack controls on nitrogen cycling and export in seasonally snow-covered catchments, *Hydrol. Process.*, 13, 2177–2190, 1999.

---

**SHIMMER (1.0):  
a novel mathematical  
model**


---

J. A. Bradley et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)

[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


- Brown, S. P. and Jumpponen, A.: Contrasting primary successional trajectories of fungi and bacteria in retreating glacier soils, *Mol. Ecol.*, 23, 481–497, 2014.
- Cannell, M. G. R. and Thornley, J. H. M.: Modelling the components of plant respiration: some guiding principles, *Ann. Bot.-London*, 85, 45–54, 2000.
- 5 Chapin, F. S., Walker, L. R., Fastie, C. L., and Sharman, L. C.: Mechanisms of Primary Succession Following Deglaciation at Glacier Bay, Alaska, *Ecol. Monogr.*, 64, 149–175, 1994.
- Cowan, A. E., Olivastro, E. M., Koppel, D. E., Loshon, C. A., Setlow, B., and Setlow, P.: Lipids in the inner membrane of dormant spores of *Bacillus* species are largely immobile, *P. Natl. Acad. Sci. USA*, 101, 7733–7738, 2004.
- 10 Darrah, P. R.: Models of the rhizosphere, 1. Microbial-population dynamics around a root releasing soluble and insoluble carbon, *Plant Soil*, 133, 187–199, 1991.
- Dessert, C., Dupre, B., Gaillardet, J., Francois, L. M., and Allegre, C. J.: Basalt weathering laws and the impact of basalt weathering on the global carbon cycle, *Chem. Geol.*, 202, 257–273, 2003.
- 15 Duc, L., Noll, M., Meier, B. E., Burgmann, H., and Zeyer, J.: High diversity of diazotrophs in the forefield of a receding Alpine glacier, *Microbial. Ecol.*, 57, 179–190, 2009.
- Ensign, K. L., Webb, E. A., and Longstaffe, F. J.: Microenvironmental and seasonal variations in soil water content of the unsaturated zone of a sand dune system at Pinery Provincial Park, Ontario, Canada, *Geoderma*, 136, 788–802, 2006.
- 20 Erhagen, B., Ilstedt, U., and Nilsson, M. B.: Temperature sensitivity of heterotrophic soil CO<sub>2</sub> production increases with increasing carbon substrate uptake rate, *Soil Biol. Biochem.*, 80, 45–52, doi:10.1016/j.soilbio.2014.09.021, 2015.
- Esperschütz, J., Pérez-de-Mora, A., Schreiner, K., Welzl, G., Buegger, F., Zeyer, J., Hagedorn, F., Munch, J. C., and Schlöter, M.: Microbial food web dynamics along a soil chronosequence of a glacier forefield, *Biogeosciences*, 8, 3283–3294, doi:10.5194/bg-8-3283-2011, 2011.
- 25 Filippelli, G. M.: The global phosphorus cycle, *Rev. Mineral. Geochem.*, 48, 391–425, 2002.
- Foeroid, B. and Yearsley, J. M.: Modelling the impact of microbial grazers on soluble rhizodeposit turnover, *Plant Soil*, 267, 329–342, 2004.
- 30 Follmi, K. B., Hosein, R., Arn, K., and Steinmann, P.: Weathering and the mobility of phosphorus in the catchments and forefields of the Rhone and Oberaar glaciers, central Switzerland: implications for the global phosphorus cycle on glacial–interglacial timescales, *Geochim. Cosmochim. Ac.*, 73, 2252–2282, 2009.

---

**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Fountain, A. G., Nylen, T. H., Tranter, M., and Bagshaw, E.: Temporal variations in physical and chemical features of cryoconite holes on Canada Glacier, McMurdo Dry Valleys, Antarctica, *J. Geophys. Res.-Biogeo.*, 113, G01S92, doi:10.1029/2007JG000430, 2008.

5 Frey, B., Rieder, S. R., Brunner, I., Plotze, M., Koetzsch, S., Lapanje, A., Brandl, H., and Furrer, G.: Weathering-associated bacteria from the Damma Glacier forefield: physiological capabilities and impact on granite dissolution, *Appl. Environ. Microb.*, 76, 4788–4796, 2010.

Frey, B., Buhler, L., Schmutz, S., Zumsteg, A., and Furrer, G.: Molecular characterization of phototrophic microorganisms in the forefield of a receding glacier in the Swiss Alps, *Environ. Res. Lett.*, 8, 015033, doi:10.1088/1748-9326/8/1/015033, 2013.

10 Garnier, P., Neel, C., Mary, B., and Lafolie, F.: Evaluation of a nitrogen transport and transformation model in a bare soil, *Eur. J. Soil Sci.*, 52, 253–268, 2001.

German, D. P., Marcelo, K. R. B., Stone, M. M., and Allison, S. D.: The Michaelis–Menten kinetics of soil extracellular enzymes in response to temperature: a cross-latitudinal study, *Global Change Biol.*, 18, 1468–1479, 2012.

15 Goebel, N. L., Edwards, C. A., Carter, B. J., Achilles, K. M., and Zehr, J. P.: Growth and carbon content of three different-sized diazotrophic cyanobacteria observed in the subtropical North Pacific, *J. Phycol.*, 44, 1212–1220, 2008.

Göransson, H., Venterink, H. O., and Baath, E.: Soil bacterial growth and nutrient limitation along a chronosequence from a glacier forefield, *Soil Biol. Biochem.*, 43, 1333–1340, 2011.

20 Göransson, H., Edwards, P., Perreijn, K., Smittenberg, R. H., and Venterink, H. O.: Rocks create nitrogen hotspots and N : P heterogeneity by funnelling rain, *Biogeochemistry*, 121, 329–338, doi:10.1007/s10533-014-0031-x, 2014.

Grant, R. F., Juma, N. G., and McGill, W. B.: Simulation of carbon and nitrogen transformations in soil – mineralization, *Soil Biol. Biochem.*, 25, 1317–1329, 1993.

25 Graversen, R. G., Mauritsen, T., Tjernstrom, M., Kallen, E., and Svensson, G.: Vertical structure of recent Arctic warming, *Nature*, 451, 53–56, doi:10.1038/nature06502, 2008.

Greenfell, T. C. and Maykut, G. A.: The optical properties of ice and snow in the Arctic basin, *J. Glaciol.*, 18, 455–463, 1977.

30 Guelland, K., Esperschutz, J., Bornhauser, D., Bernasconi, S. M., Kretzschmar, R., and Hagedorn, F.: Mineralisation and leaching of C from C-13 labelled plant litter along an initial soil chronosequence of a glacier forefield, *Soil Biol. Biochem.*, 57, 237–247, 2013a.



---

**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Guelland, K., Hagedorn, F., Smittenberg, R. H., Goransson, H., Bernasconi, S. M., Hajdas, I., and Kretzschmar, R.: Evolution of carbon fluxes during initial soil formation along the forefield of Damma glacier, Switzerland, *Biogeochemistry*, 113, 545–561, 2013b.
- Hahn, A. S. and Quideau, S. A.: Shifts in soil microbial community biomass and resource utilization along a Canadian glacier chronosequence, *Can. J. Soil Sci.*, 93, 305–318, 2013.
- Hellweger, F. L. and Bucci, V.: A bunch of tiny individuals-Individual-based modeling for microbes, *Ecol. Model.*, 220, 8–22, 2009.
- Hinsinger, P.: Bioavailability of soil inorganic P in the rhizosphere as affected by root-induced chemical changes: a review, *Plant Soil*, 237, 173–195, 2001.
- Holl, C. M. and Montoya, J. P.: Interactions between nitrate uptake and nitrogen fixation in continuous cultures of the marine diazotroph *Trichodesmium* (Cyanobacteria), *J. Phycol.*, 41, 1178–1183, 2005.
- Ingwersen, J., Poll, C., Streck, T., and Kandeler, E.: Micro-scale modelling of carbon turnover driven by microbial succession at a biogeochemical interface, *Soil Biol. Biochem.*, 40, 864–878, 2008.
- Insam, H. and Haselwandter, K.: Metabolic quotient of the soil microflora in relation to plant succession, *Oecologia*, 79, 174–178, 1989.
- Kastovska, K., Elster, J., Stibal, M., and Santruckova, H.: Microbial assemblages in soil microbial succession after glacial retreat in Svalbard (high Arctic), *Microbial. Ecol.*, 50, 396–407, 2005.
- King, A. J., Meyer, A. F., and Schmidt, S. K.: High levels of microbial biomass and activity in unvegetated tropical and temperate alpine soils, *Soil Biol. Biochem.*, 40, 2605–2610, 2008.
- Kirschke, S., Bousquet, P., Ciais, P., Saunois, M., Canadell, J. G., Dlugokencky, E. J., Bergamaschi, P., Bergmann, D., Blake, D. R., Bruhwiler, L., Cameron-Smith, P., Castaldi, S., Chevallier, F., Feng, L., Fraser, A., Heimann, M., Hodson, E. L., Houweling, S., Josse, B., Fraser, P. J., Krummel, P. B., Lamarque, J. F., Langenfelds, R. L., Le Quere, C., Naik, V., O'Doherty, S., Palmer, P. I., Pison, I., Plummer, D., Poulter, B., Prinn, R. G., Rigby, M., Ringeval, B., Santini, M., Schmidt, M., Shindell, D. T., Simpson, I. J., Spahni, R., Steele, L. P., Strode, S. A., Sudo, K., Szopa, S., Van Der Werf, G. R., Voulgarakis, A., Van Weele, M., Weiss, R. F., Williams, J. E., and Zeng, G.: Three decades of global methane sources and sinks, *Nat. Geosci.*, 6, 813–823, 2013.



---

**SHIMMER (1.0):  
a novel mathematical  
model**


---

J. A. Bradley et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Knapp, E. B., Elliott, L. F., and Campbell, G. S.: Carbon, nitrogen and microbial biomass interrelationships during the decomposition of wheat straw – a mechanistic simulation-model, *Soil Biol. Biochem.*, 15, 455–461, 1983.

5 Knelman, J. E., Legg, T. M., O'Neill, S. P., Washenberger, C. L., Gonzalez, A., Cleveland, C. C., and Nemergut, D. R.: Bacterial community structure and function change in association with colonizer plants during early primary succession in a glacier forefield, *Soil Biol. Biochem.*, 46, 172–180, 2012.

Kravchenko, L. V., Strigul, N. S., and Shvytov, I. A.: Mathematical simulation of the dynamics of interacting populations of rhizosphere microorganisms, *Microbiology+*, 73, 189–195, 2004.

10 Kuijper, L. D. J., Berg, M. P., Morrien, E., Kooi, B. W., and Verhoef, H. A.: Global change effects on a mechanistic decomposer food web model, *Global Change Biol.*, 11, 249–265, 2005.

Lancelot, C., Spitz, Y., Gypens, N., Ruddick, K., Becquevort, S., Rousseau, V., Lacroix, G., and Billen, G.: Modelling diatom and *Phaeocystis* blooms and nutrient cycles in the Southern Bight of the North Sea: the MIRO model, *Mar. Ecol.-Prog. Ser.*, 289, 63–78, 2005.

15 Laroche, J. and Breitbarth, E.: Importance of the diazotrophs as a source of new nitrogen in the ocean, *J. Sea Res.*, 53, 67–91, 2005.

Lazzaro, A., Abegg, C., and Zeyer, J.: Bacterial community structure of glacier forefields on siliceous and calcareous bedrock, *Eur. J. Soil. Sci.*, 60, 860–870, 2009.

Lazzaro, A., Brankatschk, R., and Zeyer, J.: Seasonal dynamics of nutrients and bacterial communities in unvegetated alpine glacier forefields, *Appl. Soil Ecol.*, 53, 10–22, 2012.

20 Leffelaar, P. A. and Wessel, W. W.: Denitrification in a homogeneous, closed system – experiment and simulation, *Soil Sci.*, 146, 335–349, 1988.

Lennon, J. T. and Jones, S. E.: Microbial seed banks: the ecological and evolutionary implications of dormancy, *Nat. Rev. Microbiol.*, 9, 119–130, 2011.

25 Liu, G. X., Hu, P., Zhang, W., Wu, X. K., Yang, X., Chen, T., Zhang, M. X., and Li, S. W.: Variations in soil culturable bacteria communities and biochemical characteristics in the Dongkemadi glacier forefield along a chronosequence, *Folia Microbiol.*, 57, 485–494, 2012.

Liu, Y. Y., Wu, L. H., Baddeley, J. A., and Watson, C. A.: Models of biological nitrogen fixation of legumes, a review, *Agron. Sustain. Dev.*, 31, 155–172, 2011.

30 Long, T. and Or, D.: Aquatic habitats and diffusion constraints affecting microbial coexistence in unsaturated porous media, *Water Resour. Res.*, 41, W08408, doi:10.1029/2004WR003796, 2005.

---

**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Maggi, F. and Porporato, A.: Coupled moisture and microbial dynamics in unsaturated soils, *Water Resour. Res.*, 43, W07444, doi:10.1029/2006WR005367, 2007.
- Manzoni, S. and Porporato, A.: Soil carbon and nitrogen mineralization: theory and models across scales, *Soil Biol. Biochem.*, 41, 1355–1379, 2009.
- 5 Manzoni, S., Taylor, P., Richter, A., Porporato, A., and Agren, G. I.: Environmental and stoichiometric controls on microbial carbon-use efficiency in soils, *New Phytol.*, 196, 79–91, 2012.
- Mason, C. A., Hamer, G., and Bryers, J. D.: The death and lysis of microorganisms in environmental processes, *Fems Microbiol. Lett.*, 39, 373–401, 1986.
- 10 McGill, W.: Review and classification of ten soil organic matter (SOM) models, in: *Evaluation of Soil Organic Matter Models*, edited by: Powlson, D. S., Smith, P., and Smith, J. U., Springer, Berlin, 111–132, 1996.
- McGill, W.: The physiology and biochemistry of soil organisms, in: *Soil Microbiology, Ecology and Biochemistry*, 3rd Edn., edited by: Paul, E., Elsevier, USA, 2007.
- Meola, M., Lazzaro, A., and Zeyer, J.: Diversity, resistance, and resilience of the bacterial communities at two alpine glacier forefields after a reciprocal soil transplantation, *Environ. Microbiol.*, 16, 1918–1934, doi:10.1111/1462-2920.12435, 2014.
- 15 Mindl, B., Anesio, A. M., Meirer, K., Hodson, A. J., Laybourn-Parry, J., Sommaruga, R., and Sattler, B.: Factors influencing bacterial dynamics along a transect from supraglacial runoff to proglacial lakes of a high Arctic glacier, *Fems Microbiol. Ecol.*, 59, 762–762, 2007.
- 20 Moorhead, D. L. and Sinsabaugh, R. L.: A theoretical model of litter decay and microbial interaction, *Ecol. Monogr.*, 76, 151–174, 2006.
- Moreau, M., Mercier, D., Laffly, D., and Roussel, E.: Impacts of recent paraglacial dynamics on plant colonization: a case study on Midtre Lovénbreen foreland, Spitsbergen (79° N), *Geomorphology*, 95, 48–60, 2008.
- 25 Mur, L., Skulberg, O., and Utkilen, H.: Cyanobacteria in the environment, in: *Toxic Cyanobacteria in Water: a Guide to Their Public Health Consequences, Monitoring and Management*, edited by: Chorus, I. and Bartram, J., St Edmundsbury Press, Bury St Edmunds, Suffolk, UK, 1999.
- Nemergut, D. R., Anderson, S. P., Cleveland, C. C., Martin, A. P., Miller, A. E., Seimon, A., and Schmidt, S. K.: Microbial community succession in an unvegetated, recently deglaciated soil, *Microbiol. Ecol.*, 53, 110–122, 2007.
- 30

---

**SHIMMER (1.0):  
a novel mathematical  
model**


---

J. A. Bradley et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Ohtonen, R., Fritze, H., Pennanen, T., Jumpponen, A., and Trappe, J.: Ecosystem properties and microbial community changes in primary succession on a glacier forefront, *Oecologia*, 119, 239–246, 1999.

Overland, J. E., Wang, M., and Salo, S.: The recent Arctic warm period, *Tellus A*, 60, 589–597, 2008.

Panikov, N. S. and Sizova, M. V.: A kinetic method for estimating the biomass of microbial functional groups in soil, *J. Microbiol. Meth.*, 24, 219–230, 1996.

Parton, W. J., Stewart, J. W. B., and Cole, C. V.: Dynamics of C, N, P and S in grassland soils – a model, *Biogeochemistry*, 5, 109–131, 1988.

Paul, F., Frey, H., and Le Bris, R.: A new glacier inventory for the European Alps from Landsat TM scenes of 2003: challenges and results, *Ann. Glaciol.*, 52, 144–152, 2011.

Phillips, D. A.: Efficiency of symbiotic nitrogen-fixation in Legumes, *Annu. Rev. Plant. Phys.*, 31, 29–49, 1980.

Rabouille, S., Staal, M., Stal, L. J., and Soetaert, K.: Modeling the dynamic regulation of nitrogen fixation in the cyanobacterium *Trichodesmium* sp., *Appl. Environ. Microb.*, 72, 3217–3227, 2006.

Rime, T., Hartmann, M., Brunner, I., Widmer, F., Zeyer, J., and Frey, B.: Vertical distribution of the soil microbiota along a successional gradient in a glacier forefield, *Mol. Ecol.*, 24, 1091–1108, 2014.

Sattin, S. R., Cleveland, C. C., Hood, E., Reed, S. C., King, A. J., Schmidt, S. K., Roberson, M. S., Ascarrunz, N., and Nemergut, D. R.: Functional shifts in unvegetated, perhumid, recently-deglaciated soils do not correlate with shifts in soil bacterial community composition, *J. Microbiol.*, 47, 673–681, 2009.

Schimel, J. P., Bilbrough, C., and Welker, J. A.: Increased snow depth affects microbial activity and nitrogen mineralization in two Arctic tundra communities, *Soil Biol. Biochem.*, 36, 217–227, 2004.

Schipper, L. A., Hobbs, J. K., Rutledge, S., and Arcus, V. L.: Thermodynamic theory explains the temperature optima of soil microbial processes and high Q(10) values at low temperatures, *Global Change Biol.*, 20, 3578–3586, 2014.

Schmidt, S. K., Reed, S. C., Nemergut, D. R., Grandy, A. S., Cleveland, C. C., Weintraub, M. N., Hill, A. W., Costello, E. K., Meyer, A. F., Neff, J. C., and Martin, A. M.: The earliest stages of ecosystem succession in high-elevation (5000 metres above sea level), recently deglaciated soils, *P. Roy. Soc. B-Biol. Sci.*, 275, 2793–2802, 2008.

---

**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Schulz, S., Brankatschk, R., Dümig, A., Kögel-Knabner, I., Schloter, M., and Zeyer, J.: The role of microorganisms at different stages of ecosystem development for soil formation, *Biogeosciences*, 10, 3983–3996, doi:10.5194/bg-10-3983-2013, 2013.
- Schutte, U. M. E., Abdo, Z., Bent, S. J., Williams, C. J., Schneider, G. M., Solheim, B., and Forney, L. J.: Bacterial succession in a glacier foreland of the High Arctic, *Isme J.*, 3, 1258–1268, 2009.
- Scott, E. M., Rattray, E. A. S., Prosser, J. I., Killham, K., Glover, L. A., Lynch, J. M., and Bazin, M. J.: A mathematical-model for dispersal of bacterial inoculants colonizing the wheat rhizosphere, *Soil Biol. Biochem.*, 27, 1307–1318, 1995.
- Sigler, W. V. and Zeyer, J.: Microbial diversity and activity along the forefields of two receding glaciers, *Microbial. Ecol.*, 43, 397–407, 2002.
- Smittenberg, R. H., Gierga, M., Goransson, H., Christl, I., Farinotti, D., and Bernasconi, S. M.: Climate-sensitive ecosystem carbon dynamics along the soil chronosequence of the Damma glacier forefield, Switzerland, *Global Change Biol.*, 18, 1941–1955, 2012.
- Soetaert, K. and Herman, P.: *A Practical Guide to Ecological Modelling: Using R as a Simulation Platform*, Springer, UK, 2009.
- Soetaert, K., Petzoldt, T., and Setzer, R. W.: Solving differential equations in R: package desolve, *J. Stat. Softw.*, 33, 1–25, 2010.
- Staines, K. E. H., Carrivick, J. L., Tweed, F. S., Evans, A. J., Russell, A. J., Jóhannesson, T., and Roberts, M.: A multi-dimensional analysis of pro-glacial landscape change at Sólheimajökull, southern Iceland, *Earth Surf. Proc. Land.*, 40, 809–822, doi:10.1002/esp.3662, 2014.
- Stapleton, L. M., Crout, N. M. J., Sawstrom, C., Marshall, W. A., Poulton, P. R., Tye, A. M., and Laybourn-Parry, J.: Microbial carbon dynamics in nitrogen amended Arctic tundra soil: measurement and model testing, *Soil Biol. Biochem.*, 37, 2088–2098, 2005.
- Strauss, S. L., Garcia-Pichel, F., and Day, T. A.: Soil microbial carbon and nitrogen transformations at a glacial foreland on Anvers Island, Antarctic Peninsula, *Polar Biol.*, 35, 1459–1471, 2012.
- Tamburini, F., Pfahler, V., Bunemann, E. K., Guelland, K., Bernasconi, S. M., and Frossard, E.: Oxygen isotopes unravel the role of microorganisms in phosphate cycling in soils, *Environ. Sci. Technol.*, 46, 5956–5962, 2012.
- Toal, M. E., Yeomans, C., Killham, K., and Meharg, A. A.: A review of rhizosphere carbon flow modelling, *Plant Soil*, 222, 263–281, 2000.

---

**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Turner, J., Colwell, S. R., Marshall, G. J., Lachlan-Cope, T. A., Carleton, A. M., Jones, P. D., Lagun, V., Reid, P. A., and Iagovkina, S.: Antarctic climate change during the last 50 years, *Int. J. Climatol.*, 25, 279–294, 2005.

Van Liere, L. and Walsby, A. E.: Interactions of cyanobacteria with light, in: *The Biology of Cyanobacteria*, 2nd Edn., edited by: Whitton, B. A. and Carr, N. G., Blackwell Scientific, Oxford, 1982.

Vandewerf, H. and Verstraete, W.: Estimation of active soil microbial biomass by mathematical-analysis of respiration curves – calibration of the test procedure, *Soil Biol. Biochem.*, 19, 261–265, 1987a.

Vandewerf, H. and Verstraete, W.: Estimation of active soil microbial biomass by mathematical-analysis of respiration curves – development and verification of the model, *Soil Biol. Biochem.*, 19, 253–260, 1987b.

Wieder, W. R., Bonan, G. B., and Allison, S. D.: Global soil carbon projections are improved by modelling microbial processes, *Nat. Clim. Change*, 3, 909–912, 2013.

Williams, M. W., Seibold, C., and Chowanski, K.: Storage and release of solutes from a sub-alpine seasonal snowpack: soil and stream water response, *Niwot Ridge, Colorado, Biogeochemistry*, 95, 77–94, 2009.

Xenakis, G., Ray, D., and Mencuccini, M.: Sensitivity and uncertainty analysis from a coupled 3-PG and soil organic matter decomposition model, *Ecol. Model.*, 219, 1–16, 2008.

Yoshitake, S., Uchida, M., Koizumi, H., and Nakatsubo, T.: Carbon and nitrogen limitation of soil microbial respiration in a High Arctic successional glacier foreland near Ny-Ålesund, Svalbard, *Polar Res.*, 26, 22–30, 2007.

Yoshitake, S., Uchida, M., Koizumi, H., Kanda, H., and Nakatsubo, T.: Production of biological soil crusts in the early stage of primary succession on a High Arctic glacier foreland, *New Phytol.*, 186, 451–460, 2010.

Zelenev, V. V., Van Bruggen, A. H. C., and Semenov, A. M.: “BACWAVE,” a spatial–temporal model for traveling waves of bacterial populations in response to a moving carbon source in soil, *Microbial. Ecol.*, 40, 260–272, 2000.

Zumsteg, A., Bernasconi, S. M., Zeyer, J., and Frey, B.: Microbial community and activity shifts after soil transplantation in a glacier forefield, *Appl. Geochem.*, 26, S326–S329, 2011.

Zumsteg, A., Luster, J., Goransson, H., Smittenberg, R. H., Brunner, I., Bernasconi, S. M., Zeyer, J., and Frey, B.: Bacterial, archaeal and fungal succession in the forefield of a receding glacier, *Microbial. Ecol.*, 63, 552–564, 2012.

- Zumsteg, A., Baath, E., Stierli, B., Zeyer, J., and Frey, B.: Bacterial and fungal community responses to reciprocal soil transfer along a temperature and soil moisture gradient in a glacier forefield, *Soil Biol. Biochem.*, 61, 121–132, 2013a.
- 5 Zumsteg A., Schmutz, S., and Frey, B.: Identification of biomass utilizing bacteria in a carbon-depleted glacier forefield soil by the use of <sup>13</sup>C DNA stable isotope probing, *Env. Microbiol. Rep.*, 5, 424–437, 2013b.

## GMDD

8, 1–74, 2015

### SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 1.** Outline of quantitative and predictive modelling strategy.

| Quantitative modelling  | Details   |
|---|---|
| Evaluate the extent to which initial microbial communities rely on autochthonous or allochthonous carbon.   | Whilst microbial activity is thought to be responsible for the initial build-up of labile substrate pools and facilitating the growth of higher order plant life in developed soils, it is unclear to what extent these ecosystems are self-supported by autochthonous production, or rely on external or ancient sources of carbon (Schulz et al., 2013; Bradley et al., 2014). Using SHIMMER, various carbon loading scenarios can be explored quantitatively to determine the sensitivity of microbial community development to allochthonous sources.                         |
| Assess the role of nitrogen as a key limiting nutrient.   | Nitrogen dynamics in de-glaciated forefield ecosystems are important in shaping the development of initial microbial communities (Brankatschk et al., 2011). Some evidence suggests that initial communities rely heavily on nitrogen fixation (Duc et al., 2009; Schmidt et al., 2008; Nemergut et al., 2007; Chapin et al., 1994) whereas others suggest that internal nitrogen cycling and external nutrient loading are sufficient to support initial community development (Sattin et al., 2009). Using SHIMMER, the entire nitrogen cycle can be quantified mathematically. |
| Assess how microbial diversity affects soil development.  | The diversity of microbes is likely to affect overall biogeochemical development of forefield soils (Guelland et al., 2013b; Goransson et al., 2011; Bradley et al., 2014). Initial microbial community composition is prescribed in SHIMMER, thus various community assemblages can be tested in terms of the nutrient-rich or nutrient-poor soil environment they create.   |
| Predictive modelling  | Details   |
| Identifying tipping points within the system.   | Tipping points related to population dynamics, external nutrient loading, seasonal changes, disturbance events and climate change could trigger significant changes in the dynamic behaviour of the microbial community.  |
| Assess importance of seasonality.   | The seasonal changes that affect forefield chronosequence dynamics are largely unexplored (Bradley et al., 2014). The majority of observational field data is collected during the summer (where sites are more accessible and temperatures are more amenable to working in the field). However, the dynamics of the “non-growing season” are largely under-explored, and the model representation captures the entire seasonal variation in environmental and biological components.   |
| Assess importance of disturbance events and community reset.  | Glacier forefields are extremely heterogeneous environments which are often subject to disturbances as glacial meltwater travels through the proglacial zone. This may be rich in nutrients and organic matter, but may also reset established communities and cause enhanced leaching of substrate. SHIMMER can test multiple scenarios to assess the possible effects of disturbances.  |
| Assess sensitivity of chronosequence development to future climate change.<br>Draw attention to gaps in our understanding and areas of future research. | Anthropogenic changes are likely to have a significant impact on Arctic climate and soil biogeochemistry (Graversen et al., 2008; Overland et al., 2008; ACIA, 2005; Turner et al., 2005). Models are driven primarily by data; henceforth it is likely to become apparent where future fieldwork and lab work efforts should be focussed. Models are also useful tools to highlighting gaps in current understanding of processes and systems, pressing for a quantitative consideration of system dynamics where previous attempts were more qualitative.                       |







**Table 4.** SHIMMER model formulation.

| Fundamental balance equations  |  |
|--|--|
| Rate of change of autotrophic biomass ( $A_{1-3}$ )                            | $\frac{dA_i}{dt} = U_{A_i} - G_{A_i} - X_{A_i}$  |
| Rate of change of autotrophic biomass ( $H_{1-3}$ )                            | $\frac{dH_i}{dt} = U_{H_i} - G_{H_i} - X_{H_i}$  |
| Rate of change of labile carbon substrate ( $S_1$ )                            | $\frac{dS_1}{dt} = v_{Sub} \times I_{S_1} + q \times G_{A_1} + q \times G_{H_1} + X_{A_1} + X_{H_1} - U_{S_1H_1} - W_{S_1}$  |
| Rate of change of refractory carbon substrate ( $S_2$ )                        | $\frac{dS_2}{dt} = v_{Sub} \times I_{S_2} + (1 - q) \times G_{A_1} + (1 - q) \times G_{H_1} - U_{S_2H_1} - W_{S_2}$  |
| Rate of change of dissolved inorganic nitrogen (DIN)                           | $\frac{dDIN}{dt} = v_{DIN} \times I_{DIN} - DIN_{Consumed} + DIN_{Released} - W_{DIN}$   |
| Rate of change of dissolved inorganic phosphorus (DIP)                         | $\frac{dDIP}{dt} = v_{DIP} \times I_{DIP} - DIP_{Consumed} + DIP_{Released} - W_{DIP}$   |
| Rate of change of labile organic nitrogen (ON <sub>1</sub> )                   | $\frac{dON_1}{dt} = v_{Sub} \times I_{ON_1} - ON_{1Accumulation} - ON_{1Degraded} - W_{ON_1}$  |
| Rate of change of refractory organic nitrogen (ON <sub>2</sub> )               | $\frac{dOP_2}{dt} = v_{Sub} \times I_{OP_2} - OP_{2Accumulation} - OP_{2Degraded} - W_{OP_2}$  |
| Rate of change of labile organic phosphorus (OP <sub>1</sub> )                 | $\frac{dOP_1}{dt} = v_{Sub} \times I_{OP_1} - OP_{1Accumulation} - OP_{1Degraded} - W_{OP_1}$  |
| Rate of change of refractory organic phosphorus (OP <sub>2</sub> )             | $\frac{dOP_2}{dt} = v_{Sub} \times I_{OP_2} - OP_{2Accumulation} - OP_{2Degraded} - W_{OP_2}$  |
| Carbon cycle – Biomass component   |  |
| Growth of chemo-autotrophic biomass ( $A_1$ )                                  | $U_{A_1} = A_1 \times T_1 \times d \times \rho_{sub} \times I_{maxA_1} \times \left( \frac{DIN}{DIN + (K_p \times K_{sub})} \right) \times \left( \frac{DIP}{DIP + (K_p \times K_{sub})} \right)$  |
| Growth of autotrophic biomass ( $A_2$ )  | $U_{A_2} = A_2 \times T_1 \times d \times I_{maxA_2} \times \left( \frac{L}{L + K_L} \right) \times \left( \frac{DIN}{DIN + K_C} \right) \times \left( \frac{DIP}{DIP + K_C} \right)$  |
| Growth of autotrophic N-fixing biomass ( $A_3$ )                               | $U_{A_3} = U_{A_3\_N2} + U_{A_3\_DIN}$   |
| Growth of $A_3$ with nitrogen fixation   | $U_{A_3\_N2} = A_3 \times T_1 \times d \times n_1 \times I_{maxA_3} \times \left( \frac{L}{L + K_L} \right) \times \left( \frac{DIP}{DIP + K_C} \right) \times (1 - m_{DIN})$  |
| Growth of $A_3$ with DIN   | $U_{A_3\_DIN} = A_3 \times T_1 \times d \times I_{maxA_3} \times \left( \frac{L}{L + K_L} \right) \times \left( \frac{DIN}{DIN + K_C} \right) \times \left( \frac{DIP}{DIP + K_C} \right) \times (m_{DIN})$  |
| Growth of heterotrophic subglacial biomass ( $H_1$ ) from labile substrate     | $U_{H_1\_L} = H_1 \times T_1 \times d \times \rho_{sub} \times I_{maxH_1} \times \left( J_{S_1} \times \frac{S_1}{S_1 + K_C} \right) \times \left( \frac{DIN}{DIN + (K_p \times K_{sub})} \right) \times \left( \frac{DIP}{DIP + (K_p \times K_{sub})} \right)$                  |
| Growth of heterotrophic soil biomass ( $H_2$ ) from labile substrate           | $U_{H_2\_L} = H_2 \times T_1 \times d \times I_{maxH_2} \times \left( J_{S_1} \times \frac{S_1}{S_1 + K_C} \right) \times \left( \frac{DIN}{DIN + K_C} \right) \times \left( \frac{DIP}{DIP + K_C} \right)$  |
| Growth of $H_3$ from labile substrate and nitrogen fixation                    | $U_{H_3\_L\_N2} = H_3 \times T_1 \times d \times n_1 \times I_{maxH_3} \times \left( J_{S_1} \times \frac{S_1}{S_1 + K_C} \right) \times \left( \frac{DIP}{DIP + K_C} \right) \times (1 - m_{DIN})$  |
| Growth of $H_3$ from labile substrate and DIN                                  | $U_{H_3\_L\_DIN} = H_3 \times T_1 \times d \times I_{maxH_3} \times \left( J_{S_1} \times \frac{S_1}{S_1 + K_C} \right) \times \left( \frac{DIN}{DIN + K_C} \right) \times \left( \frac{DIP}{DIP + K_C} \right) \times (m_{DIN})$  |
| Growth of heterotrophic subglacial biomass ( $H_1$ ) from refractory substrate | $U_{H_1\_R} = H_1 \times T_1 \times d \times I_{maxH_1} \times \rho_{sub} \times \left( J_{S_2} \times \frac{S_2}{S_2 + (K_p \times K_{sub})} \right) \times \left( \frac{DIN}{DIN + (K_p \times K_{sub})} \right) \times \left( \frac{DIP}{DIP + (K_p \times K_{sub})} \right)$ |
| Growth of heterotrophic soil biomass ( $H_2$ ) from refractory substrate       | $U_{H_2\_R} = H_2 \times T_1 \times d \times I_{maxH_2} \times \left( J_{S_2} \times \frac{S_2}{S_2 + K_C} \right) \times \left( \frac{DIN}{DIN + K_C} \right) \times \left( \frac{DIP}{DIP + K_C} \right)$  |
| Growth of $H_3$ from refractory substrate and nitrogen fixation                | $U_{H_3\_R\_N2} = H_3 \times T_1 \times d \times n_1 \times I_{maxH_3} \times \left( J_{S_2} \times \frac{S_2}{S_2 + K_C} \right) \times \left( \frac{DIP}{DIP + K_C} \right) \times (1 - m_{DIN})$  |
| Growth of $H_3$ from refractory substrate and DIN                              | $U_{H_3\_R\_DIN} = H_3 \times T_1 \times d \times I_{maxH_3} \times \left( J_{S_2} \times \frac{S_2}{S_2 + K_C} \right) \times \left( \frac{DIN}{DIN + K_C} \right) \times \left( \frac{DIP}{DIP + K_C} \right) \times (m_{DIN})$  |
| Growth of heterotrophic nitrogen fixers ( $H_3$ ) from labile substrate        | $U_{H_3\_L} = U_{H_3\_L\_N2} + U_{H_3\_L\_DIN}$  |
| Growth of heterotrophic nitrogen fixers ( $H_3$ ) from refractory substrate    | $U_{H_3\_R} = U_{H_3\_R\_N2} + U_{H_3\_R\_DIN}$  |
| Total growth of $H_i$  | $U_{H_i} = U_{H_i\_L} + U_{H_i\_R}$  |
| Loss of autotrophic biomass  | $G_{A_i} = T_i \times d \times \alpha_{A_i} \times A_i^2$  |
| Loss of heterotrophic biomass  | $G_{H_i} = T_i \times d \times \alpha_{H_i} \times H_i^2$  |
| Total deaths that become labile  | $G_L = q \times (G_{A_1} + G_{A_2} + G_{A_3} + G_{H_1} + G_{H_2} + G_{H_3})$   |
| Total deaths that become refractory  | $G_R = (1 - q) \times (G_{A_1} + G_{A_2} + G_{A_3} + G_{H_1} + G_{H_2} + G_{H_3})$   |
| Rate of exudate and EPS production by autotrophs                               | $X_{A_i} = \alpha_{X_{A_i}} \times U_{A_i}$  |
| Rate of exudate and EPS production by heterotrophs                             | $X_{H_i} = \alpha_{X_{H_i}} \times U_{H_i}$  |
| Total exudate production   | $X_{Total} = (X_{A_1} + X_{A_2} + X_{A_3} + X_{H_1} + X_{H_2} + X_{H_3})$  |

**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.

| Title Page               |              |
|--------------------------|--------------|
| Abstract                 | Introduction |
| Conclusions              | References   |
| Tables                   | Figures      |
| ◀                        | ▶            |
| ◀                        | ▶            |
| Back                     | Close        |
| Full Screen / Esc        |              |
| Printer-friendly Version |              |
| Interactive Discussion   |              |



## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

|                          |              |
|--------------------------|--------------|
| Title Page               |              |
| Abstract                 | Introduction |
| Conclusions              | References   |
| Tables                   | Figures      |
| ◀                        | ▶            |
| ◀                        | ▶            |
| Back                     | Close        |
| Full Screen / Esc        |              |
| Printer-friendly Version |              |
| Interactive Discussion   |              |

**Table 4.** Continued.

|  |  |
|--|--|
| Carbon cycle – Substrate component                                       |  |
| Consumption of labile substrate  | $S_{1\text{Consumed}} = \left(\frac{1}{Y_H}\right) \times U_{H_1L} + \left(\frac{1}{Y_H}\right) \times U_{H_2L} + \left(\frac{1}{n_1 \times Y_H}\right) \times U_{H_3L\_N2} + \left(\frac{1}{Y_H}\right) \times U_{H_3L\_DIN}$ |
| Consumption of refractory substrate                                      | $S_{2\text{Consumed}} = \left(\frac{1}{Y_H}\right) \times U_{H_1R} + \left(\frac{1}{Y_H}\right) \times U_{H_2R} + \left(\frac{1}{n_1 \times Y_H}\right) \times U_{H_3R\_N2} + \left(\frac{1}{Y_H}\right) \times U_{H_3R\_DIN}$ |
| Leaching   |  |
| Leaching   | $W_X = g_X \times X$   |
| Nitrogen cycle   |  |
| Labile organic nitrogen degraded   | $ON_{1\text{Degraded}} = \left(\frac{ON_1}{S_1}\right) \times (S_{1\text{Consumed}})$  |
| Refractory organic nitrogen degraded                                     | $ON_{2\text{Degraded}} = \left(\frac{ON_2}{S_2}\right) \times (S_{2\text{Consumed}})$  |
| Labile organic nitrogen accumulated                                      | $ON_{1\text{Accumulation}} = NC \times (G_L + X_T)$  |
| Refractory organic nitrogen accumulated                                  | $ON_{2\text{Accumulation}} = NC \times G_R$  |
| DIN Consumed   | $DIN_{\text{Consumed}} = NC \times \left(U_{A_1} + U_{A_2} + U_{A_3\_DIN} + U_{H_1} + U_{H_2} + U_{H_3L\_DIN} + U_{H_3R\_DIN}\right)$  |
| DIN Released   | $DIN_{\text{Released}} = ON_{1\text{Degraded}} + ON_{2\text{Degraded}}$  |
| Phosphorus cycle   |  |
| Labile organic phosphorus degraded                                       | $OP_{1\text{Degraded}} = \left(\frac{OP_1}{S_1}\right) \times (S_{1\text{Consumed}})$  |
| Refractory organic phosphorus degraded                                   | $OP_{2\text{Degraded}} = \left(\frac{OP_2}{S_2}\right) \times (S_{2\text{Consumed}})$  |
| Labile organic phosphorus accumulated                                    | $OP_{1\text{Accumulation}} = PC \times (G_L + X_T)$  |
| Refractory organic phosphorus accumulated                                | $OP_{2\text{Accumulation}} = PC \times G_R$  |
| DIP Consumed   | $DIP_{\text{Consumed}} = PC \times \left(U_{A_1} + U_{A_2} + U_{A_3} + U_{H_1} + U_{H_2} + U_{H_3}\right)$   |
| DIP Released   | $DIP_{\text{Released}} = OP_{1\text{Degraded}} + OP_{2\text{Degraded}}$  |
| Environmental and scaling equations                                      |  |
| Temperature factor response  | $T_i = \exp\left(\frac{T - T_{ref}}{10} \log_e(Q_{10})\right)$   |
| Monod expression for nitrogen fixation inhibition in the presence of DIN | $\text{if } DIN \leq DIN_t, \quad m_{DIN} = 0$ $\text{else } m_{DIN} = \left(\frac{DIN - DIN_t}{(DIN - DIN_t) + K_{iN2}}\right)$   |



**Table 5. Parameters.**

| Parameter      | Description   | Units                   | Nominal value<br>(Reference)                                    | Lower range<br>(Reference)                                   | Upper range<br>(Reference)                                   | Damma<br>Glacier,<br>Switzerland        | Athabasca<br>Glacier,<br>Canada |
|----------------|---|-------------------------|---|--|--|---|---------------------------------|
| $T_{ref}$      | Reference temperature for rates   | $^{\circ}\text{C}$      | 25<br>(Frey et al., 2010)                                       | Fixed  | Fixed  | –                                       | –                               |
| NC             | C : N ratio (mass)  | Unitless                | 0.141<br>(Bernasconi et al., 2011)                              | Fixed  | Fixed  | –                                       | –                               |
| PC             | C : P ratio (mass)  | Unitless                | 0.083<br>(Bernasconi et al., 2011)                              | Fixed  | Fixed  | –                                       | –                               |
| $Q_{10}$       | Temperature sensitivity   | Unitless                | 2.0<br>(Soetaert and Herman, 2009)                              | 1.5  | 3.1<br>(Yoshitake et al., 2010)                              | 2.2                                     | 2.2                             |
| $\alpha_A$     | Death rate (autotrophs)   | $\text{d}^{-1}$         | 0.0120<br>(German et al., 2012)                                 | 0.0060<br>(Grant et al., 1993)                               | 0.4800<br>(Scott et al., 1995)                               | –                                       | 0.0340                          |
| $\alpha_H$     | Death rate (heterotrophs)   | $\text{d}^{-1}$         | 0.0120<br>(German et al., 2012)                                 | 0.0060<br>(Grant et al., 1993)                               | 0.4800<br>(Scott et al., 1995)                               | –                                       | 0.0340                          |
| $I_{maxA}$     | Maximum growth rate (autotrophs)  | $\text{d}^{-1}$         | 1.21 (Frey et al., 2010)  | 0.30 (Mur et al., 1999)                                      | 1.40 (Mur et al., 1999)                                      | 1.40                                    | 1.4                             |
| $I_{maxH}$     | Maximum growth rate (heterotrophs)  | $\text{d}^{-1}$         | 1.21<br>(Frey et al., 2010)                                     | 0.24<br>(Ingwersen et al., 2008)                             | 4.80<br>(Darrah, 1991)                                       | 1.24<br>(Blagodatsky and Richter, 1998) | 1.24                            |
| $ex_A$         | Exudates and EPS production (autotrophs)  | Unitless                | 0.014<br>(Allison, 2005)  | 0.007<br>(Allison, 2005)                                     | 0.021<br>(Allison, 2005)                                     | –                                       | –                               |
| $ex_H$         | Exudates and EPS production (heterotrophs)  | Unitless                | 0.014<br>(Allison, 2005)  | 0.007<br>(Allison, 2005)                                     | 0.021<br>(Allison, 2005)                                     | –                                       | –                               |
| $\rho_{Sub}$   | Slow-down of subglacial microbial growth rate   | Unitless                | 0.2   | 0.1  | 1.0  | –                                       | –                               |
| $K_{Sub}$      | Lower half-saturation constants ( $K_{S_A}$ , $K_{S_H}$ and $K_P$ ) for subglacial microbes | Unitless                | 0.8   | 0.1  | 1.0  | –                                       | –                               |
| $K_L$          | Light half-saturation constant for autotrophs ( $A_2$ and $A_3$ )                           | $\text{W m}^{-2}$ (PAR) | 1.85<br>(Van Liere and Walsby, 1982)                            | 0.70<br>(Van Liere and Walsby, 1982)                         | 3.00<br>(Van Liere and Walsby, 1982)                         | –                                       | –                               |
| $K_S$          | Substrate half-saturation constant for heterotrophs   | $\mu\text{g g}^{-1}$    | 349<br>(Vandewerf and Verstraete, 1987b)                        | 50<br>(Darrah, 1991)   | 1000<br>(Knapp et al., 1983)                                 | –                                       | –                               |
| $K_N$          | DIN half-saturation constant  | $\mu\text{g g}^{-1}$    | 49.209<br>(stoichiometric)                                      | 7.050<br>(stoichiometric)                                    | 141.000<br>(stoichiometric)                                  | –                                       | –                               |
| $K_P$          | DIP half-saturation constant  | $\mu\text{g g}^{-1}$    | 28.967<br>(stoichiometric)                                      | 4.150<br>(stoichiometric)                                    | 83.000<br>(stoichiometric)                                   | –                                       | –                               |
| $\eta_i$       | Downscaling of $\gamma$ and $I_{max}$ when fixing nitrogen                                  | Unitless                | 0.50<br>(Bottomley and Myrold, 2007)                            | 0.10<br>(LaRoche and Breitbarth, 2005)                       | 0.80<br>(Breitbarth et al., 2008)                            | –                                       | –                               |
| $K_{N2}$       | Nitrogen fixation inhibition  | $\mu\text{g g}^{-1}$    | 393.672<br>(Holl and Montoya, 2005)<br>(Rabouille et al., 2006) | 56.4<br>(Holl and Montoya, 2005)<br>(Rabouille et al., 2006) | 1128<br>(Holl and Montoya, 2005)<br>(Rabouille et al., 2006) | –                                       | –                               |
| $\text{DIN}_i$ | Threshold value of DIN for nitrogen fixation inhibition                                     | $\mu\text{g g}^{-1}$    | 0   | 0  | 0  | –                                       | –                               |
| $q$            | Proportion of necromass that becomes labile ( $S_1$ )                                       | Unitless                | 0.3   | 0.1  | 0.5  | –                                       | –                               |

**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 5.** Continued.

| Parameter | Description   | Units                        | Nominal value<br>(Reference)  | Lower range<br>(Reference)               | Upper range<br>(Reference)               | Damma<br>Glacier,<br>Switzerland | Athabasca<br>Glacier,<br>Canada |
|-----------|---|------------------------------|-------------------------------|--|--|----------------------------------|---------------------------------|
| $J_{S_1}$ | Bioavailability (preference) of $S_1$                     | Unitless                     | 0.50                          | 0.50                                     | 1.00                                     | 0.68                             | 0.50                            |
| $J_{S_2}$ | Bioavailability (preference) of $S_2$                     | Unitless                     | 0.10                          | 0.10                                     | 0.50                                     | 0.15                             | 0.10                            |
| $g_{Sub}$ | Leaching of substrate                                     | $d^{-1}$                     | 0                             | 0  | 0  | –                                | –                               |
| $g_{DIN}$ | Leaching of DIN   | $d^{-1}$                     | 0                             | 0  | 0  | –                                | –                               |
| $g_{DIP}$ | Leaching of DIP   | $d^{-1}$                     | 0                             | 0  | 0  | –                                | –                               |
| $Y_A$     | Growth efficiency of autotrophs                           | $gC(gC\text{consumed})^{-1}$ | 0.200<br>(Scott et al., 1995) | 0.100<br>(Foeroid and Yearsley, 2004)    | 0.848<br>(Blagodatsky and Richter, 1998) | –                                | –                               |
| $Y_H$     | Growth efficiency of heterotrophs                         | $gC(gC\text{consumed})^{-1}$ | 0.200<br>(Scott et al., 1995) | 0.134<br>(German et al., 2012)           | 0.848<br>(Blagodatsky and Richter, 1998) | –                                | –                               |
| $d$       | Active fraction of microbial biomass                      | Unitless                     | 0.285<br>(Wang et al., 2014)  | 0.100<br>(Blagodatsky and Richter, 1998) | 0.580<br>(Blagodatsky and Richter, 1998) | 0.285                            | 0.100                           |
| $V_{Sub}$ | Proportion of allochthonous substrate deposition retained | Unitless                     | 0.60                          | 0.00                                     | 1.00                                     | –                                | 0.05                            |
| $V_{DIN}$ | Proportion of allochthonous DIN deposition retained       | Unitless                     | 0.01                          | 0.00                                     | 1.00                                     | –                                | –                               |
| $V_{DIP}$ | Proportion of allochthonous DIP deposition retained       | Unitless                     | 0.01                          | 0.00                                     | 1.00                                     | –                                | –                               |

## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




**Table 6.** Allochthonous deposition.

| Nutrient Species | Input ( $\mu\text{g g}^{-1} \text{d}^{-1}$ ) |
|------------------|--|
| $S_1$            | 1.955  |
| $S_2$            | 4.562  |
| DIN              | 6.952  |
| DIP              | 1.738  |
| $\text{ON}_1$    | 0.164  |
| $\text{ON}_2$    | 0.383  |
| $\text{OP}_1$    | 0.162  |
| $\text{OP}_2$    | 0.378  |

## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.



**Figure 1.** A conceptual overview of the main transformations and fluxes of nutrients and organic matter in and along a typical de-glaciated forefield.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




# GMDD

8, 1–74, 2015

## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.



**Figure 2.** Iterative, fully integrated interdisciplinary framework of SHIMMER, aimed at continuous model development as new datasets emerge and knowledge of the field evolves.


|                          |              |
|--------------------------|--------------|
| Title Page               |              |
| Abstract                 | Introduction |
| Conclusions              | References   |
| Tables                   | Figures      |
| ⏪                        | ⏩            |
| ◀                        | ▶            |
| Back                     | Close        |
| Full Screen / Esc        |              |
| Printer-friendly Version |              |
| Interactive Discussion   |              |





## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.



**Figure 3.** A conceptual model showing the components and transfers of SHIMMER. State variables are indicated with shading.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.



**Figure 4.** Annual meteorological forcings for SHIMMER for the Damma Glacier, Switzerland (red), and the Athabasca Glacier, Canada (black). Forcing data is provided by the WSL (Switzerland), Achuff and Coen (1980) and Staveley AAFC (Canada).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## SHIMMER (1.0): a novel mathematical model


J. A. Bradley et al.



**Figure 5.** Simulated response of autotrophic ( $A_{1-3}$ ) biomass (green line) and heterotrophic ( $H_{1-3}$ ) biomass (red line) to variation in **(a)**  $Q_{10}$  and **(b)**  $I_{maxH}$  across the entire range of plausible values, forced with meteorological data from the Damma Glacier over 75 years. The shaded segment shows the region in which model sensitivity ( $\lambda$ ) is calculated.


**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.



**Figure 6.** Model output of the Damma Glacier forefield for a 75 year time series with nominal parameter values (Table 4).





**Figure 7.** Sensitivity of model outputs ( $\lambda$ ) to individual parameter variation. The model is forced with meteorological data from the Damma Glacier (Fig. 4) over 75 years.

**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶


Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Figure 8.** Heat-map of parameter linearity and non-linearity over a range of model outputs. The model is forced with meteorological data from the Damma Glacier (Fig. 4) over 75 years.

**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**SHIMMER (1.0):  
a novel mathematical  
model**

J. A. Bradley et al.



**Figure 9.** Uncertainty of model outputs ( $\phi$ ) arising from individual parameters. The model is forced with meteorological data from the Damma Glacier (Fig. 4) over 75 years.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion







Figure 10. Model output optimised to observational data from the Damma Glacier, Switzerland.



## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.



**Figure 11.** Evolution of the microbial community composition in model output optimised from the Damma Glacier, Switzerland, over the first 20 years of soil formation.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## SHIMMER (1.0): a novel mathematical model

J. A. Bradley et al.



**Figure 12.** Evolution of **(a)** substrate dynamics, **(b)** bacterial production and **(c)** DIN assimilation of model output from the first 20 years of soil formation at the Damma Glacier, Switzerland.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)

[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Figure 13. Model output optimised to observational data from the Athabasca Glacier, Canada.