



## Strathprints Institutional Repository

**Bajaj, Deepti and Allerton, Brianne M. and Kirby, Joshua T. and Miller, Freeman and Rowe, David A. and Pohlig, Ryan T. and Modlesky, Christopher M. (2015) Muscle volume is related to trabecular and cortical bone architecture in typically developing children. BONE, 81. pp. 217-227. ISSN 8756-3282 , <http://dx.doi.org/10.1016/j.bone.2015.07.014>**

This version is available at <http://strathprints.strath.ac.uk/55256/>

**Strathprints** is designed to allow users to access the research output of the University of Strathclyde. Unless otherwise explicitly stated on the manuscript, Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Please check the manuscript for details of any other licences that may have been applied. You may not engage in further distribution of the material for any profitmaking activities or any commercial gain. You may freely distribute both the url (<http://strathprints.strath.ac.uk/>) and the content of this paper for research or private study, educational, or not-for-profit purposes without prior permission or charge.

Any correspondence concerning this service should be sent to Strathprints administrator: [strathprints@strath.ac.uk](mailto:strathprints@strath.ac.uk)

Manuscript Number: BONE-D-14-00403R2

Title: Muscle volume is related to trabecular and cortical bone architecture in typically developing children

Article Type: Original Full Length Article

Keywords: muscle, bone architecture, trabecular, cortical, femur length, MRI

Corresponding Author: Dr. Christopher Modlesky, PhD

Corresponding Author's Institution: University of Delaware

First Author: Deepti Bajaj, M.S.

Order of Authors: Deepti Bajaj, M.S.; Brianne M Allerton, M.S.; Joshua T Kirby, M.S.; Freeman Miller, MD; David A Rowe, PhD; Ryan Pohligh, PhD; Christopher Modlesky

**Abstract:** Introduction: Muscle is strongly related to cortical bone architecture in children; however, the relationship between muscle volume and trabecular bone architecture is poorly studied. The aim of this study was to determine if muscle volume is related to trabecular bone architecture in children and if the relationship is different than the relationship between muscle volume and cortical bone architecture.

**Materials and methods:** Forty typically developing children (20 boys and 20 girls; 6 to 12 y) were included in the study. Measures of trabecular bone architecture [apparent trabecular bone volume to total volume (appBV/TV), trabecular number (appTb.N), trabecular thickness (appTb.Th), and trabecular separation (appTb.Sp)] in the distal femur, cortical bone architecture [cortical volume, medullary volume, total volume, polar moment of inertia (J) and section modulus (Z)] in the midfemur, muscle volume in the midhigh and femur length were assessed using magnetic resonance imaging. Total and moderate-to-vigorous physical activity were assessed using an accelerometer-based activity monitor worn around the waist for four days. Calcium intake was assessed using diet records. Relationships among the measures were tested using multiple linear regression analysis.

**Results:** Muscle volume was moderately-to-strongly related to measures of trabecular bone architecture [appBV/TV ( $r = 0.81$ ), appTb.N ( $r = 0.53$ ), appTb.Th ( $r = 0.67$ ), appTb.Sp ( $r = -0.71$ ); all  $p < 0.001$ ] but more strongly related to measures of cortical bone architecture [cortical volume ( $r = 0.96$ ), total volume ( $r = 0.94$ ), Z ( $r = 0.94$ ) and J ( $r = 0.92$ ); all  $p < 0.001$ ]. Similar relationships were observed between femur length and measures of trabecular ( $p < 0.01$ ) and cortical ( $p < 0.001$ ) bone architecture. Sex, physical activity and calcium intake were not related to any measure of bone architecture ( $p > 0.05$ ). Because muscle volume and femur length were strongly related ( $r = 0.91$ ,  $p < 0.001$ ), muscle volume was scaled for femur length (muscle volume/femur length<sup>2.77</sup>). When muscle volume/femur length<sup>2.77</sup> was included in a regression model with femur length, sex, physical activity and calcium intake, muscle volume/femur length<sup>2.77</sup> was a significant predictor of appBV/TV, appTb.Th and appTb.Sp (partial  $r = 0.44$  to  $0.49$ ,  $p < 0.05$ ) and all measures of cortical bone architecture (partial  $r = 0.47$  to  $0.54$ ;  $p < 0.01$ ).

**Conclusions:** The findings suggest that muscle volume in the midhigh is related to trabecular bone architecture in the distal femur of children. The relationship is weaker than the relationship between muscle volume in the midhigh and cortical bone architecture in the midfemur, but the discrepancy is driven, in large part, by the greater dependence of cortical bone architecture measures on femur length

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Title: Muscle volume **is** related to trabecular **and cortical** bone architecture in **typically developing** children

Authors: Deepti Bajaj<sup>1</sup>, MS; Brianne M. Allerton<sup>1</sup>, MS, Joshua T. Kirby<sup>1</sup>, MS; Freeman Miller<sup>2</sup>, MD, David A. Rowe<sup>3</sup>, PhD, **Ryan T. Pohlig<sup>4</sup>, PhD**, Christopher M. Modlesky<sup>1</sup>, PhD

<sup>1</sup>Department of Kinesiology and Applied Physiology, University of Delaware, Newark, DE 19711

<sup>2</sup>Department of Orthopedics, Nemours AI duPont Hospital for Children, Wilmington, DE

<sup>3</sup>School of Psychological Sciences and Health, University of Strathclyde, Glasgow, UK

<sup>4</sup>**Biostatistics Core Facility, University of Delaware, Newark, DE 19713**

Abbreviations: apparent bone volume to total volume, appBV/TV; apparent trabecular number, appTb.N; apparent trabecular thickness, appTb.Th; apparent trabecular separation, appTb.Sp; section modulus, Z; polar moment of inertia, J

Address of correspondence and reprint requests:

Christopher M. Modlesky, PhD  
Department of Kinesiology and Applied Physiology  
University of Delaware  
Newark, DE 19716  
Telephone: (302) 831 4185  
Fax: (302) 831 43693  
Email: [modlesky@udel.edu](mailto:modlesky@udel.edu)

Conflicts of interest: The authors declare that they have no conflict of interest

1  
2  
3  
4 Abstract

5  
6 Introduction: Muscle is strongly related to cortical bone architecture in children; however, the  
7 relationship between muscle volume and trabecular bone architecture is poorly studied. The aim  
8 of this study was to determine if muscle volume is related to trabecular bone architecture in  
9 children and if the relationship is different than the relationship between muscle volume and  
10 cortical bone architecture.  
11  
12  
13  
14  
15  
16  
17

18 Materials and methods: Forty typically developing children (20 boys and 20 girls; 6 to 12 y)  
19 were included the study. Measures of trabecular bone architecture [apparent trabecular bone  
20 volume to total volume (appBV/TV), trabecular number (appTb.N), trabecular thickness  
21 (appTb.Th), and trabecular separation (appTb.Sp)] in the distal femur, cortical bone architecture  
22 [(cortical volume, medullary volume, total volume, polar moment of inertia (J) and section  
23 modulus (Z)] in the midfemur, muscle volume in the midthigh and femur length were assessed  
24 using magnetic resonance imaging. Total and moderate-to-vigorous physical activity were  
25 assessed using an accelerometer-based activity monitor worn around the waist for four days.  
26 Calcium intake was assessed using diet records. Relationships among the measures were tested  
27 using multiple linear regression analysis.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

43 Results: Muscle volume was moderately-to-strongly related to measures of trabecular bone  
44 architecture [appBV/TV ( $r = 0.81$ ), appTb.N ( $r = 0.53$ ), appTb.Th ( $r = 0.67$ ), appTb.Sp ( $r = -0.71$ ;  
45 all  $p < 0.001$ ] but more strongly related to measures of cortical bone architecture [cortical  
46 volume ( $r = 0.96$ ), total volume ( $r = 0.94$ ), Z ( $r = 0.94$ ) and J ( $r = 0.92$ ; all  $p < 0.001$ )]. Similar  
47 relationships were observed between femur length and measures of trabecular ( $p < 0.01$ ) and  
48 cortical ( $p < 0.001$ ) bone architecture. Sex, physical activity and calcium intake were not related  
49 to any measure of bone architecture ( $p > 0.05$ ). Because muscle volume and femur length were  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 strongly related ( $r = 0.91$ ,  $p < 0.001$ ), muscle volume was scaled for femur length (muscle  
5  
6 volume/femur length<sup>2.77</sup>). When muscle volume/femur length<sup>2.77</sup> was included in a regression  
7  
8 model with femur length, sex, physical activity and calcium intake, muscle volume/femur  
9  
10 length<sup>2.77</sup> was a significant predictor of appBV/TV, appTb.Th and appTb.Sp (partial  $r = 0.44$  to  
11  
12 049,  $p < 0.05$ ) and all measures of cortical bone architecture (partial  $r = 0.47$  to 054;  $p < 0.01$ ).  
13  
14

15  
16 Conclusions: The findings suggest that muscle volume in the midhigh is related to trabecular  
17  
18 bone architecture in the distal femur of children. The relationship is weaker than the relationship  
19  
20 between muscle volume in the midhigh and cortical bone architecture in the midfemur, but the  
21  
22 discrepancy is driven, in large part, by the greater dependence of cortical bone architecture  
23  
24 measures on femur length.  
25  
26  
27  
28  
29  
30

31 Key words: muscle, bone architecture, trabecular, cortical, femur length, MRI  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Introduction

Childhood is viewed as a critical time to maximize the mass, architecture and strength of bone and minimize fracture risk later in life [1, 2]. Areal bone mineral density (aBMD) is the single best surrogate of fracture risk widely available. However, there is a significant overlap in aBMD in those who have and have not fractured [3]. Moreover, using aBMD alone to make clinical decisions in children has been questioned [4, 5]. The development of more sophisticated imaging techniques, such as magnetic resonance imaging and computed tomography, now allow for the separation on bone into different compartments and for the evaluation of different architectural features of trabecular and cortical bone. The architecture of bone is an important contributor to its biomechanical strength [6]. Magnetic resonance imaging is particularly attractive for the study of bone in children because it does not expose them to ionizing radiation.

Due to the importance of bone architecture, identifying factors that contribute to its development could lead to an immediate and long-term reduction in fracture risk. Muscle clearly has a positive influence on bone which has led to the proposal of muscle-bone indexes that reflect the relationship between muscle and bone [7]. In theory, muscle-bone indexes can be used to identify those at greatest risk for fracture or if factors other than muscle are contributing to poor bone development [8]. Although the relationship between muscle and cortical bone architecture in children has been studied [9, 10], the relationship between muscle and trabecular bone architecture remains unclear. Because other factors may influence the development of bone architecture, such as general growth, as indicated by bone length, sex, physical activity and dietary factors, such as calcium intake, these factors must be considered when evaluating the relationship between muscle and trabecular bone architecture.

1  
2  
3  
4 The primary aim of this study was to determine if muscle volume is related to measures  
5 of trabecular bone architecture in typically developing children and if the relationship is different  
6 than the relationship between muscle volume and cortical bone architecture. We hypothesized  
7 that thigh muscle volume would be related to measures of trabecular and cortical bone  
8 architecture in the distal femur of typically developing children while controlling for femur  
9 length, sex, physical activity and calcium intake.  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20

## 21 Methods

### 22 Subjects

23  
24  
25  
26 Typically developing children 6 to 12 years of age, between the 5<sup>th</sup> and 95<sup>th</sup> percentile for  
27 height and body mass, with no history of lower extremity fracture, no intramedullary fixation in  
28 the femur or tibia, no use of chronic medications known to affect bone or muscle development  
29 and not participating in > 3 hours of organized physical activity per week were recruited from the  
30 Newark, DE, USA community by distributing flyers. The procedures followed were in  
31 accordance with the ethical standards of the institutions on human experimentation. The  
32 Institutional Review Boards at the University of Delaware and the AI duPont Hospital for  
33 Children approved the study. Written consent from parents and written assent from children, if  
34 able, were taken before testing.  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

### 50 Protocol

51  
52  
53 For each participant, height, body mass, sexual maturity, physical activity, calcium  
54 intake, measures of trabecular and cortical bone architecture, muscle volume and femur length  
55 were assessed and recorded within a two week period.  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7 Anthropometrics

8  
9 Children wore shorts, a t-shirt and no shoes during anthropometric assessment. Height  
10 was measured to the nearest 0.1 cm using a wall-mounted stadiometer (Seca model 222, Novel  
11 products, Rockton, IL). Body mass was rounded to the nearest 0.2 kg using a digital scale  
12 (Detecto 6550, Cardinal Scale, MO). Height, body mass and body mass index (BMI) percentiles  
13  
14 were calculated from the normative graphs published by the Centers for Disease Control and  
15  
16 Prevention [11].  
17  
18  
19  
20  
21  
22  
23  
24  
25

26 Tanner staging

27  
28 Sexual maturity was assessed by a physician assistant using the Tanner staging technique  
29 [12]. Signs of pubic hair growth were assessed along with testicular/penile development in boys,  
30  
31 and breast development in girls. The Tanner stages range from 1 to 5, with 1 indicating no signs  
32  
33 of sexual development (pre-pubertal), 2 indicating early sexual development and 5 indicating full  
34  
35 sexual development.  
36  
37  
38  
39  
40  
41  
42

43 Physical activity

44  
45 Physical activity was assessed using an accelerometer-based activity monitor (Actical;  
46 Philips Respironics, Sunriver, OR) that was worn on the non-dominant hip for four days (three  
47  
48 weekdays and one weekend day). Four days of data collection was chosen because it yields  
49  
50 accurate and reliable estimates of physical activity [13]. The children were asked not to remove  
51  
52 the activity monitor at any time (including sleep) except for swimming at a depth greater than  
53  
54 three feet or while bathing/showering. Activity counts were registered in 15 second epochs.  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  
4 Total physical activity was reported in average counts/d. Moderate-to-vigorous physical activity  
5 was reported in average counts/d, average minutes/d and average counts/minute/d. Physical  
6 activity was divided into moderate and vigorous levels based on the work of Puyau et al. [14]  
7 with 1500-6500 counts per minute classified as moderate activity and > 6500 counts per minute  
8 classified as vigorous activity. Only participants who wore the activity monitors for at least three  
9 complete days were included in the final analysis. This was confirmed by reviewing physical  
10 activity logs that were kept by the child with the assistance from the parent and by reviewing the  
11 graphical output generated by the activity monitor software.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

## 26 Calcium intake

27  
28 Children, with the aid of a parent, completed a diet record for two days during the week  
29 and one day on the weekend. To facilitate accurate quantification of foods, each participant and  
30 their parent received a list of serving size estimates based on comparisons to everyday objects  
31 (e.g. 3 oz of meat or poultry is approximately the size of a deck of cards). Calcium intake was  
32 estimated using the diet records and the USDA Food and Nutrient Database for Dietary Studies,  
33 v. 1.0, as previously done [15]. Only participants who recorded diet for three complete days  
34 were included in the final analysis.  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

## 48 Magnetic resonance imaging

49  
50 Trabecular bone architecture at the distal femur and cortical bone architecture and muscle  
51 volume at the level of the middle-third of the femur on the non-dominant side were estimated  
52 using MRI (GE 1.5 T; Milwaukee, WI). Children were immobilized from the waist down using  
53 the BodyFIX (Medical Intelligence, Inc., Schwabmünchen, GER), as previously described [16].  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 Different imaging protocols were used to collect images needed for the assessment of trabecular  
5  
6 bone architecture and cortical bone architecture.  
7  
8

9 Trabecular bone architecture: A phased array coil (USA Instruments; Aurora, OH) was  
10 secured to the lateral portion of the distal femur using a VacFIX system (PAR Scientific A/S;  
11 Sivlandvaenge, Denmark). The lateral distal femur was identified using a three-plane localizer.  
12  
13 Then 26 high resolution axial images were collected immediately above the growth plate in the  
14 metaphysis using a 3D fast gradient echo sequence with a partial echo acquisition (4.5 ms echo  
15 time; 30 ms repetition time; 30° flip angle; 13.89 kHz bandwidth; 9 cm field of view) and an  
16 imaging matrix of 512 x 384 reconstructed to a 175 x 175 x 700  $\mu\text{m}^3$  voxel size. Measures of  
17 trabecular bone architecture [i.e., appBV/TV, appTb.N (1/mm), appTb.Th (mm), and appTb.Sp  
18 (1/mm)] in the 20 most central images were estimated using custom software created with  
19 Interactive Data Language (IDL; Research Systems, Inc, Boulder, CO) and a procedure similar  
20 to that described by Majumdar et al. [17]. The procedure has been described in detail previously  
21 [16]. One investigator oversaw the collection of all images and one research assistant conducted  
22 all image analysis. In our laboratory, the reproducibility of the trabecular bone architecture  
23 assessment in children using this procedure ranges from 2 to 3 % [16].  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

43 Cortical bone architecture, muscle volume and femur length: Axial images (1 cm thick  
44 and 0.5 cm spacing) were collected along the length of the nondominant femur using a torso PA  
45 coil (750 ms repetition time, 14 ms echo time, 16 cm field of view and a 512 x 512 imaging  
46 matrix). Before image collection, the femur was identified using a three-plane localizer. The  
47 region of interest box was carefully positioned using the proximal end of the femoral head and  
48 the distal end of the femoral condyles to indicate the top and bottom, respectively, of the region  
49 of interest box. Using the localizer in the coronal plane, the region of interest box was rotated so  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 its midline was placed alongside the lateral portion of the femoral head and ran through the  
5  
6 center of the interchondalar notch between the femoral condyles. Using the localizer in the  
7  
8 sagittal plane, the region of interest box was positioned so its midline went through the center of  
9  
10 the femoral head and the center of the medial condyle. Images at the level of the middle third of  
11  
12 the femur were analyzed for measures of cortical bone architecture and strength and muscle  
13  
14 volume using software designed in-house with IDL. Images were first filtered with a median  
15  
16 filter to reduce pixel noise. The filtered images were then subjected to automatic segmentation  
17  
18 with a fuzzy C-means clustering algorithm. Using the software, voxels representing cortical  
19  
20 bone, the medullary cavity and muscle at the level of the mid-third of the femur were identified  
21  
22 and summed to determine their cross-sectional areas. Their volumes were quantified by  
23  
24 accounting for image thickness and spacing between images. Middle images were multiplied by  
25  
26 1.5 to account for the 1.0 cm slice thickness of each image and the 0.5 cm gap between images.  
27  
28 Images that included the proximal and distal ends of the region of interest were multiplied by an  
29  
30 appropriate correction factor ( $< 1.5$ ) so the entire image set represented the middle third of the  
31  
32 femur. Volume ( $\text{cm}^3$ ) of the total midfemur was determined by summing cortical volume and  
33  
34 medullary volume. Femur length was determined by counting the number of images and  
35  
36 adjusting for image thickness and spacing. Cross-sectional moment of inertia of the midfemur in  
37  
38 the anterior-posterior ( $\text{CSMI}_{\text{ap}}$ ;  $\text{cm}^4$ ) and medial-lateral ( $\text{CSMI}_{\text{ml}}$ ;  $\text{cm}^4$ ) directions were  
39  
40 determined using the parallel-axis theorem [18]. Polar moment of inertia ( $J$ ;  $\text{cm}^4$ ) was  
41  
42 determined by adding  $\text{CSMI}_{\text{ap}}$  and  $\text{CSMI}_{\text{ml}}$ . Section modulus ( $Z$ ;  $\text{cm}^3$ ) in the anterior-posterior  
43  
44 and medial-lateral directions was calculated by dividing  $\text{CSMI}_{\text{ap}}$  and  $\text{CSMI}_{\text{ml}}$  by the furthest  
45  
46 distance from the neutral axis in the anterior-posterior and medial-lateral directions, respectively.  
47  
48 The average  $Z$  from the two directions is reported. One investigator oversaw the collection of all  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 images and one research assistant conducted all image analysis. In our laboratory, the  
5  
6 reproducibility of the cortical bone architecture and muscle volume estimates ranges from 0.5 to  
7  
8 2.6 % [19, 20]. To assess the reproducibility of femur length measurements by MRI, five  
9  
10 children (6 to 12 years) were tested twice within 1 week. Tests were either on the same day or  
11  
12 on a separate day. Femur length estimates from the repeat tests were identical.  
13  
14  
15  
16  
17  
18

### 19 Statistical analysis

20  
21 Data were analyzed using SPSS (Version 22.0, Chicago, IL). Descriptive statistics for all  
22  
23 variables were conducted to screen for outliers and to assess normality using the Shapiro Wilk  
24  
25 test. Sex differences in physical characteristics, including muscle volume and measures of  
26  
27 trabecular and cortical bone architecture, were tested using independent t-tests for normally  
28  
29 distributed data and Mann Whitney U tests otherwise. Height, body mass and BMI percentiles  
30  
31 were compared to their respective 50<sup>th</sup> age- and sex-based percentile using a one-sample t-test.  
32  
33 Sex by Tanner stage comparisons were conducted using the Chi-square test of independence.  
34  
35  
36 Zero-order correlation analysis was used to assess relationships between muscle volume and  
37  
38 measures of trabecular and cortical bone architecture and to screen for multicollinearity among  
39  
40 the predictors of bone architecture (i.e., muscle volume, age, femur length, sex, physical activity  
41  
42 and calcium intake). Multiple linear regression was used to determine the amount of variance in  
43  
44 measures of bone architecture explained by muscle volume after controlling for femur length,  
45  
46 sex, physical activity and calcium intake. Age was excluded because it was strongly related to  
47  
48 muscle volume ( $r = 0.86$ ) and femur length ( $r = 0.92$ ) causing a high variance inflation factor  
49  
50 when it was introduced into the regression models ( $> 10$ ). In addition, although all measures of  
51  
52 trabecular and cortical bone architecture were related to age ( $r$  ranged from  $-0.57$  to  $0.66$ , and  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 0.78 to 0.85, respectively, all  $p < 0.01$ ), they were more strongly related to femur length. Muscle  
5  
6 volume was also strongly correlated with femur length ( $r = 0.91$ ,  $p < 0.001$ ), resulting in a high  
7  
8 variance inflation factor when both variables were included in regression models ( $> 6$ ).  
9  
10 Therefore, a scaling factor was created to determine the contribution of muscle volume to  
11  
12 variability in measures of trabecular and cortical bone architecture independent of femur length.  
13  
14 The scaling factor was created using a log-log regression analysis [21]. Muscle volume and  
15  
16 femur length were both natural log transformed and the transformed muscle volume was  
17  
18 regressed on transformed femur length. The regression slope (2.77) represents the power by  
19  
20 which femur length was raised to reduce collinearity. The resulting index (muscle volume/femur  
21  
22 length<sup>2.77</sup>) was unrelated to femur length ( $r = 0.03$ ) suggesting that it would be independent of  
23  
24 femur length in a regression model. In addition, when it was included in a regression model with  
25  
26 femur length, sex, MVPA and calcium intake, the variance inflation factor was low ( $< 1.5$ ).  
27  
28 Standardized ( $\beta$ ) coefficients and partial correlations were reported to express the degree of  
29  
30 variance in measures of bone architecture explained by a variable while controlling for the other  
31  
32 covariates. Alpha level was set at 0.05.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

## 43 Results

44  
45 Fifty one children ( $n = 25$  boys and 26 girls) were enrolled in the study. Four boys were  
46  
47 excluded because they did not complete diet records. One boy was excluded because he did not  
48  
49 complete MRI testing. Five girls were excluded because they had a Tanner stage  $> 2$ . One girl  
50  
51 was excluded because she did not complete the MRI testing. Twenty boys and twenty girls were  
52  
53 included in the final analysis. Physical characteristics, physical activity and calcium intake are  
54  
55 listed in Table 1. All data were normally distributed except the measures of physical activity,  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 which were log transformed before statistical analyses were conducted. However, non-  
5  
6 transformed physical activity data are reported in Table 1 for easier comparison with other  
7  
8 studies. There were no differences in age, Tanner stages, height, body mass or BMI, or age- and  
9  
10 sex-based percentiles for height, body mass or BMI between boys and girls. Percentiles for  
11  
12 height, body mass, and BMI were not different from the 50th age- and sex-based percentiles ( $p >$   
13  
14  $0.05$ ). There were also no differences in any measure of physical activity or calcium intake  
15  
16 between boys and girls ( $p > 0.05$ ). There were no differences in measures of trabecular and  
17  
18 cortical bone architecture between boys and girls ( $p > 0.05$ ), as shown in Table 2.  
19  
20  
21  
22

23  
24 Zero-order correlations between measures of trabecular and cortical bone architecture and  
25  
26 muscle volume, femur length, sex, MVPA and calcium intake are in Table 3. Muscle volume  
27  
28 was moderately-to-strongly and positively related to appBV/TV, appTb.N and appTb.Th and  
29  
30 moderately-to-strongly and negatively related to appTb.Sp (all  $p < 0.001$ ). Femur length was  
31  
32 also moderately-to-strongly and positively related to appBV/TV, appTb.N and appTb.Th and  
33  
34 moderately-to-strongly and negatively related to appTb.Sp (all  $p < 0.01$ ). These relationships are  
35  
36 demonstrated by scatter plots in Figure 1. Muscle volume and femur length were strongly and  
37  
38 positively related to cortical volume, total volume, Z and J (all  $p < 0.001$ ). These relationships  
39  
40 are demonstrated by scatter plots in Figure 2. No other variables were significantly correlated  
41  
42 with measures of trabecular or cortical bone architecture.  
43  
44  
45  
46  
47

48 Results from the multiple regression analysis with muscle volume/femur length<sup>2.77</sup>, femur  
49  
50 length, sex, MVPA and calcium intake as independent variables and measures of trabecular bone  
51  
52 architecture as dependent variables are reported in Table 4. The only significant predictors of  
53  
54 appBV/TV, appTb.Th and appTb.Sp were muscle volume/femur length<sup>2.77</sup> (all  $p < 0.05$ ) and  
55  
56 femur length (all  $p < 0.001$ ). The only significant predictor of appTb.N was femur length ( $p =$   
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 0.004); however, muscle volume/femur length<sup>2.77</sup> was only marginally insignificant (p = 0.059).  
5  
6  
7 The relationships are demonstrated by scatter plots of residuals in Figure 3.

8  
9 Results from the multiple regression analysis with muscle volume/femur length<sup>2.77</sup>, femur  
10 length, sex, MVPA and calcium intake as independent variables and measures of cortical bone  
11 architecture as dependent variables are reported in Table 5. The only significant predictors of  
12 measures of cortical bone architecture were muscle volume/femur length<sup>2.77</sup> (all p < 0.01) and  
13 femur length (all p < 0.001). The relationships are demonstrated by scatter plots of residuals in  
14 Figure 4.  
15  
16  
17  
18  
19  
20  
21  
22

23 For each of the multiple regression models, the patterns of the relationships were the  
24 same irrespective of the type of physical activity (i.e., total physical activity or MVPA) or unit of  
25 physical activity (i.e., counts/d, min/d or counts/min/d) included in the model.  
26  
27  
28  
29  
30  
31  
32

### 33 Discussion

34  
35 To our knowledge, this is the first study to examine whether muscle is related to  
36 measures of trabecular bone architecture in children. A primary finding was that mid thigh  
37 muscle volume was positively related to appBV/TV, appTb.N and appTb.Th and negatively  
38 related to appTb.Sp in the distal femur of pre and early-pubertal children. The present study is  
39 also the first to determine if the relationship between muscle volume and trabecular bone  
40 architecture is different than the relationship between muscle volume and cortical bone  
41 architecture in children. Mid thigh muscle volume was more strongly related to measures of  
42 cortical than trabecular bone architecture. However, the magnitude of the disparity in the  
43 relationships was much smaller when femur length, sex, physical activity and calcium intake  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 were statistically controlled. The findings suggest that muscle volume is an independent  
5  
6 contributor to the development of trabecular and cortical bone architecture in children.  
7  
8

9 The moderate-to-strong relationship between muscle volume and trabecular bone  
10 architecture in children that was observed in the present study is consistent with the pattern  
11 reported in animals [22, 23] and humans [24, 25]. For example, muscle paralysis induced by  
12 botulinum toxin has been shown to decrease trabecular thickness of the mandible in rats and  
13 decrease BV/TV, Tb.N and Tb.Th and increase in Tb.Sp in the proximal tibia of mice [23].  
14 Conversely, muscle contraction induced by electrical stimulation has been shown to blunt the  
15 loss of BV/TV and Tb.N associated with hindlimb suspension in rats [26]. In humans,  
16 conditions that result in a marked loss of muscle volume and strength in adults or poor accretion  
17 in children, such as spinal cord injury [25] and cerebral palsy [16], are associated with impaired  
18 trabecular bone architecture. Moreover, diminished trabecular bone architecture is associated  
19 with low muscle volume and low muscle strength [27]. In addition, relationships between  
20 appendicular muscle volume predicted by dual-energy X-ray absorptiometry and measures of  
21 trabecular bone architecture have been observed in men and women 20-97 years of age [28].  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 The positive correlations between muscle and measures of cortical bone architecture in  
42 the present study are consistent with previous reports [7, 9, 10, 29-31]. A positive moderate  
43 correlation between muscle cross sectional area and tibial cortical area was observed in 13 year  
44 olds with different ethnicities (r ranged from 0.45 – 0.60, p < 0.001) [29]. A moderate-to-strong  
45 relationship between muscle (or lean mass) and cortical bone architecture has also been reported  
46 in adult athletes and non-athletes [7], chronic stroke patients [30], patients with spinal cord injury  
47 [31], and children, adolescents and young adults [9, 10]. Despite the differences in research  
48 designs, the results of previous studies are generally consistent with ours. In the present study,  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  
4 although midthigh muscle volume was not as strongly related to measures of trabecular bone  
5  
6 architecture (r ranged from 0.53 to 0.81) as it was to measures of cortical bone architecture (r  
7  
8 ranged from 0.92 to 0.96), the relationships were still moderate-to-strong. The disparity suggests  
9  
10 that muscle volume may be more closely linked to cortical than trabecular bone. However, the  
11  
12 degree of the discrepancy was reduced substantially when femur length was statistically  
13  
14 controlled (partial r ranged from -0.44 to 0.49 for trabecular bone architecture and -0.47 to 0.56  
15  
16 for cortical bone architecture). Therefore, it appears that the discrepancy in muscle's  
17  
18 relationship with trabecular vs. cortical bone is driven, in large part, by the dependence of the  
19  
20 cortical bone architecture measures on femur length. It is also possible that the difference in the  
21  
22 strength of the relationships may be linked to the muscle and bone sites studied. Trabecular bone  
23  
24 architecture was assessed in the distal femur, whereas cortical bone architecture and muscle  
25  
26 volume were assessed at the level of the midshaft of the femur. While the shaft of the femur is  
27  
28 an attachment site for many thigh muscles (i.e., vastus intermedius, lateralis and medialis,  
29  
30 adductor longus and magnus, etc.), the distal femur is an attachment site for only one thigh  
31  
32 muscle (i.e., adductor magnus). It is also plausible that muscle volume's stronger relationship  
33  
34 with measures of cortical than trabecular bone architecture may be linked to the limited accuracy  
35  
36 of the trabecular bone measures using MRI [32]. Because the trabeculae are small and their size  
37  
38 is at the limit of the resolution than can be used to assess trabecular bone architecture with MRI,  
39  
40 the measures suffer from partial volume effects [32]. This leads to appBV/TV and appTb.N  
41  
42 overestimating actual trabecular bone volume to total volume and trabecular number,  
43  
44 respectively, and appTb.Sp underestimating actual trabecular separation. Nonetheless,  
45  
46 appBV/TV, appTb.N and appTb.Sp are correlated with their intended measures [33] and they  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 discriminate individuals who have osteoporotic fractures from those who do not have fracture  
5  
6 [34].  
7

8  
9 To our knowledge, this is also the first study to examine the relationship between a  
10 measure of bone size (i.e., bone length) and measures of trabecular bone architecture in children.  
11 Femur length, like midthigh muscle volume, was moderately-to-strongly and positively related to  
12 appBV/TV, appTb.N and appTb.Th and negatively related to appTb.Sp in the distal femur of pre  
13 and early-pubertal children. Also, similar to muscle volume, femur length was not as strongly  
14 related to measures of trabecular bone architecture as it was to measures of cortical bone  
15 architecture. Because femur length was highly correlated with muscle volume, the unique  
16 variance in trabecular bone architecture and cortical bone architecture explained by muscle  
17 volume vs. femur length is difficult to separate. In an attempt to resolve the issue of collinearity,  
18 muscle volume was scaled for femur length (i.e., muscle volume/femur length<sup>2.77</sup>). When femur  
19 length was included in a multiple regression model with muscle volume/femur length<sup>2.77</sup>, sex,  
20 physical activity and calcium intake, femur length was a significant predictor of appBV/TV,  
21 appTb.N and appTb.Sp. This finding suggests that femur length is related to measures of  
22 trabecular and cortical bone architecture independent of muscle volume. Future studies that  
23 determine if measures of trabecular and cortical bone architecture track changes in muscle due to  
24 intervention, such as resistance training, or unloading due to a neurological insult, such as spinal  
25 cord injury, would provide valuable insight into their relationship with muscle independent of  
26 bone size or general growth.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

53 Another important finding in the present study was that there was no detectable sex-effect  
54 on the relationship between muscle and bone architecture. Although boys are stronger than girls  
55 [35], previous studies have reported no sex disparity in the relationship between muscle and bone  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 in pre- and early-pubertal children [36, 37]. Sex hormones have been shown to affect the  
5  
6 muscle-bone relationship, but this influence may not be observed until puberty. In a study of 318  
7  
8 boys and girls 6 to 22 years of age, Schoenau et al. [36] reported that girls have higher cortical  
9  
10 area than boys at the onset of puberty due to the surge in estrogen. On the other hand, another  
11  
12 cross-sectional comparison of children through young adults (6 to 21 years of age) suggests that  
13  
14 appBV/TV and appTb.Th become higher and appTb.Sp becomes lower in boys during late  
15  
16  
17 appBV/TV and appTb.Th become higher and appTb.Sp becomes lower in boys during late  
18  
19 puberty and this may be related to the surge in testosterone in boys [38].  
20

21 One of the strengths of the present study was that the independent relationship between  
22  
23 muscle volume and bone architecture was examined while muscle volume was scaled for femur  
24  
25 length. This allowed us to control muscle volume for femur length while avoiding the  
26  
27 collinearity associated with the two variables. Evaluating bone architecture's relationship with  
28  
29 muscle volume while statistically controlling for physical activity was another strength of the  
30  
31 study. High levels of physical activity, especially MVPA, have a positive influence on muscle  
32  
33 [39] and bone [40-42]. Therefore, it is plausible that the relationship between muscle and bone  
34  
35 architecture is driven, at least in part, by physical activity. However, the relationship existed  
36  
37 even when total physical activity or MVPA was used as a covariate in a regression model.  
38  
39 Statistically controlling for calcium intake is another strength of the study because higher bone  
40  
41 mineral content is seen in children with a regular dietary intake of calcium [43]. Moreover, there  
42  
43 is evidence that adolescent boys retain more calcium than girls [44]. On the other hand, calcium  
44  
45 intake is not associated with increases in lean mass [45]. Although it is plausible that calcium  
46  
47 intake can cause discord in the relationship between muscle and bone, the relationships between  
48  
49 muscle volume and measures of bone architecture remained even when calcium intake was used  
50  
51 as a covariate in a regression model.  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 We acknowledge that this study has limitations not already mentioned. One is the cross-  
5  
6 sectional design which does not allow us to make inferences about cause and effect. **Although**  
7  
8 **cortical bone was strongly related to muscle volume in the present study, some studies suggest**  
9  
10 **that cortical bone is more strongly related to impact loading than the bone strain generated by**  
11  
12 **muscle forces [46]. There is also evidence that changes in cortical bone architecture occur**  
13  
14 **before changes in muscle area during growth [47]. Therefore, additional studies are needed to**  
15  
16 **determine if changes in bone architecture track changes in muscle volume that occur with growth**  
17  
18 **or with intervention. A second limitation is that the sample size was small and restricted to**  
19  
20 **Caucasians which prevents us from generalizing our findings to children of other origins. The**  
21  
22 **small sample also limits our confidence in the finding that sex did not affect bone architecture's**  
23  
24 **relationship with muscle volume and femur length. However, only children who were pre- or**  
25  
26 **early-pubertal were included in the study and the finding of no sex-effect on the muscle-bone**  
27  
28 **relationship at this stage of maturity is consistent with previous studies [36, 37]. A third**  
29  
30 **limitation is related to the physical activity data. Only children who participated in three or**  
31  
32 **fewer hours of organized physical activity per week participated in the study. Furthermore,**  
33  
34 **assessment of physical activity was limited to four days. Although we quantified MVPA, and**  
35  
36 **MVPA for the boys and girls in the present study (92 and 88 min/d) were similar to values**  
37  
38 **previously reported for a sample of 6-11 year-old Caucasian boys and girls (average MVPA = 95**  
39  
40 **and 75 min/d, respectively)[48], appropriate thresholds used to classify physical activity into**  
41  
42 **different categories have not been agreed upon [49], especially for the activity monitors used in**  
43  
44 **the present study. Nonetheless, the results were the same irrespective of the physical activity**  
45  
46 **measure used (i.e., MVPA or total physical activity) or how it was expressed (i.e., counts/d,**  
47  
48 **min/d or counts/min/d).**  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 In summary, the findings from the present study suggest that mid thigh muscle volume is  
5 related to measures of trabecular bone architecture in the distal femur and measures of cortical  
6 bone architecture in the mid femur. The findings also suggest that the relationship of bone  
7 architecture with mid thigh muscle volume is not different in pre and early-pubertal boys and  
8 girls. The relationship remained after controlling for femur length, suggesting that it may not be  
9 driven solely by general growth. Although mid thigh muscle volume is more strongly related to  
10 cortical bone architecture in the mid femur than trabecular bone architecture in the distal femur,  
11 the reason for the discrepancy is due, in large part, to the greater dependence of cortical bone  
12 architecture measures on femur length. Longitudinal studies that track changes in trabecular and  
13 cortical bone architecture, muscle volume and bone length and intervention studies that facilitate  
14 increases in muscle size and strength would help us confirm whether muscle has a unique  
15 relationship with trabecular and cortical bone architecture in children.  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

### 36 Acknowledgments

37  
38 We are thankful to the children who participated in our study and their families for their  
39 time and cooperation. This study was funded by the National Institutes of Health (HD071397  
40 and HD505030) and the National Osteoporosis Foundation.  
41  
42  
43  
44  
45  
46  
47

### 48 References

- 49  
50 [1] Heaney RP, Abrams S, Dawson-Hughes B, Looker A, Marcus R, Matkovic V, Weaver C.  
51 Peak Bone Mass. Osteoporos Int 2000;11: 985-1009.  
52  
53 [2] Modlesky CM, Lewis RD. Does exercise during growth have a long-term effect on bone  
54 health? Exerc Sport Sci Rev 2002;30: 171-6.  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4 [3] Ciarelli TE, Fyhrie DP, Schaffler MB, Goldstein SA. Variations in three-dimensional  
5 cancellous bone architecture of the proximal femur in female hip fractures and in controls. J  
6 Bone Miner Res 2000;15: 32-40.  
7  
8  
9  
10  
11 [4] Crabtree NJ, Arabi A, Bachrach LK, Fewtrell M, El-Hajj Fuleihan G, Keckskemethy HH,  
12 Jaworski M, Gordon CM, International Society for Clinical D. Dual-energy X-ray  
13 absorptiometry interpretation and reporting in children and adolescents: the revised 2013 ISCD  
14 Pediatric Official Positions. J Clin Densitom 2014;17: 225-42.  
15  
16  
17  
18  
19  
20  
21 [5] Adams JE, Engelke K, Zemel BS, Ward KA, International Society of Clinical D.  
22 Quantitative computer tomography in children and adolescents: the 2013 ISCD Pediatric Official  
23 Positions. J Clin Densitom 2014;17: 258-74.  
24  
25  
26  
27  
28  
29 [6] Jiang Y, Zhao J, Augat P, Ouyang X, Lu Y, Majumdar S, Genant HK. Trabecular bone  
30 mineral and calculated structure of human bone specimens scanned by peripheral quantitative  
31 computed tomography: relation to biomechanical properties. J Bone Miner Res 1998;13: 1783-  
32 90.  
33  
34  
35  
36  
37  
38 [7] Rittweger J, Beller G, Ehrig J, Jung C, Koch U, Ramolla J, Schmidt F, Newitt D,  
39 Majumdar S, Schiessl H, Felsenberg D. Bone-muscle strength indices for the human lower leg.  
40 Bone 2000;27: 319-26.  
41  
42  
43  
44  
45  
46 [8] Schoenau E. The "functional muscle-bone unit": a two-step diagnostic algorithm in  
47 pediatric bone disease. Pediatr Nephrol 2005;20: 356-9.  
48  
49  
50  
51 [9] Leonard MB, Elmi A, Mostoufi-Moab S, Shults J, Burnham JM, Thayu M, Kibe L,  
52 Wetzsteon RJ, Zemel BS. Effects of sex, race, and puberty on cortical bone and the functional  
53 muscle bone unit in children, adolescents, and young adults. J Clin Endocrinol Metab 2010;95:  
54 1681-9.  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4 [10] Hogler W, Blimkie CJ, Cowell CT, Inglis D, Rauch F, Kemp AF, Wiebe P, Duncan CS,  
5  
6 Farpour-Lambert N, Woodhead HJ. Sex-specific developmental changes in muscle size and bone  
7  
8 geometry at the femoral shaft. *Bone* 2008;42: 982-9.  
9
- 10  
11 [11] Kuczmarski RJ, Ogden CL, Grummer-Strawn LM, Flegal KM, Guo SS, Wei, R. M, Z.,,  
12  
13 Curtin LR, Roche AF, Johnson CL. CDC growth charts: United States. *Adv Data* 2000: 1-27.  
14  
15  
16 [12] Tanner JA. *Growth and Adolescence*: Oxford: Blackwell Scientific Publications; 1962.  
17  
18  
19 [13] Trost SG, Pate RR, Freedson PS, Sallis JF, Taylor WC. Using objective physical activity  
20  
21 measures with youth: how many days of monitoring are needed? *Med Sci Sports Exerc* 2000;32:  
22  
23 426-31.  
24  
25  
26 [14] Puyau MR, Adolph AL, Vohra FA, Zakeri I, Butte NF. Prediction of activity energy  
27  
28 expenditure using accelerometers in children. *Med Sci Sports Exerc* 2004;36: 1625-1631.  
29  
30  
31 [15] Johnson DL, Miller F, Subramanian P, Modlesky CM. Adipose tissue infiltration of  
32  
33 skeletal muscle in children with cerebral palsy. *J Pediatr* 2009;154: 715-720.  
34  
35  
36 [16] Modlesky CM, Subramanian P, Miller F. Underdeveloped trabecular bone  
37  
38 microarchitecture is detected in children with cerebral palsy using high-resolution magnetic  
39  
40 resonance imaging. *Osteoporos Int* 2008;19: 169-76.  
41  
42  
43 [17] Majumdar S, Genant HK, Grampp S, Newitt DC, Truong VH, Lin JC, Mathur A.  
44  
45 Correlation of trabecular bone structure with age, bone mineral density, and osteoporotic status:  
46  
47 in vivo studies in the distal radius using high resolution magnetic resonance imaging. *J Bone*  
48  
49 *Miner Res* 1997;12: 111-8.  
50  
51  
52 [18] Turner CH, Burr DB. Experimental techniques for bone mechanics. In: Cowen SC,  
53  
54 editor. *Bone Mechanics Handbook*. Boca Raton: CRC Press; 2001, p. 1-35.  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4 [19] Modlesky CM, Kanoff SA, Johnson DL, Subramanian P, Miller F. Evaluation of the  
5 femoral midshaft in children with cerebral palsy using magnetic resonance imaging. *Osteoporos*  
6  
7  
8  
9 Int 2009;20: 609-15.
- 10  
11 [20] Modlesky CM, Cavaola ML, Smith JJ, Rowe DA, Johnson DL, Miller F. A DXA-based  
12 mathematical model predicts midthigh muscle mass from magnetic resonance imaging in  
13  
14 typically developing children but not in those with quadriplegic cerebral palsy. *J Nutr* 2010;140:  
15  
16  
17  
18  
19 2260-5.
- 20  
21 [21] Wells JC, Cole TJ. Adjustment of fat-free mass and fat mass for height in children aged 8  
22  
23  
24  
25 y. *Int J Obes Relat Metab Disord* 2002;26: 947-52.
- 26 [22] Tsai CY, Huang RY, Lee CM, Hsiao WT, Yang LY. Morphologic and bony structural  
27 changes in the mandible after a unilateral injection of botulinum neurotoxin in adult rats. *J Oral*  
28  
29  
30  
31  
32 Maxillofac Surg 2010;68: 1081-7.
- 33 [23] Poliachik SL, Bain SD, Threet D, Huber P, Gross TS. Transient muscle paralysis disrupts  
34  
35  
36  
37 bone homeostasis by rapid degradation of bone morphology. *Bone* 2010;46: 18-23.
- 38 [24] Pollock NK, Laing EM, Taylor RG, Baile CA, Hamrick MW, Hall DB, Lewis RD.  
39  
40  
41  
42  
43  
44 Comparisons of trabecular and cortical bone in late adolescent black and white females. *J Bone*  
45  
46  
47  
48  
49  
50  
51 Miner Metab 2011;29: 44-53.
- 52 [25] Slade JM, Bickel CS, Modlesky CM, Majumdar S, Dudley GA. Trabecular bone is more  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65 deteriorated in spinal cord injured versus estrogen-free postmenopausal women. *Osteoporos Int*  
2005;16: 263-72.
- [26] Lam H, Qin YX. The effects of frequency-dependent dynamic muscle stimulation on  
inhibition of trabecular bone loss in a disuse model. *Bone* 2008;43: 1093-100.



- 1  
2  
3  
4 [27] Szulc P, Blaizot S, Boutroy S, Vilayphiou N, Boonen S, Chapurlat R. Impaired bone  
5  
6 microarchitecture at the distal radius in older men with low muscle mass and grip strength: the  
7  
8 STRAMBO study. *J Bone Miner Res* 2013;28: 169-78.  
9
- 10  
11 [28] Lebrasseur NK, Achenbach SJ, Melton LJ, 3rd, Amin S, Khosla S. Skeletal muscle mass  
12  
13 is associated with bone geometry and microstructure and serum insulin-like growth factor  
14  
15 binding protein-2 levels in adult women and men. *J Bone Miner Res* 2012;27: 2159-69.  
16  
17
- 18  
19 [29] Micklesfield LK, Norris SA, Pettifor JM. Determinants of bone size and strength in 13-  
20  
21 year-old South African children: The influence of ethnicity, sex and pubertal maturation. *Bone*  
22  
23 2011;48: 777-85.  
24  
25
- 26 [30] Pang MY, Ashe MC, Eng JJ. Tibial bone geometry in chronic stroke patients: influence  
27  
28 of sex, cardiovascular health, and muscle mass. *J Bone Miner Res* 2008;23: 1023-30.  
29  
30
- 31 [31] Modlesky CM, Slade JM, Bickel CS, Meyer RA, Dudley GA. Deteriorated geometric  
32  
33 structure and strength of the midfemur in men with complete spinal cord injury. *Bone* 2005;36:  
34  
35 331-9.  
36  
37
- 38 [32] Majumdar S, Kothari M, Augat P, Newitt DC, Link TM, Lin JC, Lang T, Lu Y, Genant  
39  
40 HK. High-resolution magnetic resonance imaging: three-dimensional trabecular bone  
41  
42 architecture and biomechanical properties. *Bone* 1998;22: 445-54.  
43  
44
- 45 [33] Liu XS, Zhang XH, Rajapakse CS, Wald MJ, Magland J, Sekhon KK, Adam MF, Sajda  
46  
47 P, Wehrli FW, Guo XE. Accuracy of high-resolution in vivo micro magnetic resonance imaging  
48  
49 for measurements of microstructural and mechanical properties of human distal tibial bone. *J*  
50  
51 *Bone Miner Res* 2010;25: 2039-50.  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4 [34] Link TM, Majumdar S, Augat P, Lin JC, Newitt D, Lu Y, Lane NE, Genant HK. In vivo  
5 high resolution MRI of the calcaneus: differences in trabecular structure in osteoporosis patients.  
6  
7 J Bone Miner Res 1998;13: 1175-82.  
8  
9  
10  
11 [35] Newman DG, Pearn J, Barnes A, Young CM, Kehoe M, Newman J. Norms for hand grip  
12 strength. Arch Dis Child 1984;59: 453-9.  
13  
14  
15 [36] Schoenau E, Neu CM, Mokov E, Wassmer G, Manz F. Influence of puberty on muscle  
16 area and cortical bone area of the forearm in boys and girls. J Clin Endocrinol Metab 2000;85:  
17 1095-8.  
18  
19  
20  
21 [37] Schoenau E, Neu CM, Beck B, Manz F, Rauch F. Bone mineral content per muscle cross-  
22 sectional area as an index of the functional muscle-bone unit. J Bone Miner Res 2002;17: 1095-  
23 101.  
24  
25  
26 [38] Kirmani S, Christen D, van Lenthe GH, Fischer PR, Bouxsein ML, McCready LK,  
27 Melton LJ, 3rd, Riggs BL, Amin S, Muller R, Khosla S. Bone structure at the distal radius during  
28 adolescent growth. J Bone Miner Res 2009;24: 1033-42.  
29  
30  
31 [39] MacDougall JD, Ward GR, Sale DG, Sutton JR. Biochemical adaptation of human  
32 skeletal muscle to heavy resistance training and immobilization. J Appl Physiol Respir Environ  
33 Exerc Physiol 1977;43: 700-3.  
34  
35  
36 [40] Fuchs RK, Bauer JJ, Snow CM. Jumping improves hip and lumbar spine bone mass in  
37 prepubescent children: a randomized controlled trial. J Bone Miner Res 2001;16: 148-56.  
38  
39  
40  
41 [41] Janz KF, Burns TL, Levy SM, Torner JC, Willing MC, Beck TJ, Gilmore JM, Marshall  
42 TA. Everyday activity predicts bone geometry in children: the Iowa bone development study.  
43 Med Sci Sports Exerc 2004;36: 1124-31.  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4 [42] Bailey DA, McKay HA, Mirwald RL, Crocker PR, Faulkner RA. A six-year longitudinal  
5 study of the relationship of physical activity to bone mineral accrual in growing children: the  
6 university of Saskatchewan bone mineral accrual study. *J Bone Miner Res* 1999;14: 1672-9.  
7  
8  
9  
10 [43] Lee WTK, Leung SSF, Lui SSH, Lau J. Relationship between long-term calcium intake  
11 and bone mineral content of children aged from birth to 5 years. *Br J Nutr* 1993;70: 235-248.  
12  
13  
14 [44] Braun M, Martin BR, Kern M, McCabe GP, Peacock M, Jiang Z, Weaver CM. Calcium  
15 retention in adolescent boys on a range of controlled calcium intakes. *Am J Clin Nutr* 2006;84:  
16 414-8.  
17  
18  
19 [45] Smilowitz JT, Wiest MM, Teegarden D, Zemel MB, German JB, Van Loan MD. Dietary  
20 fat and not calcium supplementation or dairy product consumption is associated with changes in  
21 anthropometrics during a randomized, placebo-controlled energy-restriction trial. *Nutr Metab*  
22 (Lond) 2011;8: 67.  
23  
24 [46] Schipilow JD, Macdonald HM, Liphardt AM, Kan M, Boyd SK. Bone micro-  
25 architecture, estimated bone strength, and the muscle-bone interaction in elite athletes: an HR-  
26 pQCT study. *Bone* 2013;56: 281-9.  
27  
28  
29 [47] Xu L, Nicholson P, Wang Q, Alen M, Cheng S. Bone and muscle development during  
30 puberty in girls: a seven-year longitudinal study. *J Bone Miner Res* 2009;24: 1693-8.  
31  
32  
33 [48] Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical activity  
34 in the United States measured by accelerometer. *Med Sci Sports Exerc* 2008;40: 181-8.  
35  
36  
37 [49] Guinhouya BC, Samouda H, de Beaufort C. Level of physical activity among children  
38 and adolescents in Europe: a review of physical activity assessed objectively by accelerometry.  
39 *Public Health* 2013;127: 301-11.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 Text to Figures  
5

6 **Figure 1.** Scatter plots demonstrating the **unadjusted** relationships between measures of  
7 trabecular bone architecture in the distal femur (i.e., appBV/TV, appTb.N, appTb.Th, and  
8 appTb.Sp) and mid thigh muscle volume (A-D) and femur length (E-H) in pre- and early-  
9 pubertal boys ● and girls ○. **Because no sex effect was detected, the solid line represents the**  
10 **regression line for the combined sample.**  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20

21 **Figure 2.** Scatter plots demonstrating the **unadjusted** relationships between measures of cortical  
22 bone architecture in the mid femur (i.e., cortical volume, total volume, J, and Z) and mid thigh  
23 muscle volume (A-D) and femur length (E-H) in pre- and early-pubertal boys ● and girls ○.  
24  
25 **Because no sex effect was detected, the solid line represents the regression line for the combined**  
26 **sample.**  
27  
28  
29  
30  
31  
32  
33  
34  
35

36 **Figure 3.** Scatter plots demonstrating the relationships between measures of trabecular bone  
37 architecture in the distal femur (i.e., appBV/TV, appTb.N, appTb.Th, and appTb.Sp) and 1)  
38 mid thigh muscle volume/femur length<sup>2.77</sup> while statistically controlling for femur length, sex,  
39 MVPA (min/d) and calcium intake (A-D) and 2) femur length while statistically controlling for  
40 mid thigh muscle volume/femur length<sup>2.77</sup>, sex, MVPA (min/d) and calcium intake (E-H) in pre-  
41 and early-pubertal boys ● and girls ○. **Because no sex effect was detected, the solid line**  
42 **represents the regression line for the combined sample.**  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54

55 **Figure 4.** Scatter plots demonstrating the relationships between measures of cortical bone  
56 architecture in the mid femur (i.e., cortical volume, total volume, Z and J) and 1) mid thigh  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

muscle volume/femur length<sup>2.77</sup> while statistically controlling for femur length, sex, MVPA (min/d) and calcium intake (A-D) and 2) femur length while statistically controlling for muscle volume/femur length<sup>2.77</sup>, sex, MVPA (min/d) and calcium intake (E-H) in pre- and early-pubertal boys ● and girls ○. Because no sex effect was detected, the solid line represents the regression line for the combined sample.

Table 1. Physical characteristics of pre- and early-pubertal typically developing children

	Boys (n = 20)	Girls (n = 20)	p	d
Age (y)	9.7 ± 1.6	10.0 ± 1.8	0.451	0.017
Tanner stage (1/2)				
Pubic hair	17/3	16/4	0.677	0.129
Testicular-penile/Breast	17/3	15/5	0.429	0.247
Height (m)	1.38 ± 0.12	1.39 ± 0.11	0.765	0.174
Height percentile	53 ± 29	48 ± 21	0.494	0.222
Body mass (kg)	33.2 ± 7.7	33.4 ± 7.7	0.926	0.026
Body mass percentile	57 ± 26	45 ± 24	0.195	0.437
BMI (kg/m <sup>2</sup> )	17.2 ± 1.8	17.1 ± 2.3	0.837	0.069
BMI percentile	54 ± 27	45 ± 28	0.283	0.345
Midthigh muscle volume (cm <sup>3</sup> )	891 ± 267	896 ± 250	0.945	0.019
Femur length (m)	0.378 ± 0.040	0.385 ± 0.037	0.500	0.181
Total physical activity (counts/d)	480,370 ± 205,490	458,902 ± 197,209	0.738	0.107
MVPA (counts/d)	296,220 ± 193,585	266,999 ± 167,695	0.613	0.162
MVPA (min/d)	92 ± 44	88 ± 35	0.788	0.077
MVPA (counts/min/d)	3068 ± 612	2908 ± 549	0.389	0.276
Calcium intake (mg/d)	989 ± 308	899 ± 220	0.212	0.352

Values are means ± SD. Height percentile = height relative to age- and sex-based norms; Body

mass percentile = body mass relative to age- and sex-based norms; BMI = body mass index;

BMI percentile = BMI relative to age- and sex-based norms; MVPA = moderate plus vigorous

physical activity (values are not log transformed)

Table 2. Measures of trabecular and cortical bone architecture

	Boys (n = 20)	Girls (n = 20)	p	d
<u>Trabecular bone architecture</u>				
appBV/TV	0.322 ± 0.027	0.320 ± 0.024	0.838	0.078
appTb.N (1/mm)	1.672 ± .063	1.683 ± 0.069	0.597	0.182
appTb.Th (mm)	0.192 ± 0.013	0.190 ± 0.012	0.534	0.160
appTb.Sp (mm)	0.407 ± 0.028	0.405 ± 0.030	0.856	0.069
<u>Cortical bone architecture</u>				
Cortical volume (cm <sup>3</sup> )	26.6 ± 8.8	27.0 ± 8.5	0.861	0.046
Total bone volume (cm <sup>3</sup> )	42.0 ± 13.2	42.2 ± 12.1	0.960	0.016
Z (cm <sup>3</sup> )	0.722 ± 0.234	0.716 ± 0.239	0.873	0.025
J (cm <sup>4</sup> )	1.54 ± 0.68	1.54 ± 0.68	0.978	0.001

Values are means ± SD.

Table 3. Zero-order correlations demonstrating the relationships between bone architecture and muscle volume, femur length, sex, physical activity and calcium intake

	Muscle volume (cm <sup>3</sup> )	Femur length (m)	Sex	MVPA (min/d)	Calcium Intake (mg/d)
<u>Trabecular bone architecture</u>					
appBV/TV	.81*	.72*	.03	.17	-.06
appTb.N (1/mm)	.53*	.49**	-.09	.23	-.23
appTb.Th (mm)	.67*	.57*	.10	.07	.11
appTb.Sp (mm)	-.71*	-.63*	.03	-.22	.16
<u>Cortical bone architecture</u>					
Cortical vol (cm <sup>3</sup> )	.96*	.92*	-.03	.15	-.16
Total vol (cm <sup>3</sup> )	.94*	.93*	-.01	.21	-.18
Z (cm <sup>3</sup> )	.94*	.88*	.01	.15	-.18
J (cm <sup>4</sup> )	.92*	.86*	.01	.14	-.20

Significant correlation, \*p < 0.001 and \*\*p = 0.001; MVPA = moderate to vigorous physical activity



Table 4. Predictors of trabecular bone architecture in the distal femur of typically developing children determined using multiple regression analysis

	<u>appBV/TV</u>		<u>appTb.N (1/mm)</u>		<u>appTb.Th (mm)</u>		<u>appTb.Sp (mm)</u>	
	$\beta$	P value	$\beta$	P value	$\beta$	P value	$\beta$	P value
Muscle volume/femur length <sup>2,77</sup>	.344	.001	.257	.059	.286	.020	-.348	.004
Femur length (m)	.758	.000	.410	.004	.675	.000	-.611	.000
Sex	.037	.728	-.045	.754	.075	.563	.018	.885
MVPA (min/d)	.025	.809	.149	.302	-.063	.625	-.098	.426
Calcium intake (mg/d)	.052	.627	-.200	.173	.187	.156	.057	.645

MVPA = moderate to vigorous physical activity log transformed;  $\beta$  = standardized coefficient

Table 5. Predictors of cortical bone architecture in the midfemur of typically developing children determined using multiple regression analysis

	Cortical bone volume (cm <sup>3</sup> )		Total bone Volume (cm <sup>3</sup> )		Z (cm <sup>3</sup> )		J (cm <sup>4</sup> )	
	$\beta$	P value	$\beta$	P value	$\beta$	P value	$\beta$	P value
Muscle volume/femur length <sup>2,77</sup>	.219	.000	.160	.003	.260	.001	.234	.002
Femur length (m)	.954	.000	.943	.000	.905	.000	.889	.000
Sex	.027	.640	.075	.176	.062	.384	.061	.437
MVPA (min/d)	-.055	.333	.008	.886	-.037	.597	-.045	.561
Calcium intake (mg/d)	.021	.714	-.046	.400	-.016	.816	-.033	.671

MVPA = moderate to vigorous physical activity log transformed;  $\beta$  = standardized coefficient

Figure 1  
[Click here to download high resolution image](#)




Figure 2  
[Click here to download high resolution image](#)





Figure 3

[Click here to download high resolution image](#)



**Figure 4**  
[Click here to download high resolution image](#)

