

THE UNIVERSITY OF WARWICK

Original citation:

Hill, E. M., Griffiths, Frances and House, Thomas A.. (2015) Spreading of healthy mood in adolescent social networks. Proceedings B, 282 . 20151180.

<http://dx.doi.org/10.1098/rspb.2015.1180>

Permanent WRAP url:

<http://wrap.warwick.ac.uk/70868>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work of researchers of the University of Warwick available open access under the following conditions.

This article is made available under the Creative Commons Attribution 4.0 International license (CC BY 4.0) and may be reused according to the conditions of the license. For more details see: <http://creativecommons.org/licenses/by/4.0/>

A note on versions:

The version presented in WRAP is the published version, or, version of record, and may be cited as it appears here.

For more information, please contact the WRAP Team at: publications@warwick.ac.uk

warwick**publications**wrap

highlight your research

<http://wrap.warwick.ac.uk>



Cite this article: Hill EM, Griffiths FE, House T. 2015 Spreading of healthy mood in adolescent social networks. *Proc. R. Soc. B* **282**: 20151180.
<http://dx.doi.org/10.1098/rspb.2015.1180>

Received: 20 May 2015

Accepted: 27 July 2015

Subject Areas:

health and disease and epidemiology

Keywords:

social contagion, emotional contagion, depression

Author for correspondence:

E. M. Hill

e-mail: edward.hill@warwick.ac.uk

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rspb.2015.1180> or via <http://rspb.royalsocietypublishing.org>.

Spreading of healthy mood in adolescent social networks

E. M. Hill¹, F. E. Griffiths² and T. House^{1,3}

¹Centre for Complexity Science and Warwick Infectious Disease Epidemiology Research Centre, and ²Warwick Medical School, University of Warwick, Coventry CV4 7AL, UK

³School of Mathematics, University of Manchester, Manchester M13 9PL, UK

Depression is a major public health concern worldwide. There is evidence that social support and befriending influence mental health, and an improved understanding of the social processes that drive depression has the potential to bring significant public health benefits. We investigate transmission of mood on a social network of adolescents, allowing flexibility in our model by making no prior assumption as to whether it is low mood or healthy mood that spreads. Here, we show that while depression does not spread, healthy mood among friends is associated with significantly reduced risk of developing and increased chance of recovering from depression. We found that this spreading of healthy mood can be captured using a non-linear complex contagion model. Having sufficient friends with healthy mood can halve the probability of developing, or double the probability of recovering from, depression over a 6–12-month period on an adolescent social network. Our results suggest that promotion of friendship between adolescents can reduce both incidence and prevalence of depression.

1. Introduction

Depression and other mood disorders are major and growing contributors to mortality and morbidity worldwide [1]. These mood disorders are widespread, with the World Health Organization estimating that, globally, there are currently more than 350 million people affected by depression [2]. There is evidence that social support is important for the mental well-being of adolescents [3] and that befriending can have a positive effect on mental health [4]. Recent experiments suggest that people's expression of negative or positive emotions is influenced by the level of negative or positive news from their friends and associates [5]. An improved understanding of the social processes that drive the epidemiology of these diseases therefore has the potential to bring highly significant public health benefits.

It is now very common to model infectious diseases as spreading processes on networks [6]. This approach is increasingly applied to behaviours (e.g. those related to infectious risk [7]) and non-infectious diseases that are linked to behaviours that can spread socially (e.g. obesity and smoking [8,9]).

Previous work relating to spreading of depression on social networks has generally made at least one of the following key assumptions: (i) low mood and/or depression spreads like an infectious agent; (ii) healthy mood (non-depression) does not spread like an infectious agent; (iii) the information to distinguish between transmission and no-transmission models can be found in differences in static network measures such as clustering of disease [10–12], or in coarse population-level measures such as web-search over time [13]. Here, we allow more flexibility in our model by making no prior assumption as to whether it is low mood or healthy mood that spreads. In addition, we use the dynamical behaviour of mood over time, allowing us to distinguish directly between transmission and no transmission.

2. Material and methods

(a) The data

We consider data from the in-home interview survey of the Add Health study [14], which recorded adolescents' in-school friends in addition to their CES-D (Center for Epidemiologic Studies Depression Scale) scores [15]. This was used to classify individuals as either having depressive symptoms (low mood) or not being depressed (healthy mood) according to the score cut-off associated with a clinical diagnosis of depression [16]. We took data from two time points (waves) 6 to 12 months apart, from students in saturated schools (all students in a saturated school were selected to have an in-home interview, eliminating selection bias). To be included in our study sample, for both time points the student must have provided complete answers to all the CES-D survey-related questions and be the least restricted in the number of school friends they were allowed to give (each student was either allowed to list up to five male and five female friends, or limited to listing a maximum of one male and one female friend, with students in the latter group not considered for inclusion in our study sample).

(b) Model construction, fitting and selection

We model depression status as a discrete-time Markov chain, where each individual i at time t has state $X_i(t)$, taking either the value D for depressive symptoms or N for not depressed. This model is specified by two probabilities: the probability $p = \Pr[X_i(t+1) = D \mid X_i(t) = N]$ of becoming depressed, and the probability $q = \Pr[X_i(t+1) = N \mid X_i(t) = D]$ of recovering from depression. Following Centola and Macy [17,18], we considered a model in which these probabilities depend on the number of friends of an individual who have value N or D , with this dependence taking the form of an S-shaped function. These models are referred to as N transmits and D transmits, respectively. We then fit this model with the Add Health data moving from wave 1 to wave 2, and compare with the no-transmission model that the probabilities do not depend on the moods of an individual's friends. Parameter values for our transmission and no-transmission models were found using maximum-likelihood estimation (MLE) by minimizing the negative log-likelihood $-\log(L)$ with respect to p and q using the MATLAB `fmincon()` function. Confidence intervals were obtained through calculation of the Hessian matrix at the MLE parameters and use of standard asymptotic formulae. Appendix A outlines the construction of the likelihood functions used in the fitting process. Competing models were assessed using the Akaike information criterion (AIC) [19].

(c) Simulation outline

A discrete-time Monte Carlo simulation of the no-transmission model and N -transmits model was performed on a directed network of named friends constructed from the 3084 individuals in the dataset satisfying our inclusion criteria at the first time point (wave 1). We took 10^5 independent samples from the stationary distribution for each model to calculate model quantities of interest including uncertainty. We assessed uncertainty in the observed quantities through bootstrapping. The Bonferroni method was used to account for multiple testing of statistically significant differences between models and observed data [20]. To further test the goodness-of-fit of our chosen transmission model, residual errors were analysed via a parametric bootstrap approach (see the electronic supplementary material).

3. Results

(a) Fitted parameter values

(i) No-transmission models

We obtained the no-transmission deterioration model for transitioning from healthy mood to low mood within

a year,

$$p_k = 0.088[0.075, 0.10],$$

and the no-transmission recovery model for transitioning from low mood to healthy mood within a year,

$$q_k = 0.56[0.50, 0.61].$$

(ii) N -transmits models

We obtained the N -transmits deterioration model for transitioning from healthy mood to low mood within a year,

$$p_k = \alpha + \beta \sum_{l=0}^k \binom{10}{l} \gamma^l (1-\gamma)^{10-l},$$

with $\alpha = 0.13[0.088, 0.16]$, $\beta = -0.064[-0.10, -0.025]$ and $\gamma = 0.25[0.070, 0.44]$. Note that here and elsewhere numbers such as 10 appear as the limits in the data on number of friends; k is the number of friends in the transmitting state, and the parameters estimated are a simple way to parametrize a discrete sigmoidal function as suggested by a complex contagion model. The N -transmits recovery model, for transitioning from low mood to healthy mood within a year, was

$$q_k = \alpha + \beta \sum_{l=0}^k \binom{10}{l} \gamma^l (1-\gamma)^{10-l},$$

with $\alpha = 0.53[0.46, 0.59]$, $\beta = 0.46[0.099, 0.83]$ and $\gamma = 0.71[0.45, 0.98]$.

(iii) D -transmits models

We obtained the D -transmits deterioration model for transitioning from healthy mood to low mood within a year,

$$p_k = \alpha + \beta \sum_{l=0}^k \binom{4}{l} \gamma^l (1-\gamma)^{4-l},$$

with $\alpha = 0.087[0.074, 0.10]$, $\beta = 0.021[-0.27, 0.31]$ and $\gamma = 0.84[-0.62, 2.3]$. The D -transmits recovery model, for transitioning from low mood to healthy mood within a year,

$$q_k = \alpha + \beta \sum_{l=0}^k \binom{4}{l} \gamma^l (1-\gamma)^{4-l},$$

with $\alpha = 4.1 \times 10^{-5}[-5.0, 5.0]$, $\beta = 0.64[-4.6, 5.9]$ and $\gamma = 0.050[-0.42, 0.52]$.

(b) Model comparisons

Figure 1 shows the results of fitting the different models to the $n = 2194$ data points given by our inclusion criteria. For the dependence of probabilities p and q on the number of friends with depressive symptoms (no-transmission model against D -transmits model), AIC values showed the no-transmission model was the preferred choice (figure 1a,b). When considering the no-transmission model against the N -transmits model, the N -transmits model was the preferred choice in both cases (figure 1c,d).

(c) Simulation analysis

Comparing D prevalence and edge type summary statistics to those obtained for the observed data, there were significant differences between the no-transmission model and data, while the transmission model (with the probability dependent on the number of N friends) had no statistically significant




Figure 1. Dynamical behaviour of depression status between samples as a function of N friends or D friends for observed data (and 50% CI), transmission and no-transmission models. Uncertainty in the observed values was quantified using Jeffreys intervals [21]. The ΔAIC value is calculated by subtracting the no-transmission AIC value from the transmission AIC value. (a) Probability of transitioning from healthy mood to low mood against number of D friends—transmission is not preferred to no transmission ($\Delta\text{AIC} \approx -4$). (b) Probability of recovering from low mood against number of D friends—transmission is not preferred to no transmission ($\Delta\text{AIC} \approx -0.9$). (c) Probability of transitioning from healthy mood to low mood against number of N friends—transmission is preferred to no transmission ($\Delta\text{AIC} \approx 8.4$). (d) Probability of recovering from low mood against number of N friends—transmission is preferred to no transmission ($\Delta\text{AIC} \approx 4.5$).

differences (figure 2). In particular, the $N \rightarrow N$, $N \rightarrow D$ and $D \rightarrow N$ edge statistics (where we write $A \rightarrow B$ for an individual in state A naming an individual in state B as a friend) were found to be statistically significantly different between the no-transmission model and the data (figure 2*b–d*). We also assessed goodness-of-fit and parameter identifiability through simulation, giving extra confidence to our results (see the electronic supplementary material).

4. Discussion

A major benefit of the dynamical approach that we have taken is that it avoids the problems of confounding that have been controversial in other studies of social contagion [12]. Figure 3 shows the model schematically, to provide intuitive insight into why this is the case.

In this model, there is transmission of D if the probability of event (3) happening (given the initial state) is bigger than the probability of event (1) happening (given the initial state): $\text{Pr}(\text{event (3)}) > \text{Pr}(\text{event (1)})$. There is also transmission of D if $\text{Pr}(\text{event (6)}) < \text{Pr}(\text{event (4)})$. We did not find evidence

for transmission of D based on this criterion, as shown in figure 1*a,b*. Such transmission would also be expected to lead to more $D \rightarrow D$ pairs and fewer $D \rightarrow N$ and $N \rightarrow D$ pairs than a null model. This pattern was not observed (see figure 2; electronic supplementary material, figure S1).

There is transmission of N in our model if $\text{Pr}(\text{event (2)}) < \text{Pr}(\text{event (1)})$ and also if $\text{Pr}(\text{event (5)}) > \text{Pr}(\text{event (4)})$. We found statistically significant evidence for transmission of N on the basis of this criterion, as seen in figure 1*c,d*. Such transmission would also be expected to lead to more $N \rightarrow N$ pairs and fewer $D \rightarrow N$ and $N \rightarrow D$ pairs than a null model. This pattern was observed (figure 2).

Suppose that there is homophily (similar individuals naming each other as friends) at work in the social network, either in terms of depressive symptoms, or a latent variable that is correlated with depressive symptoms. This will tend to increase the number of $D \rightarrow D$ or $N \rightarrow N$ pairs in the absence of any transmission effect, meaning that these tests (shown in figure 2) can be confounded by homophily. When working with two waves of data, however, such homophily will simply lead to fewer individuals in the initial states associated with events (3) and (5) than events (2) and (6), but in our




Figure 2. Static summary statistics for the stationary distributions of the models versus real data. Asterisks above a plot denote a significant statistical difference at the 5% level, corresponding to $p < 0.01$ using the Bonferroni method to account for multiple testing. (a) Prevalence of individuals with depressive symptoms—observed data could be plausibly generated by both transmission ($p = 0.058$) and no-transmission ($p = 0.41$) models. (b) Number of $N \rightarrow N$ edges—observed data could be plausibly generated by the transmission model ($p = 0.15$) but not by the no-transmission model ($p = 0.0014$). (c) Number of $D \rightarrow N$ edges—observed data could be plausibly generated by the transmission model ($p = 0.54$) but not by the no-transmission model ($p = 0.0035$). (d) Number of $N \rightarrow D$ edges—observed data could be plausibly generated by the transmission model ($p = 0.027$) but not by the no-transmission model ($p = 0.0067$). The fifth test and plot is for $D \rightarrow D$ edges (electronic supplementary material, figure S1). See Appendix B for the p -value calculation method, and electronic supplementary material, figure S2, for stratification by number of contacts.

approach we fit to the probability of moving to a final state given an initial state. This means that since there are still sufficient data to find a statistically significant effect, homophily cannot confound the results shown in figure 1. Our verbal argument here can be made in a more mathematically precise manner, as shown in the electronic supplementary material.

In summary, we have shown the epidemiological impact of such mood transmission in a large adolescent population, giving statistically significant evidence for spreading of healthy mood, but not for spreading of depressive symptoms. Once discovered, this behaviour is in fact plausible through a number of mechanisms. Depression has been associated with social withdrawal [22], and so depressed individuals would be expected to exert less social influence than adolescents with healthy mood. However, each individual may need sufficient exposure to others with a healthy mood in order to stay well, or become well if depressed. In support of this, there is evidence from psychology of mechanisms by which mood is transmitted between people. Automatic transmission of mood between people has been demonstrated [23].

Unconscious mimicry enhances social rapport [24], and those feeling positive towards the person with whom they are interacting socially are more likely to mimic, and so build rapport [25], and thus the opportunity for transmission of healthy mood. People who are (or have a tendency to be) depressed are less able to maintain a positive outlook from moment to moment [26], a deficit potentially compensated by interaction with healthy friends.

The static network measures provide indirect evidence of spreading of healthy mood through analysis of clustering, which shows that the no-transmission model is significantly different from the observed data, while the data and N transmits model are in agreement. Such clustering, while supportive of a transmission effect, can have other causes, and so we recommend that future empirical work measure changes in mood over time where possible.

Our results offer implications for improving adolescent mood. In particular, they suggest the hypothesis that enabling networks of friendship between adolescents has the potential to reduce both incidence and prevalence of depression. Our




Figure 3. Pictorial representation of the possible events in our model: developing or recovering from depressive symptoms; in the absence of friends, with friends with healthy mood, or with depressed friends. The changes in pairs produced are also shown.

complex contagion model suggests that adolescents with five or more healthy friends have half the probability of becoming depressed over a 6–12-month period compared with adolescents with no healthy friends, and that adolescents with 10 healthy friends have double the probability of recovering from depressive symptoms over a 6–12-month period compared with adolescents with three healthy friends. If such an effect were demonstrated in an intervention study, this would massively outperform existing interventions.

Ethics. Add Health participants provided written informed consent for participation in all aspects of Add Health in accordance with the University of North Carolina School of Public Health Institutional Review Board guidelines, which are based on the Code of Federal Regulations on the Protection of Human Subjects (45 CFR 46; see <http://www.hhs.gov/ohrp/humansubjects/guidance/45cfr46.html>).

Data accessibility. The data analysed in this study were obtained on licence from Add Health. Requests for restricted-use data access can be made to Add Health at <http://www.cpc.unc.edu/projects/addhealth/data/restricteduse/add-health-restricted-use-data-contract-1> or addhealth@unc.edu.

Authors' contributions. F.E.G. and T.H. planned the study. E.M.H. performed the simulations and numerical data analysis. E.M.H. and T.H. wrote the main paper and electronic supplementary material. All authors discussed the results and commented on the manuscript at each stage. All authors gave final approval for publication.

Competing interests. We declare we have no competing interests.

Funding. T.H. and E.M.H. are supported by the Engineering and Physical Sciences Research Council. The Add Health study was funded by

grant no. P01-HD31921 from the Eunice Kennedy Shriver National Institute of Child Health and Human Development, with cooperative funding from 23 other federal agencies and foundations. No direct support was received from grant P01-HD31921 for this analysis.

Acknowledgements. This research uses data from Add Health, a programme project directed by Kathleen Mullan Harris, and designed by J. Richard Udry, Peter S. Bearman and Kathleen Mullan Harris at the University of North Carolina at Chapel Hill. Special acknowledgement is due to Ronald R. Rindfuss and Barbara Entwisle for assistance in the original design. Information on how to obtain the Add Health data files is available on the Add Health website (<http://www.cpc.unc.edu/addhealth>). We thank Robert Goudie for R code, and Dan Sprague and Steven Hill for helpful comments, and thank the editor and two anonymous reviewers for helpful comments.

Appendix A. Likelihood functions

The following likelihood function was constructed for the development of the 'depressive symptoms' scenario, with respect to either the number of not-depressed friends or number of friends with depressive symptoms:

$$L(y|p, N) = \prod_k \binom{N_k}{y_k} p_k^{y_k} (1 - p_k)^{N_k - y_k},$$

where y_k was the number of respondents with k not-depressed friends (friends with depressive symptoms) who were classified as not depressed at the first time point and having depressive symptoms at the second time point. N_k was the total number of respondents classified as not depressed at the first time point with k not-depressed friends (friends with depressive symptoms). An equivalent likelihood function was constructed for the 'recovery from depressive symptoms' scenario:

$$L(y|q, D) = \prod_k \binom{D_k}{y_k} q_k^{y_k} (1 - q_k)^{D_k - y_k},$$

with y_k corresponding to the number of respondents with k not-depressed friends (friends with depressive symptoms) who were classified as having depressive symptoms at the first time point and not depressed at the second time point. D_k was the total number of respondents classified as having depressive symptoms at the first time point with k not-depressed friends (friends with depressive symptoms).

Appendix B. p -value calculation

To calculate the p -values in figure 2 and electronic supplementary material, figure S1, required comparison of Monte Carlo simulation output with uncertain data, and so we used the expression

$$p = \sum_x 2\pi_x \times \min(E_x, 1 - E_x),$$

where π_x is the density of the value x in the bootstrap sample from data, and E_x is the empirical cumulative distribution function of the simulation output.

References

- Moussavi S, Chatterji S, Verdes E, Tandon A, Patel V, Ustun B. 2007 Depression, chronic diseases, and decrements in health: results from the world health surveys. *Lancet* **370**, 851–858. (doi:10.1016/S0140-6736(07)61415-9)

2. World Health Organization. 2012 Depression. Fact sheet no. 369. See <http://www.who.int/mediacentre/factsheets/fs369/en/> (accessed 8 July 2014).
3. Rueger SY, Malecki CK, Demaray MK. 2010 Relationship between multiple sources of perceived social support and psychological and academic adjustment in early adolescence: comparisons across gender. *J. Youth Adolescence* **39**, 47–61. (doi:10.1007/s10964-008-9368-6)
4. Mead N, Lester H, Chew-Graham C, Gask L, Bower P. 2010 Effects of befriending on depressive symptoms and distress: systematic review and meta-analysis. *Br. J. Psychiatry* **196**, 96–101. (doi:10.1192/bjp.bp.109.064089)
5. Kramer ADI, Guillory JE, Hancock JT. 2014 Experimental evidence of massive-scale emotional contagion through social networks. *Proc. Natl Acad. Sci. USA* **111**, 8788–8790. (doi:10.1073/pnas.1320040111)
6. Danon L, Ford AP, House T, Jewell CP, Keeling MJ, Roberts GO, Ross JV, Vernon MC. 2011 Networks and the epidemiology of infectious disease. *Interdisc. Perspect. Infect. Dis.* **2011**, 1–28. (doi:10.1155/2011/284909)
7. Funk S, Gilad E, Watkins C, Jansen VAA. 2009 The spread of awareness and its impact on epidemic outbreaks. *Proc. Natl Acad. Sci. USA* **106**, 6872–6877. (doi:10.1073/pnas.0810762106)
8. Christakis NA, Fowler JH. 2007 The spread of obesity in a large social network over 32 years. *N. Engl. J. Med.* **357**, 370–379. (doi:10.1056/NEJMs066082)
9. Christakis NA, Fowler JH. 2008 The collective dynamics of smoking in a large social network. *N. Engl. J. Med.* **358**, 2249–2258. (doi:10.1056/NEJMs0706154)
10. Joiner TE, Katz J. 1999 Contagion of depressive symptoms and mood: meta-analytic review and explanations from cognitive, behavioral, and interpersonal viewpoints. *Clin. Psychol. Sci. Pr.* **6**, 149–164. (doi:10.1093/clipsy.6.2.149)
11. Hill AL, Rand DG, Nowak MA, Christakis NA. 2010 Emotions as infectious diseases in a large social network: the SISa model. *Proc. R. Soc. B* **277**, 3827–3835. (doi:10.1098/rspb.2010.1217)
12. Christakis NA, Fowler JH. 2013 Social contagion theory: examining dynamic social networks and human behavior. *Stat. Med.* **39**, 556–577. (doi:10.1002/sim.5408)
13. Bentley RA, Ormerod P. 2010 A rapid method for assessing social versus independent interest in health issues: a case study of 'bird flu' and 'swine flu'. *Soc. Sci. Med.* **71**, 482–485. (doi:10.1016/j.socscimed.2010.03.042)
14. Harris KM, Halpern CT, Whitsel E, Hussey J, Tabor J, Entzel P, Udry JR. 2009 The National Longitudinal Study of Adolescent to Adult Health: research design. See <http://www.cpc.unc.edu/projects/addhealth/design> (accessed 9 April 2015).
15. Radloff LS. 1977 The CES-D Scale: a self-report depression scale for research in the general population. *Appl. Psychiatry Meas.* **1**, 385–401. (doi:10.1177/014662167700100306)
16. Roberts RE, Lewinsohn PM, Seeley JR. 1991 Screening for adolescent depression: a comparison of depression scales. *J. Am. Acad. Child Adolescent Psychiatry* **30**, 58–66. (doi:10.1097/00004583-199101000-00009)
17. Centola D, Macy M. 2007 Complex contagions and the weakness of long ties. *Am. J. Sociol.* **113**, 702–734. (doi:10.1086/521848)
18. Centola D. 2010 The spread of behavior in an online social network experiment. *Science* **329**, 1194–1197. (doi:10.1126/science.1185231)
19. Akaike H. 1974 A new look at the statistical model identification. *IEEE Trans. Autom. Control* **19**, 716–723. (doi:10.1109/TAC.1974.1100705)
20. Bland JM, Altman DG. 1995 Multiple significance tests: the Bonferroni method. *BMJ* **310**, 170. (doi:10.1136/bmj.310.6973.170)
21. Brown LD, Cai T, Dasgupta A. 2002 Confidence intervals for a binomial proportion and asymptotic expansions. *Ann. Stat.* **30**, 160–201. (doi:10.1214/aos/1015362189)
22. Schaefer DR, Kornienko O, Fox AM. 2011 Misery does not love company: network selection mechanisms and depression homophily. *Am. Sociol. Rev.* **76**, 764–785. (doi:10.1177/0003122411420813)
23. Neumann R, Strack F. 2000 Mood contagion: the automatic transfer of mood between persons. *J. Pers. Soc. Psychol.* **79**, 211. (doi:10.1037/0022-3514.79.2.211)
24. Lakin JL, Chartrand TL. 2003 Using nonconscious behavioral mimicry to create affiliation and rapport. *Psychol. Sci.* **14**, 334–339. (doi:10.1111/1467-9280.14481)
25. Leighton J, Bird G, Orsini C, Heyes C. 2009 Social attitudes modulate automatic imitation. *J. Exp. Soc. Psychol.* **46**, 905–910. (doi:10.1016/j.jesp.2010.07.001)
26. Höhn P, Menne-Lothmann C, Peeters F, Nicolson NA, Jacobs N, Derom C, Thiery E, Van Os J, Wichers M. 2003 Moment-to-moment transfer of positive emotions in daily life predicts future course of depression in both general population and patient samples. *PLoS ONE* **8**, e75655. (doi:10.1371/journal.pone.0075655)